

Superimposed vibration on suspended push-ups (#75972)

1

First submission

Guidance from your Editor

Please submit by **19 Aug 2022** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

3 Figure file(s)

3 Table file(s)

! Custom checks

Human participant/human tissue checks



Have you checked the authors [ethical approval statement](#)?



Does the study meet our [article requirements](#)?



Has identifiable info been removed from all files?



Were the experiments necessary and ethical?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Superimposed vibration on suspended push-ups

Bernat Buscà¹, Joan Aguilera-Castells^{Corresp., 1}, Jordi Arboix-Alió^{1, 2}, Adrià Miró¹, Azahara Fort-Vanmeerhaeghe^{1, 2}, Pol Huertas¹, Javier Peña^{3, 4}

¹ Faculty of Psychology, Education Sciences, and Sport Blanquerna, Ramon Llull University, Barcelona, Spain

² School of Health Science Blanquerna, Ramon Llull University, Barcelona, Spain

³ Sport and Physical Activity Studies Centre (CEEAF), University of Vic-Central University of Catalonia, Vic, Spain

⁴ Sport performance Analysis Research Group (SPARG), University of Vic-Central University of Catalonia, Vic, Spain

Corresponding Author: Joan Aguilera-Castells

Email address: joanac1@blanquerna.url.edu

Background. Superimposition of vibration has been proposed in sports training using several devices and methods to enhance muscle activation and strength adaptations. Due to the popularity of suspension training, vibration systems have recently been developed to increase the effects of this training method. The present cross-sectional study aims to examine the effects of superimposing vibration on one of the most popular exercises in strength and conditioning programs: push-ups.

Methods. Twenty-eight physically active men and women executed push-ups in three suspended conditions (non-vibration, vibration at 25 Hz, and vibration at 40 Hz). OMNI-Res scale was registered, and surface electromyographic signals were measured for the activity of the right and left external oblique, anterior deltoid, triceps brachii, sternal, and clavicular heads of the pectoralis major.

Results. A linear mixed model indicated a significant fixed effect for vibration at 25 Hz and 40 Hz on muscle activity. Suspended push-ups with superimposed vibration (25 Hz and 40 Hz) showed a significant higher activity on left (25 Hz: $p = 0.036$, $d = 0.34$; 40 Hz: $p = 0.003$, $d = 0.48$) and right external oblique (25 Hz: $p = 0.004$, $d = 0.36$; 40 Hz: $p = 0.000$, $d = 0.59$), anterior deltoid (25 Hz: $p = 0.032$, $d = 0.44$; 40 Hz: $p = 0.003$, $d = 0.64$), and global activity (25 Hz: $p = 0.000$, $d = 0.55$; 40 Hz: $p = 0.000$, $d = 0.83$) compared to non-vibration condition. Moreover, OMNI-Res significant differences were found at 25 Hz (6.04 ± 0.32 , $p = 0.000$, $d = 4.03$, $CI = 3.27, 4.79$) and 40 Hz (6.21 ± 0.36 , $p = 0.000$, $d = 4.29$, $CI = 3.49, 5.08$) compared to the non-vibration condition (4.75 ± 0.32).

Conclusion. Superimposing vibration is a feasible strategy to enhance the muscle activity of suspended push-ups.

Superimposed vibration on suspended push-ups

Bernat Buscà¹, Joan Aguilera-Castells¹, Jordi Arboix-Alió^{1,2}, Adrià Miró¹, Azahara Fort-Vanmeerhaeghe^{1,2}, Pol Huertas¹, Javier Peña^{3,4}

¹ Faculty of Psychology, Education Sciences, and Sport Blanquerna, Ramon Llull University, Barcelona, Spain

² School of Health Science Blanquerna, Ramon Llull University, Barcelona, Spain

³ Sport and Physical Activity Studies Centre (CEEAF), University of Vic-Central University of Catalonia, Vic, Spain

⁴ Sport performance Analysis Research Group (SPARG), University of Vic-Central University of Catalonia, Vic, Spain

Corresponding Author:

Joan Aguilera-Castells¹

34 Cister Street, Barcelona, 08022, Spain

Email address: joanac1@blanquerna.url.edu

Abstract

Background. Superimposition of vibration has been proposed in sports training using several devices and methods to enhance muscle activation and strength adaptations. Due to the popularity of suspension training, vibration systems have recently been developed to increase the effects of this training method. The present cross-sectional study aims to examine the effects of superimposing vibration on one of the most popular exercises in strength and conditioning programs: push-ups.

Methods. Twenty-eight physically active men and women executed push-ups in three suspended conditions (non-vibration, vibration at 25 Hz, and vibration at 40 Hz). OMNI-Res scale was registered, and surface electromyographic signals were measured for the activity of the right and left external oblique, anterior deltoid, triceps brachii, sternal, and clavicular heads of the pectoralis major.

Results. A linear mixed model indicated a significant fixed effect for vibration at 25 Hz and 40 Hz on muscle activity. Suspended push-ups with superimposed vibration (25 Hz and 40 Hz) showed a significant higher activity on left (25 Hz: $p = 0.036$, $d = 0.34$; 40 Hz: $p = 0.003$, $d = 0.48$) and right external oblique (25 Hz: $p = 0.004$, $d = 0.36$; 40 Hz: $p = 0.000$, $d = 0.59$), anterior deltoid (25 Hz: $p = 0.032$, $d = 0.44$; 40 Hz: $p = 0.003$, $d = 0.64$), and global activity (25 Hz: $p = 0.000$, $d = 0.55$; 40 Hz: $p = 0.000$, $d = 0.83$) compared to non-vibration condition. Moreover, OMNI-Res significant differences were found at 25 Hz (6.04 ± 0.32 , $p = 0.000$ $d = 4.03$ CI = 3.27, 4.79) and 40 Hz (6.21 ± 0.36 $p = 0.00$ $d = 4.29$ CI = 3.49, 5.08) compared to the non-vibration condition (4.75 ± 0.32).

Conclusion. Superimposing vibration is a feasible strategy to enhance the muscle activity of suspended push-ups.

Introduction

Combining different strength training methods is an increasingly used strategy to reach sports performance and competitive advantages. The synergistic effect of recruiting prime movers, antagonists, and stabilizers, justifies the use of complex exercises that present instability (La Scala Teixeira et al., 2019). This effect can be even more important in sports, where perturbed tasks constitute the essence of their specificity (Behm & Anderson, 2006). The upper body muscles can benefit from instability, especially in overhead disciplines, such as handball, water polo or hockey, and gymnastic sports, continuously demanding precise, powerful, complex, and unidirectional actions. Acting as a pendulum by rotating around a singular anchor point above, suspension training uses its essential characteristics (vector resistance, stability, and pendulum) and body weight to enhance neuromuscular demands (Bettendorf, 2010).

Complex tasks involving instability have been combined with mechanical vibrations to increase its neuromuscular demands in the past (Cloak et al., 2013; Marín & Hazell, 2014; Ritzmann et al., 2014; Sierra-Guzmán et al., 2018; Aguilera-Castells et al., 2019, 2021). Vibratory training transfers vibration on the muscle to elicit the tonic vibration reflex (Cardinale & Bosco, 2003). Superimposing vibration can alter the motor unit recruitment, activating faster

and larger motor units (Martin & Park, 1997; Xu et al., 2018), thus reinforcing the possible benefits of using those devices in standard training methods (Cardinale & Wakeling, 2005). Whole-body vibration (WBV) applied through platforms is the most studied vibrating method to provoke acute neuromuscular effects (Cardinale & Lim, 2003; Rønnestad, 2009a,b) and long-term adaptations (Gollhofer, 2010; Manimmanakorn et al., 2014). Nevertheless, several devices superimposed vibration on barbells (Poston et al., 2007; Mischi & Cardinale, 2009; Moras et al., 2010; Xu, Rabotti & Mischi, 2013), dumbbells (Bosco, Cardinale & Tsarpela, 1999; Cochrane & Hawke, 2007), and cables (Issurin & Tenenbaum, 1999; Issurin et al., 2010) have also been designed to transfer vibratory stimuli to the upper body. Mischi and Cardinale (2009) reported significantly higher muscle activity in arm muscles when performing the isometric V exercise. Moras et al. (2010) have demonstrated the acute effects of superimposing vibration in a bench press in the prime movers, especially during flexion. Poston et al. (2007) showed a greater bench press average power in a vibrating condition, although they did not assess muscle activity. The authors superimposed the vibrating engine on the barbell side. Similarly, Xu et al. (2015) prototyped a bench to combine the effect of muscle tension and vibration on muscle activation, demonstrating the benefits of using an adaptive normalized least mean square algorithm to determine the real effects of superimposed vibration on the biceps brachii. Lately, vibration has been superimposed on a suspension device in lower limb exercises (Aguilera-Castells et al., 2021). When performing dynamic supine bridges and hamstring curls, surface electromyography reflected a higher activity of the muscles proximal to the straps (gastrocnemius medialis, lateralis, and semitendinosus). However, the effect on the primary movers was non-significant.

Therefore, the main objective of the present study was to examine the effects of vibration on muscle activity in dynamic suspended push-ups. It was hypothesized that the superimposed vibration on the suspension straps might obtain higher muscle activity than the suspended condition without vibration. It was also hypothesized that the OMNI-Res perceived exertion scale for resistance exercise would be higher in the vibration exercises than the non-vibrating exercises.

Materials & Methods

Design

A cross-sectional study design investigated the effects of a suspension device with superimposed vibration on upper body activation. Participants performed suspended push-ups in non-vibration, vibration at 25 Hz and 40 Hz. Surface electromyography (sEMG) was used to record and compare the activity of the right and left external oblique, anterior deltoid, triceps brachii, and sternal and clavicular heads of the pectoralis major. sEMG values were expressed as a percentage of maximum voluntary isometric contraction (% MVIC). Furthermore, the perceived exertion was assessed using the OMNI-Perceived Exertion Scale for Resistance Exercise (OMNI-Res) under all suspended push-ups conditions.

Participants

Twenty-eight physically active male ($n = 25$, mean age = 22.7 ± 3.6 years, height = 1.8 ± 0.1 m, body mass = 77.7 ± 8.4 kg, body mass index = 24.5 ± 2.1 kg·m⁻², suspension training experience = 5.2 ± 2.7 years) and female ($n = 3$, mean age = 22.6 ± 0.6 years, height = 1.6 ± 0.0 m, body mass = 56.0 ± 4.0 kg, body mass index = 21.9 ± 2.4 kg·m⁻², suspension training experience = 3.7 ± 2.5 years) voluntarily participated in the study. Participants were excluded from participating in the study if their suspension training experience was under one year, did not perform a minimum of 90 minutes of physical activity per week, or had cardiovascular, musculoskeletal, or neurological diseases. Before beginning the familiarization session, all participants were informed of all study procedures, benefits, and risks, in oral and written form, before receiving and signing the informed consent form. The Physical Activity Readiness Questionnaire (PAR-Q) was also handed out to the participants to identify any health risks related to physical exercise (Warburton et al., 2011). Three to four hours before the testing sessions participants did not ingest any stimulant substances (e.g., caffeine), food or drink. In addition, no high-intensity physical activity was performed 24 hours before the tests. The Ethics and Research Committee Board of Blanquerna Faculty of Psychology and Educational and Sport Sciences at Ramon Llull University in Barcelona, Spain (ref. number 1819005D) approved this study, and the protocols followed the principles of the Declaration of Helsinki (revised in Fortaleza, Brazil, 2013).

Procedures

A familiarization session was held one week before the testing sessions. Participants were instructed to perform the suspended push-ups with proper technique in the different conditions (non-vibration, vibration at 25 Hz and 40 Hz) in two sets of five repetitions. Anthropometric (e.g., weight, height, acromion distance) and descriptive data were collected. The test session was carried out a week later and at the same time in the morning. Researchers cleaned the electrode site with alcohol, shaving the skin area when necessary, thus placing the surface electrodes (Biopac EL504 disposable Ag-AgCl; BIOPAC System, Inc., Goleta, CA) on the external oblique (left and right), anterior deltoid, triceps brachii, sternal and clavicular portion of pectoralis major on the dominant upper limb (Criswell & Cram, 2011). A reference electrode over the iliac crest was placed and all electrodes were placed at an inter-electrode distance of 2 cm following the SENIAM guidelines (Hermens et al., 2000). Next, a standardized warm-up consisting of 10 min of dynamic upper body calisthenics and two sets of eight repetitions of strict push-ups on the floor was performed. Then, participants executed a maximal voluntary isometric contraction (MVIC) test for the right and left external oblique, anterior deltoid, triceps brachii, and the sternal and clavicular head of the pectoralis major. The MVIC values were used as a baseline to normalize the sEMG signal (Halaki & Ginn, 2012). After the normalization protocols, participants completed a set of five repetitions for each push-up condition in a randomized order. The standardized suspended push-up technique consisted of holding the legs at shoulder-width apart, the hands separated at 150% of the acromial distance, in a pronated position, and grabbing

the suspension strap handles (TRX Suspension Trainer; Fitness Anywhere, San Francisco, CA, USA). Throughout the exercise, participants were instructed to maintain the lower back natural sway. For the lower position during the suspended push-up conditions, the elbow flexion was standardized at 90° and measured using a goniometer. Customized stoppers (similar to hurdles) were used to control the elbow flexion and 150% acromial width. Participants began the suspended push-up in the upper position (elbow extension) with a plantar flexion over the plumb line between the anchor point of the suspension strap and the ground. Participants flexed their elbows to 90° (lower position) in this position, then pushed with their hands on the handles to extend their elbows and return to the upper position. The length of the suspension strap was standardized at 1.64 meters, and the inclination ranged from 20° to 33° (mean $\pm$ SD: 26.5° $\pm$ 3.5). A positional transducer (WSB 16k-200; ASM Inc., Moosinning, DEU) was used to control the range of movement in each suspended push-up condition. The positional transducer tether was attached to the chest. The measured signal was used to identify the beginning and end of each repetition and determine the eccentric phase (lower position) and the concentric phase (upper position) of the suspended push-up. The pace of the push-up repetitions was standardized using a metronome settled at 60 beats per minute. Furthermore, all participants were given two-minute rest for each suspended push-up attempt. Those repetitions that did not follow the standard technique established by the researchers were repeated with two-minute rest between attempts.

A vibration device provided the superimposed vibration on the suspension straps for suspension training settled at two frequencies (25 Hz and 40 Hz) with an amplitude of 8 mm (peak to peak). The device was attached between the ceiling anchor point and the suspension strap to transmit the vibration through the straps using a connecting rod's vertical motion caused by an electric motor's rotary motion.

Muscle Activity Assessment

Muscle activity of the analyzed muscles during suspended push-ups (non-vibration, vibration at 25 Hz and 40 Hz) was obtained using the six-channel sEMG BIOPAC MP-150 System (sampling rate: 1.0 kHz; BIOPAC System, Inc., Goleta, CA). The sEMG was processed using a bandpass filtered at 10-500 Hz with a fourth-order 50 Hz Butterworth notch filter, and then the root mean square (RMS) was calculated. Afterward, the RMS signal was normalized and expressed as a percentage of the MVIC (% MVIC). This process was done with the AcqKnowledge 4.2 software (BIOPAC System, Inc., Goleta, CA). sEMG signal normalization was established considering the best of three attempts of 5 seconds, increasing the contraction progressively for 2 seconds and maintaining the maximal isometric contraction for 3 seconds, with a 3-minute rest between MVICs trials (Jakobsen et al., 2013). The different positions to reach the MVIC followed the Konrad (2006) guidelines, thus for the sternal and clavicular head of pectoralis major the participants lay supine, with their feet on the floor and pushed with their arms (elbows 90°) against an immovable resistance (fixed bar); for the anterior deltoid, the participants sat on a bench with their feet on the floor and leaned their back against the backrest to perform a glenohumeral flexion movement by holding a fixed bar with the hand closed in a

pronated position blocking the elbows; in the previous position, the bar was adjusted to allow the participants to hold their elbows at 90°, and in this position to perform an extension movement of the elbows against the fixed bar to perform the MVIC of the triceps brachii; for the external oblique (left and right) participants laid on a bench in a side position with their legs and hips held with ratchets, in this position a manual resistance was applied against the lateral trunk flexion movement.

Perceived Subjective Exertion Measurement (OMNI-Res scale)

After performing each suspended push-up condition, participants were asked about perceived subjective exertion using the OMNI-Res scale and following Robert's (2003) protocol. During the familiarization session, a visual OMNI-Res scale was displayed to ensure that participants provided an accurate perception of the exertion. Participants were asked to report their subjective perception of exertion values ranging from 0 (extremely easy) to 10 (extremely hard). Participants were instructed that on the OMNI-Res scale, a value of 0 is equivalent to performing an unweighted exercise and a value of 10 is equivalent to lifting one repetition maximum. In the test session, the protocol mentioned above was followed, and the OMNI-Res value of each exercise condition was recorded.

Data Analysis

The data analysis consisted of recording each analyzed muscle's peak activation (%MVIC) in the different conditions of the suspended exercise (push-ups non-vibration, 25 Hz, and 40 Hz) for three intermediate repetitions. Thus, each suspended push-up condition's first and fifth repetition were discarded. In addition, the peak sEMG was analyzed for the concentric (upper position) and eccentric (lower position) phases of the three repetitions. The sEMG activation values obtained in %MVIC for both the concentric and eccentric phases were categorized as very high (>60% MVIC), high (41- 60% MVIC), moderate (21-40% MVIC), and low (<21% MVIC) activation (Escamilla et al., 2010). All recorded values after each exercise condition were analyzed as mean OMNI-Res for OMNI-Res.

Statistical Analyses

The power analysis of the sample size showed an effect size of 0.32 SD with an α level of 0.05 and power at 0.95 calculated with G*Power (version 3.1.9.6; University of Dusseldorf, Dusseldorf, Germany). The Shapiro-Wilk test was used to determine if dependent variables were normally distributed, except the OMNI-Res. The dependent variables were: i) muscle activity (external oblique, internal oblique, anterior deltoid, triceps brachii, sternal and clavicular portion of pectoralis major), ii) mean value in these muscles (global activity), and iii) the OMNI-Res values. Data from all dependent variables were shown as mean $\pm$ standard error of the mean (SE). An inferential parametric test, a linear mixed model, was carried out to determine the acute effects of suspended push-ups conditions (non-vibration, 25 Hz and 40 Hz) on each analyzed muscle and the global activity. The linear mixed model used muscle activity and global activity

as dependent variables, suspended push-ups conditions as the fixed effect, and participants as the random effect. If the linear mixed model showed a statistically significant fixed effect, post hoc comparisons were conducted. Likewise, magnitude-based inferences were calculated between non-vibration and vibration conditions (25 Hz and 40 Hz) under the concentric phase to determine whether the vibration condition had a harmful, trivial, or beneficial effect from non-vibration. Magnitude-based inferences were calculated using the spreadsheet *compatibility limits and magnitude-based decisions from p values* (Hopkins, 2007). Furthermore, the effect of suspended push-ups conditions on OMNI-Res was established using a non-parametric Friedman test. A post hoc Wilcoxon test with the Bonferroni correction was carried out in case of a significant main effect. Cohen's (1988) *d* effect size with 90% confidence intervals (CI) were calculated and interpreted as trivial (<0.2), small (from 0.2 to 0.6), moderate (from 0.6 to 1.2), large (from 1.2 to 2.0), and very large ($d>2.0$) (Hopkins et al., 2009). The SPSS statistical software (version 26; SPSS Inc., Chicago, IL, USA) was used to conduct the statistical data analyses setting the *p*-value at <0.05 .

Results

The normalized sEMG (% MVIC) values for each analyzed muscles under suspended push-ups conditions for concentric and eccentric phase are shown in Table 1, with the fixed effect of the exercise condition on muscle activity. For the concentric phase, the suspended push-ups with superimposed vibration (25 Hz and 40 Hz) showed a significant higher activity on left (25 Hz: $p = 0.036$, $d = 0.34$; 40 Hz: $p = 0.003$, $d = 0.48$) and right external oblique (25 Hz: $p = 0.004$, $d = 0.36$; 40 Hz: $p = 0.000$, $d = 0.59$), anterior deltoid (25 Hz: $p = 0.032$, $d = 0.44$; 40 Hz: $p = 0.003$, $d = 0.64$) and global activity (25 Hz: $p = 0.000$, $d = 0.55$; 40 Hz: $p = 0.000$, $d = 0.83$) compared to non-vibration condition. Superimposed vibration at 25 Hz on the suspension strap provoked a significant small increase on the sternal head of pectoralis major compared to non-vibration condition ($p = 0.007$, $d = 0.39$). For triceps brachii and clavicular head of pectoralis major a significant small increase on activity was found under suspended push-up at 40 Hz compared to non-vibration condition ($p = 0.007$, $d = 0.47$, $p = 0.000$, $d = 0.60$; respectively). For the eccentric phase, superimposed vibration (25 Hz and 40 Hz) significantly increased left (25 Hz: $p = 0.034$, $d = 0.41$; 40 Hz: $p = 0.002$, $d = 0.53$) and right external oblique (25 Hz: $p = 0.024$, $d = 0.33$; 40 Hz: $p = 0.000$, $d = 0.64$), and the sternal head of pectoralis major activity (25 Hz: $p = 0.013$, $d = 0.35$; 40 Hz: $p = 0.000$, $d = 0.51$) compared to suspended push-up without vibration. Additionally, a significant small increase of right external oblique activity was found under suspended push-up at 40 Hz, in comparison with superimposed vibration at 25 Hz ($p = 0.035$, $d = 0.29$).

*** Please, insert Table 1 near here***

For the concentric phase, magnitude-based inference analysis showed a possibly positive effect for anterior deltoid and global activity under suspended push-ups at 25 Hz compared to

non-vibration conditions. The other effects were likely trivial on the right and left external oblique, the sternal and clavicular head of pectoralis major activity, and unclear for triceps brachii while performing the suspended push-up with superimposed vibration at 25 Hz (Table 2). Suspended push-ups at 40 Hz showed a likely positive effect on the pectoralis major's right external oblique, anterior deltoid, and clavicular head. A possibly positive effect of superimposed vibration at 40 Hz on the left external oblique and triceps brachii was observed, and a very likely positive effect on global activity (Table 3). Moreover, the standardized differences at 90% CI and the chances of a superimposed vibration's beneficial, trivial, or harmful effect on suspended push-ups are represented as forest plots (Figs. 1 and 2).

Please insert Table 2 near here

Please insert Table 3 near here

Please insert Figure 1 near here

*** Please insert Figure 2 near here***

Fig. 3 shows the OMNI-Res comparison under suspended push-up conditions. A significant main effect was found on suspended push-up conditions on OMNI-Res [$X^2(2) = 26.805$, $p = 0.000$]. The perceived subjective exertion (OMNI-Res) was significantly higher for suspended push-ups at 25 Hz (6.04 ± 0.32 , $p = 0.000$, $d = 4.03$, $CI = 3.27, 4.79$) and 40 Hz (6.21 ± 0.36 , $p = 0.000$, $d = 4.29$, $CI = 3.49, 5.08$) compared to non-vibration (4.75 ± 0.32). A non-significant difference on OMNI-Res was found between suspended push-ups at 25 Hz and 40 Hz ($p = 0.867$, $d = 0.50$, $CI = 0.05, 0.94$).

*** Please insert Figure 3 near here***

Discussion

The present study showed that superimposing vibration to an upper limb suspended push-up is beneficial, increasing the analyzed muscles' global activity. This primary finding reinforces the evidence that combining different strength methods can elicit superior muscular demands (Poston et al., 2007; Mischi & Cardinale, 2009; Moras et al., 2010; Xu, Rabotti & Mischi, 2013). These results can be relevant for coaches and practitioners trying to optimize the time spent in strength and conditioning practices, especially in team-sport settings where time devoted to sessions is limited in-season.

Push-up is one of the most used upper body exercises in sports training. Moreover, it has become even more popular since several authors (Snarr et al., 2013; Calatayud et al., 2014a) have demonstrated that push-ups can be more challenging under suspension or using unstable environments (Calatayud et al., 2014b; de Araújo et al., 2020). In the present study, no comparison was performed between traditional and suspended push-ups; nevertheless, it was hypothesized that the combination of vibration and suspension increases the activation of the primary push-up movers and, probably, the stabilizers of the action. The hypothesis was mainly

confirmed in all the analyzed muscles, especially in the concentric phase at 40Hz. Several authors have demonstrated differentiated effects depending on vibration frequencies, mainly in lower body muscles (Hazell, Jakobi & Kenno, 2007; Di Giminiani et al., 2013). However, the effects on the upper body muscles are still unclear. In this vein, the addition of vibration at a higher rate of frequencies tested (30-40 Hz) has been shown as an activity enhancer in this and similar exercises performed in a vibration plate (Ashnagar et al., 2016; Grant et al., 2019). Thus, superimposing vibration seems to be a proper strategy to enhance muscle activity in a suspended push-up. However, no significant differences were found between 25Hz and 40Hz when overall muscle activity was considered.

The anterior deltoid is the most differentially demanded muscle under superimposed vibration (Grant et al., 2019) or oscillatory vibration exercises (Arora et al., 2013). Its role as a prime mover of shoulder adduction and stabilizer of the shoulder joint, together with the proximity to the vertical plane of the vibration transmission, might explain this finding (Aguilera-Castells et al., 2021). Furthermore, body inclination (20° to 33°), type of grip, and angle between the straps and the floor reinforce the role of the anterior deltoid in stabilizing the shoulder. However, this muscle is less active under unstable conditions (Calatayud et al., 2014b; Borreani et al., 2015; Youdas et al., 2020). As contributors to arm flexo-extension and shoulder adduction, these two muscle heads are close to the vertical plane and proximal to vertical vibration transmitted through the suspension strap. The activity of the anterior deltoid is probably not enough to dampen vibration. Both heads of the pectoralis major have the additional work to perform and stabilize the suspended dynamic push-up, especially the clavicular head at 40 Hz (Fig. 2) closer to the vibration point. Furthermore, the technique used in this study, with straps situated inside the grip, could explain the present findings. Indeed, this type of grip, with less distance between the two handles, makes the action more unstable, and the main involved muscles are overstimulated by the effect of vibration (Aguilera-Castells et al., 2019). This might also be the case with triceps brachii at 40 Hz. If superimposed vibration improves the quality of the strength exercises that recruit this muscle by raising the muscle activation, this effect could potentially help to reduce injuries in overhead athletes. Deltoids are synergists of the rotator cuff muscles; these muscles are typically torn by overuse when athletes present shoulder impingement (Page, 2011), one of the most frequent injuries in these disciplines.

Although the action plane of this muscle during most of the range of movement is not vertical, the action of the triceps brachii before the complete extension of the arm at the end of the concentric phase is aligned with the vertical transmission of vibration and thus overstimulated. The effect is even more apparent in the adducted technique proposed in the present study (Cogley et al., 2005). This conclusion agrees with Moras et al. (2010) comparing the vibration effects of pushing a vibratory bar and Mischi and Cardinale (2009) pushing an electromagnetic arm actuator involving biceps and triceps brachii. Both studies used frequencies around 30Hz.

According to Chen et al. (2019), in the present study, both external obliques were significantly higher stimulated with the superimposition of vibration. The external oblique is

located on the lateral and anterior parts of the abdomen. It is a broad, thin, and irregularly quadrilateral muscle whose muscular portion holds the side. Its aponeurosis is in the anterior wall of the abdomen, and the anterior internal oblique is deep below the anterior external oblique. Thus, in contrast to other studies where no effects were found for the most distal muscles from the vibration exposure point (Aguilera-Castells et al., 2021), the present findings evidenced the increased activation of abdominals (Wirth, Zurfluh & Müller, 2011). The superior fatigue of the core muscles induced by the vibration, especially in suspended exercises, might be a relevant factor (Behm et al., 2010; Mok et al., 2015). Indeed, core muscles need to use more energy to maintain posture in an unstable environment, and muscle activity increases (Cass et al., 2012; Panza et al., 2014). Again, the role of core muscles in athletic performance and injury prevention it is not negligible (Cissik, 2011). Higher activation of these muscles by means of superimposed vibration may have a superior protective effect in athletes at risk.

OMNI-Res results showed a significant increase in effort perception in both vibration frequencies for the non-vibration condition. Similarly, Marin et al. (2012b) found significant increased RPE when performing a squat + biceps curl on a WBV platform and Aguilera-Castells et al. (2021) in suspended supine bridge and hamstring curl superimposing vibration. The authors found significant OMNI-Res increases in all vibration conditions in this work. This finding suggests that the superimposition of vibration is always perceived as a more demanding condition (Marín et al., 2012a).

The present study investigated a limited variety of vibration frequencies. Although the used frequencies are the most studied, lower to 25Hz and higher than 40Hz should be considered for future research. In addition, the number of sEMG channels limited the number of muscles analyzed. Thus, one could have observed the role played by other allegedly secondary muscles during the different phases of the push-up exercise. Indeed, the cocontraction phenomenon between agonists and antagonists of an unstable upper limb task (Behm & Anderson, 2006) can be explored when adding the additional stimulus provided by the vibratory system (Rodríguez Jiménez et al., 2015). Since the exercise was dynamic, the wired system used for the sEMG assessment and its compatibility with the suspended push-up forced the investigators to choose a proper technique for avoiding electrode removals. For this reason, the effects of vibration found in the studied push-up technique might not be generalized to other types of execution. The present study was conducted with trained individuals. All of them experienced in suspension training and with total movement control, even under vibrating conditions. However, this might not happen in less experienced populations, where motion control should be guaranteed. Other protocols used accelerometers for this purpose (Buscà et al., 2020).

Conclusions

Superimposing vibration seems to be a proper strategy to enhance muscle activity in suspended push-ups. 25Hz and 40Hz frequencies provoked similar effects on global activity, and all the muscles analyzed, except in triceps brachii and anterior deltoid in the eccentric phase. Nevertheless, no differences were found between the two frequencies, except in the right

external oblique. Vibration also led to a higher value of subjective perception of exertion (OMNI-Res), but no differences were found between the two tested frequencies.

Acknowledgements

The authors thank all the study participants for their time and effort. The authors also thank Mr. Abel Folk for his collaboration in data acquisition.

References

- Aguilera-Castells J, Buscà B, Arboix-Alió J, Miró A, Fort-Vanmeerhaeghe A, Peña J. 2021. sEMG activity in superimposed vibration on suspended supine bridge and hamstring curl. *Frontiers in Physiology* 12:1–14. DOI: 10.3389/fphys.2021.712471.
- Aguilera-Castells J, Buscà B, Morales J, Solana-Tramunt M, Fort-Vanmeerhaeghe A, Rey-Abella F, Bantulà J, Peña J. 2019. Muscle activity of Bulgarian squat. Effects of additional vibration, suspension and unstable surface. *PLoS ONE* 14:e0221710. DOI: 10.1371/journal.pone.0221710.
- de Araújo RC, Nascimento VYS, Torres RJB, Trombini-Souza F, Behm D, Pitangui ACR. 2020. Can the use of unstable surfaces and instruction for conscious abdominal contraction increase the emg activity of the periscapular muscles during the dynamic push-up? *Journal of Sport Rehabilitation* 29:225–230. DOI: 10.1123/jsr.2018-0165.
- Arora S, Button DC, Basset FA, Behm DG. 2013. The effect of double versus single oscillating exercise devices on trunk and limb muscle activation. *International journal of sports physical therapy* 8:370–380.
- Ashnagar Z, Shadmehr A, Hadian M, Talebian S, Jalaei S. 2016. The effects of whole body vibration on EMG activity of the upper extremity muscles in static modified push up position. *Journal of Back and Musculoskeletal Rehabilitation* 29:557–563. DOI: 10.3233/BMR-160657.
- Behm D, Anderson K. 2006. The role of instability with resistance training. *The Journal of Strength and Conditioning Research* 20:716–722. DOI: 10.1519/R-18475.1.
- Behm DG, Drinkwater EJ, Willardson JM, Cowley PM. 2010. The use of instability to train the core musculature. *Applied Physiology, Nutrition, and Metabolism* 35:91–108. DOI: 10.1139/H09-127.
- Bettendorf B. 2010. *TRX suspension training bodyweight exercises: scientific foundations and practical applications*. San Francisco: Fitness Anywhere Inc.
- Borreani S, Calatayud J, Colado JC, Tella V, Moya-Nájera D, Martín F, Rogers ME. 2015. Shoulder muscle activation during stable and suspended push-ups at different heights in healthy subjects. *Physical Therapy in Sport* 16:248–254. DOI: 10.1016/j.ptsp.2014.12.004.
- Bosco C, Cardinale M, Tsarpela O. 1999. Influence of vibration on mechanical power and electromyogram activity in human arm flexor muscles. *European Journal of Applied Physiology* 79:306–311. DOI: 10.1007/s004210050512.
- Buscà B, Aguilera-Castells J, Arboix-Alió J, Miró A, Fort-Vanmeerhaeghe A, Peña J. 2020.

- 437 Influence of the amount of instability on the leg muscle activity during a loaded free barbell
438 half-squat. *International Journal of Environmental Research and Public Health* 17:8046.
439 DOI: 10.3390/ijerph17218046.
- 440 Calatayud J, Borreani S, Colado JC, Martín F, Rogers ME. 2014a. Muscle activity levels in
441 upper-body push exercises with different loads and stability conditions. *The Physician and*
442 *Sportsmedicine* 42:106–119. DOI: 10.3810/psm.2014.11.2097.
- 443 Calatayud J, Borreani S, Colado JC, Martín FF, Rogers ME, Behm DG, Andersen LL. 2014b.
444 Muscle activation during push-ups with different suspension training systems. *Journal of*
445 *sports science & medicine* 13:502–510.
- 446 Cardinale M, Bosco C. 2003. The use of vibration as an exercise intervention. *Exercise and sport*
447 *sciences reviews* 31:3–7.
- 448 Cardinale M, Lim J. 2003. The acute effects of two different whole body vibration frequencies
449 on vertical jump performance. *Medicina Dello Sport* 56:287–292.
- 450 Cardinale M, Wakeling J. 2005. Whole body vibration exercise: are vibrations good for you?
451 *British Journal of Sports Medicine* 39:585–589. DOI: 10.1136/bjsm.2005.016857.
- 452 Chen B, Dong Y, Guo J, Zheng Y, Zhang J, Wang X. 2019. Effects of whole-body vibration on
453 lumbar-abdominal muscles activation in healthy young adults: A pilot study. *Medical*
454 *Science Monitor* 25:1945–1951. DOI: 10.12659/MSM.912720.
- 455 Cissik JM. 2011. The role of core training in athletic performance, injury prevention, and injury
456 treatment. *Strength & Conditioning Journal* 33:10–15. DOI:
457 10.1519/SSC.0b013e3182076ac3.
- 458 Cloak R, Nevill A, Day S, Wyon M. 2013. Six-week combined vibration and wobble board
459 training on balance and stability in footballers with functional ankle instability. *Clinical*
460 *Journal of Sport Medicine* 23:384–391. DOI: 10.1097/JSM.0b013e318291d22d.
- 461 Cochrane DJ, Hawke EJ. 2007. Effects of acute upper-body vibration on strength and power
462 variables in climbers. *The Journal of Strength and Conditioning Research* 21:527. DOI:
463 10.1519/R-18505.1.
- 464 Cogley RM, Archambault TA, Fibeger JF, Koverman MM, Youdas JW, Hollman JH. 2005.
465 Comparison of muscle activation using various hand positions during the push-up exercise.
466 *The Journal of Strength and Conditioning Research* 19:628–633. DOI: 10.1519/15094.1.
- 467 Cohen J. 1988. *Statistical power analysis for the behavioral sciences*. Hilldale: Lawrence
468 Erlbaum.
- 469 Criswell E, Cram J. 2011. *Cram's introduction to surface electromyography*. Sudbury: Jones and
470 Bartlett.
- 471 Cuğ M, Ak E, Ozdemir RA, Korkusuz F, Behm DG. 2012. The effect of instability training on
472 knee joint proprioception and core strength. *Journal of sports science & medicine* 11:468–
473 474.
- 474 Escamilla RF, Lewis C, Bell D, Bramblett G, Daffron J, Lambert S, Pecson A, Imamura R,
475 Paulos L, Andrews JR. 2010. Core muscle activation during swiss ball and traditional
476 abdominal exercises. *Journal of Orthopaedic & Sports Physical Therapy* 40:265–276. DOI:

- 10.2519/jospt.2010.3073.
- Di Giminiani R, Masedu F, Tihanyi J, Scrimaglio R, Valenti M. 2013. The interaction between body position and vibration frequency on acute response to whole body vibration. *Journal of Electromyography and Kinesiology* 23:245–251. DOI: 10.1016/j.jelekin.2012.08.018.
- Gollhofer A. 2010. Functional adaptations to short and long term WBV-interventions. *Journal of Science and Medicine in Sport* 13:e101. DOI: 10.1016/j.jsams.2010.10.676.
- Grant MJ, Hawkes DH, McMahon J, Horsley I, Khaiyat OA. 2019. Vibration as an adjunct to exercise: its impact on shoulder muscle activation. *European Journal of Applied Physiology* 119:1789–1798. DOI: 10.1007/s00421-019-04168-9.
- Halaki M, Ginn K. 2012. Normalization of EMG signals: to normalize or not to normalize and what to normalize to? In: Naik G ed. *Computational intelligence in electromyography analysis - a perspective on current applications and future challenges*. London: InTech, 175–194. DOI: 10.5772/49957.
- Hazell TJ, Jakobi JM, Kenno KA. 2007. The effects of whole-body vibration on upper- and lower-body EMG during static and dynamic contractions. *Applied Physiology, Nutrition, and Metabolism* 32:1156–1163. DOI: 10.1139/H07-116.
- Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. 2000. Development of recommendations for SEMG sensors and sensor placement procedures. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology* 10:361–74.
- Hopkins WG. 2007. A spreadsheet for deriving a confidence interval, mechanistic inference and clinical inference from p value. *Sportscience* 11:16–20.
- Hopkins WG, Marshall SW, Batterham AM, Hanin J. 2009. Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise* 41:3–12. DOI: 10.1249/MSS.0b013e31818cb278.
- Issurin V, Cucca C, Temnov P, Tenenbaum G. 2010. Acute effect of strength exercises with superimposed vibration: amplitude of oscillations. *Acta Kinesiologiae Universitatis Tartuensis* 15:7–22.
- Issurin VB, Tenenbaum G. 1999. Acute and residual effects of vibratory stimulation on explosive strength in elite and amateur athletes. *Journal of Sports Sciences* 17:177–182. DOI: 10.1080/026404199366073.
- Jakobsen MD, Sundstrup E, Andersen CH, Aagaard P, Andersen L. 2013. Muscle activity during leg strengthening exercise using free weights and elastic resistance: effects of ballistic vs controlled contractions. *Human Movement Science* 32:65–78. DOI: 10.1016/j.humov.2012.07.002.
- Konrad P. 2006. *The ABC of EMG: a practical introduction of kinesiological electromyography*. Scottsdale: Noraxon INC.
- Manimmanakorn N, Hamlin MJ, Ross JJ, Manimmanakorn A. 2014. Long-term effect of whole body vibration training on jump height. *Journal of Strength and Conditioning Research* 28:1739–1750. DOI: 10.1519/JSC.0000000000000320.

- Marín PJ, Hazell TJ. 2014. Effects of whole-body vibration with an unstable surface on muscle activation. *Journal of Musculoskeletal Neuronal Interactions* 14:213–219.
- Marín PJ, Santos-Lozano A, Santin-Medeiros F, Robertson RJ, Garatachea N. 2012a. Reliability and validity of the OMNI-vibration exercise scale of perceived exertion. *Journal of sports science & medicine* 11:438–443.
- Marín PJ, Santos-Lozano A, Santin-Medeiros F, Vicente-Rodriguez G, Casajús JA, Hazell TJ, Garatachea N. 2012b. Whole-body vibration increases upper and lower body muscle activity in older adults: Potential use of vibration accessories. *Journal of Electromyography and Kinesiology* 22:456–462. DOI: 10.1016/j.jelekin.2012.02.003.
- Martin BJ, Park H-S. 1997. Analysis of the tonic vibration reflex: influence of vibration variables on motor unit synchronization and fatigue. *European journal of applied physiology and occupational physiology* 75:504–511.
- Mischi M, Cardinale M. 2009. The effects of a 28-Hz vibration on arm muscle activity during isometric exercise. *Medicine & Science in Sports & Exercise* 41:645–653. DOI: 10.1249/MSS.0b013e31818a8a69.
- Mok NW, Yeung EW, Cho JC, Hui SC, Liu KC, Pang CH. 2015. Core muscle activity during suspension exercises. *Journal of Science and Medicine in Sport* 18:189–194. DOI: 10.1016/j.jsams.2014.01.002.
- Moras G, Rodríguez-Jiménez S, Tous-Fajardo J, Ranz D, Mujika I. 2010. A vibratory bar for upper body: feasibility and acute effects on EMG activity. *Journal of Strength and Conditioning Research* 24:2132–2142. DOI: 10.1519/JSC.0b013e3181aa3684.
- Page P. 2011. Shoulder muscle imbalance and subacromial impingement syndrome in overhead athletes. *International journal of sports physical therapy* 6:51–58.
- Panza P, Vianna JM, Damasceno VO, Aranda LC, Melibe Bentes C, da Silva Novaes J, Behm DG. 2014. Energy cost, number of maximum repetitions, and rating of perceived exertion in resistance exercise with stable and unstable platforms. *Journal of Exercise Physiology Online* 17:77–87.
- Poston B, Holcomb WR, Guadagnoli MA, Linn LL. 2007. The acute effects of mechanical vibration on power output in the bench press. *Journal of Strength and Conditioning Research* 21:199–203. DOI: 10.1519/00124278-200702000-00036.
- Ritzmann R, Kramer A, Bernhardt S, Gollhofer A. 2014. Whole body vibration training - improving balance control and muscle endurance. *PLoS ONE* 9:e89905. DOI: 10.1371/journal.pone.0089905.
- Robertson R, Goss F, Rutkowski J, Brooke L, Dixon C, Timmer J, Frazee K, Dube J, Andreacci J. 2003. Concurrent validation of the OMNI perceived exertion scale for resistance exercise. *Medicine & Science in Sports & Exercise* 35:333–341. DOI: 10.1249/01.MSS.0000048831.15016.2A.
- Rodríguez Jiménez S, Benítez A, García González MA, Moras Feliu G, Maffiuletti NA. 2015. Effect of vibration frequency on agonist and antagonist arm muscle activity. *European Journal of Applied Physiology* 115:1305–1312. DOI: 10.1007/s00421-015-3108-x.

- Rønnestad BR. 2009a. Acute effects of various whole-body vibration frequencies on lower-body power in trained and untrained subjects. *The Journal of Strength & Conditioning Research* 23:1309–1315. DOI: 10.1519/JSC.0b013e318199d720.
- Rønnestad BR. 2009b. Acute effects of various whole body vibration frequencies on 1RM in trained and untrained subjects. *Journal of Strength and Conditioning Research* 23:2068–2072. DOI: 10.1519/JSC.0b013e3181b8652d.
- La Scala Teixeira C V., Evangelista AL, Pereira PE de A, Da Silva-Grigoletto ME, Bocalini DS, Behm DG. 2019. Complexity: a novel load progression strategy in strength training. *Frontiers in Physiology* 10:1–5. DOI: 10.3389/fphys.2019.00839.
- Sierra-Guzmán R, Jiménez-Díaz F, Ramírez C, Esteban P, Abián-Vicén J. 2018. Whole-body–vibration training and balance in recreational athletes with chronic ankle instability. *Journal of Athletic Training* 53:355–363. DOI: 10.4085/1062-6050-547-16.
- Snarr R, Esco M, Witte E, Jenkins C, Brannan R. 2013. Electromyographic activity of rectus abdominis during a suspension push-up compared to traditional exercises. *Journal of Exercise Physiology* 16:1–8.
- Warburton DER, Jamnik VK, Bredin SSD, McKenzie DC, Stone J, Shephard RJ, Gledhill N. 2011. Evidence-based risk assessment and recommendations for physical activity clearance: an introduction. *Applied Physiology, Nutrition, and Metabolism* 36:S1–S2. DOI: 10.1139/h11-060.
- Wirth B, Zurfluh S, Müller R. 2011. Acute effects of whole-body vibration on trunk muscles in young healthy adults. *Journal of Electromyography and Kinesiology* 21:450–457. DOI: 10.1016/j.jelekin.2010.12.007.
- Xu L, Negro F, Xu Y, Rabotti C, Schep G, Farina D, Mischi M. 2018. Does vibration superimposed on low-level isometric contraction alter motor unit recruitment strategy? *Journal of Neural Engineering* 15:066001. DOI: 10.1088/1741-2552/aadc43.
- Xu L, Rabotti C, Mischi M. 2013. Novel vibration-exercise instrument with dedicated adaptive filtering for electromyographic investigation of neuromuscular activation. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* 21:275–282. DOI: 10.1109/TNSRE.2012.2219555.
- Xu L, Rabotti C, Mischi M. 2015. On the nature of the electromyographic signals recorded during vibration exercise. *European Journal of Applied Physiology* 115:1095–1106. DOI: 10.1007/s00421-014-3091-7.
- Youdas JW, Baartman HE, Gahlon BJ, Kohnen TJ, Sparling RJ, Hollman JH. 2020. Recruitment of shoulder prime movers and torso stabilizers during push-up exercises using a suspension training system. *Journal of Sport Rehabilitation* 29:993–1000. DOI: 10.1123/jsr.2019-0381.

Table 1 (on next page)

sEMG activity for each analyzed muscle under suspended push-up conditions.

** significantly different with non-vibration condition. **significantly different with vibration at 25 Hz condition. ^aData presented as normalized muscle activity (%MVIC). ^bGlobal activity = mean of the six muscles. C = clavicular head; L = left; R = right; SE = standard error of the mean; S = sternal head.*

1

Muscles ^a	Suspended push-up			F	<i>p</i>
	Non-Vibration	Vibration at 25 Hz	Vibration at 40 Hz		
	Mean ± SE	Mean ± SE	Mean ± SE		
Concentric phase					
External oblique_R	4.30±0.40	5.08±0.43 [*]	5.60±0.44 [*]	15.81	0.000
External oblique_L	4.08±0.40	4.82±0.44 [*]	5.09±0.40 [*]	6.67	0.003
Triceps brachii	29.04±1.18	30.26±1.71	33.36±2.14 [*]	5.46	0.007
Anterior deltoid	41.18±2.10	47.05±2.86 [*]	48.84±2.39 [*]	6.51	0.003
Pectoralis major_S	24.81±2.02	29.23±2.27 [*]	27.77±2.20	5.32	0.008
Pectoralis major_C	38.67±2.47	42.26±2.44	46.16±2.28 [*]	9.99	0.000
Global activity ^b	23.68±0.85	26.45±1.05 [*]	27.80±1.02 [*]	24.15	0.000
Eccentric phase					
External oblique_R	4.46±0.35	5.11±0.40 [*]	5.72±0.40 ^{*#}	14.34	0.000
External oblique_L	4.18±0.34	4.96±0.37 [*]	5.27±0.43 [*]	7.17	0.002
Triceps brachii	26.55±1.48	28.54±1.81	27.44±1.83	1.10	0.337
Anterior deltoid	43.22±2.64	42.13±3.08	39.78±3.37	1.35	0.266
Pectoralis major_S	19.08±1.26	22.09±1.92 [*]	23.42±1.90 [*]	9.73	0.000
Pectoralis major_C	31.50±3.09	32.62±3.22	35.10±3.01	2.54	0.087
Global activity ^b	21.50±1.01	22.57±1.21	22.79±1.18	2.84	0.066

2

Table 2 (on next page)

Changes in sEMG activity for each analyzed muscle between non-vibration and 25 Hz vibration suspended push-up at concentric phase.

^aData presented as normalized muscle activity (%MVIC). ^bGlobal activity = mean of the six muscles. C = clavicular head; CI = confidence interval; L = left; R = right; SE = standard error of the mean; S = sternal head.

1

2

Muscles ^a	Suspended push-up		Percent change (90% CI)	Standardized difference (90% CI)	Chances of beneficial/ trivial/ harmful effect	Qualitative outcome
	Non- Vibration (Mean ± SE)	Vibration at 25 Hz (Mean ± SE)				
External oblique_R	4.30±0.40	5.08±0.43	18.15 (15.30; 21.00)	0.36 (0.17; 0.55)	11/88/0%	Likely trivial
External oblique_L	4.08±0.40	4.82±0.44	18.32 (15.45; 21.19)	0.34 (0.08; 0.60)	15/84/0%	Likely trivial
Triceps brachii	29.04±1.18	30.26±1.71	4.18 (3.40; 4.97)	0.16 (-0.21; 0.22)	48/3/47%	Unclear
Anterior deltoid	41.18±2.10	47.05±2.86	14.25 (12.00; 16.51)	0.44 (0.11; 0.77)	38/62/0%	Possibly positive
Pectoralis major_S	24.81±2.02	29.23±2.27	17.81 (15.02; 20.61)	0.39 (0.16; 0.62)	20/79/0%	Likely trivial
Pectoralis major_C	38.67±2.47	42.26±2.44	9.28 (7.78; 10.79)	0.28 (-0.01; 0.57)	10/89/0%	Likely trivial
Global activity^b	23.68±0.85	26.45±1.05	11.70 (9.83; 13.56)	0.55 (0.34; 0.76)	65/34/0%	Possibly positive

3

Table 3 (on next page)

Changes in sEMG activity for each analyzed muscle between non-vibration and 40 Hz vibration suspended push-up at concentric phase.

^aData presented as normalized muscle activity (%MVIC). ^bGlobal activity = mean of the six muscles. C = clavicular head; CI = confidence interval; L = left; R = right; SE = standard error of the mean; S = sternal head.

1

2

Muscles ^a	Suspended push-up		Percent change (90% CI)	Standardized difference (90% CI)	Chances of beneficial/ trivial/ harmful effect	Qualitative outcome
	Non- Vibration (Mean ± SE)	Vibration at 40 Hz (Mean ± SE)				
External oblique_R	4.30±0.40	5.60±0.44	30.36 (25.64; 35.09)	0.59 (0.37; 0.81)	75/24/0%	Likely positive
External oblique_L	4.08±0.40	5.09±0.40	24.85 (20.97; 28.72)	0.48 (0.23; 0.73)	44/55/0%	Possibly positive
Triceps brachii	29.04±1.18	33.36±2.14	14.87 (12.53; 17.22)	0.47 (0.20; 0.74)	42/57/0%	Possibly positive
Anterior deltoid	41.18±2.10	48.84±2.39	18.62 (15.70; 21.54)	0.64 (0.31; 0.97)	75/24/0%	Likely positive
Pectoralis major_S	24.81±2.02	27.77±2.20	11.95 (10.05; 13.85)	0.26 (-0.01; 0.53)	6/93/0%	Likely trivial
Pectoralis major_C	38.67±2.47	46.16±2.28	19.37 (16.34; 22.40)	0.60 (0.38; 0.82)	77/22/0%	Likely positive
Global activity^b	23.68±0.85	27.80±1.02	17.43 (14.69; 20.16)	0.83 (0.52; 1.10)	95/4/0%	Very likely positive

Figure 1

Acute effects of superimposed vibration at 25 Hz and suspended push-up without vibration on sEMG activity for each analyzed muscle at concentric phase.

Bars represent the 90% confidence interval for the effect of superimposed vibration at 25 Hz on suspended push-up. Dotted lines represent the smallest substantial threshold. Numbers show the quantitative chance (%) that the true effect of superimposed vibration at 25 Hz is harmful, trivial, or beneficial compared to non-vibration. C = clavicular head; Global activity = mean of the six muscles; L = left; R = right; S = sternal head.

Suspended push-up conditions (concentric phase)




Figure 2

Acute effects of superimposed vibration at 40 Hz and suspended push-up without vibration on sEMG activity for each analyzed muscle at concentric phase.

Bars represent the 90% confidence interval for the effect of superimposed vibration at 40 Hz on suspended push-up. Dotted lines represent the smallest substantial threshold. Numbers show the quantitative chance (%) that the true effect of superimposed vibration at 40 Hz is harmful, trivial, or beneficial compared to non-vibration. C = clavicular head; Global activity = mean of the six muscles; L = left; R = right; S = sternal head.

Suspended push-up conditions (concentric phase)




Figure 3

Perceived subjective exertion under suspended push-up conditions.

*Bars represent the mean of OMNI-Res values, and the error bars represent the standard error of the mean (SE). A.U. = arbitrary units. * significantly different from the non-vibration condition.*

