

Surviving on the edge: present and future effects of climate warming on the common frog (*Rana temporaria*) population in the Montseny massif (NE Iberia). (#75330)

1

First revision

Guidance from your Editor

Please submit by **29 Oct 2022** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

11 Figure file(s)

2 Table file(s)

4 Raw data file(s)

! Custom checks

Vertebrate animal usage checks



Have you checked the authors [ethical approval statement](#)?



Were the experiments necessary and ethical?



Have you checked our [animal research policies](#)?

Field study



Have you checked the authors [field study permits](#)?



Are the field study permits appropriate?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Surviving on the edge: present and future effects of climate warming on the common frog (*Rana temporaria*) population in the Montseny massif (NE Iberia).

Albert Montori ^{Corresp., 1}, Fèlix Amat ²

¹ Herpetology, Centre de Recerca i Estudis Ambientals de Calafell (CREAC/GREN), Calafell, Catalonia, Spain

² Herpetological Section, Natural History Museum of Granollers, Granollers, Catalonia, Spain

Corresponding Author: Albert Montori
Email address: amontori@gmail.com

The Montseny massif shelters one of the southernmost western populations of Common frogs (*Rana temporaria*) that lives in a Mediterranean climate, one which poses a challenge for the species' persistence in a scenario of rising temperatures. We evaluated the effect of climate change at three levels. Firstly, we analysed if there has been an advancement in the onset of spawning period due to the increase in temperatures. Secondly, we analysed the impact of climatic variables on the onset of the spawning period and, thirdly, how the distribution of this species could vary according to the predictions with regard to rising temperatures for the end of this century. From 2009 to 2021, we found there had been an increase in temperatures of 0,439 °C/decade, more than the 0,1°C indicated by estimates for the second half of the previous century. We found an advancement in the onset of the reproduction process of 26 days/decade for the period 2009-2022, a change that has been even more marked during the last eight years, when data were annually recorded. Minimum temperatures and the absence of frost days in the week prior to the onset of the spawning period determine the start of reproduction. Predictions on habitat availability for spawning provided by climatic niche analysis for the period 2021-2100 show a potential contraction of the species range in the Montseny and, remarkably, much isolation from the neighbouring populations.

Surviving on the edge: present and future effects of climate warming on the common frog (*Rana temporaria*) population in the Montseny massif (NE Iberia).

Albert Montori ¹, Fèlix Amat ²

¹ Centre de Recerca i Estudis Ambientals de Calafell (CREAC/GREN). Barcelona, Catalonia Spain.. amontori@gmail.com

² Herpetological Section, Natural History Museum of Granollers, C/ Palaudàries, 102, 08403 Granollers, Catalonia, Spain

Corresponding Author:

Albert Montori ¹

Passeig del Montseny, 40. 3r. 08474 Gualba. Barcelona, Catalonia, Spain.

Email address: amontori@gmail.com

Abstract

The Montseny massif shelters one of the southernmost western populations of Common frogs (*Rana temporaria*) that lives in a Mediterranean climate, one which poses a challenge for the species' persistence in a scenario of rising temperatures. We evaluated the effect of climate change at three levels. Firstly, we analysed if there has been an advancement in the onset of spawning period due to the increase in temperatures. Secondly, we analysed the impact of climatic variables on the onset of the spawning period and, thirdly, how the distribution of this species could vary according to the predictions with regard to rising temperatures for the end of this century. From 2009 to 2021, we found there had been an increase in temperatures of 0,439 °C/decade, more than the 0,1°C indicated by estimates for the second half of the previous century. We found an advancement in the onset of the reproduction process of 26 days/decade for the period 2009-2022, a change that has been even more marked during the last eight years, when data were annually recorded. Minimum temperatures and the absence of frost days in the week prior to the onset of the spawning period determine the start of reproduction. Predictions on habitat availability for spawning provided by climatic niche analysis for the period 2021-2100 show a potential contraction of the species range in the Montseny and, remarkably, much isolation from the neighbouring populations.

Keywords: Climate change, *Rana temporaria*, Onset spawn, Phenology, Habitat suitability

Introduction

Climate change due to increases in human greenhouse gas emissions since the beginning of the industrial revolution up to the present day is now considered one of the major threats to biodiversity in the twenty-first century and one which underlies several reported extinction events (Thomas et al. 2004; Parmesan 2006; Thuiller et al. 2011). Climate is rapidly changing and the same trend that points towards increases in global temperatures has been forecast for the coming decades (IPCC, 2021; Seneviratne, et al. 2014). Understanding how species respond to increases in temperatures and changes in precipitation, both at a population and species level, is one of the most important questions to be answered. With regard to amphibians, Sheridan et al. (2017) indicate that changes in body size and breeding phenology are the two major ecological consequences of climate change in the wood frog (*Lithobates sylvaticus*), but it is unclear how rising temperatures interact with these variables. Reading (2007) reported a reduction of survival, body condition, size, and egg laying in the common toad (*Bufo bufo*), and Blaustein et al. (2010), reported changes in developmental rates of eggs and larvae.

Changes in temperature or precipitation can potentially influence the timing of amphibian reproduction (Blaustein et al 2010). An earlier onset of spring breeding behaviour correlated with a warming climate has been observed among various pond-breeding frogs and toads in Europe (Beebee, 1995; Tryjanowski et al., 2003; Scott et al., 2008) and Asia (Kusano & Inoue, 2008; Primack et al., 2009). However, other species may exhibit either a trend towards a later onset of spring breeding or no trend at all (Todd et al., 2010; Klaus & Loughheed, 2013). While & Uller (2014) and Ficetola & Maiorano (2016) concluded that this group is strongly influenced by climate change after carrying out a meta-analysis with worldwide phenological data of amphibians. However, they didn't find a clear pattern for the impact climate can have on amphibian population parameters such as abundance, survival, breeding success and morphology. Both authors agreed on the low importance of rainfall changes, the effects of climate change on offspring and the strong relationship between the advancement in the onset of spawning and rising temperatures. However, in some cases, an increase in precipitation could enhance phenological advancement (Timm et al., 2007; Todd et al., 2010; Green, 2017), especially in warm and dry areas where precipitation is a limiting factor (Ficetola and Maiorano, 2016). Shifts in phenology due to climate change have been observed by many authors in several species (Beebee, 1995; Parmesan, 2006; Walther et al., 2002; While & Uller, 2014; Ficetola & Maiorano, 2016), with evidence of earlier breeding with a shifting climate and Chadwik et al. (2006) showed trends of earlier breeding associated with shifts in the climate in

Triturus species. However, findings from several independent studies reveal that the impacts of a changing climate on breeding times may be more complex. With regard to **Common Frog**, *Rana temporaria*, the most recent studies found a close relationship between rising temperatures and an advanced reproduction date (While & Uller, 2014; Ficetola & Maiorano, 2016; Bison, 2021).

One of the research areas pertaining to the effects of climate change on species entails measuring the expected shift in the ranges they will experience in the future. Changes in geographic distribution can occur because climate change affects population growth and their survival or the ability to colonise new areas (McCaffery et al., 2010), mediated by the species' physiological adaptation to a range of environmental conditions (Greenberg & Palen, 2021). Modelling habitat suitability across time using current data is commonly used to predict how amphibian ranges will change, an event shaped by the effect of climate change (see for example García et al., 2013; Duan et al., 2016; Aguelo et al., 2019; Kim et al., 2021). However, the increase in data availability on species distribution gathered over recent decades, has recently shown shifts in the species ranges. For example, Enriquez-Urzelai, et al. (2019a) demonstrated that most Iberian amphibians have changed their altitudinal and latitudinal distributions, moving up to the north or to higher altitudes in the last 20-25 years as a response to rising temperatures.

The **common frog** (~~*Rana temporaria*~~) is widely found across the temperate and cold regions in Europe, from the sea level up to 2,700m a.s.l. reaching a latitude of 70° North, above the Arctic Polar Circle. The southernmost populations of the species are located in the Rodopi mountain range in Greece, in Eastern Europe (Kuzmin et al., 2009). The second most southern rim in which the common frog is found in Western Europe, in the Montseny massif, is the southernmost foothill of the Transversal Catalan Range placed in Northeast Iberia (Figure 1). Common frogs breed in shallow, still, fresh water such as ponds, lakes or marshes. The onset of spawning will vary depending on the altitude and latitude, occurs between February to late June, but generally in March-April over the main part of the range. In the Atlantic region of Iberian Peninsula, the spawning occurs earlier, in November-December, due to the abundant rain and mild temperatures (Bea et al, 1986). Outside the breeding season, common frogs live a solitary life in damp places near ponds or marshes or in long grass.

Although the climate in the Montseny is typically Mediterranean, the massif's altitude and the northern and eastern exposure of the main hillsides provide humid and temperate conditions (Panareda-Clopés, 1979), thus allowing the persistence of populations of this cold-adapted species. Thus, this massif harbours rear edge populations of the Common Frog, where the

species experiences a small temperature buffer in comparison with most of the range, making these populations a good model to study the effects of climate change on temperate amphibians (Hampe & Petit, 2005; Habibzadeh et al., 2021).

Furthermore, the models of climatic change (González-Hidalgo et al., 2009; Peñuelas et al., 2021) for the Mediterranean area forecast more irregularity in precipitation patterns, specifically more rainfall in autumn than in spring, and concentrated over a few days in some years, while in others, there will be heavy droughts. Air temperature is rising (Vicente-Serrano et al. 2014) and in the Montseny the variation in the average air temperature in Turó de l'Home (1712m a.s.l.) for the period 1950-1999 was 0,3°C/decade (Peñuelas & Boada, 2003), and Minuartia (2016), shows a rate of increase with a similar trend (+0.24 °C/decade), for the period 1950-2014, with a higher average seasonal increase in spring and a lower one in autumn.

Our objectives were to understand how the environmental variables influence the onset of breeding phenology of the Common Frog in recent years, and to explore the changes that this species faces in the context of climate change in the edge of distribution in the Montseny massif using niche modelling. Specifically, we examined the following issues: i) we analysed whether there has been an advancement in the onset of the breeding period in recent years as a consequence of rises in global temperatures. ii) how and which meteorological variables influence this onset, and iii) whether the species will experience, in the Montseny Massif, a contraction of its distribution range in the 2100 scenario as a consequence of global warming.

Materials & Methods

Study area

A phenological survey was conducted in the Santa Fe Valley situated on the eastern side of the Montseny massif (UTM31N 5524) at 1,100m a.s.l (Figure 1). At the study site, the monthly mean temperature ranges from 3°C in January to 20°C in August and the mean annual precipitation is approximately 1000 mm, occurring mostly as rain in autumn and spring, with occasional snow in winter. The valley is primarily forested with beech (*Fagus sylvatica*) and divided by streams that produce many wetlands and ponds. We extended the study area to all the populations in the Montseny Massif to gather data for niche modelling analysis based on the known range that comprises the eastern side of the massif where the species lives associated with riparian and beech forests (Amat & Montori in press). Permits for fieldwork were granted by: Diputació de Barcelona (DIBA20140609, Servei de Fauna de la Generalitat de Catalunya: SF/0539/2019,

Onset breeding period

In 2009, and from 2014 to 2022, we recorded the day when the *Rana temporaria*'s onset breeding period began in Santa Fe del Montseny (Barcelona). This area located at 1100 m.a.s.l. is the main breeding area in the Montseny massif. During the week prior to the start of reproduction and until the day of the onset breeding, the known breeding points in the Sta. Fe Valley were surveyed. The day of year (DOY) on which the males flocked to the ponds and the first clutches were detected, was considered the day of the breeding onset. The presence of isolated males was not considered indicative of the day of breeding onset. Other discontinuous data of reproductive activity based on the observation of clutches or amplexus were obtained from reports, scientific publications, technicians and researchers in the Montseny Natural Park (Balcells, 1957; Pascual & Montori, 1982; Montori & Pascual, 1987; Campeny, 2001; Roig & Amat, 2002; <http://ornitho.cat>; <https://siare.herpetologica.es> Martí Boada com pers. and unpublished data of the authors). However, these data are not included in the analyses of the climatic factors influencing the start of the breeding period because they do not reference the day when breeding activity began, but rather, the day in which reproduction was observed. They are only considered to describe the amplitude of the breeding period in Santa Fe del Montseny.

From 2009 to 2021, daily meteorological data were obtained from "Viladrau Ws" Meteorological Station, located in Viladrau (Osona, Barcelona, at 953masl; 41.84008N; 2.41877E). Data over four weeks (28 days) prior to the spawning were taken in order to test how previous meteorological variables influence the onset of the breeding period (Suppl. Mat. Table 1). We chose four weeks because published studies reference the importance of the one to three week period prior to the onset of spawning (Bea et al. 1986; Tryjanowski et al, 2003; Loman, 2016; 2019; Bison et al. 2021). Nine meteorological variables were obtained: temperature (minimal, maximal and average in °C, relative humidity in %, rain in mm, maximum speed wind (in km/h), average of atmospheric pressure in hP, number of Days with temperature below zero <0°C and < 1°C, and Solar Irradiation in MJ/m² (Servei Meteorologic de Catalunya - <https://meteo.cat>).

To test the impact of meteorological variables on the onset of the spawning period, a four-week period prior to the start of the spawning was compared separately with ANOVA Kruskal-Wallis test using the period (week) as a grouping variable. These four periods: 1, 2, 3 and 4 correspond to 1-7, 8-14, 9-21 and 22-28 days before the start of reproduction. A non-parametric test was used in the univariate analysis since some variables did not fulfil the condition of normality (Kolmogorov-Smirnov test).

To observe if the onset of Common Frog spawning activity occurs early in recent years, we performed a linear regression between the day of the onset of reproduction (DOY: day of year) and the year. To determine which meteorological variables influence the onset of egg-spawning period Factorial Analyse, using Principal Components as an extraction method. All statistical analyses were carried out with Statistica 8.0 Software. The variables used in Factorial Analysis analyses are FD4, FD3, FD2, FD1 (Days with temperature below zero degrees four, three, two or one week before the onset of the spawning period), TX4, TX3, TX2 (Temperature Average four, three and two weeks before the onset of the spawning period) and Tm4, Tm3, Tm2, Tm1 (Minimal Temperature four, three, two and one week before the onset of the spawning period). Pluviometry, Maximal temperature and the temperature average in the first week prior to the onset of spawning has been not considered in the Factorial analyses due to low contribution to explained variance in a previous exploratory analysis.

Future distribution range contraction

We collected four-hundred georeferenced observations of *Rana temporaria* at high resolution in the northeastern Iberian range of the species (Figure 1). Nearly forty percent of localities we used originated from the Montseny massif and were recorded by us for twelve years during which we monitored amphibian populations in the natural park. We gathered nineteen bioclimatic variables at 30 seconds resolution from the WorldClimateDatabase (Hijmans et al., 2005, <https://www.worldclim.org/data/cmip6/>) of current and future climatic conditions under the coupled model intercomparison project phase 6 (CMIP6). We used the CNRM-CM6-1 model for a short (2021-2040) and long term (2081-2100) based on the lowest (126) and highest (585) emission scenarios to estimate the species habitat suitability. We selected maximum entropy approach (Phillips et al., 2006) to model habitat suitability using MaxEnt 3.4.1. because this method only uses presence data, performing well even with low sample sizes and providing reasonably accurate and interpretable models with good predictive power (Phillips & Dudík, 2008). We clipped raster layers to the same geographic extent covering the south westernmost margin of the Iberian Peninsula, including the Montseny massif, the Transversal Catalan Range and the Catalan Prepyrenean and Eastern Pyrenees. In order to minimise collinearity among bioclimatic variables, we used R usdm 1.1-8, to calculate the variance inflation factor (VIF) and removed highly correlated variables by performing a stepwise procedure. We checked the model performance using the size of the area under the receiver operating curve (AUC) which indicated high model suitability when it exceeded a threshold value of 0.75 (Pearce & Ferrier, 2000). We performed fifty replicates training the model using 25% of the presence data and quantifying variable importance by using jack-knife. Predicted

species geographic distribution based on climatic suitability under current and future scenarios was represented by means of using the logistic representation of probability of occurrence.

Results

The data on the spawning data over multiple years has been obtained from 1957 to 2022 from reports, databases and personal observations (Suppl. Mat. Table 2). This table summarizes the earliest and latest data of registered spawning for each year. During this period, the earliest amplexus was registered on February 5th, in 2021 and the latest amplexus was registered on March 27th, 1984 (Campeny, 2001). Back then, the reproductive period of *Rana temporaria* in Sta. Fe del Montseny was between early February until the second half of March. However, some sporadic spawning could occur on April (1979, per obs.).

Onset breeding period

When analysed with a linear regression on the DOY of onset spawning data recorded for the 2009 to 2022 period, we found a pattern of advancement in the beginning of reproduction (Figure 2). The onset of reproduction, quantified as day of year (DOY), was inversely correlated with the year. The linear model obtained was significant ($R = -0.7903$; $P = 0.0065$; slope = -2.5959 ; $SE = 2.4974$) and explained 62.45% of the total variation, reflecting that part of the fluctuation in the reproductive onset has advanced over time in the last years. Between 2009 to 2022 the average for an advancement of the onset of reproduction is 26 days/decade on average in the last 14 years (Figure 2). From 1997 to 2021 the average maximal temperature has risen $0.404^{\circ}\text{C}/\text{decade}$ (Suppl. Mat. Figure1 and Suppl. Mat. Table 3). This value is higher than the values published by Peñuelas & Boada (2003) and Minuartia (2016) for the Montseny massif.

The results obtained in the Kruskal-Wallis test between periods (weeks) prior to the onset of breeding activity from 2009 to 2022 related to meteorological variables analysed are shown in Table 1. Only temperature, solar irradiation and the number of days with temperature below zero degrees were significant. However, multiple comparisons indicated that significant differences only exist between the first and fourth week. Rainfall did not seem to be a determining factor (Supplementary Material Figures 2 and 3). In fact, average minimal T^a for the previous week of onset spawning is $2,09^{\circ}\text{C}$ ($STD = 0.88^{\circ}\text{C}$; $Max = 4,07^{\circ}\text{C}$; $Min = 0,9^{\circ}\text{C}$ -see meteorological data in Supplementary Material, Table1-).

The same results show the daily variation of the minimum temperature and the interannual variance average (Figure 3). In fact, the days prior to the start of reproduction, the minimal temperature stabilised and showed less variance than in previous weeks. Figure 4 shows the relationship between the DOY of the spawning period and the last DOY with one or

three consecutive days with temperatures below zero. Both correlations were significant and highlight the importance of not having frost days in the onset of reproduction.

Factorial analysis confirmed that the variables best associated with the DOY were the number of days with temperatures below zero degrees in the previous weeks. In the first factor (Table 2 and Supplementary Materials Figure 4), the variables related to the number of days those temperatures remained below zero and temperatures were more relevant, while in the second factor, we can find the variables related to temperature and frost days in the week previous to the onset of spawning. The first two factors represented 83.70% of the variance (Table 2), so these variables can be considered as indicative of the start of reproduction.

Future distribution range contraction

Habitat availability for *Rana temporaria* in North-eastern Iberia, after excluding the collinear variables, was determined by five variables which had parametric correlations between 0.695 and 0.029. Specifically, the occurrence of the species was strongly restricted to temperate conditions during the warmest month, moderate precipitation seasonality and temperatures during the driest quarter. The average test AUC for the fifty replicates was 0.896 ± 0.009 (mean $\pm$ standard deviation) indicating the model was a good fit. The predicted range of common frogs in Northeastern Iberian Peninsula was influenced by the relief as revealed by high probabilities of occurrence in the mountain areas and low probabilities at lower altitudes (Figure 5). This led to the isolation of the populations located in the southern edge of the range, especially between the Montseny and Guillerics, and between the latter and the rest of the Catalan Transversal Range. We predicted the presence of the species in the montane ranges of the Montseny except for the massif's highest peaks. After controlling for redundancy, representing habitat suitability, the analysis retained four-six variables (parametric correlations lower than 0.7). Models had AUC tests comprised between 0.867 and 0.887, indicating good performance. The Jackknife test revealed that mean temperatures in the wettest and driest quarter, maximum temperatures in the warmest month, precipitation in the warmest quarter and precipitation seasonality were the most influential variables, followed by precipitation and temperature seasonality and isothermality. Models based on scenarios ssp128 and ssp 585 (Figure 5) considerably reduced the probability of species occurrence when comparing the predictions obtained between 2040 and 2100. Our results highlighted the isolation between the populations in the southernmost edge, and between them and the Pyrenean ones. In the Montseny massif, the future predicted ranges in 2040 were similar to the ones currently occupied by the species, but occurrence probability drops and rejected an altitudinal shift towards the highest top mountains. The worst scenario, high level of emissions in 2100, predicts species occurrence

with low probability and only in the eastern side of the massif, suggesting the failure of its persistence in Montseny. Thus, in all our predicted scenarios, low and high emissions entailed a dramatic reduction in habitat availability in the Montseny massif within this century (Supplementary Material, Figure 5).

Discussion

Onset breeding period

The Common Frog in the Montseny Massif had advanced its spawning onset following the increase in temperatures due to global warming. Significantly, the four earliest dates of onset spawning periods were in 2019, 2020, 2021 and 2022 (Figure 2). Between 2009 and 2022 the average of advancement of the onset of reproduction is 26 days/decade (Figure 2). Our estimated advancement of the onset of the breeding season in Montseny Massif is high, but included within the range found by While & Uller (2014) for *Rana temporaria* (-28,1 days/decade).

Probably, the heavy Mediterranean climatic impact on the Montseny Massif. The southern distribution limit of many eurosiberian species, associated with the rapid increase in temperatures during recent years (Minuartia, 2016), could be the causes behind this great advancement. Supporting this data, (Minuartia, 2016) estimates an increase in the average annual temperature of 0.7 °C for the decade (2012-2021) in the Coastal / Pre-coastal Region (Montnegre-Corredor Natural Park), to which the Montseny Massif is close.

This earliest onset reproduction date in amphibians has been pointed out by different authors. Beebee detected, in 1995, an advancement in the beginning of reproduction of some species, which was linked to global warming. Multispecies meta-analyses have suggested that amphibians are the taxa showing the strongest phenological advancement in response to global warming, with an average breeding advancement ranging from 2.6 days per decade (Parmesan 2007) to 6.1 days per decade (While and Uller 2014). The mean advancement in *Rana temporaria*'s onset of spawning found by While & Uller (2014) was -5.95 day/decade (Stand. Dev.=11.92; Maximal advancement; -28,1 day/decade). While &Uller (2014) have clearly shown that amphibian populations consistently anticipate their breeding periods, determined by trends of global warming, particularly at high latitudes.

Our results are in accordance with Ficetola & Maiorano (2016) who conclude that temperature increase was the major factor behind phenological advancement, while the impact of precipitation on phenology was weak. In fact, we didn't obtain any significant relation between rain and DOY of onset breeding (Supplementary Material table1 and Figures 2 and 3).

For the end of the last century, other authors (Peñuelas & Boada, 2003; Minuartia, 2016), estimated an increase in air temperatures ranging from 0.24 to 0.3°C per decade in the Montseny massif. However, the data obtained in this study from 1996 to 2021 in the same weather station indicate that this increase has accelerated to 0.404 °C/decade on average of maximal temperature (Supplementary Material Figure 1), according to the observations by Peñuelas & Boada, (2003) who consider that the increase in temperatures has accelerated over recent decades.

This advancement demonstrates a great plasticity in spawning dates in relation to climate change. However, Phillimore et al. (2010) in Britain observed that although all populations exhibit a plastic response to temperature, spawning earlier in warmer years, differences within populations in their first spawning dates are dominated by local adaptation. Climate change projections for Britain in 2050-2070, will require an advancement of between 21 and 39 days for the first spawning date, but Phillimore et al. (2010) indicate that plasticity alone will only enable an advancement of 5 to 9 days, and need a microevolutionary and gene flow challenge for an earlier first spawning date of between 16-30 days over the next 50 years. However, our data demonstrates that from 2009 to 2022, the advance of the onset of spawning has been 26 days, more than the data shown by the other authors. Perhaps living and surviving in the edge has been favourable for more plastic individuals due to the suboptimal and higher variability of environmental conditions. According to this hypothesis, Lesbarrères et al. (2007) found that metamorphic size was positively correlated with individual heterozygosity and fitness in *Rana temporaria*'s Nordic distribution limit. These results suggest that genetic variability may be an important component for individual fitness in the Common Frog (Lesbarrères et al. 2007). Ruthsatz, et al. (2022) found that *R. temporaria* has a high acclimation capacity in the temperate populations, which could reflect an adaptation to increased thermal variability to enable survival in heterogeneous or suboptimal environments, especially during the larval stage. Generally, wide-ranging species, such as *R. temporaria*, exhibit a greater capacity for thermal acclimation of upper thermal tolerance than narrowing-ranging species, suggesting that selection for acclimation ability may have been a key factor that enabled geographic expansion into areas with greater thermal variability (Markle & Kozak, 2018). These authors demonstrate that young larvae of *R. temporaria* are the most temperature-sensitive life stage and highlights that early life stages might define the climate change vulnerability of amphibian populations. Therefore, this widely distributed species possesses a high capacity for developmental, morphological, and physiological plasticity. In this sense, in the Montseny Massif, *R. temporaria* has chosen, throughout the years, the week with positive minimal temperature to onset spawn (Average

minimal T^a for the previous week of onset spawning = 2,09°C; STD=0.88°C; Max=4,07; Min=0,9), with a great variability in DOY between years (Figure 2) but low variability in spawning environmental conditions.

Enriquez-Urzelai (2018) indicates that *Rana temporaria* tadpoles are an eurythermal organism and thermal niches vary across latitudinal and altitudinal clines. Furthermore, at both extremes of the latitudinal gradient, plasticity is maximal. Enriquez-Urzelai et al. (2020a) indicate that mixed signals of niche evolution and conservation at the species level and generalized conservation in thermal niches at the population level indicate that the pace of niche evolution in *R. temporaria* might be too slow to compensate climate change (mainly among terrestrial life cycle stages) according to Phillimore et al. (2010).

Previous studies identified a strong relationship between snowmelt dates and spawning dates (Corn and Muths, 2002; Corn, 2003; Bison et al, 2021), thereby confirming a high degree of plasticity in terms of breeding timing (Muir et al., 2014). Bison et al, (2021) observe in *Rana temporaria* a relationship with the DOY of snowmelt and DOY of middle spawning activity. In our case, the location of the Montseny Massif in the southern limit of distribution where the climatic conditions are Mediterranean, makes it impossible to determine the snowmelt day because the snow is not present every year and the pluviometry in the first month of the year is very scarce. Consequently, to compare the DOY of the onset of snowmelt with the theoretical snowmelt, we used the DOY of the last one or three days when there were temperatures below zero. Our results confirm one highly and significant relation between the DOY of spawning and DOY of the last frost days (Figure 4) according to the results obtained by Bison, et al. (2021).

Enriquez-Urzelai et al. (2020a) indicate that high-elevation populations had slightly wider tolerance ranges driven by increases in heat tolerance but a lower potential for acclimation. Plausibly, wider thermal fluctuations at high elevations could be extrapolated to distribution limits like Montseny populations. Then, extreme events could have more unpredictable effects on populations living in the edge of their distribution, as in the case of Montseny populations. Increases in air temperatures and more frequent extreme events (e.g., heat waves) due to climate warming will produce changes in pond temperatures (Enriquez-Urzelai, 2019b). Species could respond in situ by increasing upper thermal limits through genetic adaptation or phenotypic plasticity. However, the potential for adaptation and plastic responses to shift thermal tolerances seems limited in ectotherms, relative to predicted environmental changes (Enriquez-Urzelai et al., 2018, 2019b). Thus, species or populations living closer to their maximum thermal tolerance (e.g., southernmost populations of *R. temporaria*) might be especially vulnerable to climate change (Duarte et al., 2012; Gerick et al., 2014).

Furthermore, although the *R. temporaria* adults show great adaptability or plasticity in terms of the onset of the spawning period, the vulnerability of the larvae and adults when faced with the increase in extreme phenomena in a scenario of climate change could be high. Montori et al. (2011) observe that mortality of amphibians increases due to more frequent sudden drops in temperature, which can affect both larvae and adults (Supplementary material. Figure 6). If the advancement of reproduction continues in the coming years, the probability of post-spawning frost days will increase, given the variability of the climate in the context of an increase in extreme phenomena. Reinforcing this hypothesis, Bison et al. (2021) observed that the number of frost days during egg-development increased more quickly at high elevation. Then, early spawn could lead to an additional rise in larval or spawn mortality due to the greater number of frost days (Montori et al. 2011). Enriquez-Urzelai (2019b) predicted extensive decreases in climatic suitability in Southern Europe, which harbors a significant fraction of the species' genetic diversity. Moreover, these authors suggest that *R. temporaria*'s climatic suitability decreases at locations where water is scarce, or ponds reach temperatures of $>40^{\circ}\text{C}$. This temperature is close to the critical thermal maxima of *R. temporaria* tadpoles (Enriquez-Urzelai et al., 2019b). In the shallow ponds in which *R. temporaria* commonly breeds, thermal conditions depend on the degree of insolation and depth. Tadpoles suffer under great thermal fluctuations in ponds, and thus extreme pond temperatures may set the limits of the distribution of pond-breeding amphibians. The reproduction ponds studied by Campeny (2001) in Santa Fe del Montseny show a maximum average annual temperature between 20.83 and 26.00°C , with maximums in summer of over 32°C . In addition, these ponds are very shallow and the water has average depths oscillating from 8.19cm with SD of 4.54cm , to 15.32cm with SD of 8.19cm . In these abiotic conditions, the rising temperatures could increase the risk of larval mortality in the future.

Future distribution range contraction

The Montseny massif and the neighbouring Montnegre massif is the southernmost edge in East Iberia of ectothermic vertebrates adapted to temperate conditions (i.e. *Lacerta bilineata*, *Podarcis muralis*, *Vipera aspis* and *Zamenis longissimus*) that will likely be affected by the rise in temperatures and dryness. In this context the southwesternmost populations of Common Frog are an even more interesting species model because of their cold adaptation capabilities.

Although our results provided evidence of range contraction in the Montseny (Figure 5 and Supplementary Material, Figure 5), they failed to confidently predict the local extinction of the species. Amphibians are mostly low vagile and small vertebrates which rely on microclimatic conditions (Smith & Green, 2005; Hoffman et al., 2021), far from the scope of the climatic niche

analysis. Range contraction is frequently predicted by climatic niche modelling in analysis of the consequences of global warming in cold-adapted species (Pauli et al., 2012; Ernakovich et al., 2014). In the Montseny massif, most of the reproductive ponds are placed in the east side, more temperate and humid, whereas the species experiences the harshest conditions in the southwestern slopes exposed to more Mediterranean climate. In this area, only four reproductive sites are known and the estimated number of reproductive females is dramatically small and undoubtedly, they are the most endangered and could disappear during the coming decades.

Common frogs exhibit a large ecological plasticity, living in marshes at the sea level and high altitude lakes in the Pyrenees (Kuzmin et al., 2009). In the Montseny massif they are associated with temporal pools, in most cases in open areas within the margins of the riparian forest (F. Amat, unpublished results). The rise in temperatures increases the risk of pond desiccation leading to the failure of the reproduction because most of them depend on irrigation from neighbouring streams to maintain a long hydroperiod. Early reproduction, as our results indicated, may compensate for the shortening of the pond's hydroperiod. However, even then, it cannot outweigh extreme spring dryness that predictably could become more frequent over time in the Mediterranean region (Quintana-Seguí et al., 2016). This phenomenon could also play a negative role in the preservation of the riparian forest that provides humid refugia for the activity of adults after reproduction.

Air temperatures fluctuate more than water temperatures (Feder and Hofmann 1999). Thus, terrestrial stages are more likely to encounter more extreme temperatures than aquatic larvae. In contrast, they can use behavioural thermoregulation. This may not only allow individuals to escape unwanted temperatures (Kearney & Porter, 2009), but may also weaken directional selection based on thermal traits, through a process known as the 'Bogert effect' (Huey et al. 2003; Buckley et al. 2015). Oppositely, in water, the potential of behavioural thermoregulation to buffer heat or cold waves may be more limited.

In amphibians, metamorphic and juvenile stages can thermoregulate better than larval stages. Although most forecasts with regard to the consequences of global warming for biodiversity conservation ignore how thermal tolerance varies between life stages, the viability of the weakest link could restrict the future distribution of a species (Briscoe et al. 2012; Pincebourde and Casas 2015). Enriquez-Urzelai et al. (2020b) demonstrate that larval acclimation to high temperatures increases larval heat tolerance, but may reduce cold tolerance in immediately subsequent stages. If sudden drops in temperature are more frequent due to an

advance in the reproduction date, this can endanger the survival of some tadpole pond populations.

It is likely that wider thermal fluctuations at high elevations favour more tolerant but less plastic phenotypes, thus reducing the risk of encountering stressful temperatures during unpredictable extreme events (Enriquez-Urzelai et al. 2020b). This wider thermal fluctuation also occurs in the Montseny Massif population due its location in a Mediterranean climate, but we can't conclude if the Montseny population presents more adaptability or plasticity, and as other authors suggest, if this adaptability has a genetic basis (Phillimore et al. 2010). Future studies should address this problem in populations that survive in the southernmost limit of its wide distribution.

Acknowledgements

We are thankful to the Natural Park and Biosphere Reserve of Montseny for its financial support and logistic collaboration with the study of *Rana temporaria*. We want to thank all those people who provided us with data on the phenology of *Rana temporaria* and especially Joan Manel Roig Fernández for providing data on species phenology prior to the implementation of the current project, to Marc Franch and Adrià Jordà for assisting in some field reserach in Santa Fe and to Martí Boada, a pioneer in the annotation of phenological data from Montseny. This field data book undoubtedly has an invaluable historic utility. We also want to thank the use of data from the databases of the Institut Català d'Ornitologia (ICO: <https://ornitho.cat>) and the Asociación Herpetológica Española (AHE; <https://siare.herpetologica.es>) and Societat Catalana d'Herpetologia (CH: <https://soccatherp.org/>).

References

- Agudelo, WJ, Urbina-Cardona, N, Armenteras-Pascual, D. 2019. Critical shifts on spatial traits and the risk of extinction of Andean anurans: an assessment of the combined effects of climate and land-use change in Colombia. *Perspectives in Ecology and Conservation* 17(4):206-219. <https://doi.org/10.1016/j.pecon.2019.11.002>
- Amat, F, Montori, A. 2022. El tritó del Montseny i els Amfibis I Rèptils del Parc Natural i Reserva de la Biosfera del Montseny. Museu de Granollers Ed, (Barcelona). (in press).
- Balcells, E. 1957. Elementos nórdicos en el poblamiento de la cumbre del Montseny. *P. Inst. Biol. Apli.* 26:123-126.

475 Bea, A, Rodríguez-Teijeiro, D, Jover, L. 1986. Relations between meteorological variables and
476 the initiation of spawning period in populations of *Rana temporaria* L. in the Atlantic region of the
477 Basque Country(Northern Spain). *Amphibia-Reptilia* 7:23-31

478 Beebee, TJC. 1995. Amphibian breeding and climate. *Nature* 374(6519), 219-220
479 <https://doi.org/10.1038/374219a0>

480 Beebee, TJC. 2002. Amphibian phenology and climate change. *Conservation Biology*
481 16(6):1454-1455. DOI:10.1046/j.1523-1739.2002.02102.x

482 Bison M, Yoccoz NG, Carlson BZ, Klein G, Laigle I, Van Reeth C, Delestrade A. 2021. Earlier
483 Snowmelt Advances Breeding Phenology of the Common Frog (*Rana temporaria*) but Increases
484 the Risk of Frost Exposure and Wetland Drying. *Front. Ecol. Evol.* 9:645585.
485 <https://doi.org/10.3389/fevo.2021.645585>

486 Blaustein, AR, Walls, SC, Bancroft, BA, Lawler, JJ, Searle, CL, Gervasi, SS. 2010. Direct and
487 Indirect Effects of Climate Change on Amphibian populations. *Diversity* 2:281-313.
488 <https://doi.org/10.3390/d2020281>

489 Briscoe, NJ, Porter, WP, Sunnucks, P, Kearney, MR. 2012. Stage-dependent physiological
490 responses in a butterfly cause nonadditive effects on phenology. *Oikos*, 121:1464-1472
491 <https://doi.org/10.1111/j.1600-0706.2011.20049.x>

492 Buckley, LB, Ehrenberger, JC, & Angilletta, MJ Jr. 2015. Thermoregulatory behaviour limits
493 local adaptation of thermal niches and confers sensitivity to climate change. In R. Wilson (Ed.),
494 *Functional Ecology* 29:1038-1047 <https://doi.org/10.1111/1365-2435.12406>

495 Campeny, R. 2001. Ecologia de les larves d'amfibis anurs al Montseny. D. Phil. Thesis,
496 Universitat de Barcelona. 450pp.

497 Chadwick, EA, Slater, FM, Ormerod, SJ, 2006. Inter- and intraspecific differences in climatically
498 mediated phenological change in coexisting *Triturus* species. *Global Change Biology* 12:1069-
499 1078 <https://doi.org/10.1111/j.1365-2486.2006.01156.x>

500 Corn, PS. 2003. Amphibian breeding and climate change: importance of snow in the mountains.
501 *Conservation Biology* 17:622-625
502 <https://doi.org/10.1046/j.1523-1739.2003.02111.x>

503 Corn, P S, Muths, E 2002. Variable breeding phenology affects the exposure of amphibian
504 embryos to ultraviolet radiation. *Ecology* 83:2958-2963 [https://doi.org/10.1890/0012-9658\(2002\)083\[2958:VBPATE\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[2958:VBPATE]2.0.CO;2)

505
506 Duan, RY, Kong, XQ, Huang, MY, Varela, S, Ji, X. 2016. The potential effects of climate change
507 on amphibian distribution, range fragmentation and turnover in China. *PeerJ* 4:e2185; DOI
508 10.7717/peerj.2185.

Duarte H, Tejedo M, Katzenberger M et al. 2012. Can amphibians take the heat? Vulnerability to climate warming in subtropical and temperate larval amphibian communities. *Glob Change Biol.* 18:412-421 <https://doi.org/10.1111/j.1365-2486.2011.02518.x>

Enriquez-Urzelai, U, Palacio, AS, Merino NM, Sacco, M, Nicieza, AG. 2018. Hindered and constrained: limited potential for thermal adaptation in post-metamorphic and adult *Rana temporaria* along elevational gradients, *Journal of Evolutionary Biology* Dryad, Dataset, <https://doi.org/10.5061/dryad.2fp14fg>

Enriquez-Urzelai, U, Bernardo, N, Moreno-Rueda, G, Montori, A, Llorente, G. 2019a. Are amphibians tracking their climatic niches in response to climate warming? A test with Iberian amphibians. *Climatic Change* 154:289-301. <https://doi.org/10.1007/s10584-019-02422-9>

Enriquez-Urzelai U, Kearney MR, Nicieza AG, Tingley R. 2019b. Integrating mechanistic and correlative niche models to unravel range-limiting processes in a temperate amphibian. *Glob. Change Biol.* 2019;00:1-15. <https://doi.org/10.1111/gcb.14673>

Enriquez-Urzelai, U, Tingley, R, Kearney, M R, Sacco, M, Palacio, A S, Tejedo, M, Nicieza, A G. 2020a. The roles of acclimation and behaviour in buffering climate change impacts along elevational gradients. *J Anim Ecol.* 2020;00:1-13. <https://doi.org/10.1111/1365-2656.13222>

Enriquez-Urzelai, U, Tingley, R, Kearney, M, Sacco, M, Palacio, A S, Tejedo, M, et al. 2020b. The roles of acclimation and behaviour in buffering climate change impacts along elevation gradients. *J. Anim. Ecol.* <https://doi.org/10.1111/1365-2656.13222>

Ernakovich, JG, Hopping, KA, Berdanier, AB, Simpson, RT, Kachergis, EJ, Steltzer, H, Wallenstein, MD 2014. Predicted responses of arctic and alpine ecosystems to altered seasonality under climate change. *Global Change Biology* 20:3256-3269 <https://doi.org/10.1111/gcb.12568>

Feder, M E, Hofmann, G E. 1999. Heat-shock proteins, molecular chaperones, and the stress response: Evolutionary and ecological physiology. *Annual Review of Physiology* 61:243-282. <https://doi.org/10.1146/annurev.physiol.61.1.243>

Ficetola, G F, & Maiorano, L. (2016). Contrasting effects of temperature and precipitation change on amphibian phenology, abundance and performance. *Oecologia* 181(3):683-693 <https://doi.org/10.1007/s00442-016-3610-9>

García, A, Ortega-Huerta, MA. 2013. Potential distributional change and conservation priorities of endemic amphibians in western Mexico as a result of climate change. *Environmental Conservation* <https://doi.org/10.1017/S0376892913000349>

Gerick AA, Munshaw RG, Palen WJ, Gerick, A. A., Munshaw, R. G., Palen, W. J, Combes, S. A., O'Regan, S. M. 2014. Thermal physiology and species distribution models reveal climate

543 vulnerability of temperate amphibians. *Journal of Biogeography* 41(4):713-723.
544 <https://doi.org/10.1111/jbi.12261>

545 González-Hidalgo, J C; López-Bustins, J A; Štěpánek, P; Martín-Vide, J, de Luisa, M. 2009.
546 Monthly precipitation trends on the Mediterranean fringe of the Iberian Peninsula during the
547 second half of the twentieth century (1951-2000). *International Journal of Climatology*
548 29(10):1415-1429. <https://doi.org/10.1002/joc.1780>

549 Green, D. M. 2017. Amphibian breeding phenology trends under climate change: predicting the
550 past to forecast the future. *Global Change Biol* 23:646-656 <https://doi.org/10.1111/gcb.13390>

551 Greenberg, DA, Palen, WJ. 2021. Hydrothermal physiology and climate vulnerability in
552 amphibians. *Proceedings of Royal Society B* 288:20202273
553 <https://doi.org/10.1098/rspb.2020.2273>

554 Habibzadeh, N, Ghoddousi, A, Bleyhl, B, Kuemmerle, T. 2021. Rear-edge populations are
555 important for understanding climate change risk and adaptation potential of threatened species.
556 *Conservation Science and Practice* 2021;3:e375 <https://doi.org/10.1111/csp2.375>

557 Hampe, A, Petit, R. J. 2005. Conserving biodiversity under climate change: the rear edge
558 matters. *Ecology Letters* 8:461-467. DOI: 10.1111/j.1461-0248.2005.00739.x

559 Hijmans, RJ, Cameron, SE, Parra, JL, Jones, PG, Jarvis, A. 2005. Very high resolution
560 interpolated climate surfaces for global land areas. *International Journal of Climatology* 25:1965-
561 1978. <https://doi.org/10.1002/joc.1276>

562 Hoffmann, EP, Cavanough, KL, Mitchell, NJ. 2021. Low desiccation and thermal tolerance
563 constrains a terrestrial amphibian to a rare and disappearing microclimate niche. *Conservation*
564 *Physiology* 9(1):coab027. <https://doi.org/10.1093/conphys/coab027>

565 Huey, R B, Deutsch, C A, Tewksbury, J J, Vitt, L J, Hertz, P E, Alvarez Pérez, H J, Garland, T.
566 (2009). Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal*
567 *Society B: Biological Sciences* 276, 1939-1948. <https://doi.org/10.1098/rspb.2008.1957>

568 IPCC. 2021. IPCC report: 'Code red' for human driven global heating, warns UN chief. IPCC
569 report. <https://news.un.org/en/story/2021/08/1097362>

570 Kearney, M R, & Porter, W P. 2009. Mechanistic niche modelling: Combining physiological and
571 spatial data to predict species' ranges. *Ecology Letters* 12:334-350.
572 <https://doi.org/10.1111/j.14610248.2008.01277.x>

573 Kim, HW, Adhikari, P, Chang, MH, Seo, C. 2021. Potential distribution of amphibians with
574 different habitat characteristics in response to climate change in South Korea. *Animals*
575 11(8):2185 <https://doi.org/10.3390/ani11082185>

576 Klaus, SP, Loughheed, SC. 2013. Changes in breeding phenology of eastern Ontario frogs over
577 four decades. *Ecology and Evolution* 3(4): 835-845 <https://doi.org/10.1002/ece3.501>

578 Kusano, T, Inoue, M. 2008. Long-term trends toward earlier breeding of Japanese amphibians.
579 *Journal of Herpetology* 42:608-614. <https://doi.org/10.1670/08-002R1.1>

580 Kuzmin, S, Ishchenko, V, Tuniyev, B, Beebee, T, Andreone, F, Nyström, P, Anthony, B P,
581 Schmidt, B, Ogradowczyk, A, Ogielska, M, Bosch, J, Miaud, C, Loman, J, Cogalniceanu, D,
582 Kovács, T, Kiss, I. 2009. *Rana temporaria* (errata version published in 2016). *The IUCN Red*
583 *List of Threatened Species* 2009: e.T58734A86470817.

584 Lesbarrères, D, Schmeller, D S, Primmer, C R, & Merilä, J. 2007. Genetic variability predicts
585 common frog (*Rana temporaria*) size at metamorphosis in the wild. *Heredity* 99(1):41-46.
586 doi:10.1038/sj.hdy.6800961 <https://doi.org/10.1038/sj.hdy.6800961>

587 Loman, J. 2016. Breeding phenology in *Rana temporaria*. Local variation is due to pond
588 temperature and population size. *Ecology and Evolution* 6(17):6202-6209
589 <https://doi.org/10.1002/ece3.2356>

590 Markle, T M, Kozak, K. H. 2018. Low acclimation capacity of narrow ranging thermal specialists
591 exposes susceptibility to global climate change. *Ecology and Evolution* 8(9), 4644-4656.
592 <https://doi.org/10.1002/ece3.4006>

593 McCaffery, R M, Maxell, B A. 2010. Decreased winter severity increases viability of a montane
594 frog population. *Proceedings of the National Academy of Sciences USA* 107(19):8644-8649
595 <https://doi.org/10.1073/pnas.0912945107>

596 Minuartia, Estudis Ambientals. 2016. Estudi dels efectes del canvi climàtic en el Montseny:
597 diagnosi, impactes i vulnerabilitats. LIFE Clinomics. Acció A1. Unpublished Report, 146pp.
598 Available at [Clinomics_Canvi climatic Montseny A1_Memoria_Annexos \(lifeclinomics.eu\)](http://Clinomics_Canvi climatic Montseny A1_Memoria_Annexos (lifeclinomics.eu))

599 Montori, A, Giner, G, Béjar, X, Álvarez, D. 2011. Descenso brusco de temperaturas y nevadas
600 tardías como causas de mortalidad de anfibios durante el período reproductor. *Boletín de la*
601 *Asociación Herpetológica Española*. 22:72-74.

602 Montori, A, Pascual, X. 1987. Contribución al estudio de *Rana temporaria* L. (Amphibia,
603 Ranidae) en Santa Fe del Montseny (Barcelona). 2. Ciclo biológico. *Miscel-lania Zoologica*
604 11:299-307

605 Muir, A P, Biek, R, Mable, B K. 2014. Behavioural and physiological adaptations to low-
606 temperature environments in the common frog, *Rana temporaria*. *BMC Evol. Biol.* 14:110.
607 DOI:10.1186/1471-2148-14-110

608 Panareda-Clopés, J. 1979. Aportació a l'estudi del clima del Montseny. *Acta Geològica*
609 *Hispanica* 14, 524-528. Accesible at <https://www.recercat.cat/handle/2072/265106>

610 Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annual*
611 *Review in Ecology, Evolution and Systematics* 37:637-669.
612 <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>

613 Parmesan, C. 2007. Influences of species, latitudes and methodologies on estimates of
614 phenological response to global warming. *Global Change Biol* 13:1860-1872.
615 <https://doi.org/10.1111/j.1365-2486.2007.01404.x>

616 Pascual, X, Montori, A. 1982. Características del ciclo biológico de *Rana temporaria* L.
617 (Amphibia, Anura) en Santa Fe de Montseny (Barcelona). *P. Cent. Pir. Biol. Exp.* 13:51-54.

618 Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M, Benito Alonso JL, Coldea G, Dick
619 J, Erschbamer B, Fernández Calzado R, Ghosn D, Holten JI, Kanka R, Kazakis G, Kollár J,
620 Larsson P, Moiseev P, Moiseev D, Molau U, Molero Mesa J, Nagy L, Pelino G, Puşcaş M, Rossi
621 G, Stanisci A, Syverhuset AO, Theurillat JP, Tomaselli M, Unterluggauer P, Villar L, Vittoz P,
622 Grabherr G. 2012. Recent plant diversity changes on Europe's mountain summits. *Science*.
623 2012 Apr 20;336(6079):353-5. DOI:10.1126/science.1219033.PMID: 22517860

624 Pearce, J, Ferrier, S. 2000. Evaluating the predictive performance of habitat models developed
625 using logistic regression. *Ecological Modelling* 133, 225- 245 [https://doi.org/10.1016/S0304-](https://doi.org/10.1016/S0304-3800(00)00322-7)
626 [3800\(00\)00322-7](https://doi.org/10.1016/S0304-3800(00)00322-7)

627 Peñuelas, J, Boada, M. 2003. A global change-induced biome shift in the Montseny mountains
628 (NE Spain). *Global Change Biology* 9(2):31-140 [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2486.2003.00566.x)
629 [2486.2003.00566.x](https://doi.org/10.1046/j.1365-2486.2003.00566.x)

630 Peñuelas, J, Germain, J, Álvarez, E, Aparicio, E, Arús, P, Basnou, C, Blanché, C, Bonada, N,
631 Canals, P, Capodiferro, M, et al. 2021. Impacts of use and abuse of nature in Catalonia with
632 proposals for sustainable management. *Land* 10:144. <https://doi.org/10.3390/land10020144>

633 Phillimore, AB, Hadfield, JD, Jones, OR, Smithers, RJ. 2010. Differences in spawning date
634 between populations of Common Frog reveal local adaptation. - *Proc. Natl Acad. Sci. USA* 107:
635 8292-8297 <https://doi.org/10.1073/pnas.0913792107>

636 Phillips, SJ Dudík, M. 2008. Modeling of species distributions with Maxent: new extensions and
637 a comprehensive evaluation. *Ecography* 31:161-175 [https://doi.org/10.1111/j.0906-](https://doi.org/10.1111/j.0906-7590.2008.5203.x)
638 [7590.2008.5203.x](https://doi.org/10.1111/j.0906-7590.2008.5203.x)

639 Phillips, SJ, Anderson, RP, Schapire, RE. 2006. Maximum entropy modeling of species
640 geographic distributions. *Ecological Modelling* 190(3/4):231-259
641 <https://doi.org/10.1016/j.ecolmodel.2005.03.026>

642 Pincebourde, S, Casas, J. 2015. Warming tolerance across insect ontogeny: Influence of joint
643 shifts in microclimates and thermal limits. *Ecology* 96:986-997. [https://doi.org/10.1890/14-](https://doi.org/10.1890/14-0744.1)
644 [0744.1](https://doi.org/10.1890/14-0744.1)

645 Primack, RB, Ibáñez, I, Higuchi H, Lee, SD, Miller-Rushing, AJ, Wilson, AM, Silander, JA. 2009.
646 Spatial and interspecific variability in phenological responses to warming temperatures.
647 *Biological Conservation* 142:2569-2577 <https://doi.org/10.1016/j.biocon.2009.06.003>

648 Roig, JM, Amat, F. 2002. Fenologia reproductora de la granota roja (*Rana temporaria*) al Parc
649 Natural del Montseny. Unpublished report, Diputació de Barcelona, 14pp.

650 Quintana-Seguí, P, Martin, E, Sánchez, E, Zribi, M, Vennetier, M, Vicente-Serrano, S, Vidal, J.
651 2016. Drought: observed trends, future projections, 123-131. In: Thiebault, S, Moatti, JP eds.
652 The Meditteranena Region under climate change. A scientific update. IRD Éditions, Marseille.
653 123-131. Available at <https://memsic.ccsd.cnrs.fr/GIP-BE/hal-01401386v1>

654 Reading, CJ. 2007. Linking global warming to amphibian declines through its effects on female
655 body condition and survivorship. *Oecologia* 151:125-131 [https://doi.org/10.1007/s00442-006-](https://doi.org/10.1007/s00442-006-0558-1)
656 [0558-1](https://doi.org/10.1007/s00442-006-0558-1)

657 Ruthsatz, K, Dausmann, K H, Peck, M A, Glos, J. 2022. Thermal tolerance and acclimation
658 capacity in the European common frog (*Rana temporaria*) change throughout ontogeny. *Journal*
659 *of Experimental Zoology Part A: Ecological and Integrative Physiology* 337(5):1-14.
660 <https://doi.org/10.1002/jez.2582>

661 Scott, AW, Pithart, D, Adamson JK. 2008. Long-term United Kingdom trends in the breeding
662 phenology of the common frog, *Rana temporaria*. *Journal of Herpetology* 42(1):89-96
663 <https://doi.org/10.1670/07-022.1>

664 Seneviratne, S I, Donat, M G, Mueller, B, Alexander, LV. 2014. No pause in the increase of hot
665 temperature extremes. *Nature Climate Change* 4(3):161-163.
666 <https://doi.org/10.1038/nclimate2145>

667 Sheridan, JA, Caruso, NM, Apodaca, JJ, Rissler, LJ. 2018. Shifts in frog size and phenology:
668 Testing predictions of climate change on a widespread anuran using data from prior to rapid
669 climate warming. *Ecology and Evolution* 8:1316-1327. <https://doi.org/10.1002/ece3.3636>

670 Smith, MA, & Green, DM. 2005. Dispersal and the metapopulation paradigm in amphibian
671 ecology and conservation: are all amphibian populations metapopulations? *Ecography*
672 28(1):110128 <https://doi.org/10.1111/j.0906-7590.2005.04042.x>

673 Terhivuo, J. 1988. Phenology of spawning for the Common Frog (*Rana temporaria* L.) in Finland
674 from 1846 to 1986. *Annales Zoologici Fennici* 25, 165-175. <http://www.jstor.org/stable/23734521>

675 Thomas, CD, Cameron, A, Green, RE, Bakkenes, M, Beaumont, LJ, Collingham, YC, Erasmus,
 676 BFN, Ferreira de Siqueira, M, Grainger, AF, Hannah, L, Hughes, L, Huntley, B, van Jaarsveld,
 677 AS, Midgley, GF, Miles, L, Ortega-Huerta, MA Peterson, AT, Phillips, OL, Williams, SE. 2004.
 678 Extinction risk from climate change. *Nature* 427:145-148 <https://doi.org/10.1038/nature02121>
 679 Thuiller W, Lavergne, S, Roquet, C, Boulangeat, I, Lafourcade, B, Araujo, MA. 2011.
 680 Consequences of climate change on the tree of life in Europe. *Nature* 470:531-534
 681 <https://doi.org/10.1038/nature09705>
 682 Timm, B C, McGarigal, K, Compton, B W. (2007). Timing of large movement events of pond-
 683 breeding amphibians in western Massachusetts, USA. *Biol. Conserv* 136:442-454.
 684 <https://doi.org/10.1016/j.biocon.2006.12.015>
 685 Todd, BD, Scott, DE, Pechmann, JHK, Gibbons, JW. 2010. Climate change correlates with
 686 rapid delays and advancements in reproductive timing in an amphibian community. *Proceedings*
 687 *of the Royal Society of London B* 278:2191-2197 DOI: 10.1098/rspb.2010.1768
 688 Tryjanowski, P, Mariusz, R, Sparks, T. 2003. Changes in spawning dates of Common Frogs and
 689 Common Toads in western Poland in 1978-2002. *Annales Zoologici Fennici* 40:459-464.
 690 <https://www.jstor.org/stable/23735858>
 691 Vicente-Serrano, SM, Lopez-Moreno, JI, Beguería, S, Lorenzo-Lacruz, J, Sanchez-Lorenzo, A,
 692 García-Ruiz, JM, Azorin-Molina, C, Morán-Tejeda, E, Revuelto, J, Trigo, R, Coelho, F, Espejo,
 693 F. 2014. Evidence of increasing drought severity caused by temperature rise in southern
 694 Europe. *Environmental Research Letters* 9(4):9:044001. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/9/4/044001)
 695 [9326/9/4/044001](https://doi.org/10.1088/1748-9326/9/4/044001)
 696 Walther GR, Post, E, Convey, P, Menzel, A, Parmesan, C, Beebee, TJC, Fromentin, JM, Hoegh-
 697 Gulberg, O, Bairlein, F. 2002. Ecological responses to recent climate change. *Nature* 416:389-
 698 395 <https://doi.org/10.1038/416389a>
 699 While, G M, & Uller, T. 2014. Quo vadis amphibia? Global warming and breeding phenology in
 700 frogs, toads and salamanders. *Ecography* 37(10):921-929. <https://doi.org/10.1111/ecog.00521>

701

Figure 1

Geographic range of *Rana temporaria* in the Palearctic and in northeaster Iberia, and location of the Montseny Massif (black circle and red square).

A: Map of global distribution of Common Frog. B: Map of the Montseny massif. Green line: Administrative limit of Natural Park of Montseny. Red Line: Administrative limit of Natural Park and Biosphere Reserve of Montseny.




Figure 2

Linear regression between the day of onset reproduction from January, 1 (DOY) of *Rana temporaria* in the study site and the year, for the period 2009-2022.




Figure 3

Evolution of minimal temperature four weeks before the onset of breeding period (top) and the interannual variance (bottom) in the four weeks previous to onset of breeding period.




Figure 4

Relationship between the last day of year with air temperature below 0°C (FDOY1) previous to onset spawn and the DOY of onset egg-spawning of the Common Frog *Rana temporaria* (Left), and between the last day of year with three consecutive days with a




Figure 5

Habitat availability and categorical predictability for *Rana temporaria* in northeastern Iberia during the periods 2021-2040 and 2081-2100 predicted by MaxEnt algorithm based on CNRM-CM6 1 under low (left) and high (right) emission scenarios.




Table 1(on next page)

Kruskal-Wallis analyses results. 1st ,2nd ,3rd and 4th previous weeks to onset breeding period. Only significant variables are showed (*). Pluviometry (Rain), relative humidity, wind and Atmospheric pressure are not significant. T^a Av, Max and Min: Weakly

			Multiple Comparisons z' values between weeks						
Kruskal-Wallis test: H (3, N= 36)			Week	2		3		4	
				z	p	z	p	z	p
T ^a Av.	11.4164	0.0097*	1	0,9396	1,0000	2,2595	0,1431	3,1096	0,0112*
T ^a Max.	10.2592	0.0165*	1	0,8949	1,0000	2,0806	0,2248	2,9755	0,0176*
T ^a Min.	9.57457	0.0226*	1	1,2752	1,0000	2,1924	0,1701	2,9307	0,0203*
FD<0°C	8.97857	0.0296*	1	1,0850	1,0000	1,8569	0,3800	2,8300	0,0280*
IR	8.38596	0.0387*	1	1,7226	0,5097	2,0694	0,2311	2,7853	0,0321*

1

Table 2 (on next page)

Factor coordinates of the variables, based on correlations and Eigenvalues of correlation matrix and related statistics. FD: Number of days with temperature below 0°C. 1, 2 3, and 4: weeks before onset breeding. TX: maximal temperature. Tm: minimal temper

			Multiple Comparisons z' values between weeks						
Kruskal-Wallis test: H (3, N= 36)			Week	2		3		4	
				z	p	z	p	z	p
T ^a Av.	11.4164	0.0097*	1	0,9396	1,0000	2,2595	0,1431	3,1096	0,0112*
T ^a Max.	10.2592	0.0165*	1	0,8949	1,0000	2,0806	0,2248	2,9755	0,0176*
T ^a Min.	9.57457	0.0226*	1	1,2752	1,0000	2,1924	0,1701	2,9307	0,0203*
FD<0°C	8.97857	0.0296*	1	1,0850	1,0000	1,8569	0,3800	2,8300	0,0280*
IR	8.38596	0.0387*	1	1,7226	0,5097	2,0694	0,2311	2,7853	0,0321*

1