

Hard-bottom habitats support commercially important fish species: a systematic review for the North Atlantic Ocean and Baltic Sea

Hugo Flávio¹, Rochelle Seitz^{Corresp., 2}, David Eggleston³, Jon C. Svendsen⁴, Josianne Støttrup⁴

¹ 75 University Avenue West, Wilfrid Laurier University, Waterloo, Ontario, Canada

² Virginia Institute of Marine Science, William & Mary, Gloucester Point, Virginia, USA

³ Department of Marine, Earth & Atmospheric Sciences, North Carolina State University, Raleigh, North Carolina, USA

⁴ National Institute of Aquatic Resources, Technical University of Denmark, Lyngby, Denmark

Corresponding Author: Rochelle Seitz
Email address: seitz@vims.edu

Hard-bottom habitats span a range of natural substrates (e.g. boulders, cobble) and artificial habitats (e.g. the base of wind turbines, oil platforms). These hard-bottom habitats can provide a variety of ecosystem services, ranging from the enhancement of fish biomass and production to providing erosion control. Management decisions regarding the construction or fate of hard-bottom habitats require information on the ecological functions of these habitats, particularly for species targeted in ecosystem-based fisheries management. This study provides a systematic review of the relationships of various hard-bottom habitats to individual commercially harvested species that are managed jointly across the Atlantic by the International Council for the Exploration of the Sea (ICES). We systemically reviewed peer-reviewed publications on hard-bottom habitats including both some natural and artificial reefs, after applying various exclusion criteria. Most studies were conducted on near-shore hard-bottom habitats, and habitat importance varied according to fish species and region. We quantified the frequency with which studies demonstrate that natural and artificial hard-bottom habitats function as spawning grounds, settlement and nursery areas, and foraging grounds, as well as provide stepping-stones during migration, or new home ranges. Hard-bottom habitats generally support higher fish densities than surrounding habitat types, although not all fish species benefit from hard-bottom habitats. Of the commercially important species, cod (*Gadus morhua*) was the most frequently studied species, with enhanced biomass, density, feeding, and spawning on hard-bottom habitats compared to unstructured habitats. Moreover, hard-bottom habitats appear to be of particular importance for spawning of herring (*Clupea harengus*). Collectively, data indicate that loss of hard-bottom habitats may translate into less-favourable conditions for spawning and biomass of diverse commercial species, including

cod and herring.

Hard-bottom habitats support commercially important fish species: a systematic review for the North Atlantic Ocean and Baltic Sea

Hugo Flávio¹, Rochelle D. Seitz², David B. Eggleston³, Jon C. Svendsen⁴, Josianne G. Støttrup⁴

¹ Wilfrid Laurier University, 75 University Avenue West, Waterloo, Ontario, Canada

² Virginia Institute of Marine Science, William & Mary, Gloucester Point, Virginia 23062, USA

³ North Carolina State University, Department of Marine, Earth & Atmospheric Sciences, Raleigh, North Carolina 27695-8208, USA.

⁴ National Institute of Aquatic Resources, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

Corresponding Author:

Rochelle D. Seitz¹

Virginia Institute of Marine Science, William & Mary, Gloucester Point, Virginia 23062, USA

Email address: seitz@vims.edu

Hard-bottom habitats support commercially important fish species: a systematic review for the North Atlantic Ocean and Baltic Sea

Hugo Flávio¹, Rochelle D. Seitz², David B. Eggleston³, Jon C. Svendsen⁴, Josianne G. Støttrup⁴

¹ Wilfrid Laurier University, 75 University Avenue West, Waterloo, Ontario, Canada

² Virginia Institute of Marine Science, William & Mary, Gloucester Point, Virginia 23062, USA

³ North Carolina State University, Department of Marine, Earth & Atmospheric Sciences, Raleigh, North Carolina 27695-8208, USA.

⁴ National Institute of Aquatic Resources, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

Corresponding Author:

Rochelle D. Seitz¹

Virginia Institute of Marine Science, William & Mary, Gloucester Point, Virginia 23062, USA

Email address: seitz@vims.edu

Abstract

Hard-bottom habitats span a range of natural substrates (e.g. boulders, cobble) and artificial habitats (e.g. the base of wind turbines, oil platforms). These hard-bottom habitats can provide a variety of ecosystem services, ranging from the enhancement of fish biomass and production to providing erosion control. Management decisions regarding the construction or fate of hard-bottom habitats require information on the ecological functions of these habitats, particularly for species targeted in ecosystem-based fisheries management. This study provides a systematic review of the relationships of various hard-bottom habitats to individual commercially harvested species that are managed jointly across the Atlantic by the

International Council for the Exploration of the Sea (ICES). We systemically reviewed peer-reviewed publications on hard-bottom habitats including both some natural and artificial reefs, after applying various exclusion criteria. Most studies were conducted on near-shore hard-bottom habitats, and habitat importance varied according to fish species and region. We quantified the frequency with which studies demonstrate that natural and artificial hard-bottom habitats function as spawning grounds, settlement and nursery areas, and foraging grounds, as well as provide stepping-stones during migration, or new home ranges. Hard-bottom habitats generally support higher fish densities than surrounding habitat types, although not all fish species benefit from hard-bottom habitats. Of the commercially important species, cod (*Gadus morhua*) was the most frequently studied species, with enhanced biomass, density, feeding, and spawning on hard-bottom habitats compared to unstructured habitats. Moreover, hard-bottom habitats appear to be of particular importance for spawning of herring (*Clupea harengus*). Collectively, data indicate that loss of hard-bottom habitats may translate into less-favourable conditions for spawning and biomass of diverse commercial species, including cod and herring.

Keywords: reefs; fish abundance; spawning grounds; essential fish habitat

Introduction

Natural hard-bottom habitats are vital to the health and function of coastal ecosystems (Stål et al., 2007; Kovalenko et al., 2012; Simon et al., 2013; Lefcheck et al. 2019). These structures provide a range of services, including structural refuge for juveniles and feeding grounds for commercially important fish species. Hard-bottom habitats include a wide variety of topographic forms (e.g. high vertical profile, high complexity), substratum types (e.g. boulders, rocks, cobble, artificial materials), and degree of rugosity (e.g. smooth surfaces, perforated materials), with well-documented effects of increasing biodiversity with habitat complexity (Wilhelmsson et al., 2006; Kovalenko et al., 2012; Wehkamp and Fischer, 2013). As such, these hard substrates often comprise a three-dimensional structure, which means they can offer increased complexity in an otherwise two-dimensional landscape. Hard-bottom habitats provide substrate for the attachment of macrophytes, such as kelp and other macroalgae that further increase the complexity of the benthos and associated biodiversity, compared to unstructured habitat (Norderhaug et al., 2005; Christie et al., 2007, 2009). Reviews on the importance of coastal habitats for exploited species

identified a lack of information on how fish utilize some habitat types in the North Atlantic, particularly complex natural and artificial hard-bottom habitats (e.g. rocky shores, foundations of wind farms, oil-rigs; Seitz, 2014; Seitz et al., 2014, Fowler et al. 2018).

Artificial reefs have been used as a tool for restoration of marine habitats and as mitigation for damage to other marine habitats. Artificial reefs may be constructed using natural stone (Støttrup et al., 2017) or, more commonly, using a range of man-made materials including concrete units or scrap materials, such as decommissioned ships and oilrigs (Jensen, 2002; Soldal et al., 2002). Artificial reefs have been widely deployed to (1) enhance fisheries (Seaman and Sprague, 1991; Bombace et al., 1994; Brock, 1994), (2) mitigate damage to specific bottom habitats such as seagrass beds (*Posidonia* sp.; González-Correa et al., 2005), (3) increase the recreational value of an area (Wilhelmsson et al., 1998) or (4) as a tool for rehabilitation of coastal ecosystems (Pickering et al., 1999). Artificial reefs can be intentional (i.e. anthropogenic explicit creation of a reef) or unintentional (i.e. created as a by-product of other anthropogenic activities, such as oil platforms and wind turbines, hereafter referred to "*de facto*" reefs). Artificial reefs generally harbor higher fish densities and biomass than surrounding natural reefs (e.g. Wilhelmsson et al., 1998), which is attributed to their form and complexity (Hunter and Sayer, 2009; Degraer et al., 2020) and/or food abundance (Rilov and Benayahu, 2000; Fabi et al., 2006; Glarou et al. 2020), but artificial reefs may also increase abundance and richness of the fish community (Rilov and Benayahu, 2000). Moreover, effects of artificial reefs may vary by individual fish species (Leitão et al., 2008; Krone et al., 2013). Findings on the importance of artificial habitats for species managed by the International Council for the Exploration of the Sea (ICES) could be important for management decisions regarding habitat restoration and protection of fisheries species.

In Europe, hard-bottom habitats are one of the few marine habitat types included in the European Union's Habitat Directive (1170 Reef Habitat; EEC, 1992). For this reason, reef areas are included in several Natura 2000 networks such as the Danish Natura 2000 network. Surprisingly, the importance of temperate reefs is rarely considered in fisheries management and in fish population assessments (Lipcius et al., 2019, and references therein), and the importance of hard-bottom reefs may differ for different fishery species. Management decisions require the best available information on specific ecosystem

services that, in some cases, can be monetized (Costanza et al., 1997; Cavanagh et al., 2016). As such, the goal of this study was to summarize the existing information on ICES-managed fish species and their use of temperate hard-bottom habitats in an effort to (1) inform scientists interested in the ecological function of these habitats for some key fish species, and (2) inform managers (especially ICES managers) and conservation groups focused on coastal habitats of the potential importance of hard-bottom habitats to certain fish species. ICES is an inter-governmental marine science organization whose advisory committee translates science to management on “the sustainable use and protection of marine species,” and their work is relevant to extensive coastal areas, for example throughout Europe and the United States. Focusing our review on ICES-managed species is important because ICES-managed species are common throughout coastal areas of North America and Europe, and many commercial and recreational fisheries for these species are in poor condition, which causes concerns among conservationists as well as those dependent on fisheries as an economic activity or food source (ICES Science Plan 2013).

Materials & Methods

The overarching objective was to review the literature to quantify the ways in which hard-bottom habitats support commercially important fish species in temperate regions, particularly the North Atlantic, and Baltic and North Seas. We targeted the value of natural hard-bottom habitats and artificial reefs on exploited fish species by (1) summarizing increases or decreases in demographic variables (e.g. growth, survival, biomass, spawning rates) for individual commercial fishery species associated with hard-bottom habitats, and (2) highlighting the importance of these habitats for certain fish species so that managers and conservation groups can act on this knowledge. The spatial focus was on temperate, hard-bottom habitats (excluding subtropical and tropical areas, and the Mediterranean Sea), and we excluded some natural hard-bottom habitats such as coral reefs and biogenic habitats (see negative keywords in the search string below). We also focused on species important for management by ICES (see species-related keywords below), therefore, we did not include all commercially important species. In addition, we excluded rockfish in the Pacific Ocean, which have a separate body of literature targeted towards them, and which are not managed by ICES. Thus, our conclusions cannot be applied to all species in every area, and our focus was on non-biogenic or anthropogenically created reefs. We were most interested in

habitat enhancement from the physical structure provided by hard-bottom habitats, and thus excluded biogenic reefs, which provide additional ecological benefits such as increased nutrients (e.g. from faeces and pseudofaeces on mussel beds; Thiel and Ullrich, 2002).

Search protocol:

We followed the protocol for a systematic literature review proposed by Pullin and Stewart (2006), which set forth guidelines for planning the review to analysis of the data. We systematically reviewed the literature and evaluated papers returned by three different databases (DTU-Findit, Web of Science, and Scopus) for relevance based on our inclusion criteria (Table 1). Screening was conducted at the level of title, abstract, and full text (Pullin and Stewart, 2006) to each of several ecological study variables (see “search string” below), using the PRISMA protocol (Moher et al., 2009)(Fig. 1). The first author of this publication screened each paper by title to eliminate non-relevant papers, and a subset of articles was sent to all co-authors to compare the group’s decisions and ensure we came to consensus on what should be discarded. Then, the same procedure was followed for examining the abstracts of the remaining papers. Finally, we divided the papers for full-text examination among all authors and each person independently extracted data, with a group discussion of which papers were relevant and why, and resolving any discrepancies on what should be included. The search was run on the three different databases on April 7th, 2017. The papers returned by the three different databases were evaluated for relevance based on the inclusion criteria (Table 1), applied at three successive levels: (1) title, (2) abstract and (3) full-text. It is important to note that there are some limitations of our approach, and that we may have missed some research that did not fall into our selected categories based on the search terms we used, and results are only applicable to the species we incorporated into our search (namely ICES-managed species). Our review focused on the species for which ICES gives advice, which resulted in examining the species of importance for ICES Member Countries (i.e. Belgium, Canada, Denmark, Estonia, Finland, France, Germany, Iceland, Ireland, Latvia, Lithuania, the Netherlands, Norway, Poland, Portugal, Russia, Spain, Sweden, United Kingdom, and United States of America; US and Canadian fish stocks are not included in the advice, though these are ICES Member Countries). Therefore, interpretations of these data apply only to the species and habitats we incorporate. Additionally, we

examined the current literature (2017-2022) for more recent papers on the same ICES-managed species using hard bottom habitats (database from B. Ciotti pers. comm.), which allowed us to identify a couple of case studies to serve as a check on validity of conclusions from our systematic search.

Search string

The following search keywords were chosen for each of the ecological variables we wanted to examine: Biomass, Density, Biodiversity, Condition, Mortality, Growth, Feeding, Spawning, Nursery, Settling, Diel cycle, Site fidelity, Migration. Separate searches with all other keywords in combination with each of the ecological search keywords were conducted (Table 2). Location-related keywords and exclusion keywords were also included to narrow the search results.

Importance-related keywords (note: asterisks represent a search engine wild-card and were used to broaden the search by finding multiple words that start with the same letters):

Fish attraction, fish production, fish biomass, fish diversity, fish growth, fish survival, fish density, fish migration, demographic rate*, fish reproduction, fish mortality and fish juvenile*, feeding, nursery, spawn*.

Ecosystem-related keywords: Hard bottom, rocky reef*, stone reef*, riprap, seawall*, groin*, groyne*, offshore wind energy structure*, wind farm*, wind turbine*, artificial reef*, fish attraction device*, FAD*, shipwreck*, oil rig*, oil platform* oil pipe*, gas rig*, gas platform*, gas pipe*, offshore platform*, rig* to reef*, decommission*.

Species-related keywords: *Alopias* spp., *Amblyraja radiata*, *Ammodytes* spp., *Anguilla anguilla*, *Aphanopus carbo*, *Argentina silus*, *Beryx* spp., *Brosme brosme*, *Capros aper*, *Centrophorus squamosus*, *Centroscymnus coelolepis*, *Cetorhinus maximus*, *Chelidonichthys cuculus*, *Clupea harengus*, *Coryphaenoides rupestris*, *Dalatias licha*, *Dicentrarchus labrax*, *Dipturus batis-complex*, *Engraulis encrasicolus*, *Gadus morhua*, *Galeorhinus galeus*, *Galeus melastomus*, *Hoplostethus atlanticus*, *Lamna nasus*, *Lepidorhombus* spp., *Leucoraja circularis*, *Leucoraja fullonica*, *Leucoraja naevus*, *Limanda limanda*, *Lophius budegassa*, *Lophius piscatorius*, *Macrourus berglax*, *Mallotus villosus*, *Melanogrammus aeglefinus*, *Merlangius merlangus*, *Merluccius merluccius*, *Micromesistius poutassou*, *Molva dypterygia*,

Molva molva, *Mullus surmuletus*, *Mustelus* spp., *Pagellus bogaraveo*, *Phycis blennoides*, *Platichthys flesus*, *Pleuronectes platessa*, *Pollachius pollachius*, *Pollachius virens*, *Raja brachyura*, *Raja clavata*, *Raja microocellata*, *Raja montagui*, *Raja undulata*, *Reinhardtius hippoglossoides*, *Rostroraja alba*, *Salmo salar*, *Salmo trutta*, *Sardina pilchardus*, *Scomber scombrus*, *Scophthalmus maximus*, *Scophthalmus rhombus*, *Scyliorhinus canicula*, *Scyliorhinus stellaris*, *Sebastes* spp., *Solea solea*, *Sprattus sprattus*, *Squalus acanthias*, *Squatina squatina*, *Trachurus picturatus*, *Trachurus trachurus*, *Trachyrincus scabrus*, and *Trisopterus esmarkii*.

Location-related keywords: Belgium, Canada, Denmark, Estonia, Finland, France, Germany, Iceland, Ireland, Latvia, Lithuania, Netherlands, Norway, Poland, Portugal, Russia, Spain, Sweden, United Kingdom, United States of America, Northeast Atlantic, Atlantic Iberian, Biscay, English Channel, Bristol Channel, North sea*, Irish sea*, Cantabrian sea*, Belt sea*, Bothnian sea*, Celtic sea*, Azores, Skagerrak, Kattegat, Bornholm, Baltic, Gulf of Gdansk, Gulf of Riga, east Gotland, Faroe grounds, Iceland grounds, Greenland, Rockall, Bothnian bay, Reykjanes Ridge, Dogger bank*, Viking bank*, Bergen bank*.

Negative (Excluded) keywords: Biogenic reef*, eelgrass, mussel bed*, coral reef*, tropical, subtropical, Australia, Mediterranean, Caribbean, Indian sea.

Data extraction:

Our primary question was: “What is the habitat importance of hard-bottom natural and artificial structures for temperate fish species managed by ICES?” To answer this question, for each paper we determined the studied habitat, the ecosystem type, and species targeted, as detailed below. We broke down habitat importance into the following four main components: (1) distribution-related variables, (2) community-related variables, (3) fitness-related variables, and (4) reproduction-related variables (see details below).

Habitat importance: Fish species receive numerous benefits from benthic habitats, such as food, shelter, or spawning grounds. Such benefits may translate into measurable demographic variables (e.g. growth, survival, biomass, spawning rates), which allow researchers to assess the importance of various habitats. We initially grouped the data into the following four broad categories: (1) species distribution, (2)

community, (3) fitness, and (4) reproduction. Distribution-related variables included papers targeting diel cycles, site fidelity, or migration patterns. Community-related variables included papers targeting fish biomass, density, or biodiversity. Fitness-related variables included papers targeting condition, mortality rates, growth, and feeding. Lastly, reproduction-related variables included papers targeting spawning habitats, nursery potential, and settling (both of adults to newly formed habitats or of juveniles).

Ecosystem type: A species' ecological niche may encompass different ecosystems. Therefore, we targeted coastal marine environments and transitional areas, such as fjords and estuaries. Though we excluded biogenic reefs, we did include literature that dealt with macrophyte cover in temperate regions.

Target species: To improve future fisheries management, this review included fish species targeted for advice by the International Council for the Exploration of the Sea (ICES). Targeted species are listed above in "Species-related keywords."

Results & Discussion

Our study indicates that hard-bottom substrates may benefit multiple fish groups, such as elasmobranchs (e.g. Martin et al., 2012; Rodríguez-Cabello et al., 2008), gadoids (e.g. Reubens et al., 2014 b; Ross, Rhode, and Quattrini, 2015; van Hal et al., 2017), and other teleosts such as seabass (*Dicentrarchus labrax*; Leitão et al., 2008). Furthermore, hard substrates serve as: (1) stepping stones, where species may find shelter during migration events (Gascon and Miller, 1982), (2) spawning grounds for demersal eggs (Johannessen, 1986), (3) settlement grounds for young fish migrating out of nurseries (Santos et al., 2005), (4) foraging areas with increased prey densities (Leitão et al., 2008b) or (5) new home ranges, with newcomers developing high site fidelity rates, as observed for novel constructions such as marine wind farms (Reubens et al., 2013).

Review selection process

The search process returned an initial 2,651 papers, paring down to 2,333 papers after exclusion of duplicate references. Of those, 45 papers were returned after the subsequent screening criteria and were examined for full-text data extraction (as detailed below and in Fig. 1).

Title analysis: The title reviewing process led to the exclusion of 2036 papers (87% of the initial 2333), leaving 297 papers for abstract analysis. Most of the exclusions (54%) represented off-topic papers dealing with, for example, energy production, aquaculture, human health, oil spills, or physiology. Additionally, 35% of the excluded papers were explicitly related to non-target organisms (e.g. zebrafish, rainbow trout, invertebrates). A total of 9.5% excluded papers either studied non-target locations or other ecosystems (e.g. Hawaii or soft-bottom ecosystems, respectively). Lastly, 1.4% of the excluded titles either did not correspond to peer-reviewed papers or were not written in English.

Abstract analysis: From the 297 analysed abstracts, a total of 136 failed to comply with the inclusion criteria (46%), leaving 161 papers for full text analysis. Most of the exclusions represented papers related to non-target species (32%), followed by papers that did not relate to the target topics (25%). A total of 30% excluded papers either studied non-target locations or other ecosystems. Additionally, 13% of the analysed search results were not peer-reviewed literature.

Full text analysis: From the 161 full texts analysed, 116 (72%) did not comply with the exclusion criteria. Most of the papers were excluded based on location (49%), corresponding mostly to studies targeting rockfish in the Pacific Ocean. A total of 21% exclusions represented papers related to non-target species, while 13% studied non-target ecosystems and 7% did not relate to the target topics. A total of 11 papers were excluded for “other” reasons: four were not peer-reviewed, four were reviews, and for three it was not possible to obtain a full text version of the document. Therefore, the review focused on 45 studies distributed along the Baltic and North Seas and northern Atlantic Ocean (Fig. 2). We quantified the numbers of papers or percentage of papers reviewed as a function of the various response variables emphasized.

Papers reviewed as a function of response variables emphasized

Distribution-related variables (i.e., diel cycle, site fidelity, migration) were studied in six papers (13.4%). These targeted primarily site fidelity and migration patterns of Atlantic cod (*Gadus morhua*; e.g. Reubens et al., 2014 a, b).

Community-related variables (i.e. density, biodiversity, biomass) were the most frequently studied amongst the papers reviewed, being analysed in two thirds of the studies (Fig. 3). Most research targeted

a few species (i.e. Atlantic cod, herring [*Clupea harengus*], and saithe [*Pollachius virens*]) and a limited set of variables (i.e. biomass, density, biodiversity, and feeding). Cod was the species most commonly studied in community-related (i.e., density, biodiversity, biomass) studies (Table 2).

Fitness-related variables (i.e. condition, mortality, growth, feeding) were studied in 17 papers (38% of the total number of analysed papers). Feeding was studied primarily on cod and saithe, showing a tendency for fish use of hard-bottom habitats for foraging (Fig. 4; e.g. Johannessen, 1986; Malek et al., 2016; Wennhage and Pihl, 2002). Interestingly, with the exception of cod, little research has been carried out on the remaining fitness-related variables (Table 2).

Reproduction-related variables (i.e. spawning, settling, nursery) were studied in six papers (13.4%). Spawning was studied primarily on herring and cod, with papers on nursery and settling being almost absent for other species in the reviewed literature (Table 2). From the species targeted in our review, hard-bottom substrates appeared to be of particular importance for herring spawning (Aneer and Nellbring, 1982; Johannessen, 1986; Kääriä et al., 1997; Šaškov et al., 2014).

There was a clear predominance in the study of community-related variables in the analysed research (Fig. 3). Broadening the scope of studied variables and targeted species may prove essential to fully understand the relationships of hard-bottom reefs with commercially important species. For example, more quantitative research at the community level reveals the value of hard-bottom habitats for fish populations (essential fish habitat, *sensu* Valavanis, 2008; or effective habitat, *sensu* Dahlgren et al., 2006), and integrating this information into assessment and management may provide considerable advances towards ecosystem-based fisheries management (e.g., fish production estimates that can be monetized).

Ecosystems explored

Research was nearly equally distributed across natural and artificial reef structures, with 26 and 27 papers related to each, respectively, and eight targeting both natural and artificial structures simultaneously (Fig. 4, Table 3). Furthermore, 21 papers (46.7%) that resulted from our search performed comparisons of hard-bottom habitats with soft-bottom habitats. The review indicated that different regions of the world have different research foci. All papers on the north-western shore of the Atlantic Ocean

focused on natural reefs, with one paper also targeting an intentional artificial reef (Fig. 5). Similarly, there was a greater focus on natural reef research in the Baltic Sea, with nine out of eleven papers targeting natural reefs, as well as three papers targeting artificial *de facto* reefs, and only one paper targeting intentional artificial reefs. Conversely, in the North Sea, only 3 out of 15 papers targeted natural reefs, while 14 targeted *de facto* artificial reefs (most of which explored the effects of wind farms). Finally, on the north-eastern coast of the Atlantic Ocean, there were nearly equal numbers of papers on natural and artificial reef research (six and five papers, respectively, plus two papers that targeted both reef types simultaneously). However, of the papers targeting artificial reefs, only one studied *de facto* artificial reefs, while the remaining explored artificial intentional artificial reefs. There was a noticeable paucity of studies in the northern regions. Additionally, most of the analysed research has been conducted near-shore. Broadening the scientific knowledge on the importance of different habitats, areas, and regions along a depth gradient away from shore may provide crucial insights for future management.

Frequently studied species

There has been a consistent research effort towards multiple morphological and behavioural aspects of cod, with this species being researched twice as frequently as herring, the second-most-researched species (25 vs 12 papers, respectively; Table 2). Furthermore, while research on cod and herring is spread across the four groups of ecological variables considered in our review (i.e. species distribution, community, fitness, reproduction), research is lacking for most reviewed species on variables not related with community (e.g. condition, settling, site fidelity; Table 2).

The presence of hard-bottom habitat was associated with enhanced biomass in cod and saithe and, conversely, was negatively associated with biomass in herring and whiting (*Merlangius merlangus*; Fig. 6). Most papers compared fish densities in multiple habitats, and higher densities of cod on hard-bottom habitats were demonstrated in several papers, with only one paper suggesting fewer cod on hard-bottom compared to adjacent soft-bottom habitats. Both positive and negative relationships of hard-bottom habitats with density were demonstrated for herring, saithe, striped red mullet (*Mullus surmuletus*), whiting, pollock (*Pollachius pollachius*), plaice (*Pleuronectes platessa*), and sole (*Solea solea*). Cod, herring, saithe, striped red mullet, whiting, and pollock appeared to benefit from foraging on hard-bottom

habitat (i.e. for plaice feeding, only negative relationships of hard-bottom habitats were reported, as plaice feed less frequently over hard-bottom habitats). Few papers targeted spawning on hard-bottom habitats, and those that did reported associated benefits for cod and herring.

Changes in fish distribution

Research has demonstrated positive effects of the presence of hard-bottom substrates on cod biomass and density (e.g. Krone et al., 2013; Pihl and Wennhage, 2002). With a few exceptions, natural hard-bottom habitats and artificial reefs harbor significantly higher abundances of fish than adjacent unstructured habitats (Rodríguez-Cabello et al., 2008; Fujii, 2016). Fish movement patterns reflected relatively high site fidelity to structures (e.g. Hartman, 1987; Reubens et al., 2014 a, b), and the stomach contents of fish residing on reefs (excluding biogenic reefs) reflected the prey assemblages associated with the reefs (e.g. Reubens et al., 2014a), suggesting extended residency and foraging in the reef areas. Interestingly, catch per unit effort (CPUE) of cod was greater adjacent to wind farms than in shipwrecks in the Belgian part of the North Sea (Reubens et al., 2014b), suggesting attraction towards wind farms. It is important to note that the studied wind farms were closed to fisheries, while the shipwreck sites had no fisheries restrictions, possibly biasing the comparison.

Gadoids such as saithe, haddock (*Melanogrammus aeglefinus*) and cod were the most common species at the “Miller” oil and gas platform, in the North Sea (Fujii, 2016). Cod abundance was approximately four times higher at the platform than in nearby shipwrecks or sandy bottom. Using trawl data, Martin et al. (2012) modeled distributions of ten demersal elasmobranchs of the eastern English Channel in relation to the environment, with female small-spotted catshark (*Scyliorhinus canicula*) occurring at higher densities on hard substrate.

Within four years after the deployment of artificial reefs in an area previously closed to trawl fisheries in the Bay of Biscay, there was a rapid increase in the biomass of small-spotted catshark (Rodríguez-Cabello et al., 2008). Importantly, four to eight years after deployment, axillary sea-bream (*Pagellus acarne*), horse mackerel (*Trachurus trachurus*), and striped red mullet demonstrated the most significant increases in biomass. The observation that horse mackerel responded positively to the addition of

artificial reefs demonstrates that schooling pelagic species may also benefit from these hard-bottom habitats.

Reefs (excluding biogenic reefs) may also enhance economic returns. For example, in one location, reefs near Algarve and Faro, Portugal, raised catch rates and economic returns (Neves Santos and Costa Monteiro 1998). Specifically, CPUE at the reefs exceeded CPUE at the control sites by a factor of 2.03 – 2.28 for the protected reef and by a factor of 1.11 – 1.86 for the exploited reef. In some artificial reef systems, the economic return per unit of effort was higher at the reefs than at the control sites. Fishing on artificial reefs would be expected to yield 2.2 times the economic return compared to the control sites (Neves Santos and Costa Monteiro 1998). The productivity of reefs increased throughout the 15-year period, implying that the interaction between the hard substrate and the marine environment may take many years to produce the full effects on the carrying capacity (Whitmarsh et al., 2008). A distinct spatial pattern of fishing activity around the reefs was detected, with a concentration of fishing gear on the periphery of the reef area and a lower gear density at farther distances. A concentration of fishing gear nearby the reef would suggest that the improved fishing opportunities offered by the reefs have been recognized by fishermen. Fishermen likely take advantage of the spill-over of fish from the reefs, consistent with recent studies (Rosemond et al., 2018).

The inclusion of quantitative data on hard-bottom habitats in the monitoring of fish populations may further resolve a mismatch between fish population assessments and catches in fisheries of several fish species with a patchy distribution due to their high affinity for hard-bottom habitats (e.g. cod; Wieland et al., 2009). The mismatch between fish population assessments and catches of fish species with a patchy distribution may be further exacerbated during stock recovery, where the most suitable habitats (i.e. hard bottoms) will be populated first according to the ideal free distribution theory of MacCall (1990).

Not all fish species benefit from hard-bottom or marine constructed structures. Neutral or negative relationships with biomass and density were recorded for flatfish such as plaice (*Pleuronectes platessa*) and sole (*Solea solea*), as well as for whiting (*Merlangius merlangus*). For example, Støttrup et al. (2014) found decreasing densities of sole post restoration of a natural reef, and Bergström et al. (2013) only found whiting before the implementation of a wind farm. Neutral or negative relationships with hard-

bottom habitat for some species reflects the preferred habitat of the different species, with both flatfish and whiting occurring primarily on soft-bottom habitats (Stål et al., 2007).

The rapid growth of offshore wind farms, coupled with the high abundance and biomass of fish associated with wind farms, suggest that these *de facto* marine protected areas (MPAs) could be relatively easy to enforce, serve as broodstock sanctuaries that benefit fished populations, and optimize use of the seafloor (Fowler et al., 2018). Moreover, the economic impact of artificial reefs also appears high relative to unstructured control sites. Although the potential positive relationships of artificial reefs with fish populations are encouraging, it is important to recognize the attraction effect of artificial reefs and how such a process can facilitate overfishing (Bohnsack, 1989; Smith et al., 2015, and references therein). Artificial reefs, however, can also enhance growth and survival, for example by providing hard substrate for sessile fouling communities that provide food to fish (Todd et al., 2018). Quantitative evidence may also guide modeling efforts to assess the role of artificial reefs on population dynamics of exploited species. Lastly, given that unstructured seafloor (i.e. flat, soft-bottom habitat) can be many orders of magnitude greater in areal footprint than artificial reefs, it is important to recognize that unstructured habitats can serve as effective juvenile habitat (*sensu* Dahlgren et al., 2006) for exploited fishery species. Unstructured habitat is especially important for species for which ICES gives advice. For example, in comparing coastal habitat use among various structured (e.g. seagrass, kelp, marsh) and unstructured habitats, subtidal soft-bottom was the habitat used as a spawning or nursery area by the largest proportion of ICES species examined, with intertidal soft-bottom habitat as the next most important habitat (Seitz et al., 2014).

Habitat heterogeneity enhances biodiversity

Overall, both natural and artificial hard-bottom or reef structures tend to have a high attraction potential and house densities of fish superior to surrounding habitats (e.g. Bergström et al., 2013; Støttrup et al., 2014; Whitmarsh et al., 2008). In one study, natural structured habitats (rock, shell, rubble) compared to bare sand had positive effects on density, growth, and survival of juvenile fishes and invertebrates, with a greater positive effect for arthropod invertebrates than vertebrates (Lefcheck et al., 2019). When man-made structures are placed on surrounding soft-bottom areas (e.g. wind turbine foundations, oil

platforms), they represent beacons of habitat heterogeneity that are capable of attracting new residents (Gascon and Miller, 1982; Stenberg et al., 2015; Castège et al., 2016; Fujii, 2016). For example, cod appears to have a clear affinity for hard-bottom substrates, with increased fish densities and/or biomass near natural hard-bottom reefs (Stål et al. 2007; Støttrup et al. 2017; Kristensen et al. 2017), offshore wind farms (Bergström et al., 2013; Reubens et al., 2013; van Hal et al., 2017), oil-platforms (Fujii, 2015), shipwrecks (Wieland et al., 2009; Krone et al., 2013) and other man-made structures (Langhamer and Wilhelmsson, 2009; Wehkamp and Fischer, 2013). Similarly, pollock generally restricted its movement to a submerged reef area in Loch Ewe, Scotland, while saithe ranged more widely during daytime and returned to the reef during night-time (Sarno et al., 1994). Specific age groups of cod and pouting (*Trisopterus luscus*) can have high residency and site fidelity to the bases of marine wind turbines, feeding on the dominant epifaunal prey growing on these introduced habitats (Reubens et al., 2014a, b).

Different species react differently to the addition of hard-bottom substrates, and recent literature reports on various outcomes even within a species. For example, whiting can occur in relatively high abundance on hard substrates (e.g. shipwrecks and offshore wind farms; Krone et al., 2013; Lindeboom et al., 2011; Pihl and Wennhage, 2002), as well as on soft substrates (Stål et al. 2007; van Hal et al. 2017). Additionally, in some cases, the attraction of reef-associated demersal fish species such as bogue (*Boops boops*), horse mackerel, and axillary sea-bream may make them more vulnerable to predation, as the probability of prey–predator encounters increases compared to unstructured habitats (Leitão et al. 2008). Lastly, some commercially important species, such as dab (*Limanda limanda*) and sole, may avoid rocky substrates, with decreasing abundances whenever there is a shift from soft to hard bottom (van Hal et al. 2017). Thus, the relationship of hard-bottom structures with fish distribution and abundance patterns in the North Sea and Baltic Sea are species-dependent, a factor that must be considered during management and restoration actions.

Improved fitness and reproduction capacity

Selection by fish for complex, hard-bottom habitats generally improved fitness in those species occupying these habitats. Enhanced foraging opportunities improved condition in cod and pouting when occupying wind farm areas (De Troch et al., 2013). However, this is not always the case. For example, Mathers et al.

(1992) examined growth rates in saithe using biochemical indicators and found no differences in growth rates on an oil platform as compared to the open sea. Further, while pouting condition was enhanced in a wind farm area relative to sandy areas, condition of cod was similar in both habitat types (Reubens et al., 2014b). The similar growth rates or condition of fish on oil platforms or wind farms compared to unstructured areas suggests that although production was evident on a local scale (i.e. wind farm foundations), this may have had little effect on the larger spatial scale (i.e. the Belgian part of the North Sea). In a related example, oil and gas platforms in the North Sea provided unique feeding grounds for commercially important gadoids such as cod, saithe, and haddock (Fujii, 2016). For example, haddock predominantly consumed ophiuroids that occurred in high abundances on platforms at all depths, whereas temporal changes in the presence and abundance of saithe reflected the occurrence and availability of euphausiids (Fujii, 2016). These findings indicate that gadoids utilized the platform for foraging.

A tagging study revealed that spawning activities of cod in the Gulf of Maine occurred around specific bottom features, such as humps and ridges (Siceloff and Howell, 2013). Cod tended to move in groups after spawning, as 14 of 26 tagged and recaptured cod shifted to deep water (>100 m) within 23 to 50 days after tagging, with females tending to leave before males. While these findings indicate that the pelagic spawning of cod may be associated with large offshore hard-bottom structures, relatively shallow, complex habitats with rocks and cobble may enhance survival among the juveniles after benthic settling (Tupper and Boutilier, 1995).

A recent study highlighted the potential role of offshore oil and gas platforms as spawning sites for lump sucker (*Cyclopterus lumpus*; Todd et al., 2018). *C. lumpus* was observed during different stages of reproduction on an offshore oil and gas structure in the North Sea, providing novel evidence of spawning offshore at depths of around 40-45 m. Examples from the literature also show positive relationships of hard-bottom habitats with spawning and recruitment for several fish species, with much of the relevant literature coming from natural hard-bottom habitats. In a survey of natural hard bottoms in western Norway, spawning of herring generally occurred on hard-bottom substrates down to a depth of 10 m, with little evidence of spawning on soft-bottom habitats (Johannessen, 1986).

In the Baltic Sea, the vegetation usually attached to hard substrates seems to play an important role for successful spawning of Baltic herring (*C. harengus membras*). In a survey of natural hard bottoms in the northern Baltic Sea (Sweden), herring eggs were only found on filamentous algae (*Pilayella littoralis*) attached to hard substrates (Aneer and Nellbring 1982) and not on the surrounding hard or soft bottoms. Likewise, in the southwestern archipelago of Finland, Baltic herring spawning beds occurred in areas with vegetation growing on hard bottoms (down to a depth of 4 m); usually close to the regions' deepest zones (down to 60 m depth; Kääriä et al., 1997). The complex habitat established by the algae may provide refuge from predators, as birds and other fish are responsible for a majority of the predation of herring eggs (Aneer and Nellbring 1982). Areas without spawning beds generally had soft sediments with narrow bands of vegetation. Filamentous green algae (*Cladophora* sp.), brown algae (*Ectocarpus* sp., *Pilayella* sp.) and *Potamogeton pectinatus* made up the most common vegetation where divers observed eggs (Kääriä et al. 1997). For Baltic herring spawning on the Lithuanian coast (eastern coast of the Baltic), eggs were found on red algae *Furcellaria lumbricalis* and *Polysiphonia fucooides*, as well as hard habitat covered with mussels at 4-8 m depth (Šaškov et al. 2014). The dependence of Baltic herring on vegetation as substrate for the eggs may render this species vulnerable to pressures such as eutrophication that negatively affect macroalgal growth.

Modeling studies can be informative in assessing the effects of habitat on recruitment success. One model was used to assess the effects of MPAs on juvenile cod recruitment success (Lindholm et al., 2001). MPAs can maintain habitat complexity (i.e. more pebble-cobble substratum) compared to areas that are fished with dredges or trawls that damage the seafloor and reduce complexity (Ocean Studies Board & National Research Council 2002). Modeled survival of juvenile cod on natural reefs showed that survival was density-dependent (i.e. affected by predation or cannibalism; Wikan and Eide, 2004) and increased with seafloor habitat quality (modeled as increased size of an MPA). Juvenile cod had higher survival rates at low densities and in large MPAs, thus linking juvenile cod survival to seafloor habitat complexity (Lindholm et al. 2001). Modeled improved survival associated with complex sea-floor habitats is consistent with previous laboratory studies and may explain why various fish species prefer habitats with gravel and cobble where they can find shelter in the crevices (Gotceitas and Brown 1993; Christoffersen et al. 2018).

The review process that we used did not include all commercially important species, but rather was targeted towards species important for management by ICES, and we only examine temperate, non-biogenic reefs. The results from our study are limited to those specific species and the physical structure provided by hard-bottom, non-biogenic habitats. Our search terms were very specific and we acknowledge that this approach may have missed relevant research from habitats that we excluded. Therefore, interpretations of these data apply only to the species and habitats we incorporate.

Recent Case Studies

Although the literature for this study was not reviewed beyond those papers published in early April 2017, a related study reviewing methods that have examined the nursery role of different habitats included search terms relevant to hard-bottom habitats and fishery species of interest to ICES (Ciotti et al., in prep.). We highlight several recent case studies below that reflect this updated literature review. In the first example (Elliott et al. 2017), stereo baited remote underwater video (SBRUV) surveys were conducted in the nearshore, Scottish waters of the Firth of Clyde during summer 2013 and 2014 to determine the habitat of juvenile Atlantic cod, haddock and whiting. The fishery stocks of all three species declined in the late 20th century, and recruitment and spawning stock biomass remains relatively low in the Firth of Clyde despite efforts to re-build the stocks (Fernandes and Cook 2013, Holmes et al. 2014, ICES 2016a,b,c). Understanding juvenile gadoid habitat is particularly important given that settlement and post-settlement survival is thought to be the best means to understanding gadoid population regulation (Olsen and Moland 2011, Laurel et al. 2016). Atlantic cod were most abundant in shallow, sheltered areas composed of gravel-pebble containing maerl. Haddock and whiting predominated over deeper sand and mud. Relative abundances of Atlantic cod and whiting were positively related to the diversity of epibenthic and demersal fauna. Thus, spatial conservation measures to benefit demersal fish should consider habitat type and diversity (Elliot et al. 2017).

Another case study examined the environmental benefits of leaving de-commissioned offshore infrastructure, such as oil and gas platforms and wind turbines, in the ocean due to benefits such as biodiversity enhancement, provision of reef habitat, and protection from bottom trawling (Fowler et al. 2018). The rationale for this study was that the *in situ* ecosystem value of platforms and the negative

impacts of removal are not factored into decommissioning decisions in regions such as the North Sea, where over 80% of oil structures are more than a decade old (OSPAR Commission 2017a) and are likely integrated to at least some extent into existing ecosystems (Fowler et al. 2018). The removal policy is based on the assumption that “leaving the seabed as you found it” will minimize negative impacts on the marine environment. A total of 200 experts spanning academia, government, and private organizations were sent surveys seeking their opinions on the following: (1) appropriateness of the current removal policy, (2) identification of viable alternatives to complete removal of offshore infrastructure, (3) key environmental considerations and trade-offs for decommissioning decisions, and (4) advice on decommissioning considerations between platforms and wind turbines. Of most relevance to this review of hard-bottom habitats, 78% of the 40 respondents thought that artificial habitats with environmental value should be maintained and protected due to enhanced ecosystem services (Fowler et al. 2018). Evidence includes harboring threatened species (Bell and Smith 1999), providing reef habitat (Coolen 2020), boosting recruitment of overfished species (Love et al. 2006), producing fish biomass at a greater rate than any other marine ecosystem (by as much as a factor of ten; Claisse et al. 2014), and acting as foraging sites for top-order predators (Todd et al. 2009). Fowler et al. (2018) concluded that the traditional view that artificial structures must be removed from marine ecosystems simply because they do not “belong” there has shifted to one of environmental optimization based on comparative assessment (Fowler et al. 2015). Each decommissioning option will have positive and negative impacts that must be carefully weighed, while also accounting for site-specific characteristics and the broader environmental context of the disturbance.

Conclusions and implications for management

Our systematic review has elucidated that Atlantic cod, the most frequently studied species of the group of species managed by ICES, had increased biomass, density, feeding, and spawning associated with hard-bottom habitats compared to unstructured habitats. Spawning of herring, was increased in hard-bottom habitats, which appear to be of importance for this species. We detected that most research was conducted on Atlantic cod, herring, and saithe, and the most commonly studied variables included biomass, density, biodiversity, and feeding. Cod was the species most commonly studied, and studies

were mostly on community-related variables (i.e., density, biodiversity, biomass). However, not all responses to the presence of hard-bottom habitat were positive. For herring, saithe, striped red mullet, whiting, pollock, plaice, and sole, we quantified positive and negative relationships between fish density and hard-bottom habitats. Thus, we achieved our goal of summarizing the existing information on ICES-managed fish species and their use of temperate hard-bottom habitats. This information should help inform scientists interested in the ecological function of hard-bottom habitats for some key fish species, and inform managers and conservation groups focused on coastal habitats of the potential importance of hard-bottom habitats to certain fish species.

Natural hard-bottom reefs, if removed or destroyed, can only be restored by human intervention. Thus, it would be wise if management focuses on protecting natural reefs and finding the most cost-effective solutions for restoring reefs. In support of ecosystem-based fisheries management, research should provide additional scientific information on the value of hard-bottom habitats for biodiversity or for individual species on a local, national and regional scale. While this information is slowly being produced (e.g. Christoffersen et al., 2018; Liversage and Benkendorff, 2013; Liversage et al., 2017), the economic value of hard-bottom habitats remains largely unknown. This is in contrast to other complex marine habitats (e.g. eelgrass) where several recent studies have estimated the annual economic values per area (McArthur and Boland, 2006; Tuya et al., 2014; Cole and Moksnes, 2016). Similar economic valuation studies are needed for hard-bottom habitats, so that policy makers may assess the economic importance of protective measures related to these habitats. In areas where hard-bottom habitat has been removed or destroyed, consideration should be given to the economic and ecological benefits or disadvantages of removing decommissioned marine structures (e.g. Fowler et al., 2018). These structures could be equated to shipwrecks that become *de facto* reefs and are generally not removed.

The value of reef habitats and the need for their protection ultimately depends on the occurrence of that habitat type in the area/region and the potential for hard-bottom habitats to represent value as a resource. For example, in Denmark, although the extraction of boulders and larger stones from reef areas has ceased, extraction of smaller stones or gravel continues to take place, even though these hard bottom types are important habitats for juvenile eel and cod (Gotceitas et al., 1995; Christoffersen et al., 2018).

Research should provide the necessary information to allow for an effective restoration of the functions of destroyed reefs. Studies are increasingly investigating restoration methodologies for hard-bottom reefs (Støttrup et al., 2017; Liversage and Chapman, 2018). However, the currently available information is insufficient to conduct a restoration program that is not, in itself, explorative. From a cautionary standpoint, we recommend that anthropogenic activities degrading hard-bottom habitats should be carefully considered, and weighed against the accumulating evidence that highlights the importance of various hard-bottom habitats for several commercially important species.

Acknowledgements

This work stemmed from an ICES working group on the Value of Coastal Habitats for Exploited Species, and we thank all participants of the working group for discussions leading to insights incorporated in this research. This is contribution number XXXX from the Virginia Institute of Marine Science, William & Mary.

References

- Aneer G, Nellbring S (1982) A SCUBA-diving investigation of Baltic herring (*Clupea harengus membras* L.) spawning grounds in the Asko-Landsort area, northern Baltic proper. J Fish Biol 21:433–442. doi: 10.1111/j.1095-8649.1982.tb02849.x
- Bell N, Smith J (1999) Coral growing on North Sea oil rigs. Nature 402: 601
- Bergström L, Sundqvist F, Bergström U (2013) Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. Mar Ecol Prog Ser 485:199–210. doi: 10.3354/meps10344
- Bohnsack JA (1989) Are high densities of fishes at artificial reefs the result of habitat limitation or behavioral preference? Bull Mar Sci 44:631–645
- Bombace G, Fabi G, Fiorentini L, Speranza S (1994) Analysis of the efficacy of artificial reefs located in five different areas of the Adriatic Sea. Bull Mar Sci 55:559–580
- Brock RE (1994) Beyond fisheries enhancement: artificial reefs and ecotourism. Bull Mar Sci 55:1181–1188

- 574 Castège I, Milon E, Fourneau G, Tauzia A (2016) First results of fauna community structure and dynamics
575 on two artificial reefs in the south of the Bay of Biscay (France). *Estuar Coast Shelf Sci* 179:172–
576 180. doi: 10.1016/j.ecss.2016.02.015
- 577 Cavanagh RD, Broszeit S, Pilling GM, et al. (2016) Valuing biodiversity and ecosystem services: a useful
578 way to manage and conserve marine resources? *Proc R Soc B Biol Sci* 283:20161635. doi:
579 10.1098/rspb.2016.1635
- 580 Christie H, Jørgensen NM, Norderhaug KM (2007) Bushy or smooth, high or low; importance of habitat
581 architecture and vertical position for distribution of fauna on kelp. *J Sea Res* 58:198–208. doi:
582 10.1016/J.SEARES.2007.03.006
- 583 Christie H, Norderhaug KM, Fredriksen S (2009) Macrophytes as habitat for fauna. *Mar Ecol Prog Ser*
584 396:221–233. doi: 10.3354/meps08351
- 585 Christoffersen M, Svendsen JC, Kuhn JA, et al. (2018) Benthic habitat selection in juvenile European eel
586 *Anguilla anguilla* : implications for coastal habitat management and restoration. *J Fish Biol* 93:996–
587 999. doi: 10.1111/jfb.13807
- 588 Claisse JT, Pondella DJ, Love M et al. (2014) Oil platforms off California are among the most productive
589 marine fish habitats globally. *Proceedings of the National Academy of Sciences*, 111: 15462–67
- 590 Coolen JWP, van der Weide BB, Cuperus J, M Blomberg, Van Moorsel GWNM, Faasse MAA, Bos OG,
591 Degraer S, HJ Lindeboom (2020) Benthic biodiversity on old platforms, young wind farms, and rocky
592 reefs. *ICES Journal of Marine Science*, Vol 77, Issue 3, 250–1265
593 <https://doi.org/10.1093/icesjms/fsy092>.
- 594 Cole SG, Moksnes P-O (2016) Valuing multiple eelgrass ecosystem services in Sweden: fish production
595 and uptake of carbon and nitrogen. *Front Mar Sci* 2:121. doi: 10.3389/fmars.2015.00121
- 596 Costanza R, d'Arge R, de Groot R, et al. (1997) The value of the world's ecosystem services and natural
597 capital. *Nature* 387:253–260. doi: 10.1038/387253a0
- 598 Dahlgren C, Kellison G, Adams A, et al. (2006) Marine nurseries and effective juvenile habitats: concepts

and applications. *Mar Ecol Prog Ser* 312:291–295. doi: 10.3354/meps312291

Degraer S, Carey DA, Coolen JW, Hutchison Z., Kerckhof F, Rumes B, Vanaverbeke J (2020). Offshore wind farm artificial reefs affect ecosystem structure and functioning. *Oceanography* 33:48-57

De Troch M, Reubens JT, Heirman E, et al. (2013) Energy profiling of demersal fish: A case-study in wind farm artificial reefs. *Mar Environ Res* 92:224–233. doi: 10.1016/j.marenvres.2013.10.001

EEC (1992) Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. European Economic Community. *OJ L* 206, 22.7.1992, p. 7–50 (ES, DA, DE, EL, EN, FR, IT, NL, PT). <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A31992L0043>

Elliott SAM, Turrell WR, Heath MR, Bailey DM (2017) Juvenile gadoid habitat and ontogenetic shift observations using stereo-video baited cameras. *Marine Ecology Progress Series*, Vol. 568: 123–135, <https://doi.org/10.3354/meps12068>

Fabi G, Manoukian S, Spagnolo A (2006) Feeding behavior of three common fishes at an artificial reef in the northern Adriatic Sea. *Bull Mar Sci* 78:39–56

Fernandes PG, Cook RM (2013) Reversal of fish stock decline in the Northeast Atlantic. *Current Biology* 23: 1432–1437

Fowler AM, Macreadie P, Booth DJ (2015) Should we “reef” obsolete oil platforms? *Proceedings of the National Academy of Sciences, USA* 112: E102

Fowler AM, Jørgensen A-M, Svendsen JC, et al. (2018) Environmental benefits of leaving offshore infrastructure in the ocean. *Front Ecol Environ* 16:571–578. doi: 10.1002/fee.1827

Fujii T (2016) Potential influence of offshore oil and gas platforms on the feeding ecology of fish assemblages in the North Sea. *Mar Ecol Prog Ser* 542:167–186. doi: 10.3354/meps11534

Fujii T (2015) Temporal variation in environmental conditions and the structure of fish assemblages around an offshore oil platform in the North Sea. *Mar Environ Res* 108:69–82. doi: 10.1016/J.MARENVRES.2015.03.013

623 Gascon D, Miller RA (1982) Space utilization in a community of temperate reef fishes inhabiting small
624 experimental artificial reefs. *Can J Zool* 60:798–806

625 Glarou, M, Zrust M, Svendsen JC (2020) Using Artificial-Reef Knowledge to Enhance the Ecological
626 Function of O₂ shoreWind Turbine Foundations: Implications for Fish Abundance and Diversity. *J*
627 *Mar Sci Eng* 8:332; doi:10.3390/jmse8050332

628 González-Correa JM, Bayle JT, Sánchez-Lizaso JL, et al. (2005) Recovery of deep *Posidonia oceanica*
629 meadows degraded by trawling. *J Exp Mar Bio Ecol* 320:65–76. doi: 10.1016/J.JEMBE.2004.12.032

630 Gotceitas V, Brown JA (1993) Substrate selection by juvenile Atlantic cod (*Gadus morhua*): effects of
631 predation risk. *Oecologia* 93:31–37. doi: 10.1007/BF00321187

632 Gotceitas V, Fraser S, Brown JA (1995) Habitat use by juvenile Atlantic cod (*Gadus morhua*) in the
633 presence of an actively foraging and non-foraging predator. *Mar Biol* 123:421–430. doi:
634 10.1007/BF00349220

635 Hartman AR (1987) Movement of scorpionfishes (Scorpaenidae *Sebastes* and *Scorpaena*) in the
636 Southern California Bight. *Calif Fish Game* 73:68–79

637 Holmes SJ, Millar CP, Fryer RJ, Wright PJ (2014) Gadoid dynamics: differing perceptions when
638 contrasting stock vs. population trends and its implications to management. *ICES Journal of Marine*
639 *Science*, 71: 1433–1442

640 Hunter WR, Sayer MDJ (2009) The comparative effects of habitat complexity on faunal assemblages of
641 northern temperate artificial and natural reefs. *ICES J Mar Sci* 66:691–698. doi:
642 10.1093/icesjms/fsp058

643 International Council for the Exploration of the Sea (ICES) (2016a) Cod (*Gadus morhua*) in division VIa
644 (West of Scotland). ICES, Copenhagen. [www.ices.dk/sites/
645 pub/Publication/20Reports/Advice/2015/2015/cod-scow.pdf](http://www.ices.dk/sites/pub/Publication/20Reports/Advice/2015/2015/cod-scow.pdf)

646 International Council for the Exploration of the Sea (ICES) (2016b) Haddock (*Melanogrammus*
647 *aeglefinus*) in Subarea 4 and Divisions 6a and 3a West (North Sea, West of Scotland, Skagerrak).

648 ICES, Copenhagen. [www.ices.dk/sites/pub/Publication %20Reports/Advice/2016/2016/had-346a](http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2016/2016/had-346a)

649 International Council for the Exploration of the Sea (ICES) (2016c) Whiting (*Merlangius merlangus*) in

650 Division 6a (West of Scotland). ICES, Copenhagen. www.ices.dk/sites/pub/Publication

651 %20Reports/Advice/2016/2016/whg-scow.

652 International Council for the Exploration of the Sea (ICES). 2013. ICES Science Plan 2009-2013. ICES,

653 Copenhagen. www.ices.dk/sites/pub/Publication

654 Jensen A (2002) Artificial reefs of Europe: perspective and future. ICES J Mar Sci 59:S3–S13. doi:

655 10.1006/jmsc.2002.1298

656 Johannessen A (1986) Recruitment studies of herring (*Clupea harengus* L.) in Lindaaspollene, western

657 Norway, 1-3. Fisk Skr Ser Havundersøkelser 18:139–240

658 Kääriä J, Rajasilta M, Kurkilahti M, Soikkeli M (1997) Spawning bed selection by the Baltic herring

659 (*Clupea harengus membras*) in the Archipelago of SW Finland. ICES J Mar Sci 54:917–923

660 Kovalenko KE, Thomaz SM, Warfe DM (2012) Habitat complexity: approaches and future directions.

661 Hydrobiologia 685:1–17. doi: 10.1007/s10750-011-0974-z

662 Kristensen LD, Støttrup JG, Svendsen JC, et al. (2017) Behavioural changes of Atlantic cod (*Gadus*

663 *morhua*) after marine boulder reef restoration: Implications for coastal habitat management and

664 Natura 2000 areas. Fish Manag Ecol 24:353–360. doi: 10.1111/fme.12235

665 Krone R, Gutow L, Brey T, et al. (2013) Mobile demersal megafauna at artificial structures in the German

666 Bight - Likely effects of offshore wind farm development. Estuar Coast Shelf Sci 125:1–9. doi:

667 10.1016/j.ecss.2013.03.012

668 Langhamer O, Wilhelmsson D (2009) Colonisation of fish and crabs of wave energy foundations and the

669 effects of manufactured holes – A field experiment. Mar Environ Res 68:151–157. doi:

670 10.1016/j.marenvres.2009.06.003

671 Laurel BJ, Knoth BA, Ryer CH (2016) Growth, mortality, and recruitment signals in age-0 gadids settling

672 in coastal Gulf of Alaska. ICES Journal of Marine Science, 73: 2227–2237

- 673 Lefcheck JS, Hughes BB, Johnson AJ, et al. (2019) Are coastal habitats important nurseries? A
674 meta-analysis. *Conserv Lett* 12: e12645. doi: 10.1111/conl.12645
- 675 Leitão F, Santos MN, Erzini K, Monteiro CC (2008) Fish assemblages and rapid colonization after
676 enlargement of an artificial reef off the Algarve coast (Southern Portugal). *Mar Ecol* 29:435–448. doi:
677 10.1111/j.1439-0485.2008.00253.x
- 678 Lindeboom HJ, Kouwenhoven HJ, N Bergman MJ, et al. (2011) Short-term ecological effects of an
679 offshore wind farm in the Dutch coastal zone; a compilation. *Environ Res Lett* 6:035101
- 680 Lindholm JB, Auster PJ, Ruth M, Kaufman L (2001) Modeling the effects of fishing and implications for
681 the design of marine protected areas: Juvenile Fish Responses to Variations in Seafloor Habitat.
682 *Conserv Biol* 15:424–437. doi: 10.1046/j.1523-1739.2001.015002424.x
- 683 Lipcius RN, Eggleston DB, Fodrie FJ, et al. (2019) Modeling quantitative value of habitats for marine and
684 estuarine populations. *Front Mar Sci* 6:280
- 685 Liversage K, Benkendorff K (2013) A preliminary investigation of diversity, abundance, and distributional
686 patterns of chitons in intertidal boulder fields of differing rock type in South Australia. *Molluscan Res*
687 33:24–33. doi: 10.1080/13235818.2012.754145
- 688 Liversage K, Chapman M (2018) Coastal ecological engineering and habitat restoration: incorporating
689 biologically diverse boulder habitat. *Mar Ecol Prog Ser* 593:173–185. doi: 10.3354/meps12541
- 690 Liversage K, Cole V, Coleman R, McQuaid C (2017) Availability of microhabitats explains a widespread
691 pattern and informs theory on ecological engineering of boulder reefs. *J Exp Mar Bio Ecol* 489:36–
692 42. doi: 10.1016/J.JEMBE.2017.01.013
- 693 Love MS, Schroeder DM, Lenarz W, et al. (2006) Potential use of offshore marine structures in rebuilding
694 an overfished rockfish species, bocaccio (*Sebastes paucispinis*). *Fish B-NOAA*
695 104: 383–90
- 696 MacCall AD (1990) Dynamic geography of marine fish populations. Washington Sea Grant Program :
697 Distributed by University of Washington Press, Seattle

- 698 Malek AJ, Collie JS, Taylor DL (2016) Trophic structure of a coastal fish community determined with diet
699 and stable isotope analyses. J Fish Biol 89:1513–1536. doi: 10.1111/jfb.13059
- 700 Martin CS, Vaz S, Ellis JR, et al. (2012) Modelled distributions of ten demersal elasmobranchs of the
701 eastern English Channel in relation to the environment. J Exp Mar Bio Ecol 418–419:91–103. doi:
702 10.1016/j.jembe.2012.03.010
- 703 Mathers EM, Houlihan DF, Cunningham MJ (1992) Estimation of saithe *Pollachius virens* growth rates
704 around the Beryl oil platforms in the North Sea: a comparison of methods. Mar Ecol Prog Ser 86:31–
705 40
- 706 McArthur LC, Boland JW (2006) The economic contribution of seagrass to secondary production in South
707 Australia. Ecol Modell 196:163–172. doi: 10.1016/J.ECOLMODEL.2006.02.030
- 708 Moher D, Libersti A, Tetzla J, Altman DG, Group TP (2009) Preferred Reporting Items for Systematic
709 Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med. 6: e1000097.
- 710 Neves Santos M, Costa Monteiro C (1998) Comparison of the catch and fishing yield from an artificial reef
711 system and neighbouring areas off Faro (Algarve, south Portugal). Fish Res 39:55–65. doi:
712 10.1016/S0165-7836(98)00169-6
- 713 Norderhaug HM, Christie H, Fosså JH, Fredriksen S (2005) Fish–macrofauna interactions in a kelp
714 (*Laminaria hyperborea*) forest. J Mar Biol Assoc United Kingdom 85:1279–1286. doi:
715 10.1017/S0025315405012439
- 716 Ocean Studies Board & National Research Council (2002) Effects of trawling and dredging on seafloor
717 habitat, August 9. National Academies Press
- 718 Olsen EM, Moland E (2011) Fitness landscape of Atlantic cod shaped by harvest selection and natural
719 selection. Evolution and Ecology 25: 695–710
- 720 OSPAR Commission (2017) OSPAR inventory of offshore installations – 2015. London, UK: OSPAR
721 Commission
- 722 Pickering H, Whitmarsh D, Jensen A (1999) Artificial Reefs as a Tool to Aid Rehabilitation of Coastal

- 723 Ecosystems: Investigating the Potential. *Mar Pollut Bull* 37:505–514. doi: 10.1016/S0025-
724 326X(98)00121-0
- 725 Pihl L, Wennhage H (2002) Structure and diversity of fish assemblages on rocky and soft bottom shores
726 on the Swedish west coast. *J Fish Biol* 61:148–166. doi: 10.1111/j.1095-8649.2002.tb01768.x
- 727 Pullin AS, Stewart GB (2006) Guidelines for systematic review in conservation and environmental
728 management. *Conserv Biol* 20:1647–1656. doi: 10.1111/j.1523-1739.2006.00485.x
- 729 Reubens JT, De Rijcke M, Degraer S, Vincx M (2014a) Diel variation in feeding and movement patterns
730 of juvenile Atlantic cod at offshore wind farms. *J Sea Res* 85:214–221. doi:
731 10.1016/j.seares.2013.05.005
- 732 Reubens JT, Degraer S, Vincx M (2014b) The ecology of benthopelagic fishes at offshore wind farms: a
733 synthesis of 4 years of research. *Hydrobiologia* 727:121136. doi: 10.1007/s10750-013-1793-1
- 734 Reubens JT, Vandendriessche S, Zenner AN, et al. (2013) Offshore wind farms as productive sites or
735 ecological traps for gadoid fishes? - Impact on growth, condition index and diet composition. *Mar*
736 *Environ Res* 90:66–74. doi: 10.1016/j.marenvres.2013.05.013
- 737 Rilov G, Benayahu Y (2000) Fish assemblage on natural versus vertical artificial reefs: the rehabilitation
738 perspective. *Mar Biol* 136:931–942. doi: 10.1007/s002279900250
- 739 Rodríguez-Cabello C, Sánchez F, Serrano A, Olaso I (2008) Effects of closed trawl fishery areas on some
740 elasmobranch species in the Cantabrian Sea. *J Mar Syst* 72:418–428. doi:
741 10.1016/j.jmarsys.2007.09.010
- 742 Rosemond R, Paxton A, Lemoine H, et al. (2018) Fish use of reef structures and adjacent sand flats:
743 implications for selecting minimum buffer zones between new artificial reefs and existing reefs. *Mar*
744 *Ecol Prog Ser* 587:187–199. doi: 10.3354/meps12428
- 745 Sarno B, Glass CW, Smith GW (1994) Differences in diet and behaviour of sympatric saithe and pollack
746 in a Scottish sea loch. *J Fish Biol* 45:1–11. doi: 10.1111/j.1095-8649.1994.tb01080.x
- 747 Šaškov A, Šiaulys A, Bučas M, et al. (2014) Baltic herring (*Clupea harengus membras*) spawning grounds

- 748 on the Lithuanian coast: current status and shaping factors. *Oceanologia* 56:789–804. doi:
749 10.5697/oc.56-4.789
- 750 Seaman W, Sprague LM (1991) Artificial Habitat Practices in Aquatic Systems. In: Seaman W, Sprague
751 LM (eds) Artificial Habitats for Marine and Freshwater Fisheries. Academic Press, pp 1–29
- 752 Seitz RD (2014) Value of coastal habitats for exploited species: introduction to a theme set of articles.
753 *ICES J Mar Sci* 71:636–637. doi: 10.1093/icesjms/fst180
- 754 Seitz RD, Wennhage H, Bergström U, et al. (2014) Ecological value of coastal habitats for commercially
755 and ecologically important species. *ICES J Mar Sci* 648–665. doi: 10.1093/icesjms/fst152
- 756 Siceloff L, Howell WH (2013) Fine-scale temporal and spatial distributions of Atlantic cod (*Gadus morhua*)
757 on a western Gulf of Maine spawning ground. *Fish Res* 141:31–43. doi:
758 10.1016/j.fishres.2012.04.001
- 759 Simon T, Joyeux J-C, Pinheiro HT (2013) Fish assemblages on shipwrecks and natural rocky reefs
760 strongly differ in trophic structure. *Mar Environ Res* 90:55–65. doi:
761 10.1016/J.MARENVRES.2013.05.012
- 762 Smith JA, Lowry MB, Suthers IM (2015) Fish attraction to artificial reefs not always harmful: a simulation
763 study. *Ecol Evol* 5:4590–602. doi: 10.1002/ece3.1730
- 764 Soldal A, Svellingen I, Jørgensen T, Løkkeborg S (2002) Rigs-to-reefs in the North Sea: hydroacoustic
765 quantification of fish in the vicinity of a “semi-cold” platform. *ICES J Mar Sci* 59:S281–
766 S287. doi: 10.1006/jmsc.2002.1279
- 767 Stål J, Pihl L, Wennhage H (2007) Food utilisation by coastal fish assemblages in rocky and soft bottoms
768 on the Swedish west coast: Inference for identification of essential fish habitats. *Estuar Coast Shelf*
769 *Sci* 71:593–607. doi: 10.1016/j.ecss.2006.09.008
- 770 Stenberg C, Støttrup JG, van Deurs M, et al. (2015) Long-term effects of an offshore wind farm in the
771 North Sea on fish communities. *Mar Ecol Prog Ser* 528:257–265. doi: 10.3354/meps11261
- 772 Støttrup JG, Dahl K, Niemann S, et al. (2017) Restoration of a boulder reef in temperate waters: Strategy,

773 methodology and lessons learnt. *Ecol Eng* 102:468–482. doi: 10.1016/J.ECOLENG.2017.02.058

774 Støttrup JG, Stenberg C, Dahl K, et al. (2014) Restoration of a temperate reef: effects on the fish
775 community. *Open J Ecol* 4:1045–1059. doi: 10.4236/oje.2014.416086

776 Thiel M, Ullrich N. (2002) Hard rock versus soft bottom: the fauna associated with intertidal mussel beds
777 on hard bottoms along the coast of Chile, and considerations on the functional role of mussel
778 beds. *Helgol Mar Res* 56:21-30.

779 Todd VL, Pearse WD, Tregenza NC, et al. (2009) Diel echolocation activity of harbour porpoises
780 (*Phocoena phocoena*) around North Sea offshore gas installations. *ICES Journal of Marine Science*
781 66: 734–45

782 Todd VLG, Lavallin EW, Macreadie PI (2018) Quantitative analysis of fish and invertebrate assemblage
783 dynamics in association with a North Sea oil and gas installation complex. *Mar Environ Res* 142:69–
784 79. doi: 10.1016/J.MARENVRES.2018.09.018

785 Tupper M, Boutilier RG (1995) Effects of habitat on settlement, growth, and postsettlement survival of
786 Atlantic cod (*Gadus morhua*). *Can J Fish Aquat Sci* 52:1834–1841. doi: 10.1139/f95-176

787 Tuya F, Haroun R, Espino F (2014) Economic assessment of ecosystem services: Monetary value of
788 seagrass meadows for coastal fisheries. *Ocean Coast Manag* 96:181–187. doi:
789 10.1016/J.OCECOAMAN.2014.04.032

790 Valavanis VD (2008) Preface: European Commission’s “Scientific Support to Policies” Action EnviEFH:
791 Environmental Approach to Essential Fish Habitat Designation. *Hydrobiologia* 612:1–3. doi:
792 10.1007/s10750-008-9494-x

793 van Hal R, Griffioen AB, van Keeken OA (2017) Changes in fish communities on a small spatial scale, an
794 effect of increased habitat complexity by an offshore wind farm. *Mar Environ Res* 126:26–36. doi:
795 10.1016/j.marenvres.2017.01.009

796 Wehkamp S, Fischer P (2013) Impact of coastal defence structures (tetrapods) on a demersal hard-
797 bottom fish community in the southern North Sea. *Mar Environ Res* 83:82–92. doi:

798 10.1016/j.marenvres.2012.10.013

799 Wennhage H, Pihl L (2002) Fish feeding guilds in shallow rocky and soft bottom areas on the Swedish
800 west coast. *J Fish Biol* 61:207–228. doi: 10.1111/j.1095-8649.2002.tb01772.x

801 Whitmarsh D, Santos MN, Ramos J, Monteiro CC (2008) Marine habitat modification through artificial
802 reefs off the Algarve (southern Portugal): An economic analysis of the fisheries and the prospects
803 for management. *Ocean Coast Manag* 51:463–468. doi: 10.1016/j.ocecoaman.2008.04.004

804 Wieland K, Pedersen EMF, Olesen HJ, Beyer JE (2009) Effect of bottom type on catch rates of North Sea
805 cod (*Gadus morhua*) in surveys with commercial fishing vessels. *Fish Res* 96:244–251. doi:
806 10.1016/J.FISHRES.2008.11.010

807 Wikan A, Eide A (2004) An analysis of a nonlinear stage-structured cannibalism model with application to
808 the Northeast Arctic cod stock. *Bull Math Biol* 66:1685–1704. doi: 10.1016/j.bulm.2004.03.005

809 Wilhelmsson D, Öhman MC, Ståhl H, Shlesinger Y (1998) Artificial Reefs and Dive Tourism in Eilat,
810 Israel. *Ambio* 27:764–766

811 Wilhelmsson D, Yahya SAS, Öhman MC (2006) Effects of high-relief structures on cold temperate fish
812 assemblages: A field experiment. *Mar Biol Res* 2:136–147. doi: 10.1080/17451000600684359

813

814

815

Table 1 (on next page)

Admission/exclusion criteria

Admission/exclusion criteria for hard-bottom review

1 Table 1: Admission/exclusion criteria

Criteria	Include	Exclude
Peer-reviewing	Peer-reviewed	Everything else
Text Language	English	Everything else
Ecosystem	Natural and artificial hard-bottom reefs in marine or contiguous salt water environments (i.e. fjords and estuaries)	Biogenic reefs or other structures placed at non-marine environments
Target species	Fish species managed by ICES	No fish species managed by ICES present
Location	North hemisphere, temperate zone (not including Mediterranean Sea)	Mediterranean Sea, everything else
Type of study	Field studies	Everything else

2

Table 2 (on next page)

Total number of papers (no parentheses) and percentage of papers in relation to the total of papers (in parentheses) for each of the top eight fish species that were most commonly selected in our review, in relation to the different ecological variables o

The groups of variables are delineated by the grey horizontal lines: community-, fitness-, reproduction- and distribution-related (from top to bottom; terms defined in the Materials & Methods section). In the “Total” column, the percentage of papers is shown in relation to the total number of papers reviewed (i.e. 45). Note that any given paper may simultaneously target more than one species and/or variable. Most research has targeted few species (i.e. cod, herring, and saithe) and a limited set of variables (i.e. biomass, density, biodiversity, and feeding). Full genus names given in “Species-related keywords” section.

1 Table 1: Total number of papers (no parentheses) and percentage of papers in relation to the total of papers (in parentheses) for each of the top
 2 eight fish species that were most commonly selected in our review, in relation to the different ecological variables of interest in the review. The
 3 groups of variables are delineated by the grey horizontal lines: community-, fitness-, reproduction- and distribution-related (from top to bottom;
 4 terms defined in the Materials & Methods section). In the “Total” column, the percentage of papers is shown in relation to the total number of
 5 papers reviewed (i.e. 45). Note that any given paper may simultaneously target more than one species and/or variable. Most research has
 6 targeted few species (i.e. cod, herring, and saithe) and a limited set of variables (i.e. biomass, density, biodiversity, and feeding). Full genus
 7 names given in “Species-related keywords” section.

	<i>G. morhua</i>	<i>C. harengus</i>	<i>P. virens</i>	<i>M. surmu.^a</i>	<i>M. merlangus</i>	<i>P. pollachius</i>	<i>P. platessa</i>	<i>S. solea</i>	Total
Biomass	3(12)	1(8.3)	1(9.1)	3(30)	3(30)	2(22.2)	1(11.1)	0(0)	6(13.3)
Density	15(60)	6(50)	5(45.5)	7(70)	7(70)	6(66.7)	7(77.8)	5(62.5)	24(53.3)
Biodiversity	6(24)	4(33.3)	5(45.5)	7(70)	6(60)	5(55.6)	5(55.6)	7(87.5)	15(33.3)
Condition	3(12)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	3(6.7)
Mortality	2(8)	2(16.7)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	3(6.7)
Growth	1(4)	0(0)	1(9.1)	0(0)	0(0)	0(0)	0(0)	0(0)	2(4.4)
Feeding	8(32)	2(16.7)	4(36.4)	1(10)	1(10)	2(22.2)	2(22.2)	0(0)	12(26.7)
Spawning	1(4)	4(33.3)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	5(11.1)
Nursery	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)
Settling	1(4)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	1(2.2)
Diel cycle	4(16)	1(8.3)	2(18.2)	1(10)	1(10)	2(22.2)	1(11.1)	1(12.5)	4(8.9)
Site fidelity	3(12)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	3(6.7)
Migration	1(4)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	1(2.2)
Total	25(55.6)	12(26.7)	11(24.4)	10(22.2)	10(22.2)	9(20)	9(20)	8(17.8)	45

8 ^a *Mullus surmuletus*

Table 3(on next page)

Distribution of papers per fish species and different hard-bottom structures.

Artificial reefs are further split into intentional and de facto reefs, with de facto reefs further subdivided into the various structures that form de facto reefs. “Other” includes wave-energy foundations and coastal-protection structures (i.e. riprap and breakwater tetrapods). The percentage of papers in relation to the total of papers for each species is shown in the brackets. In the “Total” column and row, the proportion of papers is shown in relation to the total number of papers reviewed (i.e. 45). Note that any given paper may simultaneously target more than one species and/or more than one hard-bottom structure. Overall, data shows an even distribution of research between natural and artificial reefs, despite the spatial patterns noted in Figure 4. Full genus names given in “Species-related keywords” section.

1 Table 1: Distribution of papers per fish species and different hard-bottom structures. Artificial reefs are further split into intentional and *de facto*
 2 reefs, with *de facto* reefs further subdivided into the various structures that form *de facto* reefs. “Other” includes wave-energy foundations and
 3 coastal-protection structures (i.e. riprap and breakwater tetrapods). The percentage of papers in relation to the total of papers for each species is
 4 shown in the brackets. In the “Total” column and row, the proportion of papers is shown in relation to the total number of papers reviewed (i.e. 45).
 5 Note that any given paper may simultaneously target more than one species and/or more than one hard-bottom structure. Overall, data shows an
 6 even distribution of research between natural and artificial reefs, despite the spatial patterns noted in Figure 4. Full genus names given in
 7 “Species-related keywords” section.

	G. morhua	C. harengus	P. virens	M. surmu. ^a	M. merlangus	P. pollachius	P. platessa	S. solea	Total
Natural Reefs	12(48)	9(75)	9(81.8)	5(50)	5(50)	7(77.8)	5(55.6)	3(37.5)	26(57.8)
Artificial Reefs	18(72)	4(33.3)	7(63.6)	6(60)	7(70)	4(44.4)	6(66.7)	7(87.5)	27(60)
Intentional	3(12)	1(8.3)	3(27.3)	2(20)	1(10)	3(33.3)	1(11.1)	2(25)	8(17.8)
<i>De facto</i>	15(60)	3(25)	4(36.4)	4(40)	6(60)	1(11.1)	5(55.6)	5(62.5)	19(42.2)
Wind Farms	10(40)	3(25)	1(9.1)	3(30)	6(60)	0(0)	5(55.6)	5(62.5)	12(26.7)
Shipwrecks	3(12)	0(0)	1(9.1)	1(10)	1(10)	0(0)	1(11.1)	0(0)	4(8.9)
Platforms	2(8)	0(0)	3(27.3)	0(0)	0(0)	0(0)	0(0)	0(0)	3(6.7)
Other	3(12)	0(0)	0(0)	1(10)	0(0)	1(11.1)	0(0)	0(0)	4(8.9)
Total	25(55.6)	12(26.7)	11(24.4)	10(22.2)	10(22.2)	9(20)	9(20)	8(17.8)	45

8 ^a *Mullus surmuletus*

Figure 1

PRISMA flow diagram for our hard-bottom systematic review showing numbers of papers at different stages in the review.

PRISMA flow diagram for our hard-bottom systematic review showing numbers of papers at different stages in the review.




Figure 2

Geographical distribution of the papers that were included in the review.

Geographical distribution of the papers that were included in the review.




Figure 3

Number of papers targeting each of the ecological response variables recorded. Note that each paper may target more than one variable. Variables were grouped by colors, showing a clear dominance of community-related studies.

Number of papers targeting each of the ecological response variables recorded. Note that each paper may target more than one variable. Variables were grouped by colors, showing a clear dominance of community-related studies.




Figure 4

Number of papers targeting each of the studied hard-bottom substrates. Note that papers may target more than one topic simultaneously.

Number of papers targeting each of the studied hard-bottom substrates. Note that papers may target more than one topic simultaneously.




Figure 5

Distribution of papers with grouping of studied reef structures.

Distribution of papers with grouping of studied reef structures: natural reefs only (downward pointing green triangles); artificial reefs only (upward pointing blue triangles); both natural and artificial reefs (orange diamonds).




Figure 6

Effects of the presence of hard-bottom habitats on the most commonly studied variables.

Effects of the presence of hard-bottom habitats on the most commonly studied variables: biomass (B), density (D), feeding (F) and spawning (S), for the species most commonly targeted in recent research. Given the variety of methods used in literature the effects are listed as positive, neutral or negative. When no comparison was performed with a different substrate, the effect was considered unknown.

