

# ZFP36 ring finger protein like 1 significantly suppresses human coronavirus OC43 replication

Tooba Momin<sup>1</sup>, Andrew Villasenor<sup>1</sup>, Amit Singh<sup>1</sup>, Mahmoud Darweesh<sup>2</sup>, Aditi Singh<sup>1</sup>, Mrigendra Rajput<sup>Corresp. 1</sup>

<sup>1</sup> Department of Biology, University of Dayton, Dayton, Ohio, United States

<sup>2</sup> Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Uppsala, Sweden

Corresponding Author: Mrigendra Rajput  
Email address: mrajput1@udayton.edu

CCCH-type Zinc finger proteins (ZFP) are small cellular proteins that are structurally maintained by zinc ions. Zinc ions coordinate the protein structure in a tetrahedral geometry by binding to cystine- cystine or cysteines - histidine amino acids. ZFP's unique structure enables it to interact with a wide variety of molecules including RNA; thus, ZFP modulates several cellular processes including the host immune response and virus replication. CCCH-type ZFPs have shown their antiviral efficacy against several DNA and RNA viruses. However, their role in the human coronavirus is little explored. We hypothesized that ZFP36L1 also suppresses the human coronavirus. To test our hypothesis, we used OC43 human coronavirus (HCoV) strain in our study. We overexpressed and knockdown ZFP36L1 in HCT-8 cells using lentivirus transduction. Wild type, ZFP36L1 overexpressed, and ZFP36L1 knockdown cells were each infected with HCoV-OC43, and the virus titer in each cell line was measured over 96 hours post-infection (p.i.). Our results show that HCoV-OC43 replication was significantly reduced with ZFP36L1 overexpression while ZFP36L1 knockdown significantly enhanced virus replication. ZFP36L1 knockdown HCT-8 cells started producing infectious virus at 48 hours p.i. which was an earlier timepoint as compared to wild -type and ZFP36L1 overexpressed cells. Wild-type and ZFP36L1 overexpressed HCT-8 cells started producing infectious virus at 72 hours p.i.. Overall, the current study showed that overexpression of ZFP36L1 suppressed human coronavirus (OC43) production.

# ZFP36 Ring Finger Protein Like 1 significantly suppresses Human coronavirus OC43 replication

Tooba Momin<sup>1</sup>, Andrew Villasenor<sup>1</sup>, Amit Singh<sup>1</sup>, Mahmoud Darweesh<sup>2</sup>, Aditi Singh<sup>1</sup>, Mrigendra Rajput<sup>1</sup>

<sup>1</sup> Department of Biology, University of Dayton, Dayton, Ohio, USA 45469

<sup>2</sup> Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden, 75105

Corresponding Author:

Mrigendra Rajput

SC 234, Department of Biology, 300 College Street, University of Dayton, Dayton, Ohio, 45469, USA

# **Abstract**

CCCH-type Zinc finger proteins (ZFP) are small cellular proteins that are structurally maintained by zinc ions. Zinc ions coordinate the protein structure in a tetrahedral geometry by binding to cystine- cystine or cysteines - histidine amino acids. ZFP's unique structure enables it to interact with a wide variety of molecules including RNA; thus, ZFP modulates several cellular processes including the host immune response and virus replication. CCCH-type ZFPs have shown their antiviral efficacy against several DNA and RNA viruses. However, their role in the human coronavirus is little explored. We hypothesized that ZFP36L1 also suppresses the human coronavirus. To test our hypothesis, we used OC43 human coronavirus (HCoV) strain in our study. We overexpressed and knockdown ZFP36L1 in HCT-8 cells using lentivirus transduction. Wild type, ZFP36L1 overexpressed, and ZFP36L1 knockdown cells were each infected with HCoV-OC43, and the virus titer in each cell line was measured over 96 hours post-infection (p.i.).

Our results show that HCoV-OC43 replication was significantly reduced with ZFP36L1 overexpression while ZFP36L1 knockdown significantly enhanced virus replication. ZFP36L1 knockdown HCT-8 cells started producing infectious virus at 48 hours p.i. which was an earlier timepoint as compared to wild -type and ZFP36L1 overexpressed cells. Wild-type and ZFP36L1 overexpressed HCT-8 cells started producing infectious virus at 72 hours p.i.. Overall, the current study showed that overexpression of ZFP36L1 suppressed human coronavirus (OC43) production.

**Keywords:** CCCH type Zinc finger protein, ZFP36L1, RNA binding protein, human coronavirus OC43

# Introduction

ZFPs are small cellular proteins that are structurally maintained by zinc ions. Zinc ions coordinate the protein structure in a tetrahedral geometry (Abbehausen, 2019; Hajikhezri, 2020). There are over 40 different types of ZFPs that have been annotated so far (Hajikhezri, 2020). ZFP's unique structure enables it to interact with a wide variety of molecules such as DNA, RNA, PAR (poly-ADP-ribose), and cellular proteins and thus modulate several cellular processes including host immune response and virus replication (Müller et al., 2007; Cassandri et al., 2017; Takata et al., 2017; Tang, Wang & Gao, 2017; Meagher et al., 2019; Lal, Ullah & Syed, 2020; Nchioua et al., 2020; Gonzalez-Perez et al., 2021; Wang & Zheng, 2021). Among various ZFPs, the CCCH-type ZFP family contains zinc ions that coordinate protein structure by binding to cystine-cystine or cysteines-histidine amino acids (Abbehausen, 2019; Hajikhezri, 2020). The CCCH-type ZFP family has also been characterized for its antiviral (Hajikhezri, 2020; Tang, Wang & Gao, 2017; Zhang, et al., 2020; Guo et al., 2004; Zhao et al., 2019; Gao, Guo & Goff, 2002; Zhu et al., 2020; Musah, 2004; Chen, Jeng & Lai, 2017; Scozzafava et al., 2003; Schito et al., 2006; Angiolilli et al., 2021) and immune modulator activity (Wang et al., 2015; Tu et al., 2019; Haneklaus et al., 2017; Lv et al., 2021; Matsushita et al., 2009; Wawro, Kochan & Kasza, 2016; Uehata, & Akira, 2013; Chen et al., 2018; Mino et al., 2015; Fu & Blackshear, 2017; Stumpo, Lai & Blackshear, 2010; Shrestha, Pun & Park, 2018; Kontoyiannis, 2018; Chiu et al., 2022).

CCCH-type ZFPs show their antiviral efficacy against several RNA viruses including Influenza A virus (Tang, Wang & Gao, 2017), retrovirus (Gao, Guo & Goff, 2002; Zhu et al., 2011; Zhu et al., 2017) filoviruses (Müller, 2007), and alphavirus such as Sindbis virus, Semliki Forest virus, Ross River virus, and Venezuelan equine encephalitis virus (Bick et al., 2003). However, CCCH-type ZFP's role on the human coronavirus is little explored. The current study is designed to evaluate the effect of ZFP36L1, a CCCH-type ZFP, on human coronavirus (HCoV)-OC43 replication.

# Materials & Methods

## Cells, Virus Strains and Virus Propagation

HCT-8 cells (ATCC, Manassas, VA) were cultured in Roswell Park Memorial Institute (RPMI) 1640 Medium (Gibco BRL, Grand Island, NY) and supplemented with 10% heat-inactivated fetal bovine serum (FBS), (ATCC, Manassas, VA), and antibiotic-antimycotic: penicillin 100 units /ml, streptomycin 0.10 mg /ml and amphotericin B 0.25 µg /ml (Sigma-Aldrich, St. Louis, MO). During virus culture, HCT-8 cells were adapted to 1% FBS. HCT-8 cells cultured with RPMI 1640 medium supplemented with 1% FBS were used to grow and subsequently titrate the OC43 human coronavirus (HCoV) stain (ATCC, Manassas, VA).

## Overexpression and knockdown of ZFP36L1

HCT-8 cells were stably overexpressed for ZFP36L1 (NCBI reference sequence: NM\_001244701.1) with a green fluorescent protein (GFP) marker using a lentivirus vector. The ZFP36L1 gene containing both tandem zinc finger domains (TZFD) such as TZFD1 and TZFD2 were cloned in a pLV-eGFP plasmid with the help of Vector Builder Inc, IL. To make the lentivirus, pLV-eGFP plasmids containing our gene of interest were co-transfected with VSV-G and packaging plasmids encoding Gag/Pol and Rev in HEK293T cells. After 48 hours, the supernatant containing the lentivirus was collected and purified by centrifugation followed by filtration. Purified lentivirus was concentrated using a sucrose gradient ultracentrifugation and this concentrated, purified lentivirus was used in the study.

Similar to ZFP36L1 overexpression, HCT-8 cells were knockdown for ZFP36L1 using ZFP36L1 specific shRNA (GTAACAAGATGCTCAACTATA). The ZFP36L1 shRNA was stably expressed using a lentivirus by cloning it in a pLV-mCherry plasmid. Lentivirus for ZFP36L1-

shRNA was prepared as per the above-mentioned method by co-transfection of pLV-mCherry containing ZFP36L1 shRNA with VSV-G and packaging plasmids in HEK293T cells. The prepared lentiviruses were used to either overexpress or knock down ZFP36L1. Successful lentivirus transduction was measured through GFP or mCherry expression for ZFP36L1 overexpression (GFP) or ZFP36L1 knockdown (mCherry), respectively. Transduced HCT-8 cells were selected with an increased concentration of puromycin (2-3  $\mu\text{g/ml}$ ) over 7 days. Selected cells were further characterized for ZFP36L1 overexpression or knockdown using a western blot with ZFP36L1-specific antibodies.

### **Western blot analysis for ZFP36L1 expression**

To confirm ZFP36L1 overexpression or ZFP36L1 knockdown; wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were individually seeded in T25 flasks. When cells reached 75-80% confluency, cells were lysed using a radioimmunoprecipitation assay buffer (RIPA buffer) (Cell Signaling Technology, Danvers, MA) supplemented with protease-phosphatase inhibitor (Cell Signaling Technology, Danvers, MA). Lysates were then centrifuged at 3000 X g for 15 minutes at 4 °C. The supernatant was collected and the protein concentration in each supernatant was measured using the Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific, Waltham, MA). 40  $\mu\text{g}$  cell lysates were separated through 12% resolving SDS PAGE gel. After separation, proteins were transferred onto a polyvinylidene difluoride (PVDF) membrane (Thermo Fisher Scientific, Waltham, MA). The PVDF membrane was blocked with 5% skimmed milk (Sigma-Aldrich, St. Louis, MO) in Tris-buffered saline (TBS) for 1 hour at room temperature followed by incubation with anti- ZFP36L1 antibody (1:1000) (Thermo Fisher Scientific, Waltham, MA) and anti-  $\beta$ actin antibody (1:4000) (Cell Signaling Technology, Danvers, MA) overnight at 4 °C. After overnight incubation, membranes

were washed with tris-buffered saline +0.1% Tween 20 (TBST) and incubated with HRP conjugated secondary antibodies (1:2000) for 1 hour at room temperature. After washing, membranes were developed using the Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific, Waltham, MA). Images of the western blot were taken by the Odyssey XF Imaging System (LI-COR Biosciences, Lincoln, NE). Band intensity for ZFP36L1 proteins was normalized with  $\beta$  actin using ImageJ software (Schneider, Rasband & Eliceiri, 2012). A significant difference in ZFP36L1 expression in ZFP36L1 overexpressed and knockdown cells compared to wild-type cells was estimated using a paired T-test.

#### **Determining ZFP36L1's effect on HCT-8 cells viability**

The effect of ZFP36L1 overexpression and its knockdown on cell viability was measured by trypan blue exclusion assay (Strober, 2015). Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown cells were individually seeded in 6 well plates ( $1.5 \times 10^6$  cell/well) in triplicate. 96 hours post-seeding, cells were washed with sterilized phosphate-buffered saline (PBS) and detached with 0.25% trypsin-EDTA (ATCC, Manassas, VA). Detached cells were washed with PBS by centrifugation at 500x g for 5 minutes at 4 °C, and then cells were stained with 0.4% trypan blue for 3 minutes and examined for cell. Changes in cell viability following ZFP36L1 overexpression or its knockdown compared to wild-type cells was estimated by paired t-test.

#### **Measuring the effect of ZFP36L1 expression on virus titration**

Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were infected with HCoV-OC43 with 0.1 multiple of infection (MOI) individually. The supernatant from these cells was collected at 24 hours, 48 hours, 72 hours, and 96 hours p.i. Collected cell supernatants were then centrifuged at 1000Xg at 4°C for 15 minutes to remove cell debris and stored at -80 °C until used. Once samples from all time points were collected, the HCoV-OC43 virus titer was

determined as per the aforementioned method (Reed & Muench, 1938). Changes in virus titer in ZFP36L1 overexpressed or ZFP36L1 knockdown cells were compared to wild-type cells and statistically analyzed by a paired T-test.

# **Measuring the effect of ZFP36L1 expression on HCoV-OC43 replication.**

To measure the effect of ZFP36L1 overexpression or ZFP36L1 knockdown on HCoV-OC43 replication, we infected ZFP36L1 overexpressed, ZFP36L1 knockdown or wild type HCT-8 cells with HCoV-OC43 (MOI: 0.1). Infected cells were collected at 72 and 96 hours p.i. Viral RNA was isolated from infected cells using the QIAamp Viral RNA Mini kit (Qiagen, Valencia, CA, USA). The viral nucleocapsid was quantified using qPCR (Stratagene MX3000P Real-Time Thermocycler, Stratagene Inc., La Jolla, USA) in 25 µl reaction using syber green dye. Primer sequence for nucleocapsid (F: 5'-: GCTGTT TWTGTAAAG TCYAAA GT-3', R: 5'-ATTCTGATAGAGAGTGCYTAT Y-3') were used (Al-Khannaq, et al., 2016) with qPCR amplification cycle at 95°C/ 2 minutes, 40 cycles of (95 °C/15 and 60 °C/ 1 minutes) followed by melting curve cycle at: 95°C/ 15 seconds, 60°C/ 1 minute and 95°C/ 15 seconds. Fold change in HCoV nucleocapsid expression in each cell was estimated by paired -test.

# **Statistical analysis**

The significant change in HCoV-OC43 titer and virus replication in wild-type, ZFP36L1 overexpressed, or ZFP36L1 knockdown cells was estimated using a paired T-test with 95% degree of freedom. Virus titer in wild-type, ZFP36L1 overexpression or ZFP36L1 knockdown cells was repeated at least three times with calculations for average, standard deviation, and standard error.

# **Results**

**ZFP36L1 was overexpressed or knockdown in HCT-8 cells.**



A stable ZFP36L1 overexpression with an upstream GFP marker in HCT-8 cells was generated using a lentivirus system. GFP expression in HCT-8 cells was considered positive for ZFP36L1 overexpression (Figure 1B), which was further confirmed by western blot (Figure 2 and Figure 3). Similarly, ZFP36L1 was knockdown using ZFP36L1-specific shRNA. The shRNA was located downstream to mCherry and expression of ZFP36L1-specific shRNA was determined by mCherry expression (Figure 1C) and ZFP36L1 knockdown was confirmed by western blot analysis (Figure 2 and Figure 3). Our results showed that lentivirus significantly overexpressed or knockdown ZFP36L1 in HCT-8 cells ( $p < 0.05$ ) (Figure 2 and Figure 3).

#### **ZFP36L1 overexpressing or its knockdown did not affect HCT-8 cells' viability**

The effect of ZFP36L1 overexpression or its knockdown was measured on HCT-8 cells' viability using trypan blue exclusion assay. Results showed that overexpression or knockdown of ZFP36L1 in HCT-8 cells did not affect its viability. Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown cells showed viability as  $94.83 \pm 1.01\%$ ,  $94.16 \pm 0.71\%$ , and  $95.83 \pm 0.43\%$  at 96 hours post seeding, respectively. These values were non-significant different to each other ( $p < 0.05$ ) (Figure 4). Additionally, no apparent morphological changes were observed among these cells.

#### **ZFP36L1 overexpression significantly suppressed while ZFP36L1 knockdown significantly enhanced the HCoV-OC43 production.**

Wild type, ZFP36L1 overexpressed, and ZFP36L1 knockdown HCT-8 cells were infected individually with HCoV-OC43 with MOI of 0.1. Cell supernatants were collected at 24 hours, 48 hours, 72 hours, and 96 hours p.i. and analyzed for virus titer.

Results showed that ZFP36L1 overexpression in HCT-8 cells significantly reduced virus titer ( $p < 0.05$ ) (Figure 5). Virus titer in ZFP36L1 overexpressed cells was  $2.24 \pm 1.28 \log_{10}/\text{ml}$  and  $4.32 \pm 0.00 \log_{10}/\text{ml}$  at 72 hours and 96 hours p.i. respectively. These titer values were significantly lower than virus titers in wild-type cells at same time points, such as 72 hours p.i. ( $4.08 \pm 0.11 \log_{10}/\text{ml}$ ) and 96 hours p.i. ( $5.42 \pm 0.10 \log_{10}/\text{ml}$ ) ( $p < 0.05$ ) (Figure 5).

Results with ZFP36L1 knockdown HCT-8 cells showed that ZFP36L1 knockdown significantly enhanced virus titer ( $p < 0.05$ ) (Figure 5). Knocking down ZFP36L1 facilitated the infectious virus production as early as 48 hours p.i. while wild-type cells produced infectious viruses at 72 hours p.i. The virus titer in ZFP36L1 knockdown cells was recorded as  $0.00 \pm 0.00 \log_{10}/\text{ml}$ ,  $2.86 \pm 0.00 \log_{10}/\text{ml}$ ,  $4.52 \pm 0.22 \log_{10}/\text{ml}$ , and  $5.85 \pm 0.01 \log_{10}/\text{ml}$  at 24 hours, 48 hours, 72 hours and 96 hours p.i., respectively. While wild-type HCT-8 cells have virus titer of  $0.00 \pm 0.00 \log_{10}/\text{ml}$ ,  $0.00 \pm 0.00 \log_{10}/\text{ml}$ ,  $4.08 \pm 0.11 \log_{10}/\text{ml}$ , and  $5.42 \pm 0.10 \log_{10}/\text{ml}$  at 24 hours, 48 hours, 72 hours and 96 hours p.i., respectively. Virus titer in ZFP36L1 knockdown cells was significantly higher at 48 hours and 96 hours p.i. compared to wild-type cells ( $p < 0.05$ ) (Figure 5). Results also showed a lower cytopathic effect in ZFP36L1 overexpressed or wild-type HCT-8 cells compared to ZFP36L1 knockdown cells at 72 hours p.i. (Figure 6)

**ZFP36L1 overexpression significantly suppressed while ZFP36L1 knockdown significantly enhanced the HCoV-OC43 RNA replication.**

To further confirm ZFP36L1's effect on HCoV-OC43 RNA replication, wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were individually infected with HCoV-OC43 (MOI: 0.1). Infected cells were collected at 72 and 96 hours p.i. Viral RNA was isolated from infected cells and viral nucleocapsid transcription (RNA concentration) was analyzed using qPCR. Results did not show any significant difference in HCoV-OC43 nucleocapsid RNA

concentration among these cells at 72 hours p.i. However, at 96 hours p.i. ZFP36L1 knockdown HCT-8 cells showed significantly higher HCoV-OC43 nucleocapsid transcription compared to wild type HCT-8 cells ( $p<0.05$ ). ZFP36L1 knockdown HCT-8 cells displayed an  $11.14\pm2.21$ -fold increase in HCoV-OC43 nucleocapsid RNA compared to wild-type HCT-8 cells. While ZFP36L1 overexpressed cells displayed a significantly lower HCoV-OC43 nucleocapsid RNA ( $0.37\pm0.13$  fold) compared to wild-type cells at 96 hours p.i. ( $p<0.05$ ) (Figure 7).

## Discussion

The current study was designed to determine the role of ZFP36L1 (a CCCH type ZFP) on HCoV-OC43 replication. Our results showed that overexpression of ZFP36L1 significantly reduced infectious HCoV-OC43 production while ZFP36L1 knockdown significantly enhanced virus titer compared to wild-type cells. ZFP36L1 overexpression also reduced the RNA replication of HCoV-OC43 and suppressed the apparent cytopathic effect in infected cells. ZFPs are one of the most abundant proteins in humans which can make up to 5% of total human proteins (Vilas et al., 2018). ZFPs have an extremely high binding ability. They can bind to cellular DNA, RNA, lipids, proteins, and PAR (poly-ADP-ribose); therefore, ZFPs can modulate several cellular types of machinery (Müller et al., 2007; Cassandri et al., 2017; Takata et al., 2017; Tang, Wang & Gao, 2017; Vilas et al., 2018; Meagher et al., 2019; Lal, Ullah & Syed, 2020; Nchioua et al., 2020, Wang & Zheng, 2021; Gonzalez-Perez et al., 2021). The diverse binding properties of ZFPs make it difficult to characterize their functional effect in cells (Vilas et al., 2018). However, such a challenge is overcome by classifying the ZFPs and then identifying their functional characteristics (Cassandri et al., 2017). Classification of ZFP is based on zinc ion, zinc ion interaction with specific amino acids, and the protein's folded structure (Krishna, Majumdar & Grishin, 2003). Based on such classification, CCCH-type ZFP is characterized to interact with

RNA and thus modulate RNA metabolism in the cell (Maeda & Akira, 2017). including interfering with RNA virus replication (Gao, Guo & Goff, 2002; Cassandri et al., 2017).

The known mechanisms by which CCCH-type ZFPs exhibit these antiviral or immunomodulatory activities is by limiting the total mRNA turnover in the cell. CCCH-type ZFPs such as ZFP36L1 have two tandem zinc finger (TZF) domains that are known to bind with adenyl and uracyl nucleotides-rich (AU-rich) elements (AREs) in mRNA. This interaction facilitates RNA degradation by CCR4-NOT complex-mediated deadenylation, followed by 5' decapping and exonuclease-mediated nucleotide cleaving (Blackshear, 2002; Lai, Kennington & Blackshear, 2003; Lykke-Andersen & Wagner, 2005; Suk et al., 2018; Lai et al., 2019; Lai et al., 2000; Chiu et al., 2022).

Coronavirus genome, including HCoV-OC43's genome is 5'-capped with a 3' poly(A) tail of variable length (Fehr & Perlman, 2015). The length of the poly (A) tail varies at different stages of the virus replication cycle and viruses with longer poly (A) tails replicate at a faster rate (Wu et al., 2013). Therefore, the effect of ZFP36L1 on viral poly (A) may explain reduced virus production with ZFP36L1 overexpression in the current study. Our study not only showed that ZFP36L1 suppressed the infectious HCoV-OC43 production, but also reduced HCoV-OC43 nucleocapsid transcription indicating that ZFP36L1 mediates its antiviral effect by limiting the viral RNA in infected cells.

However, there is the possibility that ZFP36L1 can reduce virus replication with different mechanisms other than poly A tail interaction. A study showed that CCCH Type ZFP also targets the non-ARE sequence of 3' and 5' (untranslated region) UTR in mRNA (Li et al., 2015). While another study showed that CCCH Type ZFP targets CG-rich viral sequences (Meagher et al., 2019). The study also showed that ZFP36 (ZFP36L1) suppressed the virus production (influenza

A virus) by interfering with viral protein translation/ export from the nucleus to the cytoplasm without affecting viral RNA replication (Lin et al., 2020). Therefore, a detailed study to determine ZFP36L1's mechanism of action for suppressing coronavirus replication needs to be explored.

## Conclusions

The current study showed that overexpression of ZFP36L1, a CCCH type ZFP significantly reduced HCoV-OC43 RNA (nucleocapsid) and infectious virus production. A reduced viral production was in correlation with reduced cytopathic effect in the infected cells. Furthermore, ZFP36L1 knockdown significantly enhanced the HCoV-OC43 replication and infectious virus production. However, additional mechanisms employed to reduce virus replication still need to be explored.

## Acknowledgments

The author thanks the Department of Biology, University of Dayton, Ohio, USA, for providing support to conduct the current research.

## References

1. Abbehausen C. 2019. Zinc finger domains as therapeutic targets for metal-based compounds - an update. *Metallomics* 11(1): 15-28.
2. Al-Khannaq MN, Ng KT, Oong XY, Pang YK, Takebe Y, Chook JB, Hanafi NS, Kamarulzaman A, Tee KK. 2016. Molecular epidemiology and evolutionary histories of human coronavirus OC43 and HKU1 among patients with upper respiratory tract infections in Kuala Lumpur, Malaysia. *Virology Journal*. 25;13:33. doi: 10.1186/s12985-016-0488-4. PMID: 26916286; PMCID: PMC4766700.
3. Angiolilli C, Leijten EFA, Bekker CPJ, Eeftink E, Giovannone B, Nordkamp MO, van der Wal M, Thijs JL, Vastert SJ, van Wijk F, Radstake TRDJ and van Loosdregt J. 2021. ZFP36 family members regulate the pro-inflammatory features of psoriatic dermal fibroblasts. *Journal of Investigative Dermatology*. 142(2):402-413
4. Bick MJ, Carroll JW, Gao G, Goff SP, Rice CM, MacDonald MR. 2003. Expression of the zinc-finger antiviral protein inhibits alphavirus replication. *Journal of Virology*. 77(21):11555-62. doi: 10.1128/jvi.77.21.11555-11562.2003. PMID: 14557641; PMCID: PMC229374.

5. Blackshear, PJ. 2002. Tristetraprolin and other CCCH tandem zinc-finger proteins in the regulation of mRNA turnover. *Biochemical Society Transactions*. 30(6): 945-52.
6. Cassandri M, Smirnov A, Novelli F, Pitolli C, Agostini M, Malewicz M, Melino G, Raschellà G. 2017. Zinc-finger proteins in health and disease. *Cell Death Discovery*. 13;3:17071. doi: 10.1038/cddiscovery.2017.71. PMID: 29152378; PMCID: PMC5683310.
7. Cassandri MA, Smirnov F, Novelli C, Pitoll M, Agostini M, Malewicz GG., Raschellà 2017. Zinc-finger proteins in health and disease. *Cell Death Discovery*. 3(1): 17071.
8. Chen SC., Jeng KS, and Lai MMC. 2017. Zinc Finger-Containing Cellular Transcription Corepressor ZBTB25 Promotes Influenza Virus RNA Transcription and Is a Target for Zinc Ejector Drugs. *Journal of Virology*. 91(20).
9. Chen XF, Wu J, Zhang YD, Zhang CX, Chen XT, Sun JH, Chen TX. 2018. Role of Zc3h12a in enhanced IL-6 production by newborn mononuclear cells in response to lipopolysaccharide. *Pediatrics & Neonatology*. 59(3):288-295. doi: 10.1016/j.pedneo.2017.09.006. Epub 2017 Sep 21. PMID: 29054363.
10. Chiu H, Chiu HP, Yu HP, Lin LH, Chen ZP, Lin YL, Lin RJ. 2022. Zinc Finger Protein ZFP36L1 Inhibits Flavivirus Infection by both 5'-3' XRN1 and 3'-5' RNA-Exosome RNA Decay Pathways. *Journal of Virology*. 12;96(1):e0166521. doi: 10.1128/JVI.01665-21. Epub 2021 Oct 13. PMID: 34643435; PMCID: PMC8754223.
11. Fehr AR and Perlman S. 2015. Coronaviruses: an overview of their replication and pathogenesis. *Methods in Molecular Biology*, 1282: 1-23.
12. Fu M and Blackshear PJ. 2017. RNA-binding proteins in immune regulation: a focus on CCCH zinc finger proteins. *Nature Reviews Immunology*, 17(2): 130-143.
13. Gao G., Guo X, and Goff SP. 2002. Inhibition of retroviral RNA production by ZAP, a CCCH-type zinc finger protein. *Science*, 297(5587): p. 1703-6.
14. Gonzalez-Perez, AC., Stempel M, Wyler E, Urban C, Piras A, Hennig T, Ganskih S, Wei Y, Heim A, Landthaler M, Pichlmair A, Dölken L, Munschauer M, Erhard F and Brinkmann MM. 2021. The Zinc Finger Antiviral Protein ZAP Restricts Human Cytomegalovirus and Selectively Binds and Destabilizes Viral UL4/UL5 Transcripts. *mBio*. 12(3).
15. Guo X, Carroll JW, Macdonald MR, Goff SP and Gao G. 2004. The zinc finger antiviral protein directly binds to specific viral mRNAs through the CCCH zinc finger motifs. *Journal of Virology*. 78(23): 12781-12787.
16. Hajikhezri Z, Darweesh M, Akusjärvi G, Punga T. 2020. Role of CCCH-Type Zinc Finger Proteins in Human Adenovirus Infections. *Viruses*. 18;12(11):1322. doi: 10.3390/v12111322. PMID: 33217981; PMCID: PMC7698620.
17. Haneklaus M, O'Neil JD, Clark AR, Masters SL, O'Neill LAJ. 2017. The RNA-binding protein Tristetraprolin (TTP) is a critical negative regulator of the NLRP3 inflammasome. *Journal of Biological Chemistry*. 28;292(17):6869-6881. doi: 10.1074/jbc.M116.772947. Epub 2017 Mar 16. PMID: 28302726; PMCID: PMC5409458.
18. Kontoyiannis DL. 2018. An RNA checkpoint that keeps immunological memory at bay. *Nature Immunology*. 19(8): 795-797.
19. Kozaki T, Takahama M, Misawa T, Matsuura Y, Akira S, Saitoh T. 2015. Role of zinc-finger anti-viral protein in host defense against Sindbis virus. *International*

- Immunology*. 27(7):357-64. doi: 10.1093/intimm/dxv010. Epub 2015 Mar 10. PMID: 25758257; PMCID: PMC4565983.
20. Krishna SS., Majumdar I., and Grishin NV. 2003. Structural classification of zinc fingers: survey and summary. *Nucleic Acids Research* 31(2): 532-50.
21. Lai WS, Carballo E, Thorn JM, Kennington EA, Blackshear PJ. 2000. Interactions of CCCH zinc finger proteins with mRNA. Binding of tristetraprolin-related zinc finger proteins to Au-rich elements and destabilization of mRNA. *Journal of Biological Chemistry* 9;275(23):17827-37. doi: 10.1074/jbc.M001696200. PMID: 10751406.
22. Lai WS, Kennington EA and Blackshear PJ. 2003. Tristetraprolin and its family members can promote the cell-free deadenylation of AU-rich element-containing mRNAs by poly(A) ribonuclease. *Molecular and Cellular Biology*, 23(11): p. 3798-812.
23. Lai WS, Stumpo DJ, Wells ML, Gruzdev A, Hicks SN, Nicholson CO, Yang Z, Faccio R, Webster MW, Passmore LA, Blackshear PJ. 2019. Importance of the Conserved Carboxyl-Terminal CNOT1 Binding Domain to Tristetraprolin Activity *In Vivo*. *Molecular and Cellular Biology*. 39(13):e00029-19. doi: 10.1128/MCB.00029-19. PMID: 31036567; PMCID: PMC6580703.
24. Lal S, Ullah BA, and Syed S. 2020. The Role of Zinc-Finger Antiviral Proteins in Immunity against Viruses. *Molecular Genetics, Microbiology and Virology*, 35(2): 78-84.
25. Li M, Yan K, Wei L, Yang J, Lu C, Xiong F, Zheng C, Xu W. 2015. Zinc finger antiviral protein inhibits coxsackievirus B3 virus replication and protects against viral myocarditis. *Antiviral Research*. 123:50-61. doi: 10.1016/j.antiviral.2015.09.001. Epub 2015 Sep 2. PMID: 26343012.
26. Lin RJ, Huang CH, Liu PC, Lin IC, Huang YL, Chen AY, Chiu HP, Shih SR, Lin LH, Lien SP, Yen LC, Liao CL. 2020. Zinc finger protein ZFP36L1 inhibits influenza A virus through translational repression by targeting HA, M and NS RNA transcripts. *Nucleic Acids Research*. 27;48(13):7371-7384. doi: 10.1093/nar/gkaa458. PMID: 32556261; PMCID: PMC7367194.
27. Lv L, Qin T, Huang Q, Jiang H, Chen F, Long F, Ren L, Liu J, Xie Y, Zeng M. 2021. Targeting Tristetraprolin Expression or Functional Activity Regulates Inflammatory Response Induced by MSU Crystals. *Frontiers in Immunology*. 16;12:675534. doi: 10.3389/fimmu.2021.675534. PMID: 34335573; PMCID: PMC8322984.
28. Lykke-Andersen J. and Wagner E. Recruitment and activation of mRNA decay enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-1. *Genes & Development*, 19(3): p. 351-61.
29. Maeda K. and Akira S. 2017. Regulation of mRNA stability by CCCH-type zinc-finger proteins in immune cells. *International Immunology*, 29(4): 149-155.
30. Matsushita K, Takeuchi O, Standley DM, Kumagai Y, Kawagoe T, Miyake T, Satoh T, Kato H, Tsujimura T, Nakamura H, Akira S. 2009. Zc3h12a is an RNase essential for controlling immune responses by regulating mRNA decay. *Nature*. 30;458(7242):1185-90. doi: 10.1038/nature07924. Epub 2009 Mar 25. PMID: 19322177.
31. Meagher J., Takata M, Gonçalves-Carneiro D, Keane SC, Rebendenne A, Ong H, Orr VK, MacDonald MR, Stuckey JA, Bieniasz PD and Smith JL. 2019. Structure of the zinc-finger antiviral protein in complex with RNA reveals a mechanism for

- selective targeting of CG-rich viral sequences. *Proceedings of the National Academy of Sciences*. 116(48): 24303-24309.
32. Mino T, Murakawa Y, Fukao A, Vandenbon A, Wessels HH, Ori D, Uehata T, Tartey S, Akira S, Suzuki Y, Vinuesa CG, Ohler U, Standley DM, Landthaler M, Fujiwara T, Takeuchi O. 2015. Regnase-1 and Roquin Regulate a Common Element in Inflammatory mRNAs by Spatiotemporally Distinct Mechanisms. *Cell*. 21;161(5):1058-1073. doi: 10.1016/j.cell.2015.04.029. PMID: 26000482.
33. Müller S, Möller P, Bick MJ, Wurr S, Becker S, Günther S and Kümmerer BM. 2007. Inhibition of filovirus replication by the zinc finger antiviral protein. *Journal of virology*. 81(5): 2391-2400.
34. Musah RA 2004, The HIV-1 nucleocapsid zinc finger protein as a target of antiretroviral therapy. *Current Topics in Medicinal Chemistry*, 4 (15): 1605-22.
35. Nchioua R, Kmiec D, Müller JA, Conzelmann C, Groß R, Swanson CM, Neil SJD, Stenger S, Sauter D, Münch J, Sparrer KML and Kirchhoff F. 2020. SARS-CoV-2 Is Restricted by Zinc Finger Antiviral Protein despite Preadaptation to the Low-CpG Environment in Humans. *mBio*. 11(5).
36. Reed LJ and Muench J. 1938. A simple method of estimating fifty percent endpoints.. *American Journal of Epidemiology*.27(3): 493-497.
37. Schito ML, Soloff AC, Slovit D, Trichel A, Inman JK, Appella E, Turpin JA and Barratt-Boyes SM. 2006. Preclinical evaluation of a zinc finger inhibitor targeting lentivirus nucleocapsid protein in SIV-infected monkeys. *Current HIV Research*. 4(3): 379-386.
38. Schneider C., Rasband W. and Eliceiri K. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9, 671–675 (2012). <https://doi.org/10.1038/nmeth.2089>.
39. Scozzafava A., Owa T, Mastrolorenzo A and Supuran CT . 2003. Anticancer and antiviral sulfonamides. *Current Medicinal Chemistry*. 10(11): 925-953.
40. Shrestha A, Pun NT and Park PH. 2018.ZFP36L1 and AUF1 Induction Contribute to the Suppression of Inflammatory Mediators Expression by Globular Adiponectin via Autophagy Induction in Macrophages. *Biomolecules & Therapeutics*.26(5): 446-457.
41. Strober W. 2015. Trypan Blue Exclusion Test of Cell Viability. *Current Protocols in Immunology*. 2;111:A3.B.1-A3.B.3. doi: 10.1002/0471142735.ima03bs111. PMID: 26529666; PMCID: PMC6716531
42. Stumpo DJ., Lai WS, and Blackshear PJ. 2010. Inflammation: cytokines and RNA-based regulation. *Wiley Interdisciplinary Reviews:RNA*, 1(1): 60-80.
43. Suk FM, Chang CC, Lin RJ, Lin SY, Liu SC, Jau CF, Liang YC. 2018. ZFP36L1 and ZFP36L2 inhibit cell proliferation in a cyclin D-dependent and p53-independent manner. *Scientific Reports*. 9;8(1):2742. doi: 10.1038/s41598-018-21160-z. Erratum in: *Sci Rep*. 2019 Nov 20;9(1):17457. PMID: 29426877; PMCID: PMC5807420.
44. Takata MA, Gonçalves-Carneiro D, Zang TM, Soll SJ, York A, Blanco-Melo D and Bieniasz PD. 2017. CG dinucleotide suppression enables antiviral defence targeting non-self RNA. *Nature*. 550(7674): 124-127.Tu Y, Wu X, Yu F, Dang J, Wang J, Wei Y, Cai Z, Zhou Z, Liao W, Li L, Zhang Y. 2019. Tristetraprolin specifically regulates the expression and alternative splicing of immune response genes in HeLa cells. *BMC Immunology*. 2;20(1):13. doi: 10.1186/s12865-019-0292-1. PMID: 31046669; PMCID: PMC6498542.




45. Tang Q, Wang X, and Gao G. 2017. The Short Form of the Zinc Finger Antiviral Protein Inhibits Influenza A Virus Protein Expression and Is Antagonized by the Virus-Encoded NS1. *Journal of Virology*, 91(2): p. e01909-16.
46. Uehata T. and Akira S. 2013. mRNA degradation by the endoribonuclease Regnase-1/ZC3H12a/MCPIP-1. *Biochimica et Biophysica Acta*, 1829 (6-7): p. 708-13.
47. Vilas CK, Emery LE, Denchi EL, Miller KM. 2018. Caught with One's Zinc Fingers in the Genome Integrity Cookie Jar. *Trend in Genetics*. 34(4):313-325. doi: 10.1016/j.tig.2017.12.011. Epub 2018 Jan 19. PMID: 29370947; PMCID: PMC5878116.
48. Villa N, Do A, Hershey JW, Fraser CS. 2013. Human eukaryotic initiation factor 4G (eIF4G) protein binds to eIF3c, -d, and -e to promote mRNA recruitment to the ribosome. *Journal of Biological Chemistry*. 15;288(46):32932-40. doi: 10.1074/jbc.M113.517011. Epub 2013 Oct 3. PMID: 24092755; PMCID: PMC3829144.
49. Wang G. and Zheng C. 2021. Zinc finger proteins in the host-virus interplay: multifaceted functions based on their nucleic acid-binding property. *FEMS Microbiology Reviews*. 45(3).
50. Wang KT, Wang HH, Wu YY, Su YL, Chiang PY, Lin NY, Wang SC, Chang GD, Chang CJ. 2015. Functional regulation of Zfp361l and Zfp3612 in response to lipopolysaccharide in mouse RAW264.7 macrophages. *Journal of Inflammation*. 16;12:42. doi: 10.1186/s12950-015-0088-x. PMID: 26180518; PMCID: PMC4502546.
51. Wawro M, Kochan J, and Kasza A, The perplexities of the ZC3H12A self-mRNA regulation. *Acta Biochimica Polonica*, 63(3): p. 411-5.
52. Wu HY, Ke TY, Liao WY, Chang NY. 2013. Regulation of coronaviral poly(A) tail length during infection. *PLoS One*. 29;8(7):e70548. doi: 10.1371/journal.pone.0070548. PMID: 23923003; PMCID: PMC3726627.
53. Zhang B, Goraya MU, Chen N, Xu L, Hong Y, Zhu M and Chen JL .2020. Zinc Finger CCCH-Type Antiviral Protein 1 Restricts the Viral Replication by Positively Regulating Type I Interferon Response. *Frontiers in microbiology*. 11: 1912-1912.
54. Zhao Y, Song Z, Bai J, Liu X, Nauwynck H and Jiang P. 2019. ZAP, a CCCH-Type Zinc Finger Protein, Inhibits Porcine Reproductive and Respiratory Syndrome Virus Replication and Interacts with Viral Nsp9. *Journal of Virology*. 93(10).
55. Zhu M, Ma X, Cui X, Zhou J, Li C, Huang L, Shang Y, Cheng Z. 2017. Inhibition of avian tumor virus replication by CCCH-type zinc finger antiviral protein. *Oncotarget*. 19;8(35):58865-58871. doi: 10.18632/oncotarget.19378. PMID: 28938603; PMCID: PMC5601699.
56. Zhu M., Zhou J, Liang Y, Nair V, Yao Y and Cheng Z. 2020. CCCH-type zinc finger antiviral protein mediates antiviral immune response by activating T cells. *Journal of Leukocyte Biology*. 107(2): 299-307.
57. Zhu Y, Chen G, Lv F, Wang X, Ji X, Xu Y, Sun J, Wu L, Zheng YT, Gao G. 2011. Zinc-finger antiviral protein inhibits HIV-1 infection by selectively targeting multiply spliced viral mRNAs for degradation. *Proceedings of the National Academy of Sciences of the United States of America*. 20;108(38):15834-9. doi: 10.1073/pnas.1101676108. Epub 2011 Aug 29. PMID: 21876179; PMCID: PMC3179061.

58. Zhu Y, Wang X, Goff SP, Gao G. 2012. Translational repression precedes and is required for ZAP-mediated mRNA decay. *The EMBO Journal*. 5;31(21):4236-46. doi: 10.1038/emboj.2012.271. Epub 2012 Sep 28. PMID: 23023399; PMCID: PMC3492732.

# Figure 1

## Overexpression and knockdown of ZFP36L1 in HCT-8 cells


Wild type HCT8 wells (A), ZFP36L1 overexpressed HCT-8 cells with GFP marker (B), and ZFP36L1 knockdown HCT-8 cells with mCherry marker (C). Overexpression and knockdown of ZFP36L1 were performed by lentivirus transduction.



# Figure 2

Western blot for confirming ZFP36L1 overexpression and knockdown in HCT-8 cells


Cell lysate for wild type HCT-8 cell (A), ZFP36L1 overexpressed (B) and ZFP36L1 knockdown (C) were separated with 12% resolving SDS PAGE gel and transferred to PVDF membrane. Proteins on the membrane were detected with an anti-ZFP36L1 antibody and anti-  $\beta$ actin antibody with HRP-conjugated secondary antibodies.



# Figure 3

Relative quantification of ZFP36L1 expression in HCT-8 cell following its overexpression and knockdown


Cell lysate for wild-type HCT-8 cell, ZFP36L1 overexpressed and ZFP36L1 knockdown was analyzed for ZFP36L1 and  $\beta$  actin using western blot. Band intensity for ZFP36L1 proteins was normalized with  $\beta$  actin using ImageJ software. A significant difference in ZFP36L1 expression in ZFP36L1 overexpressed and knockdown cells compared to wild-type cells was estimated using a paired T-test. Asterisks are showing significant differences in ZFP36L1 expression.



# Figure 4

## Effect of ZFP36L1 on HCT-8 cells viability


The effect of ZFP36L1 overexpression and its knockdown on cell viability was measured by trypan blue exclusion assay. Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown cells were individually seeded in 6 well plates. After 96 hours post-seeding, cells were detached and stained with 0.4% trypan blue to determine the percent viability. Changes in cell viability following ZFP36L1 overexpression or its knockdown compared to wild-type cells was estimated by paired T-test.



# Figure 5

Human coronavirus-OC43 titer in HCT-8 cells.


Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were infected individually with HCoV-OC43 with 0.1 MOI. Supernatant from these cells was collected at 24 hours, 48 hours, 72 hours, and 96 hours p.i. and analyzed for virus titer. Asterisks are showing significant differences in virus titer.



# Figure 6

Effect of ZFP36L1 expression on Human coronavirus-OC43 induced cytopathic effect in HCT-8 cells

Wild type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were infected individually with HCoV-OC43 with 0.1 MOI. Cytopathic effect was observed at 72 hours p.i. at 40X magnification





# Figure 7

## Effect of ZFP36L1 expression on Human coronavirus-OC43 replication

Wild-type, ZFP36L1 overexpressed and ZFP36L1 knockdown HCT-8 cells were infected individually with HCoV-OC43 with 0.1 MOI. Viral RNA was isolated from infected cells at 72 and 96 hours p.i. Isolated RNA was quantified using qPCR (for viral nucleocapsid). Fold change in nucleocapsid RNA in ZFP36L1 overexpressed and knockdown cells as compared to wild-type HCT-8 cells were estimated using paired T-test. Asterisks are showing significant differences in viral RNA.

