

The first record of chigutisaurid amphibian from the Late Triassic Tiki Formation and the probable Carnian pluvial episode in central India

Sanjukta Chakravorti ^{Corresp., 1}, Dhurjati Prasad Sengupta ¹

¹ Geological Studies Unit, Indian Statistical Institute, Kolkata, West Bengal, India

Corresponding Author: Sanjukta Chakravorti
Email address: chirpymoni2009@gmail.com

A new, partially preserved skull of chigutisaurid amphibian (*temnospondyli*) has been reported for the first time from the Late Triassic Tiki Formation of India. Chigutisaurids are now known to occur in Australia's Early and Late Triassic, the Late Triassic in India, Argentina, and Brazil, the Jurassic of South Africa and Australia, and the Cretaceous of Australia. In India, the first appearance of chigutisaurids marks the Carnian – middle Carnian / Norian Boundary. This work also attempts to correlate, again for the first time, the advent of chigutisaurids and the occurrence of Carnian Pluvial Episodes (CPE) in the Late Triassic Maleri and Tiki Formation of Central India. The new specimen belongs to the genus *Compsocerops* prevalent in the Late Triassic Maleri Formation occurring 700 km south. However, the chigutisaurid specimen recovered from the Tiki Formation is a new species when compared to that of the Maleri Formation. It has the presence of an inward curved process of the quadratojugal as opposed to the straight downward trending process of the quadratojugal, the presence of vomerine foramen, shorter and wider interpterygoid vacuities, wider subtemporal vacuities, the base of the interpterygoid vacuities at the same level with the base of the subtemporal vacuity, it proves that the Tiki Formation is coeval with the Lower Maleri Formation and a part of Upper Maleri.

The first record of chigutisaurid amphibian from the Late Triassic Tiki Formation and the probable Carnian Pluvial Episode in Central India.

Sanjukta Chakravorti ^{1*}, Dhurjati Prasad Sengupta¹

¹ Geological Studies Unit, Indian Statistical Institute, 203, B.T.Road, Kolkata- 700108, West-Bengal, India.

Corresponding Author

Sanjukta Chakravorti^{1*}

*Email address: chirpymoni2009@gmail.com

Abstract

A new, partially preserved skull of chigutisaurid amphibian (temnospondyli) has been reported for the first time from the Late Triassic Tiki Formation of India. Chigutisaurids are now known to occur in Australia's Early and Late Triassic, the Late Triassic in India, Argentina, and Brazil, the Jurassic of South Africa and Australia, and the Cretaceous of Australia. In India, the first appearance of chigutisaurids marks the Carnian – middle Carnian / Norian Boundary. This work also attempts to correlate, again for the first time, the advent of chigutisaurids and the occurrence of Carnian Pluvial Episodes (CPE) in the Late Triassic Maleri and Tiki Formation of Central India. The new specimen belongs to the genus *Compsocerops* prevalent in the Late Triassic Maleri Formation occurring 700 km south. However, the chigutisaurid specimen recovered from the Tiki Formation is a new species when compared to that of the Maleri Formation. It has the presence of an inward curved process of the quadratojugal as opposed to the straight downward trending process of the quadratojugal, the presence of vomerine foramen, shorter and wider interpterygoid vacuities, wider subtemporal vacuities, the base of the interpterygoid vacuities at the same level

with the base of the subtemporal vacuity, it proves that the Tiki Formation is coeval with the Lower Maleri Formation and a part of Upper Maleri.

Introduction

Temnospondyls are a very diverse and widespread group of extinct amphibians that thrived from the Carboniferous to the Cretaceous reaching their peak in the Triassic ([Konietzko-Meier et al. 2018](#); [Konietzko-Meier et al. 2019](#)). Capable of flourishing in both land and water in a variety of ecological niches, the fossils of these extinct amphibians are found in almost all places from Antarctica in the South to Greenland in the North, India being no exception ([Schoch et al. 2014](#)). The temnospondyl fossils in India are found in various Gondwana deposits throughout the country ([Chakravorti & Sengupta 2019](#); [Chowdhury 1965](#); [Sengupta 1995](#); [Sengupta 2002](#); [Tripathi 1969](#)). Chigutisaurids are by far one of the most important temnospondyl amphibians having the longest temporal range ([Marsicano 1999](#)) ranging from the Early Triassic to the Cretaceous ([Dias-da-Silva et al. 2012](#)). However, the majority of them originated in the Upper Carnian to Lower Norian of the Late Triassic Period. The family Chigutisauridae is much less diversified than the other temnospondyl families. Falling within the superfamily Brachyopoidea which comprises parabolic, brevirostrine skulled temnospondyls, Chigutisauridae forms a single monophyletic family in the phylogenetic position ([Warren & Marsicano 2000](#)). The spatial and temporal distribution of the chigutisauridae is schematically represented in Table -1 along with their general habitat ([Bandyopadhyay & Ray 2020b](#); [Bonaparte 1975](#); [Cabrera 1944](#); [DIAS-da-SILVA et al. 2012](#); [Pledge 2013](#); [Rusconi 1949](#); [Rusconi 1951](#); [Sengupta 1995](#); [Warren 1981](#); [Warren & Marsicano 2000](#)). The earliest origin of the family chigutisauridae is *Keratobrachyops australis* in the Early Triassic of Australia ([Warren 1981](#)) with the latest being *Koolasuchus cleelandi* ([Warren et al. 1997](#)) in the Cretaceous of Australia as well.

So far, the origin and diversification of chigutisaurids remain restricted only to the Gondwana countries.

The Gondwana successions of India are exposed in four discrete basins coinciding with some of the major river valleys throughout the Indian subcontinent ([Pascoe 1973](#); [Robinson 1970](#); [Veevers & Tewari 1995](#)). Of these, the Late Triassic Maleri and the Tiki Formations of the Pranhita-Godavari Valley Basin and the Son Valley Basin respectively are long known to be coeval ([Chatterjee 1974](#); [Chatterjee & Roy-Chowdhury 1974](#); [Mukherjee & Ray 2014](#); [Robinson 1970](#)). Both Formations are known for the metoposaurid *Panthesaurus maleriensis* ([Arche & Lopez-Gomez 2014](#); [Chakravorti & Sengupta 2019](#); [Sengupta 2002](#)). In the Late Triassic Maleri Formation *P. maleriensis* is thought to be restricted within the Carnian and the chigutisaurids appear in the mid-Carnian to early Norian ([Chakravorti & Sengupta 2019](#); [Sengupta 1995](#)). Though a considerable amount of work has been done on the microvertebrates ([Bhat 2017](#); [Hussain 2018](#); [Ray et al. 2016a](#)), rhynchosaurs ([Mukherjee & Ray 2012](#); [Mukherjee & Ray 2014](#)) and phytosaurs ([Datta et al. 2021a](#); [Datta et al. 2019a](#); [Datta et al. 2021b](#)) of the Tiki Formation; no comprehensive work has been done in the last decade on its temnospondyl faunal contents. [Chakravorti & Sengupta \(2019\)](#) in their taxonomic revision of the Indian metoposaurids, included the metoposaurids of the Tiki Formation and grouped them into a new genus *Panthesaurus maleriensis* based on morphometric and phylogenetic approaches. However, the biostratigraphic implications of the Tiki Formation based on its temnospondyl contents have not been attempted so far, though, the taphonomic aspects of metoposaurids from the Tiki Formation have been discussed by [Rakshit & Ray \(2020\)](#). Also, to date, no chigutisaurid remains were reported from the Late Triassic Tiki Formation though the same is widely prevalent in the Late Triassic Maleri Formation ([Sengupta 1995](#)). Therefore, the finding of a chigutisaurid amphibian from the Late Triassic Tiki Formation is very important in the context of correlating

the Late Triassic Maleri and Tiki Formation, India, and their position concerning global biostratigraphic correlation. This paper will subsequently highlight a brief geological setting of the Tiki Formation followed by the taxonomic status of chigutisaurids from the Tiki Formation and subsequently its role in demarcating the Carnian Pluvial Episode in India.

The Carnian Pluvial Episode – A Global Climatic Consequence

The Carnian Pluvial Episode (CPE) can be defined as a geologically short-lived (234-232 Million Years Ago) monsoonal period of extreme rainfall that brought about significant changes in several depositional environments ([Arche & Lopez-Gomez 2014](#); [Dal Corso et al. 2015](#); [Furin et al. 2006](#); [Schlager & Schöllnberger 1974](#); [Simms & Ruffell 1990](#)). The Carnian Pluvial Episode (CPE) was a global phenomenon. Geochemical data suggest that global warming involved environmental and biotic changes. Radioisotopic ages coupled with biostratigraphic correlation suggest a possible link to the eruption of the Wrangellia Large Igneous Province (LIP) ([Dal Corso et al. 2020](#)). CPE was a significant (but previously neglected) time of extinction linked to the Carnian explosive diversification of many key modern groups of plants and animals ([Dal Corso et al. 2020](#)). The CPE marks a distinct change in the hydrological cycle during which the climate shifted from arid to humid conditions and back again to arid conditions ([Bernardi et al. 2018](#)). It is represented by remarkable enhancement of the hydrological cycle demarcated by four episodes of increased rainfall indicated by diverse sedimentary and paleontological data, repeated Carbon Cycle perturbations, evidenced by sharp negative C-isotope excursions, coincided with global environmental changes and climate warming all of which suggest a cause-and-effect relationship ([Dal Corso et al. 2015](#)).

The Carnian is the earliest part of the Late Triassic and its base or lower boundary is dated at approximately 237 million years based on U-Pb radiometric dating of a single crystal zircon from a tuff layer within a section having strong biostratigraphic constraints ([Dal Corso et al. 2015](#); [Dal Corso et al. 2012](#); [Maron et al. 2019](#)). The upper boundary of the Carnian is constrained at approximately 227 million years based on magnetostratigraphic correlations between the marine successions of Tethys and the astrochronological time scale of the continental Newark Basin ([Kent et al. 2017](#)). The Carnian is subdivided into Julian (Early Carnian) and Tuvanian (Late Carnian) substages. The Julian – Tuvanian boundary occurs at approximately 233 million years ([Dal Corso et al. 2015](#); [Kent et al. 2017](#)). The beginning of the onset of CPE is well defined from ammonoid, conodont, and sporomorph biostratigraphic dating and is synchronous in several geological settings. It coincides with the first appearance of the ammonoid genus *Austrotrachyceras* in the Julian ([Dal Corso et al. 2020](#); [Dal Corso et al. 2012](#); [Roghi et al. 2010](#); [Simms & Ruffell 1990](#); [Sun et al. 2016](#)). However, the upper boundary or the end of CPE is poorly defined in most locations. It is usually placed at the base or within the Tuvanian 2 based on sedimentological (e.g., end of terrigenous sediment supply) and chemostratigraphic (last C-isotope excursion) evidence ([Dal Corso et al. 2020](#); [Dal Corso et al. 2015](#); [Dal Corso et al. 2018](#)). The total duration of this pluvial episode is variable. Cyclostratigraphy of marine successions of the South China Block and continental successions of the Wessex Basin (United Kingdom) gives a duration of the CPE of approximately 1.2 ma but this is variable and longer than 1.6-1.7 million years as indicated by integrated stratigraphy (biostratigraphy and magnetostratigraphy).

Temnospondyl amphibians in the Carnian Pluvial Episode

The CPE facilitated the Dinosaur Diversification Event (DDE) ([Bernardi et al. 2018](#)). However, the role of CPE on the temnospondyls has not much been discussed barring a few papers ([Buffa et al.](#)

2019; Fortuny et al. 2019; Gee & Jasinski 2021a; Lucas 2020a). The amphibious temnospondyls living both on land and water were the most sensitive to the changes in climate. Two dominant groups of temnospondyls, in this context, were the metoposaurids and the chigutisaurids. According to Fortuny et al. (2019) the gigantism of the metoposaurids might have been linked to the Carnian Pluvial Episode. Buffa et al. (2019) also stated that the diversification of the metoposaurids might have been linked to the CPE and the post-CPE aridification led to the extinction of the metoposaurids during the Rhaetian. Gee & Jasinski (2021a) have also commented on the fact that the physiological variation of the metoposauridae and their palaeoclimatic range also corroborates a palaeo-environmental barrier. Finally, Lucas (2020b) concluded that climate change that occurred during CPE played an important part in the metoposaurid evolution. According to Lucas (2020b), Metoposaurids appeared during the CPE, attained their highest diversity and cosmopolitan distribution during this time and had reduced diversity and showed endemism in the post-CPE climate.

Thus, the presence of *Compsocerops* in both Maleri and Tiki Formation enhances the scope to discuss the palaeoenvironment of these two Late Triassic basins in India and to compare the possible reason for faunal turnover from Carnian to Norian concerning the amphibious temnospondyls (Sengupta 1995).

Geological setting of the Tiki Formation and its comparison to the Maleri Formation

The Tiki Formation named after the small village of Tiki in the district of Shahdol, Madhya Pradesh has been an interest to scientists for decades. Reports on the geology and palaeontology of the Tiki Formation date back to as early as 1877 when Hughes (1877) noticed reptilian fossils near this village. Cotter (1917) noticed several other such fossils and finally Fox (1931) formally

designated the area as the “Tiki stage”. [Aiyengar \(1937\)](#) first divided the “Tiki stage” into lithostratigraphic units viz. the lower unit being fossiliferous and composed dominantly of red and green mudstones, proportionately lesser number of sandstones and mud-galls while the upper unit is composed of ferruginous sandstones and shales. [Robinson \(1970\)](#) in her memoir kept the Tiki Formation to be coeval with the Late Triassic Maleri Formation. However, [Dutta & Ghosh \(1993\)](#) did not recognize the separate entity of the Tiki Formation and placed Tiki rocks in the upper part of the Pali Formation forming the “Pali-Tiki Formation”. [Roychowdhury et al. \(1975\)](#) based on the megaflorea assemblage noted the age of the Nidpur beds is Anisian and separated the upper part of the Tiki Formation to be Carnian – Rhaetian in age. [Maheshwari et al. \(1976\)](#) separated the Tiki Formation to be a separate entity (including the Nidpur beds) and based on the mega flora and faunal assemblages suggested the age of the Tiki (including Nidpur beds) Formation to be ranging from Anisian to Norian with a possible extension to Rhaetian. [Mukherjee et al. \(2012\)](#) revised the stratigraphy of the Rewa Basin and put the Tiki Formation with the coeval Carnian Lower Maleri Formation. [Ray et al. \(2016b\)](#) in the study of vertebrate faunal assemblage of the Tiki formation also suggested Tiki Formation be of Carnian in age but they narrowed the range to Otischalkian to early Adamanian. The common conclusions of all this literature are that the Late Triassic Tiki Formation is dominantly Carnian and its fauna can be correlated with the Lower Maleri fauna. So far, no evidence of a Norian age was assigned to any part of the Tiki Formation. As stated earlier, based on the faunal pieces of evidence and correlating it with the Late Triassic Maleri Formation of the Pranhita – Godavari valley a Carnian age was assigned to the Tiki Formation ([Dutta & Ghosh 1993](#); [Kutty et al. 1987](#); [Sengupta 1992](#)). Henceforth, through the Decades, the Tiki formation was considered to be coeval with the Carnian Maleri Formation ([Mukhopadhyay et al. 2010](#); [Sengupta 1992](#); [Veevers & Tewari 1995](#)). Only recently, [Datta et al. \(2021b\)](#) while describing a new phytosaur

from the Tiki Formation, commented that the age of the Tiki Formation may range from Carnian to Early/Middle Norian.

To date, the faunal assemblage of the Tiki Formation includes fishes belonging to the family Ceratodontidae, Hybodontidae and new undescribed forms of Xenacanthidae ([Ray et al. 2016a](#)), temnospondyl amphibians belonging to Metoposauridae; reptilian belonging to families Rhynchosauridae, Rauisuchidae, Acrodonta, basal Saurischia, Dromatheridae, and Traversodontidae. Mammaliaformes are also reported from the Tiki Formation (Tables 2a and 2b) ([Bandyopadhyay & Ray 2020b](#); [Ray et al. 2016a](#)).

The appearance of chigutisaurids in India is noted with the demise of the metoposaurs, rhynchosaurs, and primitive phytosaurs. Prosauropods, large in size also appeared during that time ([Novas et al. 2010](#)). These faunal turnovers were thought to demarcate India's Carnian – Norian boundary ([Datta et al. 2021b](#)). However, recent signatures ([Dal Corso et al. 2015](#)) of the pluvial event and its role in extinction might shift this boundary to Carnian – mid-Carnian/Norian.

The Maleri Formation starts with a 250-meter-thick mudstone ([Dasgupta et al. 2017](#); [Kutty & Sengupta 1989](#)). At the top of the mudstone, a sandy zone initiates the sand–mud alternations of Upper Maleri ([Kutty & Sengupta 1989](#)). This sandy zone contains a maximum number of rhynchosaur fossils, abundant metoposaurids, and unionids. The chigutisaurids in Maleri appear just above this sandy zone ([Sengupta 1995](#)) and no rhynchosaurs or metoposaurids are known from that level (or above that). The occurrence of chigutisaurids in upper part of Tiki Formation is also restricted within a sandy zone which do not contain metoposaurids or rhynchosaurs. Unionids are also present there but in lesser abundance than Maleri. Phytosaur teeth are also present. This sandy horizon noticed in Maleri and Tiki has been stratigraphically placed below the Carnian – mid-Carnian/ Norian boundary and may indicate the Carnian Pluvial Episodes (CPE) in India.

Materials used with an overview of their preservation and methods

The new skull material along with the clavicle (ISI A 202) excavated by the authors and the specimen RH01/Pal/CHQ/Tiki/15 previously described as metoposaurid ([Kumar & Sharma 2019](#)) has been studied in detail and referred to in this paper. The map of the temnospondyl-bearing localities of the Tiki Formation has been modified here with faunal boundaries (hypothetical faunal boundary demarcated in a red dotted line, after [Mukherjee & Ray \(2012\)](#)) (Fig: 1). The temnospondyl bearing (metoposaurid and chigutisaurid) localities of the Maleri Formation have also been extensively mapped and modified (after [Dasgupta et al. \(2017\)](#); [Kutty & Sengupta \(1989\)](#)) (Fig: 2). Some distinct sections have been logged in the Tiki Formation and has been compared with the existing and modified logs of the Late Triassic Tiki and Maleri Formation.

Preservation of specimen ISI A 202 - The skull along with a fragmentary clavicle, ISI A 202, is poorly preserved (Figs: 3,4,5,6). Only the left half of the skull is preserved and the specimen is heavily eroded. Thus, the ornaments are not well observed in all the areas. The upper part of the parietal and postfrontal have coarse ridges and grooves preserved in them. The skull, its fragments and the clavicle, all have been excavated from the same point in the location and were present together with the same individual as the skull.

Preservation of specimen RH01/Pal/CHQ/Tiki/15-

Only the picture of the palate published as *Metoposaurus* in the paper published by [Kumar & Sharma \(2019\)](#)(Fig: 7) could be studied. As mentioned in the paper ([Kumar & Sharma 2019](#)) the material could not be excavated from the field. The photograph of the said publication was reproduced with permission of the journal editor and a higher resolution image was reproduced for better clarity

and study. The palate is dorsoventrally elongated and slightly sheared. The edges of the palate are not well preserved.

The new specimens with ISI numbers, ISI A 202 and the published specimen of [Kumar & Sharma \(2019\)](#) RH01/Pal/CHQ/Tiki/15 were recovered from mudrocks at a distance of about 100 meters from each other from the village of Tenduadh in the Late Triassic Tiki Formation. Thus, ISI A 202 and RH01/Pal/CHQ/Tiki/15 (previously published by [Kumar & Sharma \(2019\)](#) as metoposaurid) are now the two chigutisaurid individuals that are being reported for the first time from the Tiki Formation.

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is urn:lsid:zoobank.org:pub:1B45D1E1-9FFE-421D-8060-4174334A7EF4. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central SCIE, and CLOCKSS.

Chigutisauridae from the Tiki Formation

Systematic Palaeontology

Temnospondyli [Zittel 1887](#)

Stereospondyli [Zittel 1888](#)

Chigutisauridae [Rusconi 1951](#)

Compsocerops [Sengupta 1995](#)

Compsocerops tikiensis sp. nov. (ISI A 202/1, ISI A 202/2, ISI A 202/3, ISI A 202/4, and ISI A 202/5, RH01/Pal/CHQ/Tiki/15) (Fig: 3)

Type material: ISI A 202/1 which comprises the left half of a skull roof, ISI A 202/2 – a nearly complete clavicle and ISI A 202/3, ISI A 202/4, and ISI A 202/5 – broken parts of the skull are the holotype. The holotype materials are housed in Geological Studies Unit, Indian Statistical Institute, Kolkata, India (Fig: 3).

Paratypes: ISI A 202/2, ISI A 202/3, ISI A 202/4 and ISI A 202/5, RH01/Pal/CHQ/Tiki/15.

Referred material: A palate (RH01/Pal/CHQ/Tiki/15) previously assigned to a metoposaurid by (Kumar & Sharma 2019) has been referred (Fig: 7). The parabolic skull outline, vaulted pterygoid, shape and proportion of the interpterygoid vacuities, wide and folded palatine ramus of the pterygoid, and comparatively narrow cultriform process of the parasphenoid in RH01/Pal/CHQ/Tiki/15 indicate that it is not a metoposaurid. The cultriform process is wider than *Compsocerops cosgriffi* but narrower than any of the metoposaurids. This palate is comparatively well preserved and bears definite characteristics of a chigutisaur as it appears from the field photograph (Fig: 7). Only the photograph of the palate is available for study.

Locality: Southwest of the Village of Tenduadh (23°59'41" N; 81°25'02" E), just next to the Barakachh - Beohari Road in the district of Shahdol, Madhya Pradesh, Central India.

Etymology: The new species of chigutisaur is named after the Late Triassic Tiki Formation from where the specimen has been excavated and studied.

Diagnosis of genus: A chigutisaur temnospondyl identified as *Compsocerops* by the presence of the following combination of characters: skull outline parabolic in shape, orbits anteriorly placed, raised rim of the naris, presence of a well preserved conspicuous quadratojugal projection, presence of squamosal horn (which though broken and eroded is recognizable), and ill preserved postparietal horn (though ill preserved the presence of horns can be clearly identified), well preserved vaulted pterygoid, long and narrow cultriform process of the parasphenoid, dentigerous area restricted to the anterior portion of the palate, short and restricted palatine dentition not reaching to the middle of the choana, wide ramus of the pterygoid with a gentle fold, walls like quadrate ramus of the pterygoid, presence of postpterygoid process, typical shape of the ramus of the pterygoid and that of the subtemporal vacuities, long dorsal process of the clavicle with a distinct groove and bean-shaped cross section.

Diagnosis of species: The new species of *Compsocerops* is identified by the presence of an inward curved process of the quadratojugal as opposed to the straight downward trending process of the quadratojugal in *Compsocerops cosgriffi*, presence of vomerine foramen, shorter and wider interpterygoid vacuities, wider subtemporal vacuities, the base of the interpterygoid vacuities at the same level with the base of the subtemporal vacuity as opposed to *Compsocerops cosgriffi* (where the base of the interpterygoid vacuity is lower than the base of the subtemporal vacuity making the interpterygoid vacuities longer and slenderer in *Compsocerops cosgriffi*) and wider cultriform process of the parasphenoid.

Description and comparative anatomy

The skull roof (Fig: 3)

The dorsal part of the skull roof can only be studied in ISI A 202.

269 The skull roof is parabolic in shape with a broad and concave posterior part of the skull table. Even
 270 though the anterior part of the skull roof is mostly broken, a major portion of the left orbit is
 271 preserved. The orbit is subcircular in shape and bordered by the prefrontal, jugal, postfrontal and
 272 postorbital. The orbit is located in the anterior half of the skull and is laterally placed. The posterior
 273 and posterolateral part of the left external nares is also preserved. It can be understood from the
 274 posterior outline of the external nares that they are oval. The external nares are located very close
 275 to the anterior border of the skull roof. The posterior part of the external nares is thick and raised.
 276 This character is noted in chigutisaurids and is found in *Pelorocephalus tenax* ([Marsicano 1999](#))
 277 and *Compsocerops cosgriffi* ([Sengupta 1995](#)). The supraorbital sensory canal in the region of the
 278 naris is unusually deep around the posterolateral border of the naris which is responsible for the
 279 thick and raised posterior part of the naris in the ISI A 202. This characteristic feature has also
 280 been noted in *Compsocerops cosgriffi* ([Sengupta 1995](#)) where anteriorly the naris is flushed with
 281 the skull roof. The entire disposition of the sensory sulci is not well preserved in the specimen.
 282 Apart from the supraorbital sulcus, the presence of postorbital dermal sensory sulcus through the
 283 postfrontal can be recognized by the deep continuous canal like structure in these two bones. The
 284 infraorbital sulcus is visible in the maxilla but gradually becomes less prominent as it enters the
 285 jugal to form the jugal lateral dermal sensory sulcus. Just like other comparable chigutisaurids
 286 like *Keratobrachyops australis* ([Warren 1981](#)), *Siderops kehli* ([Warren & Hutchinson 1983](#)),
 287 *Pelorocephalus tenax* ([Marsicano 1999](#); [Rusconi 1949](#)), *Compsocerops cosgriffi* ([Sengupta 1995](#)) the
 288 lacrimal is absent in ISI A 202 and the maxilla enters the border of the external nares. However,
 289 the anterior part of the skull is fragmentary and heavily eroded. The better-preserved left side of
 290 the dorsal part of the skull roof consists of partially preserved prefrontal, postfrontal, postorbital
 291 and supratemporal. The squamosal is broken at the posterior part, and parietal, postparietal,

tabular, jugal and quadratojugals are also partially preserved. The surfaces of the bones are eroded in most places and ornamentations are poorly preserved. The parietal is comparatively large, rectangular and broken along the midline. Coarse ridges and grooves can be recognized from the anterior part of the parietal. Just as in *Compsocerops* the postparietal of ISI A 202 (Fig: 3) is much shorter in length than the parietal. The pineal foramen is not preserved in the parietal. The suture of the postparietal with the tabular is obliterated. The postparietal is broken and eroded along the midline and at its posterior part in the region of the postparietal horn. The postparietal horn is broken in ISI A 202 (Fig: 3). However, there is clear evidence that the horns exist. Postparietal horns are the most unambiguous synapomorphy of *Compsocerops*. It is preserved in *C. cosgriffi* (Sengupta 1995), *C. sp.* (Dias-da-Silva et al. 2012) and *C. tikiensis* sp. nov. These horns are not preserved in any other chigutisaur (the relevant area is not preserved in *Siderops* (Warren & Hutchinson 1983) and *Koolasuchus* (Warren et al. 1997). The tabular is most likely to be in contact with the parietal though that part is not very well preserved. The broad tabular–parietal contact is considered to be a diagnostic character of *Compsocerops* (Sengupta 1995). The tabular horn is broken. This post-quadratojugal process is robust and despite the very poor preservation of the skull in general, the posterior quadratojugal process is well preserved. The shapes and sutural patterns of the posterior left side of the skull are very similar to *Compsocerops cosgriffi*.

The Palate (Fig: 4)

Kumar and Sharma 2019, (Fig: 7) described the palate (RH01/Pal/CHQ/Tiki/15) as a metoposaurid palate (Fig:7). However, no detailed osteological description or identifying characters were described by the authors as to why the specimen was identified to be a metoposaurid. The authors only described the palate to be the ‘dorsal’ part of a metoposaurid as it has conical teeth present on the anterior part. However, this description is vague and of no taxonomical significance

whatsoever as all temnospondyls have conical teeth and both chigutisaurids and metoposaurids have teeth and tusks in the anterior part of the skulls. Again, dentition restricted to the anterior margin of the skull is a characteristic of all temnospondyls with parabolic skulls. Additionally, [Kumar & Sharma \(2019\)](#) grouped the palate collected from the village of Tenduadh with the specimens of metoposaurid clavicle collected from the village of Jora. This grouping is not viable as the two villages are approximately 12 kilometres apart from each other and there is a probability that these two villages might be parts of the Upper and Lower parts of the Tiki Formation and may even be of different ages as discussed later. The specimen, as said in [Kumar & Sharma \(2019\)](#), was too friable and could not be excavated by them. Thus, there is no option to study the specimen first-hand. Henceforth, the image of [Kumar & Sharma \(2019\)](#), has been replicated into a high-resolution photograph with the required permission from the editor of the Journal of the Palaeontological Society of India to study the detail of the described specimen (RH01/Pal/CHQ/Tiki/15) (Fig: 7).

The studied specimen is the palate of a temnospondyl and not the dorsal view of the skull as erroneously stated by [Kumar & Sharma \(2019\)](#). The palate showed in the picture (RH01/Pal/CHQ/Tiki/15) (Fig:7) has a distinct vaulted pterygoid, parabolic skull outline, and comparatively narrow cultriform process than metoposaurids. The specimen (RH01/Pal/CHQ/Tiki/15), ([Kumar & Sharma 2019](#)) (Fig:7) has a parabolic skull, thickening of the pterygoid, presence of vaulted pterygoid, presence of post-pterygoid process and concave vertical wall of pterygoid that are characteristic of chigutisaurids as written repeatedly above. The specimen is indeed friable with dense networks of fractures that obscured the clear identification of the sutures. The specimen (RH01/Pal/CHQ/Tiki/15) is partly eroded along the lateral margins

as well as anteriorly and posteriorly. The right half of the palate is slightly compressed, deformed and curved (Fig: 7).

The anterior portion of the palate is considerably broken both anteriorly and anterolaterally. Though the sutures cannot be delineated, the presence of vomer is very apparent. The vomer is broken anteriorly and the anterior palatal vacuity is not preserved (RH01/Pal/CHQ/Tiki/15) (Fig: 7). The posterior part of the preserved vomer includes the anterior tongue of the cultriform process of the parasphenoid. The left half of a possible vomerine cavity is preserved. The vomerine cavity is present only in Jurassic chigutisaur *Siderops kehli* ([Warren & Hutchinson 1983](#)) and it is absent in *Compsocerops* ([Sengupta 1995](#)) or *Pelorocephalus* ([Marsicano 1999](#)). The left lateral margin of the right choana is aligned to the left lateral margin of the right interpterygoid vacuity. The ectopterygoids are exposed on the anterolateral margins of the interpterygoid vacuities and are preserved on both sides. The ectopterygoid borders the anterior portion of the subtemporal vacuity inwards. The subtemporal vacuity is wide and broad bordered by the ectopterygoid and the parasphenoid on the inward margin and the quadratojugal, the alar process of the jugal on the outward lateral margin. The dentigerous area is restricted to the anterior region of the palate. The anterior and the anterolateral margins of the palate are broken, and all teeth are not preserved. However, two broken ectopterygoid teeth can be seen preserved at the anterolateral corner of the ectopterygoid in contact with the palatine in the left part of the palate. The palatine teeth row in the left half of the skull is also preserved partially. Like other chigutisaurids, the dentigerous area of the palate is remarkably short. The palatine row of teeth is not continuous up to the middle of the choana. This character has been considered to be a synapomorphy of *Compsocerops cosgriffi* ([Sengupta 1995](#)). Conical, inward curved four complete palatine teeth are preserved in the margin of the left palatine bone of the palate. Since, the dentigerous area is restricted to the anterior part

of the skull the posterior part is longer in proportion and covered by large and wide subtemporal
vacuity (Fig: 4).

In (RH01/Pal/CHQ/Tiki/15) (Fig:7) ([Kumar & Sharma 2019](#)) both the interpterygoid vacuities are
well preserved. The interpterygoid vacuities are quadrangular in shape, shorter and wider
compared to *Compsocerops cosgriffi* ([Sengupta 1995](#)). The borders of the interpterygoid vacuities
are approximately parallel-sided. The interpterygoid vacuities are bordered dominantly by the
cultriform process along the inward margin as well as the vomer. Anteriorly, it is bordered by the
vomer and the palatine. The pterygoid forms the dominant margin of the interpterygoid vacuities
laterally with a small area being occupied by the ectopterygoid. Posteriorly, these are formed by
the parasphenoid. In ISI A 202/1, the interpterygoid vacuities are not completely preserved. In
both, the specimens ISI A 202 and (RH01/Pal/CHQ/Tiki/15) ([Kumar & Sharma 2019](#)), the
interpterygoid vacuities are shorter and broader than *Compsocerops cosgriffi* where the base levels
of the interpterygoid vacuities are lower than that of the subtemporal vacuities. The subtemporal
vacuity extends anteriorly to the level higher than the centre of the interpterygoid vacuities.

In (RH01/Pal/CHQ/Tiki/15) ([Kumar & Sharma 2019](#)) (Fig:7) both the pterygoids are preserved.
They are deep and vaulted. The vertical lateral wall of the pterygoid projects posteriorly possibly
up to the posterior level of the occipital condyles which are broken. The palatal ramus of the
pterygoid is visible on both sides in (RH01/Pal/CHQ/Tiki/15). The palatal ramus of the pterygoid
is longitudinally concave with a gentle fold which is again a character of some chigutisaurids
specially *Compsocerops*. The quadrate ramus of the pterygoid is better preserved on the right side
of the palate (RH01/Pal/CHQ/Tiki/15). The quadrate ramus of the pterygoid looks like a wall as
they are deeply vaulted. The ascending ramus of the pterygoid is not visible in
(RH01/Pal/CHQ/Tiki/15). A broken post pterygoid process that is a projection from the posterior

border of the pterygoid corpus is visible on the right side of the palate (RH01/Pal/CHQ/Tiki/15). This area on the left side of the palate of (RH01/Pal/CHQ/Tiki/15) is broken. The postpterygoid process is considered to be an apomorphic character for *Compsocerops cosgriffi* (Marsicano 1999). The suture of the quadrate and pterygoid is present on the outer side of the downturned part of the quadrate ramus of the pterygoid. In ISI A 202, (Fig: 4) only the right pterygoid is ill-preserved but the bone surface is crushed. However, a distinct post pterygoid process characteristic of *Compsocerops* is present. Though the bone is crushed and compressed, the vaulted nature of the pterygoid can be made out because of the concavity of the vertical wall of the pterygoid. In both, the specimens (RH01/Pal/CHQ/Tiki/15) (Kumar & Sharma 2019) and ISI A 202, the palatine ramus of the pterygoid is much broader and wider than that in *Compsocerops cosgriffi*.

Just like other chigutisaurids, the base of the parasphenoid is almost hexagonal with a long extension in the form of the cultriform process placed between two interpterygoid vacuities in (RH01/Pal/CHQ/Tiki/15) (Kumar & Sharma 2019). The parasphenoid has a long suture with the pterygoid laterally and the exoccipitals posteriorly. A distinct raised longitudinal keel is present on the ventral surface of the cultriform process in this specimen. The presence of this keel in the cultriform process has been noted by Marsicano (1999) as a distinguishing character present only in *Pelorocephalus mendozensis*. However, first-hand studies reveal that this longitudinal keel of the cultriform process is also present in *Compsocerops cosgriffi* from the Maleri Formation of Pranhita - Godavari Valley Basin. The cultriform process of parasphenoid of this specimen is comparatively narrower than all other specimens of *Compsocerops cosgriffi*. The cultriform process of the *Compsocerops* species from Tiki is wider than *C. cosgriffi*, the cultriform process is also comparatively broader than *Siderops kehli*, more comparable to the width of the cultriform process in the specimen previously denoted as *Kuttycephalus triangularis* (Sengupta 1995). The cultriform

process preserved in (RH01/Pal/CHQ/Tiki/15) ([Kumar & Sharma 2019](#)) is thin and constricted in the middle part of the interpterygoid vacuities and gets broader as it progresses to the anterior part of the process. This type of cultriform process is unique among the chigutisaurids. In the specimen photographed by [Kumar & Sharma \(2019\)](#), the anterior tongue of the cultriform process is in contact with the vomer and lies posterior to the level of the anterior margin of the interpterygoid vacuities. The cultriform process is not preserved in ISI A 202. The occipital condyles are broken as well. In the earliest known chigutisaur *Keratobrachyops*, the cultriform process of the parasphenoid is also narrower than ISI A 202.

The Occiput (Fig: 5)

The occiput is very ill preserved only in ISI A 202 (Fig: 5). The occiput could not be prepared due to the extremely fragile nature of the skull. Removing the matrix load from the occiput would result in the sagging of the entire specimen. However, from the little that could be studied, it can be said that in occipital view, the quadrate ramus of the pterygoid is deeply downturned. The vagus nerve foramen is preserved on the left exoccipital lateral to the broken occipital condyle. The ascending process of the exoccipital is wide and inclined and meets the descending process of the postparietal. A sub-circular, matrix filled, paraquadrate foramen is present in the quadratojugal. The quadrate is partially preserved in the occipital view. It is bounded by the squamosal, quadratojugal and the downturned pterygoid. The absence of occiput makes the comparison of ISI A 202 difficult with the different species of *Pelorocephalus* as different species of the genus are differentiated, to a great extent, by their occipital characters ([Marsicano 1999](#)).

Clavicle (Fig: 6)

An almost complete left clavicle (ISI A 202/2) (Fig: 6) was found associated with the skull (ISI A 202/1) during excavation. The clavicle has a flat eye-drop shaped blade and a long straight dorsal process that ascends almost straight, nearly at ninety degrees with the plate. The cross-section of the process at the dorsal end is bean shaped as a feeble furrow runs along the process. This is very similar to the clavicle of *Compsocerops cosgriffi* (Sengupta 1995), *Siderops kehli* (Warren & Hutchinson 1983) and *Koolasuchus cleelandi* (Warren et al. 1997).

Results

Tiki Formation:

To date, no temnospondyl fauna has been recorded from the upper part Tiki Formation. Excavation taken up in 2018 by the authors revealed the first chigutisaurid from the Tiki Formation in the Tenduadh locality (Fig: 8) in the upper part of the Tiki Formation. Several vertebrae and postcranial bones of metoposaurid have been excavated from the Jora and Tiki Nala sections which have been assigned to the lower Tiki Formation from our field studies. Based on the changes in the faunal assemblage in the lower and the upper Tiki Formation (Tables 2a and 2b) and considering the lithostratigraphy, a boundary between the basal and upper Tiki Formation has been assigned and the zone demarcating the Carnian to Norian faunal turnover in the Tiki Formation has been approximated and marked in red dotted line in the map (Fig: 1). The lithological logs modified after Kumar & Sharma (2019) and Mukherjee & Ray (2012) reveal that just like the Maleri Formation, the basal Tiki Formation is dominated by a large band of red mudrock intercalated with peloidal calcirudite-calcarenite (Sarkar 1988) (Fig: 9). The Jora Nala section in the Carnian basal Tiki has been logged in detail in this work (Fig:9a). This shows the dominance of greenish to

reddish siltstones and mudstones in the Jora Nala section with sparse deposition of trough cross-bedded sandstones in between.

Maleri Formation

In the revised map of the Maleri Formation and (after [Dasgupta et al. 2017](#); [Kutty & Sengupta 1989](#)) a boundary between the Carnian basal Maleri and Middle Carnian / Norian Upper Maleri has been established from both lithological and faunal contents (Fig: 2 – faunal boundary indicated by a green broken line). From our field studies and maps it is evident that though sandstone - mudstone alternation is present throughout the Maleri Formation, the Carnian basal Maleri is abundant in red mudrocks and calcirudites (Fig: 2,10) and moving towards Upper Maleri there is a sudden increase in the deposition on siliciclastic sediments leading to the more frequent occurrence of sandstone bands alternating with red mudstone (Fig: 10).

Discussion

The CPE had a significant impact on the terrestrial environment and ecosystem globally. The evidence of CPE has never been worked upon or mentioned in India because of the lack of proper age constraints present in the Late Triassic Maleri and Tiki Formations in India.

Tiki Formation

No detailed sedimentological or geochemical studies have been carried out in the Late Triassic Tiki Formation in India to analyse the associated changes from Carnian to Norian through the humid phase of the Carnian Pluvial Episode. Though, [Ahmed & Ray \(2010\)](#) presented a geochemical analysis of 42 nodular carbonates confirming their pedogenic origin, no details of the localities of collection in terms of lower and upper Tiki have been provided.

The terrestrial influx of sediments is significantly low at the period denoted by the sparse occurrence of sandstones in the basal Tiki Formation (Fig: 9). The presence of *Unio* beds in between the basal thick layers of mud reflects a stagnant quiet and well-watered environment. This basal mud encompasses areas like the Jora and Tiki River sections. Abundant postcranial fragments of metoposaurids and rhynchosaurs have been collected from these sections. Moving upwards in the direction of the dip of the beds, there is a sudden influx of siliciclastic sediments marked by thick sandstone units with little intermittent mudstone. This could be a demarcation of the rapid influx of siliciclastic sediments that took place during CPE in the Tiki Formation. Only two dominant sand bodies are observed in Tiki before the recurrence of a thick horizon of mud and subsequently sand-mud alternations indicating the onset of seasonality and aridity in the Norian. The Norian of Tiki Formation is demarcated by red mudstones, whitish sandstones and sparse calcirudites. The Norian Upper Tiki Formation is exposed in sections near Tenduadh as shown in the map (Fig: 1) and an estimated approximate faunal boundary between the Carnian and the Norian in the Tiki Formation is also furnished as in Fig: 1. Tiki has a long history of yielding fossil vertebrates ([Bandyopadhyay & Ray 2020b](#); [Chatterjee & Roy-Chowdhury 1974](#)) (Table 1). It has a rich Late Triassic faunal association as shown in Table 2a and 2b ([Bandyopadhyay & Ray 2020a](#); [Chatterjee & Majumdar 1987](#); [Mukherjee & Ray 2014](#)). The Tiki faunal assemblage was thought to be coeval to the Lower Maleri faunal assemblage ([Datta 2005](#); [Kutty & Sengupta 1989](#)). However, [Datta et al. \(2019b\)](#) stated that the Tiki fauna ranges from Middle Carnian to Early Norian and is younger than Lower Maleri Fauna. The mid Carnian / Norian Upper Maleri fauna has chigutisaurids. The discovery of a chigutisaurid from the upper part of the Tiki Formation confirms the views of [Datta et al. \(2019b\)](#) regarding the presence of Middle Carnian / Norian fauna in the Tiki Formation.

492 *Maleri Formation*

493 The abundance of red mudstone in the basal Maleri Formation (Fig: 11) with a sudden increase in
 494 the frequency of sandstone bands in the upper part can be correlated with the advent of the Carnian
 495 Pluvial Episode (CPE) in India. The overall palaeoenvironment and sedimentology of the Maleri
 496 formation have been worked upon by earlier workers ([Dasgupta & Ghosh 2018](#); [Sarkar 1988](#)). Most
 497 of these studies were done on the Maleri Formation as a whole without distinguishing its basal and
 498 upper parts. Also, no detailed analyses about the changes in sedimentology or geology or
 499 geochemistry have been done to study the changes in the pattern of sedimentation from the Carnian
 500 basal Maleri to the Norian Upper Maleri.

501 Evidence from detailed geological mapping, logs and associated faunal turnover in the Late
 502 Triassic Maleri Formation all point toward the distinctive effect of CPE in India. The overall
 503 thickness of the Maleri Formation is about 350-600m which is variable in different places. From
 504 the map of the Maleri Formation and log of the same near Achlapur and Nalapur (Fig: 10), it is
 505 evident that the basal part of the Maleri Formation has a large stretch or band of red mudrock-
 506 dominated unit. This unit of red mudrock stretches from about 100m-200m in thickness in basal
 507 Maleri and is the thickest band of red mudrock encountered in the entire Maleri Formation (Fig:
 508 12). It encompasses villages namely Aigerrapalli (19°15'22.4" N; 79°27'23.4" E), Achlapur
 509 (19°09'34" N; 79°31'51" E) and Nambala (19°13'47" N; 79°26'07" E), areas slightly north of
 510 village Gampalpalli (19°10'11" N; 79°30'53" E) and is rich in vertebrate fossils, the most
 511 significant among them being rhynchosaurs and metoposaurids. This significantly thick basal
 512 Maleri mudstone has sporadic carbonate grainstones [*sensu* ([Dunham 1962](#)); calcarenites–
 513 calcirudites of [Sarkar \(1988\)](#)] and the presence of palaeosols with no significant siliciclastic
 514 deposition of sandstone present within this mud. The upper part of the Formation is approximately

200–350 m thick and has three to four multi-storeyed sheet sandstone bodies (each 10–35 m thick) vertically separated by mudrock-dominated intervals (15–55 m thick). The mudrock intervals in the upper part are lithologically similar to those occurring in the lower part of the Formation. The mudrock units comprise both stratified and massive mudstones ([Dasgupta et al. 2017](#)) with sporadic carbonate grainstones (calcirudite of [Sarkar \(1988\)](#)). Moving upwards from the thick stretch of mudrock, the influx of siliciclastic sedimentation increases as evident from the increase in the deposition of frequent sandstone units. The beginning of these sandstone units is marked by the presence of metoposaurids and rhynchosaurs and unionid fossils in a sandy zone. In the upper part, apart from the chigutisaurids, there are basal sauropods like *Jaklapallisaurus*, and sauropodomorphs like *Nambalia*, probable Guaibasaurids-and-Herrerasaurus-like forms ([Novas et al. 2010](#)). The authors in the same work, also mentioned that early theropods are known from the Norian-Rhaetian time from North and South America, India, South Africa, and Europe and the demise of members of Lower Maleri fauna like rhynchosaurs together with the global extinction of *Chiniquodon* (cynodont) and Proterocampsidae (archosauriform) mark the Carnian – Norian boundary and also the North Tethyan Pluvial Episode of end Carnian (CPE). [Benton et al. \(2018\)](#) have argued that CPE triggered the diversification of early dinosaurs. It has been discussed earlier those indications of CPE are present in Maleri and Tiki Formations of India. The sudden appearance of several basal dinosaurs like *Jaklapallisaurus*, *Nambalia*, probable Guaibasauridae and *Herrerasaurus* ([Novas et al. 2010](#)) in the Norian Upper Maleri fauna also corresponds to that.

Thus, the palaeoenvironment of the Maleri Formation shifted from a comparatively arid and dry climate in the Julian at the basal substage of Carnian to a high competence fluvial-lacustrine environment with the presence of small, ephemeral and vegetated swamps or ponds along the flow path of the channels at the time of Carnian Pluvial Episode from the end of Julian to Tuvanian and

back again to fluvial deposition in the Norian ([Dasgupta et al. 2017](#)). The episodes of increased rainfall during the Carnian Pluvial Episode demarcated by increased frequency of sandstone deposition are intervened by seasonality as evident from the red mudrock alternations between sandstones. Similar climatic shifts are seen from the coeval Santa Maria to Caturrita formations of Brazil ([Dal Corso et al. 2015](#)). These shifts indicating a major variation of the hydrological regime in terrestrial depositional settings suggest an enhancement of the hydrological cycle during the CPE. Recently, [Lucas \(2010\)](#) stated that the demise of metoposaurids in most parts of the world during the Carnian is related to the end of the enhanced hydrological cycle at the dying phase of CPE. The disappearance of key herbivorous groups such as dicynodonts and rhynchosaurs of Carnian and their places taken up by giant sauropodomorphs seems to be linked to CPE which is not documented in India so far.

Also, the *Hyperodepadon* Assemblage Zone (HAZ) is characterized by the presence of rhynchosaur *Hyperodepadon* and is present in the lower part of the Ischigualasto Formation of Argentina, the Lossiemouth Sandstone Formation of Scotland, and the Lower Maleri Formation of India ([Langer et al. 2010](#)). The HAZ is dated as late Carnian to early Norian, approx. 228–224 Ma by some authors ([Benton et al. 2018](#); [Brusatte et al. 2010](#); [Ezcurra et al. 2017](#)). Most metoposaurids in the Gondwana deposits are considered to be Carnian in age ([Chakravorti & Sengupta 2019](#); [Gee & Jasinski 2021b](#); [Sengupta 2002](#)). The demise of the metoposaurids *Panthsaurus maleriensis* ([Chakravorti & Sengupta 2019](#)) in India along with the demise of *Hyperodepadon* ([Chatterjee 1974](#); [Mukherjee et al. 2012](#)) in both Late Triassic Maleri and Tiki Formation also points to the presence and effect of CPE in India. The demise of the metoposaurids left a vacant niche to be occupied by the chigutisaurids in the Norian suggesting short-lived aridity in post Carnian stage.

The Carnian of Argentina has its age radiometrically constrained between 231.4 ± 0.3 and 225.9 ± 0.9 Ma ([Martínez et al. 2016](#)) and similar reports are present from Santa Maria and Caturrita Formations of Brazil (233.2 ± 0.7 and 225.4 ± 0.4 Ma) ([Langer et al. 2018](#)). The onset of CPE is well constrained in stratigraphic sections like the Southern Alps of Italy, Northern Calcareous Alps of Austria, Transdanubian Range of Hungary, and the Nanpanjiang Basin of the South China block and is placed at the substages Julian 1 – Julian 2 boundary of the Carnian ([Gallet et al. 1994](#)). Due to the lack of any radiometric dating, Late Triassic Maleri and Tiki Formations are poorly constrained and pose difficulty in global correlation, their correlation is based only on available fauna. The CPE has always been dated as mid-Carnian ([Ruffell et al. 2016](#)) but this is not a unanimous viewpoint. The Italian Dolomites occur between the Aonoides/Austriacum interval (about Julian) and the base of the Subbullatus Zone (Tuvalian), dated at 234–232 Ma ([Dal Corso et al. 2015](#); [Roghi et al. 2010](#)). Further constraint has been documented in borehole successions in the southwest UK, which indicates a maximum duration of 1.09 MYA ([Miller et al. 2017](#)). The precise radiometric dating to constrain the Maleri and Tiki Formations and to denote the beginning of CPE in India will shed further light on the pattern of faunal diversification post CPE event in the subcontinent and help in the global stratigraphic correlation. A continental carbon isotope record in southwest England shows multiple carbon cycle perturbations during CPE ([Miller et al. 2017](#)). The CPE is not only the time interval of increased humidity but also a major carbon perturbation. Unfortunately, no carbon isotope data is noted from the Maleri and the Tiki Formations of India.

Conclusion

1. In the current work a new species of chigutisarid amphibian, *Compsocerops tikiensis* sp. nov. from the Late Triassic Tiki Formation of the Rewa Gondwana Basin has been described in detail. The presence of chigutisaurid *Compsocerops tikiensis* sp. nov. in the

Upper part of the Tiki Formation is the first evidence of the Norian chigutisaurid amphibian from the said Formation and is important for correlation of the Late Triassic basins worldwide.

2. The presence of *Compsocerops* ([Sengupta 1995](#)) in Tiki, for the first time, confirms the presence of the Upper Maleri faunal element in Tiki. [Lucas \(2020b\)](#) thought that the demise of metoposaurids in most parts of the world was at the end of Carnian and that tallies with the last appearance datum of the metoposaurids of Maleri. The chigutisaurids, both in Maleri and Tiki have their first appearance datum at the onset of the Middle Carnian or Early Norian.
3. Along with the extinction of the rhynosaurs and *Parasuchus* (primitive phytosaur) ([Chatterjee 1978](#)), chiniquodontids (cynodonts), the Carnian – Norian Extinction Event (CNEE) also caused the extinction of the metoposaurids in India. Chigutisaurids appeared in Middle Carnian / Norian and India is the only place which accommodates definite metoposaurids and chigutisaurids within the same formations (the Late Triassic Maleri and Tiki Formations) the former being replaced by the latter. Incidentally, among the phytosaurs, the *Parasuchus* of Lower Maleri fauna is replaced by the *Leptosuchus* like forms of Upper Maleri and *Volcanosuchus statisticae* ([Datta et al. 2021b](#)) in the upper part of the Tiki Formation.
4. The post-CNEE empty niche left by the metoposaurids in the Late Triassic Gondwana deposits of India (controversially Brazil as well, (see [Dias-da-Silva et al. 2011](#)) was occupied by the chigutisaurids in the Norian. The availability of phytosaur teeth along with *C. tikiensis* sp. nov. only indicated their co-existence in the same aquatic niche but does not necessarily point towards any prey-predatory relationship between the phytosaurids and

the chigutisaurids. However, detailed studies on histology and growth pattern of the chigutisaurids might shed light on the gigantism of these amphibious animals in the post-CNEE and recovery of the temnospondyls.

5. The presence of both metoposaurids and chigutisaurids and the faunal turnover from the Carnian to the Norian along with the extinction of the rhynchosaurs (*Hyperodapedon*) ([Mukherjee et al. 2012](#)) and *Parasuchus* ([Chatterjee 1978](#)) in the Carnian of both the Late Triassic Maleri and Tiki Formation and the presence of prosauropods in the Upper Maleri Formation and undescribed dinosauriformes including theropod-like forms ([Bandyopadhyay & Ray 2020b](#)) sheds light and documents for the first time the existence and effect of the Carnian Pluvial Episode in India.
6. The finding of *Compsocerops tikiensis* sp. nov. from the Tiki Formation and assessing its importance in global palaeoclimatic and palaeoclimatic correlation paves the way for future scope of works in finding the effect of CPE in India and comparing its faunal diversification at a global scale. Also, finding any new and better-preserved specimen of *Compsocerops tikiensis* sp. nov. from the Tiki Formation would help to establish the phylogenetic relationship of the other chigutisaurids around the world with that of the Indian counterparts. As the specimens are deformed broken and have poor preservation potential, phylogenetic analysis is beyond the scope of this paper as it will add more missing and misinterpreted data.

Acknowledgements

The author SC would like to thank Ms. Aindrila Roy, Project Linked Person (PLP) at the Geological Studies Unit, Indian Statistical Institute, Kolkata for her help in editing and formatting the manuscript. The authors' SC and DPS would like to acknowledge Mr Lakshman

Mahankur for helping in the preparation of the poorly preserved material. We acknowledge the editor (Surendra Kumar) of the Journal of Palaeontology of India for providing us with permission to reproduce the published picture published on page 159, (Plate IV. number 1) of the journal volume 64(2), December 31, 2019. We are also very grateful to Dr Claudia Marsicano, Dr Valentin Buffa and the anonymous reviewer who through their insightful comment and meticulous suggestions has enriched this work to a great extent.

Reference

- Ahmed F, and Ray S. 2010. Paleoclimate of Upper Triassic India: new observations from the Rewa Gondwana basin. *Geochimica et Cosmochimica Acta*: PERGAMON-ELSEVIER SCIENCE LTD THE BOULEVARD, LANGFORD LANE, KIDLINGTON p A6-A6.
- Aiyengar N. 1937. A note on the Maleri beds of Hyderabad State (Deccan) and the Tiki beds of south Rewa. *Records of the Geological Survey of India* 71:401-406.
- Arche A, and Lopez-Gomez J. 2014. The Carnian pluvial event in Western Europe: new data from Iberia and correlation with the Western Neotethys and eastern North America–NW Africa regions. *Earth-Science Reviews* 128:196-231.
- Bandyopadhyay S, and Ray S. 2020a. Gondwana vertebrate faunas of India: their diversity and intercontinental relationships. *Episodes* 43:438-460.
- Bandyopadhyay S, and Ray S. 2020b. Gondwana vertebrate faunas of India: their diversity and intercontinental relationships. *Episodes Journal of International Geoscience* 43:438-460.
- Bandyopadhyay and Ray 2020
- Benton MJ, Bernardi M, and Kinsella C. 2018. The Carnian Pluvial Episode and the origin of dinosaurs. *Journal of the Geological Society* 175:1019-1026.
- Bernardi M, Gianolla P, Petti FM, Mietto P, and Benton MJ. 2018. Dinosaur diversification linked with the Carnian Pluvial Episode. *Nature Communications* 9:1-10.
- Bhat MS. 2017. Techniques for systematic collection and processing of vertebrate microfossils from their host mudrocks: a case study from the Upper Triassic Tiki Formation of India. *Journal of the Geological society of India* 89:369-374.
- Bonaparte JF. 1975. *Sobre la Presencia Del Laberintodonte Pelorocephalus en la Formación de Ischigualasto Y Su Significado Estratigráfico (Brachypoidea-Chigutisauridae)*.
- Brusatte SL, Nesbitt SJ, Irmis RB, Butler RJ, Benton MJ, and Norell MA. 2010. The origin and early radiation of dinosaurs. *Earth-Science Reviews* 101:68-100.
- Buffa V, Jalil NE, and Steyer JS. 2019. Redescription of Arganasaurus (Metoposaurus) azerouali (Dutuit) comb. nov. from the Upper Triassic of the Argana Basin (Morocco), and the first phylogenetic analysis of the Metoposauridae (Amphibia, Temnospondyli). *Papers in Palaeontology* 5:699-717.
- Cabrera Á. 1944. *Sobre un estegocéfalo de la Provincia de Mendoza*: Instituto del Museo de la Universidad Nacional de La Plata.
- Chakravorti S, and Sengupta DP. 2019. Taxonomy, morphometry and morphospace of cranial bones of Panthasaurus gen. nov. maleriensis from the Late Triassic of India. *Journal of Iberian Geology* 45:317-340.

- Chatterjee S. 1974. A rhynchosaur from the Upper Triassic Maleri Formation of India. *Philos Trans R Soc Lond B Biol Sci* 267:209-261. 10.1098/rstb.1974.0001
- CHATTERJEE S. 1978. A PRIMITIVE PARASUCHID (PHYTOSAUR) REPTILE FROM THE UPPER TRIASSIC MALERI FORMATION OF INDIA.
- Chatterjee S, and Majumdar PK. 1987. Tikisuchus romeri, a new rauisuchid reptile from the Late Triassic of India. *Journal of Paleontology* 61:787-793.
- Chatterjee S, and Roy-Chowdhury T. 1974. Triassic Gondwana vertebrates from India. *Indian Journal of Earth Sciences* 1:96-112.
- Chowdhury TR. 1965. A new metoposaurid amphibian from the Upper Triassic Maleri Formation of Central India. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences* 250:1-52.
- Cotter GdP. 1917. A Revised Classification of the Gondwana System." *Rec. Geol. Surv. India*, Vol. XLVIII.
- Dal Corso J, Bernardi M, Sun Y, Song H, Seyfullah LJ, Preto N, Gianolla P, Ruffell A, Kustatscher E, and Roghi G. 2020. Extinction and dawn of the modern world in the Carnian (Late Triassic). *Science Advances* 6:eaba0099.
- Dal Corso J, Gianolla P, Newton RJ, Franceschi M, Roghi G, Caggiati M, Raucsik B, Budai T, Haas J, and Preto N. 2015. Carbon isotope records reveal synchronicity between carbon cycle perturbation and the "Carnian Pluvial Event" in the Tethys realm (Late Triassic). *Global and Planetary Change* 127:79-90.
- Dal Corso J, Mietto P, Newton RJ, Pancost RD, Preto N, Roghi G, and Wignall PB. 2012. Discovery of a major negative $\delta^{13}\text{C}$ spike in the Carnian (Late Triassic) linked to the eruption of Wrangellia flood basalts. *Geology* 40:79-82.
- Dal Corso J, Ruffell A, and Preto N. 2018. The Carnian pluvial episode (Late Triassic): new insights into this important time of global environmental and biological change. *Geological Society of London*.
- Dasgupta S, and Ghosh P. 2018. Freshwater carbonates within a Late Triassic siliciclastic fluvial system in a Gondwana rift basin: The Maleri Formation, India. *Sedimentary geology* 373:254-271.
- Dasgupta S, Ghosh P, and Gierlowski-Kordesch EH. 2017. A discontinuous ephemeral stream transporting mud aggregates in a continental rift basin: the Late Triassic Maleri Formation, India. *Journal of Sedimentary Research* 87:838-865.
- Datta D, Kumar N, and Ray S. 2021a. Taxonomic identification of isolated phytosaur (Diapsida, Archosauria) teeth from the Upper Triassic of India and their significances. *Historical Biology* 33:272-282.
- Datta D, Mukherjee D, and Ray S. 2019a. Taphonomic signatures of a new Upper Triassic phytosaur (Diapsida, Archosauria) bonebed from India: Aggregation of a juvenile-dominated paleocommunity. *Journal of Vertebrate Paleontology* 39:e1726361.
- Datta D, Ray S, and Bandyopadhyay S. 2019b. Cranial morphology of a new phytosaur (Diapsida, Archosauria) from the Upper Triassic of India: implications for phytosaur phylogeny and biostratigraphy. *Papers in Palaeontology*.
- Datta D, Ray S, and Bandyopadhyay S. 2021b. Cranial morphology of a new phytosaur (Diapsida, Archosauria) from the Upper Triassic of India: implications for phytosaur phylogeny and biostratigraphy. *Papers in Palaeontology* 7:675-708.
- Datta P. 2005. Earliest mammal with transversely expanded upper molar from the Late Triassic (Carnian) Tiki Formation, South Rewa Gondwana Basin, India. *Journal of Vertebrate Paleontology* 25:200-207.
- Dias-da-Silva S, Cabreira SF, and Da Silva LR. 2011. Occurrence of giant stereospondyl remains in the Santa Maria Formation (Middle/Upper Triassic of southern Brazil). *Alcheringa* 35:11-19.

- DIAS-da-SILVA S, Sengupta DP, Cabreira SF, and Da Silva LR. 2012. The presence of Compsocerops (Brachyopoidea: Chigutisauridae)(Late Triassic) in southern Brazil with comments on chigutisaurid palaeobiogeography. *Palaeontology* 55:163-172.
- Dunham RJ. 1962. Classification of carbonate rocks according to depositional textures.
- Dutta PK, and Ghosh SK. 1993. The century-old problem of the Pali-Parsora-Tiki stratigraphy and its bearing on the Gondwana classification in Peninsular India. *Journal of Geological Society of India (Online archive from Vol 1 to Vol 78)* 42:17-31.
- Ezcurra MD, Fiorelli LE, Martinelli AG, Rocher S, von Baczko MB, Ezpeleta M, Taborda JR, Hechenleitner EM, Trotteyn MJ, and Desojo JB. 2017. Deep faunistic turnovers preceded the rise of dinosaurs in southwestern Pangaea. *Nature Ecology & Evolution* 1:1477-1483.
- Fortuny J, Arbez T, Muijal E, and Steyer JS. 2019. Reappraisal of 'Metoposaurus hoffmani' Dutuit, 1978, and description of new temnospondyl specimens from the Middle–Late Triassic of Madagascar (Morondava Basin). *Journal of Vertebrate Paleontology* 39:e1576701.
- Fox SCS. 1931. *The Gondwana System and related formations*: Government of Indian Central Publication Branch.
- Furin S, Preto N, Rigo M, Roghi G, Gianolla P, Crowley JL, and Bowring SA. 2006. High-precision U-Pb zircon age from the Triassic of Italy: Implications for the Triassic time scale and the Carnian origin of calcareous nannoplankton and dinosaurs. *Geology* 34:1009-1012.
- Gallet Y, Besse J, Krystyn L, Théveniaut H, and Marcoux J. 1994. Magnetostratigraphy of the Mayerling section (Austria) and Erenkolu Mezarlik (Turkey) section: Improvement of the Carnian (late Triassic) magnetic polarity time scale. *Earth and Planetary Science Letters* 125:173-191.
- Gee BM, and Jasinski SE. 2021a. Description of the metoposaurid *Anaschisma browni* from the New Oxford Formation of Pennsylvania. *Journal of Paleontology* 95:1061-1078.
- Gee BM, and Jasinski SE. 2021b. Description of the metoposaurid *Anaschisma browni* from the New Oxford Formation of Pennsylvania. *Journal of Paleontology*:1-18.
- Hughes TW. 1877. *The Wardha Valley coal-field*: Government of India.
- Hussain S. 2018. Study of Late Triassic Vertebrate Fossils from Tiki Formation, Shahdol District, Madhya Pradesh. *Masters' Dissertation, Knowledge Repository, Central University of Punjab*.
- Kent DV, Olsen PE, and Muttoni G. 2017. Astrochronostratigraphic polarity time scale (APTS) for the Late Triassic and Early Jurassic from continental sediments and correlation with standard marine stages. *Earth-Science Reviews* 166:153-180.
- Konietzko-Meier D, Gruntmeier K, Marcé-Nogué J, Bodzioch A, and Fortuny J. 2018. Merging cranial histology and 3D-computational biomechanics: a review of the feeding ecology of a Late Triassic temnospondyl amphibian. *PeerJ* 6:e4426. 10.7717/peerj.4426
- Konietzko-Meier D, Werner JD, Wintrich T, and Martin Sander P. 2019. A large temnospondyl humerus from the Rhaetian (Late Triassic) of Bonenburg (Westphalia, Germany) and its implications for temnospondyl extinction. *Journal of Iberian Geology* 45:287-300.
- Kumar J, and Sharma KM. 2019. Micro and mega-vertebrate fossils from the Late Triassic Tiki Formation, South Rewa Gondwana basin, India: palaeoenvironmental and palaeobiogeographic implications. *Journal of the Palaeontological Society of India* 64:151-168.
- Kutty T, Jain S, and Chowdhury TR. 1987. Gondwana sequence of the northern Pranhita-Godavari Valley: its stratigraphy and vertebrate faunas.
- Kutty T, and Sengupta D. 1989. Late Triassic formations of the Pranhita-Godavari valley and their vertebrate faunal sequence-a reappraisal: *Indian Journal of Earth Science*, v. 16.
- Langer MC, Ezcurra MD, Bittencourt JS, and Novas FE. 2010. The origin and early evolution of dinosaurs. *Biological Reviews* 85:55-110.
- Langer MC, Ramezani J, and Da Rosa ÁA. 2018. U-Pb age constraints on dinosaur rise from south Brazil. *Gondwana Research* 57:133-140.

- Lucas SG. 2010. The Triassic timescale based on nonmarine tetrapod biostratigraphy and biochronology. *Geological Society, London, Special Publications* 334:447-500.
- Lucas SG. 2020a. Biochronology of Late Triassic Metoposauridae (Amphibia, Temnospondyli) and the Carnian pluvial episode. *Annales Societatis Geologorum Poloniae*.
- Lucas SG. 2020b. Biochronology of Late Triassic Metoposauridae (Amphibia, Temnospondyli) and the Carnian pluvial episode. *Annales Societatis Geologorum Poloniae*. p 409-418, doi: 410.14241/asgp. 12020.14229.
- Maheshwari HK, Kumaran K, and Bose M. 1976. The age of the Tiki Formation: with remarks on the miofloral succession in the Triassic Gondwanas of India.
- Maron M, Muttoni G, Rigo M, Gianolla P, and Kent DV. 2019. New magnetobiostratigraphic results from the Ladinian of the Dolomites and implications for the Triassic geomagnetic polarity timescale. *Palaeogeography, Palaeoclimatology, Palaeoecology* 517:52-73.
- Marsicano CA. 1999. Chigutisaurid amphibians from the Upper Triassic of Argentina and their phylogenetic relationships. *Palaeontology* 42:545-565.
- Martínez RN, Apaldetti C, Correa GA, and Abelín D. 2016. A Norian lagerpetid dinosauroform from the Quebrada del Barro Formation, northwestern Argentina. *Ameghiniana* 53:1-13.
- Miller CS, Peterse F, Da Silva A-C, Baranyi V, Reichert GJ, and Kürschner WM. 2017. Astronomical age constraints and extinction mechanisms of the Late Triassic Carnian crisis. *Scientific Reports* 7:1-7.
- Mukherjee D, and Ray S. 2012. Taphonomy of an Upper Triassic vertebrate bonebed: a new rhynchosaur (Reptilia; Archosauriformes) accumulation from India. *Palaeogeography, Palaeoclimatology, Palaeoecology* 333:75-91.
- Mukherjee D, and Ray S. 2014. A new Hesperodapedon (Archosauriformes, Rhynchosauria) from the Upper Triassic of India: implications for rhynchosaur phylogeny. *Palaeontology* 57:1241-1276.
- Mukherjee D, Ray S, Chandra S, Pal S, and Bandyopadhyay S. 2012. Upper Gondwana succession of the Rewa basin, India: understanding the interrelationship of lithologic and stratigraphic variables. *Journal of the Geological society of India* 79:563-575.
- Mukhopadhyay G, Mukhopadhyay S, Roychowdhury M, and Parui P. 2010. Stratigraphic correlation between different Gondwana basins of India. *Journal of the Geological society of India* 76:251-266.
- Novas FE, Ezcurra MD, Chatterjee S, and Kuttu T. 2010. New dinosaur species from the Upper Triassic Upper Maleri and Lower Dharmaram formations of central India. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 101:333-349.
- Pascoe EH. 1973. *A Manual of the Geology of India and Burma*: Controller of Publications.
- Pledge NS. 2013. First South Australian Labyrinthodont: A Possible Chigutisaurid Stereospondyl Amphibian from the Late Triassic at Leigh Creek. *Transactions of the Royal Society of South Australia* 137:127-134.
- Rakshit N, and Ray S. 2020. Mortality dynamics and fossilisation pathways of a new metoposaurid-dominated multitaxic bonebed from India: a window into the Late Triassic vertebrate palaeoecosystem. *Historical Biology*:1-23. doi.org/10.1080/08912963.2020.1777550
- Ray S, Bhat MS, Mukherjee D, and Datta P. 2016a. Vertebrate fauna from the Late Triassic Tiki Formation of India: new finds and their biostratigraphic implications.
- Ray S, Bhat MS, Mukherjee D, and Datta P. 2016b. Vertebrate fauna from the Late Triassic Tiki Formation of India: new finds and their biostratigraphic implications. *The Palaeobotanist* 65:47-59.
- Robinson PL. 1970. The Indian Gondwana formations-a review. *1st international symposium on gondwana stratigraphy*:201-268.


- Roghi G, Gianolla P, Minarelli L, Pilati C, and Preto N. 2010. Palynological correlation of Carnian humid pulses throughout western Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology* 290:89-106.
- Roychowdhury M, Sastry M, Shah S, Singh G, and Ghosh S. 1975. Triassic floral succession in the Gondwana of Peninsular India. *Gondwana Geology Australia Natural University Press, Canberra*:149-158.
- Ruffell A, Simms M, and Wignall P. 2016. The Carnian Humid Episode of the late Triassic: a review. *Geological Magazine* 153:271-284.
- Rusconi C. 1949. Otra especie de Labyrinthodonte del Triásico de Mendoza. *Revista del Museo de Historia Natural de Mendoza* 3:93-94.
- Rusconi C. 1951. Laberintodontes triásicos y pérmicos de Mendoza. *Revista del Museo de Historia Natural de Mendoza* 5:33-158.
- Sarkar S. 1988. Petrology of caliche-derived peloidal calcirudite/calcarenite in the Late Triassic Maleri Formation of the Pranhita-Godavari valley, South India. *Sedimentary geology* 55:263-282.
- Schlager W, and Schöllnberger W. 1974. Das Prinzip stratigraphischer Wenden in der Schichtfolge der Nördlichen Kalkalpen. *Mitt Geol Ges Wien* 66:165-193.
- Schoch RR, Milner AR, and Witzmann F. 2014. Skull morphology and phylogenetic relationships of a new Middle Triassic plagiosaurid temnospondyl from Germany, and the evolution of plagiosaurid eyes. *Palaeontology* 57:1045-1058.
- SENGUPTA DP. 1992. Metoposaurus maleriensis Roychowdhury from the Tiki Formation of Son-Mahanadi Valley of Central India.
- Sengupta DP. 1995. Chigutisaurid temnospondyls from the late triassic of India and review of the family chigutisauridae.
- Sengupta DP. 2002. Indian metoposaurid amphibians revised. *Paleontological Research* 6:41-65.
- Simms MJ, and Ruffell AH. 1990. Climatic and biotic change in the late Triassic. *Journal of the Geological Society* 147:321-327.
- Sun Y, Wignall PB, Joachimski MM, Bond DP, Grasby SE, Lai XL, Wang L, Zhang ZT, and Sun S. 2016. Climate warming, euxinia and carbon isotope perturbations during the Carnian (Triassic) Crisis in South China. *Earth and Planetary Science Letters* 444:88-100.
- Tripathi C. 1969. *Fossil labyrinthodonts from the Panchet Series of the Indian Gondwanas*: Manager of Publications, Civil Lines.
- Veevers JJ, and Tewari R. 1995. *Gondwana master basin of peninsular India between Tethys and the interior of the Gondwanaland province of Pangea*: Geological Society of America.
- von Zittel KA. 1887. *Handbuch der Paläontologie. 1. Paläozoologie: 3. Vertebrata:(pisces, amphibia, reptilia, aves)*: Oldenbourg.
- Warren A. 1981. A horned member of the labyrinthodont super-family Brachyopoidea from the Early Triassic of Queensland. *Alcheringa* 5:273-288.
- Warren A, and Hutchinson MN. 1983. The last labyrinthodont? A new brachyopoid (Amphibia, Temnospondyli) from the early Jurassic Evergreen Formation of Queensland, Australia. *Philosophical Transactions of the Royal Society of London B, Biological Sciences* 303:1-62.
- Warren A, and Marsicano C. 2000. A phylogeny of the Brachyopoidea (Temnospondyli, Stereospondyli). *Journal of Vertebrate Paleontology* 20:462-483.
- Warren A, Rich TH, and Vickers-Rich P. 1997. The last last labyrinthodonts. *PALAEONTOGRAPHICA ABTEILUNG A-STUTTGART*- 247:1-24.

Table 1(on next page)

Spatial and temporal range of the family Chigutisauridae along with their paleoecology and paleoenvironment.

Abbreviations Max ma = Maximum age million years ago, Min ma = Minimum age million years ago, Fm. = Geological Formations from which the specimen had been excavated. The table has been modified after data was acquired from the Paleobiology Database on 10th May 2022, using the family name “Chigutisauridae”. Data for the table was modified after Bonaparte (1957); Marsicano (1999); Dias – da- Silva and Sengupta (2012); Warren (2006); Pledge (2013); Cabrera (1944); Rusconi (1949); Rusconi (1951); Sengupta (1995); Warren and Hutchinson (1983); Warren (1981); Warren and Marsicano (2000).

1 **Table 1**

Name	Early_interval	Late_interval	Max_ma	Min_ma	Country	Fm.	Lithology	Palaeoenvironm
<i>Koolasuchus cleelandi</i>	Late Aptian	—	122.46	112.03	Australia	Eumeralla Fm.	Sandstone	
<i>Koolasuchus cleelandi</i>	Late Aptian	—	122.46	112.03	Australia	Eumeralla Fm.	Sandstone	
<i>Koolasuchus cleelandi</i>	Late Aptian	—	122.46	112.03	Australia	Eumeralla Fm.	Sandstone	
<i>Koolasuchus cleelandi</i>	Late Aptian	—	122.46	112.03	Australia	Eumeralla Fm.	Sandstone	
<i>Koolasuchus cleelandi</i>	Aptian	—	125	113	Australia	Eumeralla Fm.	Sandstone	
<i>Siderops kehli</i>	Pliensbachian	Toarcian	190.8	174.1	Australia	Evergreen Fm.	Ironstone/Sandstone	
<i>Chigutisauridae indet</i>	Mid Carnian	—	228	—	Australia	Leigh Creek Fm. Santa Maria	Siltstone	
<i>Compsocerops</i>	Mid Carnian	Norian	228	208.5	Brazil	Fm. Upper Maleri	Mudstone	
<i>Compsocerops cosgriffi</i>	Mid Carnian	Norian	228	208.5	India	Fm. Upper Maleri	Mudstone	
<i>Kuttycephalus triangularis</i>	Mid Carnian	Norian	228	208.5	India	Fm.	Mudstone	
<i>Compsocerops tikiensis</i>	Mid Carnian	Norian	228	208.5	India	Upper Tiki Fm	Mudstone	
<i>Pelorocephalus mendozensis</i>	Carnian	Norian	237	208.5	Argentina	Cacheuta Fm.	"Siliciclastic"	The overall palaeoenvironm is fluviolacustr
<i>Pelorocephalus tenax</i>	Carnian	Norian	237	208.5	Argentina	Cacheuta Fm.	"Siliciclastic"	
<i>Pelorocephalus cacheutensis</i>	Carnian	Norian	237	208.5	Argentina	Cacheuta Fm.	"Siliciclastic"	
<i>Pelorocephalus ischigualastensis</i>	Carnian	Norian	237	208.5	Argentina	Ischigualasto Fm.	Siliciclastic	
<i>Keratobrachyops australis</i>	Induan	—	252.17	251.2	Australia	Arcadia Fm.	Mudstone	

2 Table -1: The table shows the spatial and temporal range of the family Chigutisauridae along with their paleoecology and paleoenvironment.
3 Abbreviations Max ma = Maximum age million years ago, Min ma = Minimum age million years ago, Fm. = Geological Formations from which the
4 specimen had been excavated. The table has been modified after data was acquired from the Paleobiology Database on 10th May 2022, using
5 the family name "Chigutisauridae". Data for the table was modified after Bonaparte 1957; Marsicano 1999; Dias – da- Silva and Sengupta 2012;
6 Warren 2006; Pledge 2013; Cabrera 1944; Rusconi 1949; Rusconi 1951; Sengupta 1995; Warren and Hutchinson 1983; Warren 1981; Warren and
7 Marsicano 2000)

Table 2(on next page)

Fossils excavated from the Late Triassic Tiki Formation (modified after Bandopadhyay and Ray 2020)

2a. Fish fossils excavated from the Late Triassic Tiki Formation (modified after Bandopadhyay and Ray 2020).

2b. Vertebrate fossil assemblage (tetrapod content) of the Late Triassic Tiki Formation of the Rewa Basin, India (modified after Bandopadhyay and Ray 2020).

1 **Table 2a**

Order/Family	Genus and Species	Order/Family	Genus and Species
Chondrichthyes		Osteichthyes	
Lonchididae	<i>Lonchidion estesi</i>	Ptychoceratodontidae	<i>Ceratodus sp.</i>
	<i>Lonchiodon incumbens</i>		<i>Ptychoceratodus oldhami</i>
	<i>Pristrisodus tikiensis</i>	Gnathorhizidae	<i>Gnathorhiza sp.</i>
Xenacanthidae	<i>Mooreodontus indicus</i>	Actinopterygii	Undescribed
	<i>Mooreodontus jaini</i>		
	<i>Tikiodontus asymmetricus</i>		

2 **Table- 2a:** Fish fossils excavated from the Late Triassic Tiki Formation (modified after Bandopadhyay and Ray 2020).

3 **Table 2b**

Order/Family	Genus and Species	Order/Family	Genus and Species
Amphibia		Diapsida	
Metoposauridae	<i>Panthesaurus maleriensis</i>	Archosauriformes	<i>Galtonia sp.</i> , <i>Protecovasaurus sp.</i> , and other intermediate forms
Chigutisauridae	<i>Compsocerops tikiensis</i>	Dinosauriformes	Undescribed Theropod-like (?) forms
Diapsida		Aetosauria	Undescribed
Phytosauria	<i>Volcanosuchus statisticae(?) leptosuchomorph</i>	Synapsida	

Rhynchosauria	<i>Hyperodapedon tikiensis</i>	Cynodontia	<i>Ruberodon roychowdhurii</i>
Rauisuchidae	<i>Tikisuchus romeri</i>	Mammaliaformes	<i>Tikitherium copei</i>
Rhynchocephalia	<i>Undescribed</i>		<i>Gondwanadon tapani</i>

4

5 **Table- 2b:** Vertebrate fossil assemblage (tetrapod content) of the Late Triassic Tiki Formation of the Rewa Basin, India (modified
6 after Bandopadhyay and Ray 2020).

Figure 1


Palatal surface of the skull photograph (RH01/Pal/CHQ/Tiki/15) (Kumar and Sharma 2019). Recreated with permission from the editor of the Palaeontological Society of India.

a. Shows the field photograph published in Kumar and Sharma (2019). b. Line drawing showing the disposition of the bones in the palatal surface of the skull published in Kumar and Sharma (2019). The abbreviation stated in the figure are as follows: cf = cultriform process, ect= ectopterygoid, itv = interpterygoid vacuity, pal= palatine, ps = parasphenoid, q = quadrate, stf = subtemporal foramen, v = vomer.



15 cm

Fig:1a



15 cm

Fig:1b

Figure 2

Dorsal surface of the skull roof of ISI A 202 *Compsoceroops tikiensis* sp. nov.

a. Reconstructed photograph of the dorsal surface of the skull roof of ISI A 202. Scale bar = 5 cm. b. Line drawing showing the disposition of the preserved bones in the dorsal part of the skull roof in ISI A 202. The abbreviation stated in the figure are as follows: j=jugal, mx=maxilla, n=nasal, p = parietal, po = postorbital, pof = postfrontal, pp = postparietal, prf = prefrontal, qj = quadratojugal, sq = squamosal, st = supratemporal, t = tabular. Scale bar = 5 cm.




Fig: 2a





Fig: 2b

Figure 3


Palatal surface of the skull of ISI A 202 *Compsocerops tikiensis* sp. nov.

a. Reconstructed photograph of the palatal surface of the skull of ISI A 202. Scale bar = 10 cm. b. Line drawing showing the disposition of the preserved bones in the dorsal part of the skull roof in ISI A 202. The abbreviation stated in the figure are as follows: ect = ectopterygoid, eo = eoccipital, pal = palatine, ps= parasphenoid, pt= pterygoid, pv= palatine vacuity, q= quadrate, stf = subtemporal foramen.



10 cm

Fig: 3a



10 cm

Fig: 3b

Figure 4

The poorly preserved occiput of ISI A 202 *Compsocerops tikiensis* sp. nov.




Figure 5

Dorsal view of poorly preserved left clavicle of ISI A 202 *Compsocerops tikiensis* sp. nov.




Figure 6

Geological Map showing the Tiki Formation, Rewa Basin, India.

Geological map of the Tiki Formation, Rewa Basin, India, modified after Mukherjee et al. (2012). The red dotted line shows the hypothetical faunal boundary.




Figure 7

Field photograph showing the exposure of *Compsocerops tikiensis* sp. nov. embedded in mudstone in the Upper part of the Tiki Formation. Image taken by the authors during field season 2019.



Figure 8

Lithologs of the Tiki Formation.

a. Litholog in the Jora Nala Section in the Lower Tiki Formation. b. Litholog in the Lower part of Tiki Formation modified after Mukherjee et al. (2012). c. Litholog of the Tiki Formation modified after Kumar and Sharma (2019).




Figure 9

Field photograph of the sand-mud alternation in the Tiki Formation near Tenduadh. Photograph taken by the authors during field season 2019.

a. Trough cross- bedded sandstone in the Tiki Formation b. Extensive mudstone in the Tiki Formation.



Figure 10

Geological map of the Maleri Formation, Pranhita-Godavari Valley Basin, India

Geological map modified after Kutty and Sengupta (1989), Dasgupta et al. (2017) showing the sand- mud alternations in the Maleri Formation, Pranhita- Godavari Valley Basin, India. The green line represents the faunal boundary that occurred due to the faunal turnover from the Carnian Lower Maleri to the Norian Upper Maleri Formation.




Figure 11

Litholog of the Maleri Formation modified after Kutty and Sengupta (1989).




Figure 12

Field photograph showing the abundance of red mudstone in the basal part of the Maleri Formation. Image taken by the authors during field season 2015 at Pranhita- Godavari Valley.

