

Adaptive changes in bodybuilders in conditions of different energy supply modes and intensity of training load regimes using machine and free weight exercises

Andrii Chernozub¹, Veaceslav Manolachi^{2,3}, Anatolii Tsos¹, Vladimir Potop^{3,4,5}, Georgiy Korobeynikov^{6,7}, Victor Manolachi^{2,3}, Liudmyla Sherstiuk⁸, Jie Zhao⁶, Ion Mihaila^{4,5}

¹ Lesya Ukrainka Volyn National University, Lutsk, Ukraine

² Dunarea de Jos University of Galati, Galati, Romania

³ State University of Physical Education and Sport, Chisinau, Republic of Moldova

⁴ Department of Physical Education and Sport, University of Pitesti, Pitesti, Romania

⁵ Doctoral School of Sports Science and Physical Education, University of Pitesti, Pitesti, Romania

⁶ National University of Physical Education and Sport, Kyiv, Ukraine

⁷ German Sport University Cologne, Institute of Psychology, Cologne, Germany

⁸ Petro Mohyla Black Sea National University, Mikolaiv, Ukraine

Corresponding Author:

Veaceslav Manolachi

Str. Domneasca no. 47, Galati, 800008, Romania

Email address: manolachivsciences@yahoo.com

Abstract

Background. The research was aimed at comparing the effect of using two types of training load_s different in intensity and energy supply. We studied the influence of the proposed load variations (machine and free weight exercises) on long-term adaptation of the body at the stage of specialized basic training in bodybuilding.

Methods. 64 athletes aged 18-20 years were examined. The research participants were divided into 4 groups, 16 athletes in each group. Athletes of group_s 1 and 3 used a complex of free weight exercises. Group 2 and 4 participants performed machine exercises. Bodybuilders of group 1 and 2 were trained in conditions of medium intensity training load ($R_a=0.58$) in the anaerobic-glycolytic mode of energy supply. Athletes of the 3rd and 4th groups used high

intensity load ($R_{a_} = 0.71$) in the anaerobic-alactate mode of energy supply. We managed to determine the nature of adaptation processes using methods of control testing of strength capabilities, bioimpedansometry, anthropometry, biochemical analysis of blood serum (LDH, creatinine).

Results. The study showed that the difference in the dynamics of the participants' maximum strength development (on example of chest muscles) did not depend on the content of machine or free weight exercises, but on the features of training load regimes. Thus, the controlled indicator of strength capabilities in athletes of groups 3 and 4 increased by 5.1% compared to groups 1 and 2. During all stages of the study, the indicators of the projectile working mass in athletes of groups 3 and 4 exceeded the results observed in groups 1 and 2 by 25.9%. At the same time, the amount of load in a set is on average 2 times higher in athletes of groups 1 and 2. Group 4 athletes, who used machine exercises and high intensity training load, increased the circumferential body measurements by 3.8 times (the chest), compared to the results recorded in group 1 athletes. Athletes of group 1 and 2 showed increasing in body fat by 3.4% compared to the initial level on the background of large load volume. The basal creatine level in bodybuilders of groups 3 and 4 increased by 3.7 times after 12 weeks of study, which indicates an accelerated growth of muscle mass.

Conclusion. The most pronounced adaptive body changes in bodybuilders at the stage of specialized basic training occurred during high intensity training load and anaerobic-alactate energy supply mode. Machine exercises contributed to increasing the morpho functional indicators of athletes more than free weight exercises.

Introduction

Improving the training process by optimizing loads is one of the main issues in bodybuilding. There is a constant search for effective combination of strength exercises and training programs. The issue of adaptation to a stressful stimulus in bodybuilding remains relevant. Adaptation changes are of great interest among leading specialists (*Aerenhouts & D'Hondt, 2020; Coratella et al., 2020; Heidel, Novak & Dankel, 2022*). This problem is especially acute at the stage of specialized basic training in bodybuilding. The complexity of the process of training optimization is associated with a high level of body resistance to physical exertion. Even if the load parameters contribute to the accelerated growth of muscle mass and strength (*Chernozub et al., 2020; Marshall et al., 2020; Schwanbeck et al., 2020*). At the same time, using a wide range of means, principles and methods of training in the previous stages complicates the mechanisms of increasing adaptation reserves in the shortest possible time and requires the search for new ways of improving the training process (*Chernozub et al., 2018; Saeterbakken et al., 2022*).

The impact of training loads different in volume and intensity on adaptation processes in untrained people is of constant interest among scientists. Scientists in sports physiology and fitness have been paying attention to the research of this problem in recent years (*Titova et al., 2018; Saeterbakken et al., 2019; Schott, Johnen & Holfelder, 2019*). The development of the integral method of quantitative estimation of load capacity in power fitness (*Chernozub et al.,*

2020, 2022) allowed to solve the long-standing problem of quantitative estimation of power load parameters depending on the conditions of muscle activity and the level of functional capabilities. The load factor (R_a) reflects the indicator of the load regime intensity in power sports. Indicators of maximum strength, duration of work in a set, amplitude of movement, duration of concentric and eccentric phases of movement, number of repetitions in a set, allow to determine this coefficient. Indicators of projectile working mass (m) and load volume in the set (Wn) demonstrate the magnitude of the physical stimulus depending on the training load regime. The effectiveness of using quantitative indicators for assessing the intensity of load regimes was studied in fitness and MMA (Chernozub *et al.*, 2020, 2022).

The complex use of morphometric and physiological indicators allows to evaluate the effectiveness of training loads and their adequacy to body capabilities. Blood biochemical markers helped to determine the nature of adaptive and compensatory reactions to a physical stimulus in different regimes of training loads. At the same time, the control of biochemical blood markers shows the level of resistance to training loads in certain modes of energy supply. It is known that the growth of maximum strength and muscle mass occurs after loads in anaerobic modes of energy supply (alactate and glycolytic). These changes occur due to hypertrophy of fast-twitch muscle fibers, growth of intra-muscular and inter-muscular coordination. The alactate and glycolytic anaerobic modes of energy supply differ in the mechanism of action, sources, duration and power (Shaner *et al.*, 2014; Parks *et al.*, 2020; Wilke, Stricker & Usedy, 2020; Wilk, Zajac & Tufano 2021). At the same time, the question of finding an effective combination of training load (machine or with free weight exercises) in bodybuilding that differ in terms of energy supply and intensity has not been investigated by scientists yet.

For many years, the question of determining the effectiveness of using machine or free weight exercises in bodybuilding, fitness, powerlifting has caused conflicting views among scientists, coaches and athletes (Wirth *et al.*, 2016; Wilke, Coratella *et al.*, 2020; Wilke, Stricker & Usedy, 2020). In recent years, a number of specialists have paid close attention to the study of this issue not only in athletes, but also in people who use complexes of similar strength exercises in the process of muscle activity during rehabilitation (Schott, Johnen & Holfelder, 2019; Marshall *et al.*, 2020). At the same time, the issue of determining the nature of changes in the concentration of testosterone, somatropin, and cortisol in the blood serum in response to training during machine and free weight exercises has been actively investigated (Shaner *et al.*, 2014; Schwanbeck *et al.*, 2020; González-Hernández *et al.*, 2022). The results of these studies contribute to the in-depth observation of the features of body adaptive and compensatory reactions to physical stimuli when using similar complexes of training exercises (Stajer, Vranes & Ostojic, 2018; Chernozub *et al.*, 2020; Saeterbakken *et al.*, 2022). However, despite the sufficient number of studies in this direction, there is no clear definition of informative biochemical blood markers for assessing the growth of muscle mass, increasing the functional capabilities of bodybuilders.

Thus, the main problem of bodybuilding is the lack of physiologically based mechanisms for the ratio of external stimulus indicators and adaptation processes. Another pressing issue at the stage of specialized basic training is the effective combination of training means and load modes because athletes have high level of resistance to power loads. The in-depth study of these problems will allow to improve the training process in bodybuilding.

The purpose of the study. To determine the impact of training loads different in intensity and energy supply on performance in bodybuilding; to study the peculiarities of adaptive body changes using machine and free weight exercises at the stage of specialized basic training.

Materials & Methods

Participants

64 athletes aged 18-20 years were examined. Their experience of training in bodybuilding is 4.2 ± 0.4 years. A series of studies was conducted over 12 weeks in 2021 on the basis of fitness centers Gold Gym, Fight House, Septem Fitness, Gym Style (Ukraine). The research participants were divided into 4 groups, 16 athletes in each group. Athletes of groups 1 and 3 used a complex [training regime which included](#) free weight exercises. Participants of groups 2 and 4 used machine exercises [for all training](#). However, bodybuilders of groups 1 and 2 trained in conditions of medium intensity training load ($R_a = 0.58$) in the anaerobic-glycolytic mode of energy supply. Athletes of the 3rd and 4th groups used high intensity training load ($R_a = 0.71$) in the anaerobic-alactate mode of energy supply.

The algorithm, structure and methods of the study were approved by the ethical committee for biomedical research of Lesya Ukrainka Volyn National University (ecbr18.08.2022) in accordance with the ethical standards of the Declaration of Helsinki. Participants gave written informed consent for the research in accordance with the recommendations of the biomedical research ethics committees (WHO, 2000).

Diagnostic equipment of the university medical and biological laboratory was used for conducting medical examination, assessment of functional status, indicators of body composition and biochemical control of creatinine concentration and LDH activity in the blood serum of the participants of the examined groups.

Maximal muscle strength

Monitoring the development of chest muscles maximum strength (1 RM) in the examined bodybuilders took place in four stages: at the beginning of the study and every 4 weeks. The control exercise “barbell press lying on a horizontal bench” was used to determine the maximum strength of the chest muscles of athletes performing free weight exercises. To determine the chest muscles maximum strength, we used the control exercise “bench press on the Smith simulator”. The procedure of control testing of the studied indicator 1 RM was carried out according to the generally accepted methodology for power sports with the observance of the appropriate exercise technique (e.g. Loturco et al., 2018).

Training load parameters

Using the integral method of quantitative estimation of load capacity in power fitness (e.g. Chernozub, 2018) was determined the load factor (R_a) at the beginning of the study. Its

parameters reflect the intensity level of the power load regime. At each of the four stages of control (at the beginning of the study and after every 4 weeks), average group indicators of projectile working mass (m) and load volume (W_n) were determined in each set.

Circumferential body measurements

Body circumference measurements (on the example of chest) were made using a centimeter tape following the generally accepted methodology. Controlled indicators were measured at the beginning and during the next 12 weeks of the study with an interval of 30 days.

Body composition

Determination of body composition indicators was carried out using the method of ~~bioimpedansometry~~ bio electrical impedance (BIA) followed by computer processing of the obtained results. Body composition parameters were recorded at the beginning and during the next 12 weeks of the study with an interval of 30 days. The method of ~~bioimpedansometry~~ BIA allowed to determine a large number of parameters, but during the research we controlled the content of fat-free mass (FFM, kg) and body fat (BF, %) in the study participants. The KM-AR-01 Diamond-AST diagnostic computerized hardware and software complex (VYUSK. 941118.001 RE) (e.g. *Martyrosov, Nikolayev & Rudnev, 2006*) was used to determine the studied indicators of body composition.

Biochemical parameters

The creatinine concentration and the activity of lactate dehydrogenase (LDH) in the blood serum of bodybuilders were determined by the kinetic method on the equipment of the company "High Technology Inc" (USA) with a set of reagents PRESTIGE 24i LQ LDH (Poland). Blood sampling was performed by a medical worker in compliance with the internationally accepted requirements for medical and biological research (e.g. *Tietz, Finley, & Pruden, 1995*). Control of changes in the parameters of the studied biochemical blood indicators took place at the beginning and after 12 weeks of training at rest (before exercise) and at the end of the training session.

Experimental design

The research was conducted in several stages:

At the *first stage*, machine and free weight exercises were selected to achieve the study objective. These complexes consisted of strength training exercises, which optimally engage the working muscle groups and allow to achieve maximum muscle fatigue in the required period of time. The machine exercises (using the example of basic exercises) were the following: Smith machine bench press, head off bench press, block thrust behind the head, standing cinder block triceps extension, Larry Scott curl, lying leg press. The free weight exercises (on the example of basic exercises) included: barbell chest press, seated barbell (dumbbell) bench press, bent-over row, French bench press, cross body hammer curl, barbell squat. Using the integral method of quantitative estimation of load capacity in power fitness (e.g. *Chernozub, 2018*), load modes of high ($R_a = 0.71$) and medium ($R_a = 0.58$) intensity were developed. Taking into account the duration of motor activity until complete fatigue of working muscle groups in a separate set, training load were performed in conditions of anaerobic-alactate or anaerobic-glycolytic modes of energy supply. During 12 weeks of research, 36 training sessions were conducted. The

training duration was 30-33 minutes. Not more than 3-4 muscle groups were loaded during one training. We used 1 basic and 2 isolated exercises for each group of muscles.

At the *second stage*, we studied the peculiarities of changes in maximum muscle strength development (using the example of chest muscles), circumferential body measurements (the chest), parameters of body composition (BF and FFM), indicators of projectile working mass (m) and load volume (W_n) in a set. The analysis of the studied morpho functional indicators allowed to clearly determine the nature of adaptive body changes in bodybuilders at this stage of training in conditions of different energy supply and intensity of training load regimes using machine and free weight exercises. We studied the character of changes in lactate dehydrogenase activity and creatinine concentration in the blood serum of athletes in response to a stress stimulus (training load) in order to determine the nature of the adaptive and compensatory body reactions.

At the *third stage*, the results of circumference measurements, indicators of fat-free mass and body fat were compared during 12 weeks of using the proposed machine and free weight exercises in conditions of different energy supply modes and intensity training load regimes, according to the nature of changes in creatinine basal level. We compared the results of changes in the concentration of LDH activity in the blood serum of athletes who used anaerobic-glycolytic and anaerobic-alactate modes of energy supply for muscle activity. The obtained results will allow to determine the optimal combination of using machine or free weight exercises on the background of high ($R_a=0.71$) or medium ($R_a=0.58$) training load intensity, which will contribute to the most effective increase in functional capabilities and muscle mass growth in bodybuilders at the stage of specialized basic training.

Statistical analysis

Statistical analysis of the research results was performed using the IBM *SPSS*Statistics 26 program package (StatSoftInc., USA). Median, lower and upper quartiles, interquartile range (IQR) were determined. Kruskal-Wallis H test was used for testing whether samples originate from the same distribution. Friedman's two-way analysis of variance by ranks was used to compare indicators of the same sample of subjects during the control period. Kendall's W (the Kendall's coefficient of concordance) is an effect size index for Friedman test. The Kolmogorov-Smirnov test was used to determine the normal distribution (e.g. Nasledov, 2013). The G-Power 3.1.96 program was used to calculate statistical power (determining the smallest sample size for the study). The sample size was evaluated using statistical tests: Wilcoxon signet-rank test (one sample case); ANOVA: Repeated measures, between factors. Statistical results are reported according to the APA style (e.g. Yağın et al., 2021).

Results

Table 1 presents the dynamics of the chest muscles maximum strength development (1 RM) in athletes of all 4 groups during 12 weeks of the study.

Insert here Table 1

Table 1 shows that during the period of research, there was positive dynamics in the chest muscles maximum strength development in bodybuilders of all groups. The most pronounced growth of the studied indicator was found in athletes of the 4th group (by 21.3%), who used

machine exercises with high intensity training load in conditions of anaerobic-alactate mode of energy supply. The smallest development of the chest muscle maximum strength was observed in group 2 athletes (by 15.1%) using free weight exercises in conditions of medium intensity training load and anaerobic-glycolytic mode of energy supply. At the same time, in groups of athletes in the conditions of using training load identical in intensity, but different sets of exercises, the average-group difference between the dynamics of the studied indicator is not significant.

Training load parameters

The results of the average group changes in the parameters of projectile working mass, used by bodybuilders during 12 weeks of the study, are graphically presented in Fig. 1.

Figure 1 demonstrates that despite almost identical parameters of the maximum muscle strength development in all groups at the beginning of the study (Table 1), the controlled indicators of the projectile working mass was by 22.7% higher in athletes of groups 3 and 4 who used high intensity training load ($R_a=0.71$) regardless of the type of exercises (Fig. 1). After 12 weeks of training, the bodybuilders of groups 3 and 4 showed a similar difference in the projectile working mass parameters compared to group 1 and 2 athletes, who used medium intensity training load ($R_a=0.58$).

Insert here Figure 1

Figure 2 depicts the results of average group changes in the load volume in a set. Due to the peculiarities of medium-intensity training load mode, group 1 and 2 athletes used 2 times higher volume of training load from the very beginning of the study. The last 4 weeks of research showed that group 1 and 3 representatives, who used the same type of free weight exercises on the background of different load modes, had a 175.4% difference between the groups in the load volume in a set. However, group 2 and 4 athletes, who used machine exercises in conditions of medium ($R_a=0.58$) and high ($R_a=0.71$) intensity training load, had an identical difference in the studied indicator.

Insert here Figure 2

Circumferential body measurements

The dynamics of the chest circumference measurements in the bodybuilders of examined groups is shown in Table 2.

Insert here Table 2

Table 2 clearly depicts that chest circumferential measurements were positively growing in athletes of all 4 groups during the study. But the largest increase in the studied indicator among the examined bodybuilders was in group 4 athletes (by 4.9%) using machine exercises with high intensity training load in the anaerobic-alactate mode of energy supply. At the same time, group 1 athletes, who used free weight exercises in conditions of anaerobic-glycolytic mode of energy supply and medium-intensity training load ($R_a=0.58$), showed an increase in the studied indicator by 1.3% for a similar period of time, comparing with participants of other groups.

Body composition

The changes in the indicators of fat-free mass and body fat of athletes who took part in 12-week research are given in Table 3.

Insert here Table 3

The results of bioimpedansometry showed that body fat index decreased greatly (by 4.0%) in group 1 athletes compared to the initial data. The index of fat-free mass in the same group of bodybuilders increased by 3.1% compared to the results recorded in representatives of other groups.

The fat-free mass index in group 4 athletes showed the largest increase of 5.7% among the examined bodybuilders over a period of 12 weeks of training. However, their body fat index did not change significantly over the same period of time.

Biochemical parameters

Fig. 3 graphically presents the changes in lactate dehydrogenase activity in the blood serum of all group participants using complexes machine and free weight exercises during all stages of the research.

Insert here Figure 3

Figure 3 shows that, LDH activity increased in the blood serum of bodybuilders in the following way: group 1 - by 16.6%, and group 2 - by 14.5%, compared to the state of rest. It happened in response to a stress stimulus at the beginning of the study. However, the activity of this enzyme in group 3 and 4 athletes in response to high intensity training load during machine exercises did not change significantly.

After 12 weeks of training, LDH activity in the blood serum increased with a smaller progression, and only in athletes of group 1 (by 13.1%) and 2 (by 6.8%) in response to medium intensity training load in anaerobic-glycolytic energy supply mode. At the same time, the studied enzyme activity in group 3 and 4 athletes did not change in response to training load in the conditions of anaerobic-alactate energy supply mode, regardless of using the type of strength exercises.

Fig. 4 displays the results of changes in the creatinine concentration in the blood serum of the examined bodybuilders in response to different modes of anaerobic energy supply performing machine and free weight exercises.

Insert here Figure 4

The results analysis shows that the initial measurements of creatinine concentration in the blood serum decreased in athletes of all groups in response to a stress stimulus. Thus, in representatives of groups 3 and 4, the concentration of the studied biochemical indicator shows an average 2-fold decrease in response to high intensity training load in the conditions of anaerobic-alactate energy supply mode, compared to the results recorded in bodybuilders of groups 1 and 2. At the same time, there was no significant difference between the effect of using machine or free weight exercises on the parameters of the studied biochemical indicator.

After 12 weeks of training, the creatinine concentration demonstrated changes in athletes of all examined groups in response to a stress stimulus. Thus, its concentration increased by 7.6%

($p < 0.05$) on average in athletes of all groups. The basal creatinine level was 3.7 times higher in athletes of groups 3 and 4. These changes in the basal level of creatinine indicate an accelerated growth of muscle mass in athletes of groups 3 and 4.

Discussion

This research describes the influence of training loads of different intensity and energy supply on performance in bodybuilding. It studies the process of adaptation to training loads (machine and free weight exercises) at the stage of specialized basic training. The study results indicate that there is no significant difference between machine and free weight exercises at this stage of training. ~~We have identified that it~~ is the high-intensity loads in combination with the anaerobic-alactate mode of energy supply that ~~contributes-contributed~~ to maximum adaptive body changes. The results of this study will contribute to the optimization of training loads at the stage of specialized basic training. The obtained results will also help to better understand the processes of adaptation in bodybuilding while using high and medium intensity training loads on the background of anaerobic modes of energy supply.

The lack of uniform mechanisms for training activity optimization is one of the unsolved problems in the modern training system of bodybuilding. The combination of effective load variations, principles, methods and tools is one of the most controversial scientific directions. In-depth study of adaptive body changes in bodybuilders is one of the priority areas of research in sports physiology (Tang, Chan & Kuo, 2014; Chernozub et al., 2020; Saeterbakken et al., 2022). Determining the peculiarities of adaptive body changes of bodybuilders using machine and free weight exercises in different modes of energy supply and their effectiveness is one of the directions of this problem. However, the majority of studies are specifically aimed at the comparative assessment of the effectiveness of using machine and with free weight exercises due to the dynamics of morpho functional indicators and changes in certain biochemical blood indicators (Wochyński & Sobiech, 2017; Aerenhouts & D'Hondt, 2020; Wilke, Stricker & Usedly, 2020). At the same time, leading scientists in this direction paid only partial attention to the course of adaptive changes in conditions of using these exercises on the background of different intensity of training load and modes of anaerobic energy supply allowing to increase the effectiveness in the shortest possible time while reducing the risk of adaptation failure.

The greatest increase in maximum muscle strength and circumferential measurements of bodybuilders occurred during high-intensity training loads ($R_a = 0.71$) in the anaerobic-alactate mode of energy supply. The rate of morpho functional indicators growth does not depend on type of exercises (machine or free weight). These adaptive changes are associated with accelerated hypertrophy of fast-twitch type B muscle fibers, which actively participate during exercises in anaerobic-alactate mode of energy supply (Joannis et al., 2020; Lopez et al., 2021). At the same time, the increase in the maximum muscle strength indicators is associated with growing indicators of intra-muscular and inter-muscular coordination in conditions of the anaerobic-alactate mode of energy supply (Wilk, Zajac & Tufano, 2021; Carvalho et al., 2022).

The results of bioimpedance measurement indicate that using free weight exercises and medium intensity training load significantly decreased the level of body fat with a minimal

increase in fat-free mass. Such changes in this body composition indicator highlights significant energy expenditure due to the large amount of work performed and the use of a significant number of stabilizer muscle groups during free weight exercises, as well as the possible manifestation of compensatory reactions to a physical stimulus (e.g. *Chernozub, et al., 2020*). The level of body fat of athletes performing machine exercises in the anaerobic-alactate mode of energy supply did not change during the research. However, the fat-free mass increased two times compared to the results of the participants using anaerobic-glycolytic energy supply mode for the same period, which indicates long-term adaptation on the background of pronounced hypertrophy of fast-twitch muscle fibers (e.g. *Moesgaard et al., 2022*).

The indicator of the projectile working mass was 25% higher in conditions of high intensity training load, and the parameters of load volume in a series of 4 sets were twice smaller compared to the results found during medium intensity training load despite identical initial parameters for the development of maximum muscle strength of the examined groups. At the same time, the use of machine or free-weight exercises did not affect the parameters of these load indicators. The characteristic difference in the projectile working mass values and performed volume load depends entirely on the power load regime and the variability of the parameters of its main components. (e.g. *Chernozub et al., 2018*). Thus, doubling the duration of concentric and especially eccentric phases of movement significantly reduces not only the level of inertia during exercise, but also the number of repetitions in a set, which is necessary to ensure the fatigue of the working muscles within the limits of the predominantly anaerobic-alactate mode of energy supply (*Chernozub et al., 2020; Kubo, Ikebukuro & Yata, 2021*).

The increase in lactate dehydrogenase activity in the blood serum of bodybuilders in response to medium-intensity training load are confirmed by the results of a series of studies in power fitness (e.g. *Titova et al., 2018*). A significant increase in LDH activity within the physiological norm in response to strength training indicates the accumulation of lactate in the blood of athletes during muscle activity and the activation of the predominantly anaerobic-glycolytic mechanism of ATP resynthesis (e.g. *Wochyński & Sobiech, 2017*). The level of LDH activity in response to high intensity training load during strength exercises, especially machine exercises, did not change both in conditions of short-term and long-term adaptation. This is due to the fact that in these conditions of muscle activity, energy supply occurs due to the creatine phosphokinase mechanism of ATP resynthesis and the absence of lactate in the blood (*Stajer, Vranes & Ostojic, 2018; González-Hernández et al., 2022*). The obtained results also indicated that there was no need to involve additional muscle groups during machine exercises, which are extremely necessary for stabilizing the body position in free weight exercises (e.g. *Chernozub et al., 2018*).

The changes in creatinine concentration did not depend on the type of exercises. ~~Rather, t~~The change of this blood biochemical marker ~~depends-depended upon~~ the intensity of loads and the mode of energy supply. The obtained data confirm the results of research in fitness (e.g. *Chernozub et al., 2020*). However, the increase in the basal level of creatinine concentration by 6 times indicate an increase in the adaptive reserves for the creatine phosphokinase mechanism of

ATP resynthesis during anaerobic-alactate mode of energy supply and high intensity training load (e.g. *Tang, Chan & Kuo, 2014*).

Thus, the obtained results indicate the expediency of using high intensity load ($R_a=0.71$) and anaerobic-alactate energy supply mode in combination with machine exercises at the stage of specialized basic training in bodybuilding. The study proved that adaptive body changes were the most pronounced on the background of accelerated growth of strength and muscle mass with minimal reduction in body fat (*Coratella et al., 2020; Heidel, Novak & Dankel, 2022*). Biochemical indicators of LDH activity and creatinine concentration in the blood serum are not only markers for evaluating adaptive body changes in conditions of strength load of various intensity (*Stajer, Vranes & Ostojic, 2018; Chernozub et al., 2020; González-Hernández et al., 2022*), but they also contribute to the development of new mechanisms for improving the training process. The optimization of training process takes place due to the correction of load volume components taking into account those processes that will take place in the body depending on the involvement of certain mechanisms of energy supply.

Conclusions

Using sets of machine exercises in combination with high intensity training load contribute to the growth of strength capabilities and muscle mass of bodybuilders at the stage of specialized basic training in the shortest possible time. The absence of additional involvement of stabilizer muscles in the process of performing machine exercises allows to reduce energy consumption in most cases, and using high intensity training load in the conditions of anaerobic-alactate energy supply mode will allow to increase the reserves of the creatine phosphokinase mechanism of ATP resynthesis and thus the efficiency of the long-term adaptation process as a whole. The implementation of the study results in the training activity will help to find effective ways of optimizing training load taking into account the adaptation potential of the body, the peculiarities of the training stages and muscle activity.

The studied blood biochemical markers were used for evaluating adaptive changes in bodybuilders in response to high and medium intensity training load while using machine and free weight exercises. A clear distinction of changes in LDH activity and creatinine concentration in the blood serum in response to this stress stimulus allows to determine the level of involvement of certain energy supply modes and energy recovery mechanisms during muscle activity, which in the perspective will help with developing innovative systems for load correction and optimization of the training process as a whole.

Acknowledgements

The article is a part of the planned scientific work “Development and implementation of innovative technologies and correction of the functional state of a person during physical activity in sports and physical therapy”, (state registration number 0117U007145) and Ministry of Education and Science of Ukraine project number 0118U000809.

References

- Aerenhouts D, D'Hondt E. 2020. Using Machines or Free Weights for Resistance Training in Novice Males? A Randomized Parallel Trial. *International Journal of Environmental Research and Public Health* 26;17(21):7848 DOI:10.3390/ijerph17217848.
- Carvalho L, Junior RM, Barreira J, Schoenfeld BJ, Orazem J, Barroso R. 2022. Muscle hypertrophy and strength gains after resistance training with different volume-matched load: a systematic review and meta-analysis. *Applied Physiology, Nutrition, and Metabolism* 47(4):357-368 DOI:10.1139/apnm-2021-0515.
- Chernozub A, Titova A, Dubachinskiy O, Bodnar A, Abramov K, Minenko A, Chaban I. 2018. Integral method of quantitative estimation of load capacity in power fitness depending on the conditions of muscular activity and level of training. *Journal of Physical Education and Sport* 18(1):217–221 DOI:10.7752/jpes.2018.01028.
- Chernozub A, Potop V, Korobeynikov G, Timnea OC, Dubachinskiy O, Ikkert O, Briskin Y, Boretsky Y, Korobeynikova L. 2020. Creatinine is a biochemical marker for assessing how untrained people adapt to fitness training load. *PeerJ* 8:e9137 DOI:10.7717/peerj.9137.
- Chernozub A, Manolachi V, Korobeynikov G, Potop V, Sherstiuk L, Manolachi V, Mihaila I. 2022. Criteria for assessing the adaptive changes in mixed martial arts (MMA) athletes of strike fighting style in different training load regimes. *PeerJ* 10:e13827 DOI:10.7717/peerj.13827.
- González-Hernández JM, Jiménez-Reyes P, Cerón JJ, Tvarijonaviciute A, Llorente-Canterano FJ, Martínez-Aranda LM, García-Ramos A. 2022. Response of Muscle Damage Markers to an Accentuated Eccentric Training Protocol: Do Serum and Saliva Measurements Agree? *Journal of Strength and Conditioning Research* 1;36(8):2132-2138 DOI:10.1519/JSC.0000000000003777.
- Coratella G, Tornatore G, Longo S, Esposito F, Cè E. Specific prime movers' excitation during free-weight bench press variations and chest press machine in competitive bodybuilders. *Eur J Sport Sci*. 2020 Jun;20(5):571-579. doi: 10.1080/17461391.2019.1655101
- Heidel KA, Novak ZJ, Dankel SJ. 2022. Machines and free weight exercises: a systematic review and meta-analysis comparing changes in muscle size, strength, and power. *The Journal of Sports Medicine and Physical Fitness* 62(8):1061-1070 DOI:10.23736/S0022-4707.21.12929-9.
- Joanisse S, Lim C, McKendry J, Mcleod JC, Stokes T, Phillips SM. 2020. Recent advances in understanding resistance exercise training-induced skeletal muscle hypertrophy in humans. *F1000Research* 9:F1000 Faculty Rev-141 DOI:10.12688/f1000research.21588.1.
- Kubo K, Ikebukuro T, Yata H. 2021. Effects of 4, 8, and 12 Repetition Maximum Resistance Training Protocols on Muscle Volume and Strength. *Journal of Strength and Conditioning Research* 35(4):879-885 DOI: 10.1519/JSC.0000000000003575.
- Lopez P, Radaelli R, Taaffe DR, Newton RU, Galvão DA, Trajano GS, Teodoro JL, Kraemer WJ, Häkkinen K, Pinto RS. 2021. Resistance Training Load Effects on Muscle

- Hypertrophy and Strength Gain: Systematic Review and Network Meta-analysis. *Medicine & Science in Sports & Exercise* 53(6):1206-1216 DOI: 10.1249/MSS.0000000000002585.
- Loturco I, Suchomel T, Kobal R, Arruda AS, Guerriero A, Pereira LA, Pai CN. 2021. Force-Velocity Relationship in three Different Variations of Prone Row Exercises. *Journal of Strength and Conditioning Research* 35(2):300–309
DOI:10.1519/JSC.0000000000002543.
- Marshall EM, Parks JC, Tai YL, Kingsley JD. 2020. The Effects of Machine-Weight and Free-Weight Resistance Exercise on Hemodynamics and Vascular Function. *International Journal of Exercise Science* 13(2):526-538
- Martyrosov EG, Nikolayev DV, Rudnev SG. 2006. *Technologies and methods of defining body composition*. Moscow: Nauka.
- Moesgaard L, Beck MM, Christiansen L, Aagaard P, Lundbye-Jensen J. 2022. Effects of Periodization on Strength and Muscle Hypertrophy in Volume-Equated Resistance Training Programs: A Systematic Review and Meta-analysis. *Sports Medicine* 52(7):1647-1666 DOI:10.1007/s40279-021-01636-1.
- Nasledov AD. 2013. *IBM SPSS statistics 20 and AMOS: professional statistical data analysis*. St. Petersburg: Piter.
- Parks JC, Marshall EM, Tai YL, Kingsley JD. 2020. Free-weight versus weight machine resistance exercise on pulse wave reflection and aortic stiffness in resistance-trained individuals. *European Journal of Sport Science* 20(7):944-952
DOI:10.1080/17461391.2019.1685007.
- Saeterbakken AH, Olsen A, Behm DG, Bardstu HB, Andersen V. 2019. The short- and long-term effects of resistance training with different stability requirements. *PLoS One* 14(4):e0214302 DOI:10.1371/journal.pone.0214302.
- Saeterbakken AH, Stien N, Pedersen H, Andersen V. 2022. Core Muscle Activation in Three Lower Extremity Exercises With Different Stability Requirements. *Journal of Strength and Conditioning Research* 36(2):304-309. DOI:10.1519/JSC.0000000000003465.
- Schott N, Johnen B, Holfelder B. 2019. Effects of free weights and machine training on muscular strength in high-functioning older adults. *ExpGerontol* 122:15-24
DOI:10.1016/j.exger.2019.03.012.
- Schwanbeck SR, Cornish SM, Barss T, Chilibeck PD. 2020. Effects of training with free weights versus machines on muscle mass, strength, free testosterone, and free cortisol levels. *Journal of Strength and Conditioning Research* 34(7):1851-1859
DOI:10.1519/JSC.0000000000003349.
- Shaner AA, Vingren JL, Hatfield DL, Budnar RG, Duplanty AA, DW Hill. 2014. The acute hormonal response to free weight and machine weight resistance exercise. *Journal of Strength and Conditioning Research* 28(4):1032-40 DOI:10.1519/JSC.0000000000000317.

- Stajer V, Vranes M, Ostojic SM. 2018. Correlation between biomarkers of creatine metabolism and serum indicators of peripheral muscle fatigue during exhaustive exercise in active men. *Research in Sports Medicine* 20:1–8 DOI:10.1080/15438627.2018.1502185.
- Tang FC, Chan CC, Kuo PL. 2014. Contribution of creatine to protein homeostasis in athletes after endurance and sprint running. *European Journal of Nutrition* 53(1):61–71 DOI:10.1007/s00394-013-0498-6.
- Tietz NW, Finley PR, & Pruden EL. 1995. *General clinical tests. Clinical guide to laboratory tests*. 3rd ed. Philadelphia: WB Saunders, 578.
- Titova AV, Chorniy OG, Dolgov AA, Gladyr TA. 2018. Parameters of biochemical control as a criteria of adaptive changes in the organism of athletes with various fitness levels engaged in the conditions of power fitness. *Ukrainian Journal of Medicine, Biology and Sports* 3(2):278–283 DOI:10.26693/jmbs03.02.278.
- WHO (2000) *World Health Organization. WHO Regional Office for Europe*; Copenhagen: 2000. Air quality guidelines for Europe, 2nd edition.
- Wilk M, Zajac A, Tufano J. 2021. The Influence of Movement Tempo During Resistance Training on Muscular Strength and Hypertrophy Responses: A Review. *Sports Medicine* 51(8):1629-1650 DOI:10.1007/s40279-021-01465-2.
- Wilke J, Stricker V, Usedly S. 2020. Free-Weight Resistance Exercise Is More Effective in Enhancing Inhibitory Control than Machine-Based Training: A Randomized, Controlled Trial. *Brain Sciences* 10(10):702 DOI:10.3390/brainsci10100702.
- Wirth K, Keiner M, Hartmann H, Sander A, Mickel C. 2016. Effect of 8 weeks of free-weight and machine-based strength training on strength and power performance. *Journal of Human Kinetics* 53:201-210 DOI:10.1515/hukin-2016-0023.
- Wochyński Z, Sobiech K. 2017. Impact of special aviation gymnastics instruments training on selected hormones in cadets' blood serum and plasma. *International Journal of Occupational Medicine and Environmental Health* 30(4):655-664 DOI:10.13075/ijomeh.1896.00904.
- Yağın B, Yağın F, Gözükar H, Çolak C. 2021. A web-based software for reporting guidelines of scientific researches. *Journal of Cognitive Systems* 6(1):39–43 DOI:10.52876/jcs.916182.