

Sodic stress differently influences physiological traits and anti-oxidant enzymes in pear and peach cultivars

Anshuman Singh^{Corresp., 1, 2}, Ashwani Kumar³, Parbodh Chander Sharma³, Raj Kumar³, Rajender Kumar Yadav³

¹ ICAR- Central Soil Salinity Research Institute, Karnal, Karnal, Haryana, India

² ICAR- Central Institute for Subtropical Horticulture, Lucknow, Uttar Pradesh, India

³ ICAR- Central Soil Salinity Research Institute, Karnal, Haryana, India

Corresponding Author: Anshuman Singh

Email address: anshumaniari@gmail.com

Background. The growth and physiological responses to sodic stress of pear and peach are poorly understood. Insights into how sodic stress alters tree physiology remain vital to developing salt tolerant scion and rootstock cultivars.

Methods. The effects of sodic stress (soil pH_s ~8.8) on tree growth and physiological traits of field grown trees of pear cultivars Punjab Beauty and Patharnakh, and peach cultivars Partap and Shan-e-Punjab were recorded using standard procedures. Sodic-induced changes in oxidative stressors, proline, anti-oxidant enzymes and leaf ions were measured to draw inferences.

Results. Sodic-induced reductions in vegetative growth were particularly marked in Patharnakh pear and Partap peach compared with other cultivars. Although sodic stress triggered a significant increase in leaf malondialdehyde (MDA) and hydrogen peroxide (H₂O₂), their levels relative to controls were much higher in peach than in pear; reflecting that peach suffered from greater oxidative stress. Interestingly, MDA and H₂O₂ levels did not seem to be deleterious enough to trigger proline-induced osmotic adjustment in pears. The activities of anti-oxidant enzymes strongly varied with the cultivar; specifically, the sodic-induced increases in CAT and SOD activities were much higher in Punjab Beauty pear and Shan-e-Punjab peach. Principal Component Analysis revealed an explicit convergence between CAT and SOD activities in Punjab Beauty and Shan-e-Punjab cultivars in response to sodic-induced oxidative stress. Correlation analysis revealed that leaf Na⁺ strongly inhibited tree growth in peach than in pear. Leaf K⁺ and proline were found to be the major osmolytes in sodic-stressed pear and peach cultivars, respectively.

Conclusions. We have for the first time studied the effects of sodic stress on important tree growth and physiological traits of commercially important pear and peach cultivars. Our findings revealed a marked suppressive effect of sodic stress on tree growth in peach than in pear. The sodic-induced upticks in leaf malondialdehyde, hydrogen peroxide and Na⁺ seemed to induce proline-mediated osmotic adjustment in peach but not in pear. The overall better sodic tolerance in pear compared to peach was ascribed to increased activities of anti-oxidant enzymes catalase and superoxide dismutase enzymes together with restricted Na⁺ uptake and better leaf K⁺ levels. Further investigations are needed to elucidate the effects of sodic stress on genetic and transcriptional changes, and on fruit yield and quality.

Sodicity stress differently influences physiological traits and anti-oxidant enzymes in pear and peach cultivars

Anshuman Singh^{1,2,*}, Ashwani Kumar¹, P.C. Sharma¹, Rajkumar¹, and R.K. Yadav¹

¹ICAR – Central Soil Salinity Research Institute, Karnal

²ICAR – Central Institute for Subtropical Horticulture, Lucknow

Corresponding author: Anshuman Singh

Email address: Anshuman.Singh@icar.gov.in

Abstract

Background. The growth and physiological responses to sodicity stress of pear and peach are poorly understood. Insights into how sodicity stress alters tree physiology remain vital to developing salt tolerant scion and rootstock cultivars.

Methods. The effects of sodicity stress (soil pH_s ~8.8) on tree growth and physiological traits of field grown trees of pear cultivars Punjab Beauty and Patharnakh, and peach cultivars Partap and Shan-e-Punjab were recorded using standard procedures. Sodicity-induced changes in oxidative stressors, proline, anti-oxidant enzymes and leaf ions were measured to draw inferences.

Results. Sodicity-induced reductions in vegetative growth were particularly marked in Patharnakh pear and Partap peach compared with other cultivars. Although sodicity stress triggered a significant increase in leaf malondialdehyde (MDA) and hydrogen peroxide (H₂O₂), their levels relative to controls were much higher in peach than in pear; reflecting that peach suffered from greater oxidative stress. Interestingly, MDA and H₂O₂ levels did not seem to be deleterious enough to trigger proline-induced osmotic adjustment in pears. The activities of anti-oxidant enzymes strongly varied with the cultivar; specifically, the sodicity-induced increases in CAT and SOD activities were much higher in Punjab Beauty pear and Shan-e-Punjab peach. Principal Component Analysis revealed an explicit convergence between CAT and SOD

activities in Punjab Beauty and Shan-e-Punjab cultivars in response to sodicity-induced oxidative stress. Correlation analysis revealed that leaf Na^+ strongly inhibited tree growth in peach than in pear. Leaf K^+ and proline were found to be the major osmolytes in sodicity-stressed pear and peach cultivars, respectively.

Conclusions. We have for the first time studied the effects of sodicity stress on important tree growth and physiological traits of commercially important pear and peach cultivars. Our findings revealed a marked suppressive effect of sodicity stress on tree growth in peach than in pear. The sodicity-induced upticks in leaf malondialdehyde, hydrogen peroxide and Na^+ seemed to induce proline-mediated osmotic adjustment in peach but not in pear. The overall better sodicity tolerance in pear compared to peach was ascribed to increased activities of anti-oxidant enzymes catalase and superoxide dismutase enzymes together with restricted translocation of Na^+ uptake and better leaf K^+ levels. Further investigations are needed to elucidate the effects of sodicity stress on genetic and transcriptional changes, and on fruit yield and quality.

INTRODUCTION

Presently, about 1 billion hectare global land area is affected by salinity and sodicity problems to varying extents (Hopmans et al., 2021). Out of this, ~76 million hectare area is impacted by the human-induced secondary salinization and sodification (Hossain, 2019). The severity of the problem is evidenced by the fact that approximately 1.5 million hectares of land become unsuitable for agricultural production every year because of salinity and related problems (Hossain, 2019). Available evidence suggests that salt sensitive horticultural crops including pear (Ahmad et al., 2022) and peach (Hernández et al., 2021) are particularly adversely affected under saline and sodic conditions. In contrast to the commonly accepted threshold of 8.5 for the pH of

saturated soil paste (pH_s), $pH_s > 8.2$ seems to be more realistic for classifying the soils as sodic under Indian conditions: soil pH_s of 8.2 is often associated with an exchangeable sodium percentage (ESP) of 15; a limit above which the soils are generally considered to be sodic (Abrol et al., 1988). Generally, soil pH_s of 8.5 roughly corresponds to ESP of ~ 50 , high enough to suppress the crop growth (Sharma et al., 2016; Abrol et al., 1988). Although alkali salts such as sodium carbonate (Na_2CO_3) are often more detrimental to plant growth than the neutral salts such as sodium chloride ($NaCl$) (Yang et al., 2009), even in fairly salt tolerant crops (Abbas et al., 2021), alkali stress research continues to receive little attention. Sodic soils in the Trans-Gangetic Plains of India (study area) mostly have predominance of highly soluble Na_2CO_3 and $NaHCO_3$ salts, and are thus prone to the abrupt increases in the soil pH (Mandal, 2012).

In sodic soils, excessive Na^+ causes the clay dispersion, surface crusting and deterioration in the soil physical properties (Qadir et al., 2007). Besides poor physical properties, high pH, osmotic and ionic stresses, and nutrient deficiencies are other limitations to plant growth in the sodic soils (Qadir & Schubert, 2002). Additionally, calcium carbonate ($CaCO_3$) concretions in the sub-soil also hamper plant establishment (Sharma et al., 2016). Considerable spatial variations in soil pH are also frequently seen in sodic soils; the more sodic parts of the field are often less congenial for crop growth (Samra et al., 1988). In fruit crops, the adverse effects of sodicity stress on plant growth (Saxena & Gupta, 2006; Krishnamoorthy, 2009) are ascribed to altered plant water relations (Li et al., 2020), reduced levels of photosynthetic pigments (Krishnamoorthy, 2009; Li et al., 2020), lipid peroxidation and oxidative stress (Ahmad et al., 2014), and ionic stress (Singh et al., 2018b). Sodicity-stressed plants accumulate osmolytes such as proline for the osmotic adjustment (Krishnamoorthy, 2009; Ahmad et al., 2014; Singh et al.,

2018a), and also activate the antioxidant enzymes for scavenging the free radicals (Ahmad et al., 2014).

Although sodic soils display noticeable improvements in the physico-chemical properties following amendment application and salt leaching with the fresh water, such improvements are mostly transient and confined to the top soil (<15 cm) (Sharma & Singh, 2019) such that sub-soil constraints continue to persist (Kumar et al., 2019). Under such conditions, agronomic practices such as planting into amendment-treated auger-holes often give better results (Gill & Abrol, 1991; Saxena & Gupta, 2006). As a majority of fruit crops are mostly highly sensitive to salinity and sodicity stresses (Singh et al. 2018a), development of salt tolerant scion and rootstock cultivars is absolutely essential to sustain the fruit production in salt-affected soils (Mahmoud et al., 2020). Although salt tolerance is a complex polygenic trait, and is greatly influenced by the genetic and environmental cues (Flowers, 2004), there exists ample genotypic variation that needs to be explored for identifying the salt tolerant genotypes (Mahmoud et al., 2020; Singh et al. 2018a).

Although some studies have shown the adverse effects of salinity stress on tree growth and physiology of pears (Myers et al., 1995; Musacchi et al., 2006; Oron et al., 2002) and peaches (Boland et al., 1993; Karakas et al., 2000; Soliman et al., 2017), their responses to sodicity stress remain elusive; excluding some preliminary observations that high soil pH and the related sub-soil constraints may suppress the plant growth (Elkins et al., 2012; Abd-Elmegeed et al., 2013; Mestre et al., 2015; Tagliavini et al., 1995). However, the soil pH levels in these studies were rather low to draw reasonable inferences, and the plausible physiological changes accounting for the reduced plant growth were also not investigated. Plant responses to salt stress often strikingly vary with the experimental conditions; for instance, depending on

experimental conditions, pears are either highly sensitive (Ebert, 1999) or moderately tolerant (Musacchi et al., 2006) to the salinity stress. Interestingly, most of the aforementioned studies had used NaCl as the sole salinizing agent in relatively controlled short-term experiments; the results may be altogether different when a different salt is used to induce the salt stress (Grieve et al., 2012). While an interplay among the oxidative stressors such as malondialdehyde, osmolytes, anti-oxidant enzymes and the leaf ions is known to greatly influence the plant responses to salinity stress in pears and peaches (Erturk et al., 2007; Dejampour et al., 2012; Wu & Zou, 2009; Yousefi et al., 2019), such responses remain uninvestigated under sodic conditions. Insights from other fruit crops [e.g., *Malus halliana* (Jia et al., 2019) and *Morus alba* (Hui-Hui et al., 2019)] also suggest that plant physiological responses to salinity and sodicity stresses remarkably vary with each other: the effects of salinity cannot be used as a reliable proxy for the effects of sodicity.

The subtropical areas in the Trans-Gangetic Plains of India (study region) usually have a frost free spring and adequate chilling availability for the cultivation of low chill cultivars of temperate fruits such as pear and peach. Pear cultivars Patharnakh [*Pyrus pyrifolia* (Burm F.) Nakai] and Punjab Beauty (*Pyrus communis* L.) (Singh et al., 2015), and peach cultivars Partap and Shan-e-Punjab [*Prunus persica* (L.) Batsch] (Bhatnagar and Kaul, 2002) are particularly suitable for cultivation in this area given their commercial potential, low chilling requirement and the ability to tolerate the high summer temperatures typical of the region. To our knowledge, systematic studies have not yet been carried out to evaluate the effects of sodicity stress on tree growth and physiological relations of pears and peaches. This study intended to delineate sodicity-induced changes in tree growth, leaf oxidative stress markers,

proline, enzymatic anti-oxidants and leaf ions in pear and peach cultivars, and how these changes influence the overall cultivar-specific tree responses to sodicity stress.

MATERIALS & METHODS

Location

The experiment was conducted between January, 2015 and April, 2019 at the experimental farm of Indian Council of Agricultural Research-Central Soil Salinity Research Institute, Karnal, India (29° 42' 20.6" N, 76° 57' 19.80" E, 243 m above mean sea level). The region has a semi-arid subtropical climate with hot summers and dry winters. The long-term average annual rainfall is ~700 mm. A sodic field was used to evaluate the pear and peach cultivars.

Experimental material

The pear cultivars Punjab Beauty (*Pyrus communis* L.) and Patharnakh [*Pyrus pyrifolia* (Burm F.) Nakai] grafted on Kainth (*Pyrus pashia* Buch.-Ham. Ex D. Don); and peach cultivars Partap and Shan-e-Punjab [*Prunus persica* (L.) Batch] on Sharbati rootstock [*P. persica* (L.) Batch] were evaluated. The bare root plants were planted on 17th January 2015 in auger-holes (diameter 20 cm, depth 120 cm) in a sodic field, keeping the graft joint ~15 cm above the surface. The auger-holes were refilled with a mixture of the original soil and 5 kg of gypsum (Ca₂SO₄.2H₂O) before planting for better plant establishment (Sharma et al., 2014). The square system of planting was used, with between- and within-row spacings of 6 m each in both pears (Sangwan et al., 2015) and peaches (Kanwar and Singh, 2004). Trees were trained to the modified leader system, and the recommended management practices were followed for better

tree growth. Irrigation water was applied through 1 m wide channels. Plants were irrigated at the weekly intervals during summer months and at fortnightly intervals during July- August.

Treatments

Soil samples were collected from 24 random points of the field, from four depths (0-15, 15-30, 30-60 and 60-100 cm) using an auger. After air drying, the samples were ground and sieved (2.0 mm sieve) for determining the pH of saturated soil paste (pH_s) and soil saturation extract electrical conductivity (EC_e) using digital pH and conductivity meters (Eutech, Singapore). Based on soil analysis, the experimental plants were grouped into control (mean soil $pH_s = 8.22$, $EC_e = 0.71$ dS m^{-1}) and sodic ($pH_s = 8.80$, $EC_e = 0.94$ dS m^{-1}) treatments for recording the observations. Both soil pH_s and EC_e increased with depth, and there were significant differences between control and sodicity treatments (pH_s $F = 41.94$, $p < 0.001$; EC_e $F = 27.52$, $p < 0.001$) (*Table S1*). The groundwater used in irrigation had the following composition: electrical conductivity- 0.65 dS m^{-1} , pH- 8.04, Na^+ - 2.41 me L^{-1} , K^+ - 0.15 me L^{-1} , $Ca^{2+} + Mg^{2+}$ - 4.17 me L^{-1} , Cl^- - 0.98 me L^{-1} and HCO_3^- - 4.31 me L^{-1} .

Tree growth

Four trees of each pear and peach cultivar representing the control and sodic treatments were randomly tagged for recording the observations. Tree height and trunk diameter were measured during the last week of April, 2019. Trunk diameter readings, recorded using a digital Vernier caliper (Mitutoyo, Japan) 15 cm above the graft union, were converted into trunk cross sectional area (TCSA) by the formula: $TCSA = \pi(d/2)^2$; where d= mean of east-west and north-south trunk diameters. The canopy volume (CV) was computed by the formula: $CV = (w^2 \times h)/2$; where w= canopy diameter in east-west and north-south directions; and h= tree height.

157 Leaf physiological traits

158 The fully expanded leaves from the middle of the shoots were collected from all the
 159 directions of each replicate tree (n= 4), pooled, packed in zip lock bags inside an ice-box, and
 160 immediately brought to the laboratory. The total leaf chlorophyll (TC) was estimated by
 161 overnight incubation of 200 mg chopped leaves in 80% acetone (Arnon, 1949) using a
 162 spectrophotometer (UV 3200, Lab India Analytical, India). Lipid peroxidation, in terms of
 163 malondialdehyde (MDA) content, was estimated by the method of Heath & Packer, (1968). One
 164 g of leaf tissue was homogenized in 5 ml of 0.1 per cent trichloroacetic acid (TCA, w/v) and
 165 centrifuged at 8000 g for 15 min. Supernatant (1 ml) was precipitated in 4 ml of 20% (w/v) TCA
 166 containing 0.5% (w/v) 2-thiobarbituric acid. The reaction mixture was heated in a water bath (at
 167 95°C for 30 min) with constant stirring and quickly cooled in the ice bath. Sample was
 168 centrifuged at 8000 x g for 10 min, absorbance of the supernatant was measured at 532 nm
 169 against distilled water, and the value for non-specific absorption at 600 nm was subtracted. The
 170 MDA concentration was calculated by using the molar extinction coefficient of 155 mM⁻¹ cm⁻¹.
 171 Hydrogen peroxide (H₂O₂) content was estimated using the procedure described in Sinha,
 172 (1972). The leaf tissue was macerated in 5 ml of ice cold 0.01 M phosphate buffer (pH 7.0) and
 173 centrifuged at 8000 g for 10 min. 1.95 ml of 0.01 M potassium phosphate buffer (pH 7.0)
 174 followed by 5% potassium dichromate (2 ml) and glacial acetic acid (1:3; v/v) were added to 50
 175 µl supernatant. The sample tubes were kept in the boiling water bath for 10 min and then cooled.
 176 The absorbance was read at 570 nm against the reagent blank without sample extract and the
 177 H₂O₂ content was calculated from the standard calibration curve (10 to 160 µmol of H₂O₂).
 178 Proline was extracted using 200 mg leaf tissue homogenized in 10 ml of 3% sulphasalicylic
 179 acid (Bates et al., 1973). Two ml of extract was reacted with 2 ml each of acid-ninhydrin and

glacial acetic acid for 1 hour at 100°C, and the reaction was terminated in water bath. The reaction mixture was extracted with 4 ml toluene, mixed vigorously and the chromophore containing toluene was aspirated from the aqueous phase. The absorbance was measured at 520 nm using toluene as blank.

Anti-oxidant enzymes

Fresh leaf samples (~250 mg) were homogenized in 0.1 M phosphate buffer (pH 7.5) containing 5% polyvinyl polypyrrolidone (w/v), 1 mM ethylenediamine tetraacetic acid, and 10 mM b-mercapto-ethanol for assaying the ascorbate peroxidase (APX, EC 1.11.1.11) and superoxide dismutase (SOD, EC 1.15.1.1) activities. The APX activity was assayed as described in Nakano & Asada, (1981); enzyme activity was calculated using extinction coefficient of 2.8 mM⁻¹ cm⁻¹. The SOD assay was performed following the methodology given in Beauchamp & Fridovich, (1971). The absorbance of the solution was measured at 560 nm with a UV-VIS spectrophotometer (UV 3200, Lab India Analytical, India). One unit of SOD was calculated as the amount of enzyme required to inhibit the photo-reduction of one mmol of nitroblue tetrazolium. Catalase (CAT, EC 1.11.1.6) and peroxidase (POX, EC 1.11.1.7) were extracted in 0.01 M phosphate buffer (pH 7.5) with 3% polyvinyl polypyrrolidone (w/v) by homogenizing the fresh leaf tissue (1.0 g). The homogenate was centrifuged at 4°C for 15 min at 10,000 x g and the clear supernatant was used for the assay. The CAT activity was determined as the disappearance of H₂O₂ at 240 nm (25°C) for 1 min (Aebi, 1984). The POX was assayed by determining the rate of guaiacol oxidation in the presence of H₂O₂ at 470 nm (Rao et al., 1998). One unit of POX activity was defined as the amount of enzyme required to oxidize one nmol of guaiacol min⁻¹ ml⁻¹.

Leaf Na⁺ and K⁺

The leaf samples were first dried to a constant weight at 60° C in a hot air oven (Narang Scientific Works, India). The samples were then finely ground using a hammer mill. One hundred mg of the powdered sample was digested in the di-acid [nitric acid (HNO₃) and perchloric acid (HClO₄); 3:1] mixture for determining the Na⁺ and K⁺ contents (mg g⁻¹ DW) using a flame photometer (Systronics India).

Statistical analysis

The experiment was laid out in a randomized block design. The main and interaction effects of independent factors (treatment and cultivar) on each dependent variable were examined by a two-way Analysis of Variance (ANOVA). The assumptions of equality of variances (Levene's test) and normality (quantile-quantile plot) were checked prior to the ANOVA, and some variables (tree height, proline and peroxidase in case of peach data) were log-transformed to improve the ANOVA assumptions. The effect size measure omega squared (ω^2) was computed to estimate the variance in the response variable(s) accounted for by the explanatory variables. The use of sample size independent effect size measures such as ω^2 improves the statistical power by reducing the risk of Type II error (Altman, 2000). Tukey's test ($p < 0.05$) was used for mean comparisons (JASP v. 0.15). The data are expressed as mean (n=4) \pm standard deviation (SD). Principal Component Analysis (PCA) (Bartlett's test of sphericity, $p < 0.001$) was applied to reduce the dimensionality and to detect the key patterns in data (Jamovi v. 2.2). Pearson's bivariate correlations between the measured traits were computed (Julkowska et al., 2019).

RESULTS

Analysis of Variance

The results for the Analysis of Variance (ANOVA) revealed the strong repressive effects of sodicity stress on tree growth and physiological traits in both pear and peach. Although sodicity-induced reductions in tree height (TH), trunk cross sectional area (TCSA) and canopy volume (CV) were highly significant ($p < 0.001$) in both the crops, a perusal of the effect-size measure (ω^2) values implied that TCSA was far less sensitive to sodicity stress than were both TH and CV; regardless of the crop (*Table 1*). Likewise, based on ω^2 values, a more adverse effect of sodicity stress was apparent on peach than on pear growth. The ω^2 values were low-to-moderate (< 0.600) for the most leaf physiological traits, but relatively high (> 0.700) for leaf proline, ascorbate peroxidase (APX) and Na^+ in pear. This suggested that explanatory variable (i.e., sodicity stress) accounted for a reasonably high variance in leaf proline, APX and Na^+ . Pear cultivars differed markedly with each other for all the traits except leaf proline, APX and Na^+/K^+ ratio (*Table 1*). The sodicity-triggered increases in leaf malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) levels were far greater in peach compared to pear in terms of ω^2 values. This, together with more or less similar values of ω^2 for the leaf proline, implied the greater sensitivity to oxidative stress of peaches than of pears. Contrarily, the values of ω^2 evinced moderate-to-strong increases in APX, peroxidase (POX) and catalase (CAT) activities in pear but not in peach; while upregulation in the superoxide dismutase (SOD) activity was quite similar in both the cases. This again indicated a better anti-oxidant system to cope with the free radicals in pear than in peach (*Table 1*).

Tree growth

There were strong differences between the cultivars for sodicity-induced reductions in tree growth. For instance, the reductions in trunk cross sectional area (TCSA) and canopy volume (CV) relative to controls were much lower in Punjab Beauty pear (16.49% and 44.50%,

respectively) than in Patharnakh (41.89% and 69.28%, respectively) (*Table 2*). Sodicity stressed Partap peach trees displayed much higher reductions in tree height (TH, 36.24%), TCSA (74.28%) and CV (90.33%) than corresponding decreases of 29.23%, 11.94% and 58.58% in the cultivar Shan-e-Punjab (*Table 2*).

Leaf physiological traits

Sodicity stress caused appreciable reductions in total leaf chlorophyll (TC) in pear (Punjab Beauty- 26.95%, Patharnakh- 21.43%). Leaf malondialdehyde (MDA), hydrogen peroxide (H_2O_2) and proline levels were invariably higher in sodic (pH_s 8.8) than in control (pH_s 8.2) treatment. Leaf MDA increased marginally (9.61%) in Punjab Beauty and moderately (19.92%) in Patharnakh (*Table 3*). Both the cultivars showed identical increases (~12.0%) in leaf H_2O_2 . Leaf proline accumulation in sodic treatment (pH_s 8.8) was considerably higher in Punjab Beauty (33.86%) compared to Patharnakh (20.15%). Partap and Shan-e-Punjab peaches had 30.80 and 18.09% less TC, respectively, in sodic soils (pH_s 8.8) than in controls (pH_s 8.2) (*Table 3*).

Leaf anti-oxidant enzymes

Sodicity-triggered increases in ascorbate peroxidase (APX) and catalase (CAT) activities relative to controls were more pronounced in pear cultivar Patharnakh (34.49 and 35.97%, respectively) than in Punjab Beauty (28.44 and 25.16%, respectively) (*Table 4*). Contrarily, peroxidase (POX) and superoxide dismutase (SOD) activities were 2.6- and 1.4-folds higher, respectively, in sodicity-stressed Punjab Beauty leaves than in Patharnakh (*Table 4*). However, in absolute terms, only POX activity was higher in sodicity-stressed Patharnakh while both CAT and SOD activities much higher in Punjab Beauty. In case of peach, the sodicity-triggered upregulation in leaf APX and POX activities relative to controls were markedly higher in Shan-

e-Punjab (29.41 and 22.99%, respectively) than in Partap (13.49 and 6.24%, respectively). Conversely, CAT and SOD activities were more noticeable in the cultivar Partap (38.82 and 16.19%, respectively) compared to the cultivar Shan-e-Punjab (8.37 and 14.79%, respectively). Nonetheless, Shan-e-Punjab significantly outperformed Partap for the absolute leaf CAT and SOD levels in the sodic soils (pH_s 8.8) (*Table 4*).

Leaf Na⁺ and K⁺

As expected, sodicity stress caused an increase in leaf Na⁺ and a decrease in K⁺, regardless of the cultivar. Pear cultivars Punjab Beauty and Patharnakh displayed considerably higher leaf Na⁺ (43.91 and 74.57%, respectively) in sodic (pH_s 8.8) than in control (pH_s 8.2) soils. In contrast, leaf K⁺ declined significantly in sodicity stressed Punjab Beauty (17.76%) and Patharnakh (28.69%) trees (*Table 5*). In sodic soils (pH_s 8.8), peach cultivars Partap and Shan-e-Punjab had 71.68 and 58.24% more leaf Na⁺, respectively, and 31.97 and 24.21% less K⁺, respectively (*Table 5*).

Principal component analysis

The Principal Component Analysis (PCA) was quite efficient in reducing the dimensionality: the first two Principal Components (PCs) (Eigen value >1.0) alone explained 90.95% of the cumulative variance in data in pear (*Table S2*; Figure 1), and 94.61% in peach (*Table S2*; Figure 2). In case of pear, leaf Na⁺, K⁺ and Na⁺/K⁺ ratio alongside proline, APX and CV were the highly weighted variables on PC1; and TH, TCSA, HP, MDA and CAT on PC2 (*Table S2*). Likewise, for peach, TH, HP, proline and Na⁺/K⁺ ratio were the best represented variables on PC1, and TCSA, APX, POX, CAT and K⁺ on PC2 (*Table S2*). Interestingly, PCA achieved a clear-cut discrimination for the cultivar- and treatment-specific effects in data: while

PC1 unambiguously distinguished the control and sodicity treatments, PC2 clearly separated the tested cultivars from each other (Figure 1, 2). The PCA also unveiled some interesting patterns in data. For pear, tight clustering of tree growth attributes, K and TC in the upper left quadrant implied a straightforward role of leaf K^+ in osmotic adjustment in the sodicity-stressed pears (Figure 1). Likewise, there was an apparent cultivar-specific upregulation in CAT and SOD activities in response to the oxidative (H_2O_2 and MDA) and ionic (Na^+) stresses in Punjab Beauty trees; increased POX activity alone seemed to have alleviated salt stress in cultivar Patharnakh (Figure 1). In the case of peach, leaf K^+ did not have any obvious effect in alleviating the salt stress. Contrarily, osmoregulation through proline apparently increased in response to increasing HP, MDA and Na^+ levels. Of the enzymatic anti-oxidants, APX and POX were rather specific to the cultivar Partap, and CAT and SOD to the cultivar Shan-e-Punjab (Figure 2).

Correlation analysis

In pear, tree growth traits and total leaf chlorophyll (TC) had strong positive correlations with leaf K^+ ($r > 0.800$, $p = 0.000$). Expectedly, leaf Na^+/K^+ ratio had moderate-to-strong negative correlations with TH ($r = -0.623$, $p = 0.010$), TC ($r = -0.602$, $p = 0.014$) and CV ($r = -0.788$, $p = 0.000$) (Table S3, Figure 3). A strong positive correlation between leaf MDA and H_2O_2 ($r = 0.914$, $p = 0.000$) reflected their synergistic adverse effects on pear tree growth and physiology. Leaf proline had moderate positive correlations with MDA ($r = 0.578$, $p = 0.019$) and H_2O_2 ($r = 0.519$, $p = 0.039$). Similarly, strong positive relationships ($r > 0.800$, $p = 0.000$) were found between antioxidants (CAT and SOD) and oxidative stress indicators (MDA and H_2O_2) (Table S3, Figure 3). In peach, leaf K^+ had strong positive correlations with TC ($r = 0.887$, $p = 0.000$), APX ($r = 0.732$, $p = 0.001$) and POX ($r = 0.791$, $p = 0.000$), strong negative correlations with CAT ($r = -0.987$, $p = 0.000$) and SOD ($r = -0.839$, $p = 0.000$), and a moderate negative correlation with

H_2O_2 ($r = -0.655$, $p = 0.006$) (Table S4, Figure 4). In contrast to pear, elevated leaf Na^+ had strong inhibitory effects ($r > -0.880$, $p = 0.000$) on tree growth in peach. The adverse effects of both MDA and HP on peach tree growth were also apparent ($r > -0.800$) (Table S4, Figure 4).

DISCUSSION

Sodic stress is a serious constraint on pear and peach productivity (Elkins et al., 2012; Mestre et al., 2015). The observation that sodic stress impairs pear and peach growth (Thind and Mahal, 2020) is anecdotal; key physiological traits underpinning sodic tolerance remain elusive. In this backdrop, we aimed to understand the effects of sodic stress on tree growth and physiological traits in commercially important pear and peach cultivars.

We found that peaches in general were more adversely affected than pears. Moreover, sodic stress had a greater adverse effect on Patharnakh pear and Partap peach than other tested cultivars. The experimental soils used in this study still have $\text{pH} > 8.5$, particularly below 30 cm depth. Additionally, a clayey sub-soil further limits the water flux and nutrient availability; hampering the plant growth (Kumar et al., 2019). This may partly explain the differential responses to sodic stress of the crops and cultivar tested by us (Samra et al., 1988). Excess salts suppress the vegetative growth and biomass production in pears (Matsumoto et al., 2006; Okubo et al., 2000) and peaches (Massai et al., 1997, 2004); albeit in a genotype-specific manner such that some cultivars are more adversely affected than others (Krishnamoorthy, 2009). The depletion of leaf chlorophyll may be caused by the lime-induced chlorosis when pears (Ma et al., 2005) and peaches (Thomidis & Tsipouridis, 2005) are grown in the high pH soils. The sodic soils of the study area are often deficient in iron (Kaledhonkar et al., 2019), a key element in chlorophyll synthesis. Furthermore, high soil pH may inhibit the uptake of Mg^{2+} and Fe^{2+} ions

needed for chlorophyll synthesis (Jia et al., 2019). The presence of bicarbonate (NaHCO_3 , predominant salt in the study area soils; Mandal, 2012) suppresses iron availability to the peach (Molassiotis et al., 2005) and pear (Valipour et al., 2020) plants. The increase in apoplastic pH in the presence of bicarbonates limits iron transport to the root stele and restricts its uptake (Molassiotis et al., 2005).

Sodic stress triggered a significant increase in leaf MDA and H_2O_2 accumulation, the major indicators of oxidative injury (Sorkheh et al., 2012; Shen et al., 2021); regardless of the crop and cultivar tested. Nonetheless, the increases in leaf MDA in sodic stress-stressed plants relative to controls were much higher in peach (~37.0-42.0%) than in pear (9.61-19.91%) as were the increases in leaf H_2O_2 (~22.0-27.0% and ~12.0%, respectively), suggesting that peaches in general were more adversely affected by the oxidative stress (Shahvali et al., 2020). Although comparative studies are lacking, the *Prunus* spp. including peach are mostly highly sensitive to the salt-induced oxidative stress and lipid peroxidation (Erturk et al., 2007; Shen et al., 2021; Toro et al., 2021). Salt-induced osmotic and oxidative stresses more adversely affect osmotic-sensitive than osmotic-tolerant genotypes; the latter show relatively efficient reactive oxygen species (ROS) scavenging when exposed to these stresses (Rajabpoor et al., 2014).

Although direct evidence is lacking, insights from a related species (*Pyrus betulaefolia* Bunge) suggest that salt-triggered increases in leaf MDA may not be deleterious enough to cause the lipid peroxidation in pears (Wu & Zou, 2009). The overexpression of genes '*Pp14-3-3*' [from *P. pyrifolia* (Burm F.) Nakai] and '*PbrNHX2*' (from *P. betulaefolia* Bunge) dramatically improved the salt tolerance in transgenic tobacco and *P. ussuriensis* Maxim by upregulating the activity of the antioxidant enzymes (Li et al., 2014; Dong et al., 2019). Specifically, increased APX activity may account for a better redox homeostasis in pear; '*14-3-3*' proteins interact with

APX for scavenging the reactive oxygen species implicated in the oxidative damage (Li et al., 2014). Our results showed that sodicity-induced increases in APX activity were noticeably higher in pears (28.44-34.49%) than in peaches (13.49-29.41%). The cultivar differences for both MDA and H₂O₂ were highly significant ($p < 0.001$) in pear (Wu & Zou, 2009), but only weekly significant (MDA: $p = 0.07$, H₂O₂: $p = 0.002$) in peach (Toro et al., 2021); implying quite distinct and somewhat shared responses to sodicity stress of pear and peach cultivars, respectively, in terms of lipid peroxidation. Genotypic differences for these oxidative stress markers are either pronounced or subtle in different *Pyrus* and *Prunus* species (Sorkheh et al., 2012; Rajabpoor et al., 2014; Zarafshar et al., 2014; Li et al., 2015; Toro et al., 2021).

Salt-stressed plants accumulate compatible osmolytes such as proline that, in addition to osmoregulation, also reduce the cellular damage by scavenging the ROS (Dejampour et al., 2011; Rahnesan et al., 2018; Yousefi et al., 2019). The tested cultivars did not differ significantly in leaf proline under both control and sodic treatments (Wen et al., 2011). Furthermore, the increase in leaf proline in response to MDA and H₂O₂ accumulation was moderate in pear ($r = 0.578$ and 0.519 , respectively; Fig. 3) but quite strong in peach ($r = 0.870$ and 0.930 , respectively; Fig. 4). Leaf proline levels may not always be sufficient enough to induce the osmotic adjustment (Larher et al., 2009; Singh et al., 2022; Wen et al., 2011). Despite being a typical osmolyte, proline may not essentially lessen the osmotic potential of pear leaves (Larher et al., 2009), and other organic osmolytes such as glycine betaine) may also be involved in the osmotic adjustment. The fact that MDA and H₂O₂ levels may not necessarily be toxic enough to upregulate the proline activity (Wen et al., 2011) also supports our findings as leaf MDA and H₂O₂ were two-three folds higher in peach than in pear (Table 2). Plausibly, a lower

than expected increase in proline activity (Regni et al., 2019) might reflect higher sodicity tolerance in pear than in peach (Ebert, 1999; Musacchi et al., 2006).

All the tested cultivars displayed increased activities of leaf antioxidant enzymes in response to sodicity stress. Enzymatic anti-oxidants efficiently protect salt-stressed plants from ROS induced oxidative stress, and are considered to be the reliable markers for discriminating the salt-tolerant and salt-sensitive genotypes (Sorkheh et al., 2012; Yousefi et al., 2019; Aazami et al., 2021). The cultivar differences for anti-oxidant activities (Regni et al., 2019) can be explained by the complex nature of anti-oxidant expression in plant cells (Racchi, 2013) and the cell organelle-specific activities of anti-oxidant enzymes (Niu & Liao, 2016). The *P. pashia* rootstock was found to better protect the Flemish Beauty scions than clonal (Quince A and C) rootstocks against oxidative damage via enhanced CAT, POX and SOD activities (Sharma & Sharma, 2008). Similarly, peach seedling and clonal rootstocks differed considerably in leaf antioxidant levels in the presence of NaHCO_3 (Molassiotis et al., 2005). The SOD constitutes the first line of defense in alleviating the oxidative stress in plants; it dismutates the superoxide anion (O_2^-) to produce molecular oxygen (O_2) and H_2O_2 . The CAT then decomposes H_2O_2 into O_2 and H_2O (Cavalcanti et al., 2004). Obviously, a balance between SOD and CAT activities and not their relative levels *per se*, seem crucial to maintaining the O_2^- and H_2O_2 levels in a steady-state (Azarabadi et al., 2017). The CAT and SOD activities were not only much higher (Table 3) but also had a clear synergistic effect (Fig. 1, 2) in the sodicity-stressed Punjab Beauty pear and Shan-e-Punjab peach; enabling them to better adapt to the sodicity stress (Sorkheh et al., 2012). The decreased activity of CAT often comes at the expense of greater oxidative damage-characterized, for instance, by the increased accumulation of H_2O_2 (Molassiotis et al., 2005).

Sodicity-stressed pear and peach trees had significantly higher leaf Na^+ and lower K^+ than controls. In the sodic treatment, Punjab Beauty pear showed considerably lower increase in leaf Na^+ than Patharnakh; helping it maintain a higher leaf K^+ . A more or less similar response was also seen in peach. Restricted translocation of Na^+ to aerial plant parts (Matsumoto et al., 2006), achieved for example by Na^+ exclusion in common pears (Musacchi et al., 2006) prevents the xylem loading and translocation of Na^+ to the leaves. The differential accumulation of leaf Na^+ and K^+ in response to the salt stress has also been observed in both own-rooted and grafted peaches (Massai et al., 1997, 2004) and interspecific *Prunus* hybrids (Dejampour et al., 2011), with low Na^+ accumulators showing better salt tolerance (Massai et al., 1997). Reduced accumulation of leaf Na^+ , achieved by root exclusion (Musacchi et al., 2006) or partitioning into the basal leaves (Massai et al., 2004) together with the maintenance of adequate leaf K^+ (Massai et al., 2004) improves the salt tolerance.

In this study, the PCA was highly efficient in reducing the dimensionality, and in differentiating the cultivar- and sodicity-specific effects in data. Specifically, PCA delineated the putative traits linked to sodicity stress tolerance in the pear and peach cultivars. Previously, PCA has been used to unveiling the key responses to the salt stress in other fruit crops (Sorkheh et al., 2012; Abid et al., 2020). Multivariate techniques such as PCA are usually more suitable for detecting the key patterns in data having complex (multicollinear) variables (Julkowska et al., 2019). Additionally, graphical visualization of PCA loadings provides an easier and intuitive means to understanding the shared and contrasting physiological responses to the salt stress (Sorkheh et al., 2012; Singh et al., 2022). Based on the correlation analysis, MDA, H_2O_2 and leaf Na^+ were found to have a greater repressive effect on tree growth in peaches than in pears.

Furthermore, a strong correlation between leaf K^+ and growth traits and leaf chlorophyll in pear, but not in peach, was indicative of leaf K^+ mediated osmotic adjustment in pears.

CONCLUSIONS

Although sodicity stress suppressed tree growth regardless of the cultivar, strong genotypic differences were quite apparent: Punjab Beauty pear and Shan-e-Punjab peach exhibited better tolerance to sodicity stress. We found that sodicity-triggered increases in leaf malondialdehyde, hydrogen peroxide and Na^+ had a greater repressive effect on tree growth in peaches than in pears, and induced the proline-mediated osmotic adjustment in the former. The higher activities of catalase and superoxide dismutase enzymes coupled with restricted Na^+ uptake and the maintenance of adequate leaf K^+ are the plausible explanations for overall better sodicity tolerance in pear. Further studies should aim to understand the effects of sodicity stress on other biochemical and transcriptional changes, and how such changes influence the fruit yield and quality in the sodicity stressed pear and peach cultivars.

ABBREVIATIONS

ANOVA: Analysis of Variance; APX: Ascorbate peroxidase; CAT: Catalase; CV: Canopy volume; DW: Dry weight basis; EC_e : Soil saturation extract electrical conductivity; H_2O_2 : hydrogen peroxide; MDA: Malondialdehyde; $me\ L^{-1}$: Milli equivalent per liter; PCA: Principal Component Analysis; PCs: Principal Components; POX: Peroxidase; pH_s : pH of saturated soil paste; ROS: Reactive oxygen species; SOD: Superoxide dismutase; TCSA: Trunk cross sectional area; TH: Tree height

Acknowledgments: Head, Department of Fruit Science, Punjab Agricultural University, Ludhiana, India is thanked for providing the pear and peach plants. Mr. Dheeraj Kumar is appreciated for his technical help.

REFERENCES

Aazami MA, Rasouli F, Panahi TR. 2021. Influence of salinity stress on morphological, nutritional and physiological attributes in different cultivars of *Prunus amygdalus* L. *Journal of Plant Nutrition* **44**: 1758–1769.

Abbas G, Amjad M, Saqib M, Murtaza B, Asif Naeem M, Shabbir A, Murtaza G. 2021. Soil sodicity is more detrimental than salinity for quinoa (*Chenopodium quinoa* Willd.): A multivariate comparison of physiological, biochemical and nutritional quality attributes. *Journal of Agronomy and Crop Science* **207**: 59–73.

Abd-Elmegeed NA, Nabeel WA, Nasr MM. 2013. Effect of foliar application of some nutrients on " Le-Conte" pear trees grown under calcareous soil conditions. *Egyptian Journal of Horticulture* **40**: 277–293.

Abid M, Zhang YJ, Li Z, Bai DF, Zhong YP, Fang JB. 2020. Effect of salt stress on growth, physiological and biochemical characters of four kiwifruit genotypes. *Scientia Horticulturae* **271**: 109473.

Abrol IP, Yadav JSP, Massoud FI. 1988. Salt-Affected Soils and their Management. <https://agris.fao.org/agris-search/search.do?recordID=XF2016024038>.

Aebi H. 1984. Catalase. *In*: Methods in Enzymology, (L. Packer Ed.). Academic Press, Orlando, **105**: 121–126.

Ahmad M, Alabd A, Gao Y, Yu W, Jamil W, Wang X, Wei J, Ni J, Teng Y, Bai S. 2022. Three stress-responsive NAC transcription factors, Pp-SNACs, differentially and synergistically regulate abiotic stress in pear. *Scientia Horticulturae* **305**: 1-13.

Ahmad P, Ozturk M, Sharma S, Gucel S. 2014. Effect of sodium carbonate-induced salinity–alkalinity on some key osmoprotectants, protein profile, antioxidant enzymes, and lipid peroxidation in two mulberry (*Morus alba* L.) cultivars. *Journal of Plant Interactions* **9**: 460–467.

Altman D, 2000. A review of experiment wise Type I error: implications for univariate Post Hoc and for multivariate testing. p. 20. Paper presented at the Annual Meeting of the Southwest Educational Research Association (Dallas, January 27-29, 2000). <https://files.eric.ed.gov/fulltext/ED438322.pdf>.

- 478 **Arnon DI. 1949.** Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*.
479 *Plant Physiology* **24**: 1.
- 480 **Azarabadi S, Abdollahi H, Torabi M, Salehi Z, Nasiri J. 2017.** ROS generation, oxidative
481 burst and dynamic expression profiles of ROS-scavenging enzymes of superoxide dismutase
482 (SOD), catalase (CAT) and ascorbate peroxidase (APX) in response to *Erwinia amylovora* in
483 pear (*Pyrus communis* L). *European Journal of Plant Pathology* **147**: 279–294.
- 484 **Bates L, Waldren R, Teare I. 1973.** Rapid determination of free proline for water-stress
485 studies. *Plant and Soil* **39**: 205–207.
- 486 **Beauchamp C, Fridovich I. 1971.** Superoxide dismutase: improved assays and an assay
487 applicable to acrylamide gels. *Analytical Biochemistry* **44**: 276–287.
- 488 **Bhatnagar P, Kaul MK. 2002.** Growth and development studies in peach. *Journal of Eco-*
489 *Physiology* **5**: 93-95.
- 490 **Boland AM, Mitchell PD, Jerie PH. 1993.** Effect of saline water combined with restricted
491 irrigation on peach tree growth and water use. *Australian Journal of Agricultural Research* **44**:
492 799–816.
- 493 **Cavalcanti FR, Oliveira JTA, Martins-Miranda AS, Viégas RA, Silveira JAG. 2004.**
494 Superoxide dismutase, catalase and peroxidase activities do not confer protection against
495 oxidative damage in salt-stressed cowpea leaves. *New Phytologist* **163**: 563–571.
- 496 **Dejampour J, Aliasgarzad N, Zeinalabedini M, Niya MR, Hervan EM. 2012.** Evaluation of
497 salt tolerance in almond [*Prunus dulcis* (L.) Batsch] rootstocks. *African Journal of*
498 *Biotechnology* **11**: 11907–11912.
- 499 **Dejampour J, Aliasgarzadb N, Grigorianc V, Majidid E. 2011.** Salt tolerance comparison of
500 four interspecific *Prunus* hybrids. *Acta Horticulturae* **96**: 81–86.
- 501 **Dong H, Wang C, Xing C, Yang T, Yan J, Gao J, Li D, Wang R, Blumwald E, Zhang S.**
502 **2019.** Overexpression of PbrNHX2 gene, a Na⁺/H⁺ antiporter gene isolated from *Pyrus*
503 *betulaefolia*, confers enhanced tolerance to salt stress via modulating ROS levels. *Plant Science*
504 **285**: 14–25.
- 505 **Ebert G. 1999.** Salinity problems in (sub-) tropical fruit production. *Acta Horticulturae* **531**: 99–
506 106.
- 507 **Elkins RB, Bell R, Einhorn T. 2012.** Needs assessment for future US pear rootstock research
508 directions based on the current state of pear production and rootstock research. *Journal of the*
509 *American Pomological Society* **66**: 153–163.

- Erturk U, Sivritepe N, Yerlikaya C, Bor M, Ozdemir F, Turkan I. 2007.** Responses of the cherry rootstock to salinity in vitro. *Biologia Plantarum* **51**: 597–600.
- Flowers TJ. 2004.** Improving crop salt tolerance. *Journal of Experimental Botany* **55**: 307–319.
- Gill HS, Abrol IP. 1991.** Salt affected soils, their afforestation and its ameliorating influence. *International Tree Crops Journal* **6**: 239–260.
- Grieve CM, Grattan SR, Maas EV. 2012.** Plant salt tolerance. *ASCE Manual and Reports on Engineering Practice* **71**: 405–459.
- Heath RL, Packer L. 1968.** Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics* **125**: 189–198.
- Hernández JA, Diaz-Vivancos P, Acosta-Motos JR, Barba-Espín G. 2021.** Where biotic and abiotic stress responses converge: Common patterns in response to salinity and Plum pox virus infection in pea and peach plants. *Annals of Applied Biology* **178**: 281-292.
- Hopmans JW, Qureshi AS, Kisekka I, Munns R, Grattan SR, Rengasamy P, Ben-Gal A, Assouline S, Javaux M, Minhas PS, Raats PAC, Skaggs TH, Wang G, De Jong van Lier Q, Jiol H, Lavado RS, Lazarovitch N, Taleisnik E. 2021.** Critical knowledge gaps and research priorities in global soil salinity. *Advances in Agronomy* **169**: 1-191.
- Hossain MS. 2019.** Present scenario of global salt affected soils, its management and importance of salinity research. *International Research Journal of Biological Sciences* **1**: 1-3.
- Hui-Hui Z, Guang-Liang S, Jie-Yu S, Xin L, Ma-Bo L, Liang M, Nan X, Guang-Yu S. 2019.** Photochemistry and proteomics of mulberry (*Morus alba* L.) seedlings under NaCl and NaHCO₃ stress. *Ecotoxicology and Environmental Safety* **184**: 109624.
- Jia X, Wang H, Svetla S, Zhu Y, Hu Y, Cheng L, Zhao T, Wang Y. 2019.** Comparative physiological responses and adaptive strategies of apple *Malus halliana* to salt, alkali and saline-alkali stress. *Scientia Horticulturae* **245**: 154–162.
- Julkowska MM, Saade S, Agarwal G, Gao G, Pailles Y, Morton M, Awlia M, Tester M. 2019.** MVApp–multivariate analysis application for streamlined data analysis and curation. *Plant Physiology* **180**: 1261–1276.
- Kaledhonkar MJ, Meena BL, Sharma PC. 2019.** Reclamation and nutrient management for salt-affected soils. *Indian Journal of Fertilizers* **15**: 566–575.
- Kanwar JS, Singh H. 2004.** Scope of high density plantings of peach in the subtropics of India. *Acta Horticulturae* **662**: 221-224.

- 541 **Karakas B, Bianco RL, Rieger M. 2000.** Association of marginal leaf scorch with sodium
542 accumulation in salt-stressed peach. *HortScience* **35**: 83–84.
- 543 **Krishnamoorthy V. 2009.** Studies on the performance of different varieties of aonla,
544 pomegranate and annona under sodic soil. *Asian Journal of Horticulture* **4**: 149–151.
- 545 **Kumar S, Narjary B, Kumar K, Jat HS, Kamra SK, Yadav RK. 2019.** Developing soil matric
546 potential based irrigation strategies of direct seeded rice for improving yield and water
547 productivity. *Agricultural Water Management* **215**: 8–15.
- 548 **Larher FR, Lugan R, Gagneul D, Guyot S, Monnier C, Lespinasse Y, Bouchereau A. 2009.**
549 A reassessment of the prevalent organic solutes constitutively accumulated and potentially
550 involved in osmotic adjustment in pear leaves. *Environmental and Experimental Botany* **66**:
551 230–241.
- 552 **Li H, Liu D, Rao J, Liu Y, Ge F, Chen C. 2014.** Overexpression of Pp14-3-3 from *Pyrus*
553 *pyrifolia* fruit increases drought and salt tolerance in transgenic tobacco plant. *Biologia* **69**: 880–
554 887.
- 555 **Li X, Dong W, Lin J, Wang Z, Yang Q, Chang Y, Zhang Z. 2015.** Structural, physiological
556 and biochemical responses of *Pyrus calleryana* offspring to salt stress. *European Journal of*
557 *Horticultural Science* **80**: 306–315.
- 558 **Li Y, Liu C, Sun X, Liu B, Zhang X, Liang W, Huo L, Wang P, Ma F, Li C. 2020.**
559 Overexpression of MdATG18a enhances alkaline tolerance and GABA shunt in apple through
560 increased autophagy under alkaline conditions. *Tree Physiology* **40**: 1509–1519.
- 561 **Ma C, Tanabe K, Itai A, Tamura F, Chun JP, Teng Y. 2005.** Tolerance to lime-induced iron
562 chlorosis of Asian pear rootstocks (*Pyrus spp.*). *Journal of the Japanese Society for Horticultural*
563 *Science* **74**: 419–423.
- 564 **Mahmoud LM, Dutt M, Vincent CI, Grosser JW. 2020.** Salinity-induced physiological
565 responses of three putative salt tolerant citrus rootstocks. *Horticulturae* **6**: 90.
- 566 **Mandal AK. 2012.** Delineation and characterization of salt affected and waterlogged areas in the
567 Indo-Gangetic plain of central Haryana (District Kurukshetra) for reclamation and management.
568 *Journal of Soil Salinity and water quality* **4**: 21–25.
- 569 **Massai R, Gucci R, Tattini M. 1997.** Salinity tolerance in four different rootstocks for
570 peach. *Acta Horticulturae* **465**: 363–370.
- 571 **Massai R, Remorini D, Tattini M. 2004.** Gas exchange, water relations and osmotic adjustment
572 in two scion/rootstock combinations of *Prunus* under various salinity concentrations. *Plant and*
573 *Soil* **259**: 153–162.

574 **Matsumoto K, Chun JP, Tamura F, Kamamoto Y, Tanabe K. 2006.** Salt tolerance in *Pyrus*
575 species is linked to levels of Na and Cl translocation from roots to leaves. *Journal of the*
576 *Japanese Society for Horticultural Science* **75**: 385–391.

577 **Mestre L, Reig G, Betran JA, Pinochet J, Moreno MÁ. 2015.** Influence of peach–almond
578 hybrids and plum-based rootstocks on mineral nutrition and yield characteristics of ‘Big
579 Top’ nectarine in replant and heavy-calcareous soil conditions. *Scientia Horticulturae* **192**: 475–
580 481.

581 **Molassiotis AN, Diamantidis GC, Therios IN, Tsirakoglou V, Dimassi KN. 2005.** Oxidative
582 stress, antioxidant activity and Fe (III)-chelate reductase activity of five *Prunus* rootstocks
583 explants in response to Fe deficiency. *Plant Growth Regulation* **46**: 69–78.

584 **Musacchi S, Quartieri M, Tagliavini M. 2006.** Pear (*Pyrus communis*) and quince (*Cydonia*
585 *oblonga*) roots exhibit different ability to prevent sodium and chloride uptake when irrigated
586 with saline water. *European Journal of Agronomy* **24**: 268–275.

587 **Myers BA, West DW, Callinan L, Hunter CC. 1995.** Long term effects of saline irrigation on
588 the yield and growth of mature Williams pear trees. *Irrigation Science* **16**: 35–46.

589 **Nakano Y, Asada K. 1981.** Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in
590 spinach chloroplasts. *Plant and Cell Physiology* **22**: 867–880.

591 **Niu L, Liao W. 2016.** Hydrogen peroxide signaling in plant development and abiotic responses:
592 crosstalk with nitric oxide and calcium. *Frontiers in Plant Science* **7**: 230.

593 **Okubo M, Furukawa Y, Sakuratani T. 2000.** Growth, flowering and leaf properties of pear
594 cultivars grafted on two Asian pear rootstock seedlings under NaCl irrigation. *Scientia*
595 *Horticulturae* **85**: 91–101.

596 **Oron G, DeMalach Y, Gillerman L, David I, Lurie S. 2002.** SW—soil and water: effect of
597 water salinity and irrigation technology on yield and quality of pears. *Biosystems Engineering*
598 **81**: 237–247.

599 **Qadir M, Oster JD, Schubert S, Noble AD, Sahrawat KL. 2007.** Phytoremediation of sodic
600 and saline-sodic soils. *Advances in Agronomy* **96**: 197–247.

601 **Qadir M, Schubert S. 2002.** Degradation processes and nutrient constraints in sodic soils. *Land*
602 *Degradation & Development* **13**: 275–294.

603 **Racchi, ML. 2013.** Antioxidant defenses in plants with attention to *Prunus* and *Citrus* spp.
604 *Antioxidants* **2**: 340–369.

- 605 **Rahneshan Z, Nasibi F & Moghadam AA. 2018.** Effects of salinity stress on some growth,
606 physiological, biochemical parameters and nutrients in two pistachio (*Pistacia vera* L.)
607 rootstocks. *Journal of Plant Interactions* **13**: 73–82.
- 608 **Rajabpoor S, Kiani S, Sorkheh K, Tavakoli F. 2014.** Changes induced by osmotic stress in the
609 morphology, biochemistry, physiology, anatomy and stomatal parameters of almond species
610 (*Prunus* L. spp.) grown in vitro. *Journal of Forestry Research* **25**: 523–534.
- 611 **Rao MV, Watkins CB, Brown SK, Weeden NF. 1998.** Active oxygen species metabolism in
612 'White Angel' x 'Rome Beauty' apple selections resistant and susceptible to superficial scald.
613 *Journal of the American Society for Horticultural Science* **123**: 299–304.
- 614 **Regni L, Del Pino AM, Mousavi S, Palmerini CA, Baldoni L, Mariotti R, Mairech H, Gardi**
615 **T, D'Amato R, Proietti P. 2019.** Behavior of four olive cultivars during salt stress. *Frontiers in*
616 *Plant Science*. **10**: 867.
- 617 **Samra JS, Singh VP, Sharma KNS. 1988.** Analysis of spatial variability in sodic soils: 2.
618 Point-and block-kriging. *Soil Science* **145**: 250–256.
- 619 **Sangwan AK, Dhillon WS, Singh H, Singh NP, Chohan SK, Gill PPS. 2015.** Influence of
620 horti-silviculture combinations on pre-bearing growth and physiological parameters of
621 pear. *Indian Journal of Horticulture* **72**: 21-27.
- 622 **Saxena CK, Gupta SK. 2006.** Effect of soil pH on the establishment of litchi (*Litchi chinensis*)
623 plants in an alkali environment. *Indian journal of Agricultural Sciences* **76**: 547-549.
- 624 **Shahvali R, Shiran B, Ravash R, Fallahi H, DJeri BB. 2020.** Effect of symbiosis with
625 arbuscular mycorrhizal fungi on salt stress tolerance in GF677 (peach x almond) rootstock.
626 *Scientia Horticulturae* **272**: 1-11.
- 627 **Sharma DK, Chaudhari SK, Singh A. 2014.** In salt-affected soils agroforestry is a promising
628 option. *Indian Farming* **63**: 19–22.
- 629 **Sharma DK, Singh A, Sharma PC, Dagar JC, Chaudhari SK. 2016.** Sustainable management
630 of sodic soils for crop production: opportunities and challenges. *Journal of Soil Salinity and*
631 *Water Quality* **8**: 109–130.
- 632 **Sharma PC, Singh A. 2019.** Reviving the productivity of salt-affected lands: Technological
633 options, constraints and research needs. *Research Developments in Saline Agriculture* **591–627**.
- 634 **Sharma S, Sharma N. 2008.** Effect of rootstocks on leaf water potential, water relations,
635 antioxidant activities and drought tolerant in Flemish Beauty pear under water stress conditions.
636 *Indian Journal of Plant Physiology* **13**: 266–271.

- 637 **Shen W, Liu D, Zhang H, Zhu W, He H, Li G, Liu J. 2021.** Overexpression of β -cyanoalanine
638 synthase of *Prunus persica* increases salt tolerance by modulating ROS metabolism and ion
639 homeostasis. *Environmental and Experimental Botany* **186**: 1-12.
- 640 **Singh A, Kumar A, Kumar R, Sheoran P, Yadav RK, Sharma PC. 2022.** Multivariate
641 analyses discern shared and contrasting eco-physiological responses to salinity stress of Ziziphus
642 rootstocks and budded trees. *South African Journal of Botany* **146**: 573–584.
- 643 **Singh A, Sharma DK, Kumar R, Kumar A, Yadav RK, Gupta SK. 2018a.** Soil salinity
644 management in fruit crops: a review of options and challenges. In: Engineering Practices for
645 Management of Soil Salinity- Agricultural, Physiological, and Adaptive Approaches (Gupta,
646 S.K., Goyal, M.R., Singh, A., Eds.). Apple Academic Press Inc. p. 39–85.
- 647 **Singh A, Kumar A, Sheoran P, Yadav RK. 2018b.** Effects of sodic water irrigation on growth,
648 physiological relations and ion uptake in two Ziziphus rootstocks. *Indian Journal of Agricultural*
649 *Sciences* **88**: 1413–1418.
- 650 **Singh D, Dhillon WS, Singh NP, Gill PPS. 2015.** Effect of girdling on leaf nutrient levels in
651 pear cultivars Patharnakh and Punjab Beauty. *Indian Journal of Horticulture* **72**: 319-324.
- 652 **Sinha AK. 1972.** Colorimetric assay of catalase. *Analytical biochemistry* **47**: 389–394.
- 653 **Soliman MA, Abo-Ogiela HM, El-Saedony NA. 2017.** Reducing adverse effects of salinity in
654 peach trees grown in saline clay soil. *Alexandria Science Exchange Journal* **38**: 800–809.
- 655 **Sorkheh K, Shiran B, Rouhi V, Khodambashi M, Sofo A. 2012.** Salt stress induction of some
656 key antioxidant enzymes and metabolites in eight Iranian wild almond species. *Acta*
657 *Physiologiae Plantarum* **34**: 203–213.
- 658 **Tagliavini M, Masia A, Quartieri M. 1995.** Bulk soil pH and rhizosphere pH of peach trees in
659 calcareous and alkaline soils as affected by the form of nitrogen fertilizers. *Plant and Soil* **176**:
660 263–271.
- 661 **Thind SK, Mahal JS. 2020.** Package of Practices for Cultivation of Fruits. Punjab Agricultural
662 University, Ludhiana. p. 176. https://www.pau.edu/content/ccil/pf/pp_fruits.pdf. Retrieved 26th
663 February, 2021.
- 664 **Thomidis T, Tsipouridis C. 2005.** Influence of rootstocks, pH, iron supply (in nutrient solutions)
665 and *Agrobacterium radiobacter* on chlorophyll and iron concentration in leaves of a peach
666 variety. *Journal of Plant Nutrition* **28**: 1833–1842.
- 667 **Toro G, Pimentel P, Salvatierra A. 2021.** Effective categorization of tolerance to salt stress
668 through clustering Prunus Rootstocks according to their physiological performances.
669 *Horticulturae* **7**: 542.

- 670 **Valipour M, Baninasab B, Khoshgoftarmanesh AH, Gholami M. 2020.** Oxidative stress and
671 antioxidant responses to direct and bicarbonate-induced iron deficiency in two quince rootstocks.
672 *Scientia Horticulturae* **261**: 1-9.
- 673 **Wen XP, Ban Y, Inoue H, Matsuda N, Kita M, Moriguchi T. 2011.** Antisense inhibition of a
674 spermidine synthase gene highlights the role of polyamines for stress alleviation in pear shoots
675 subjected to salinity and cadmium. *Environmental and Experimental Botany* **72**: 157–166.
- 676 **Wu QS, Zou YN. 2009.** Adaptive responses of birch-leaved pear (*Pyrus betulaefolia*) seedlings
677 to salinity stress. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* **37**: 133–138.
- 678 **Yang CW, Xu HH, Wang LL, Liu J, Shi DC, Wang DL. 2009.** Comparative effects of salt-
679 stress and alkali-stress on the growth, photosynthesis, solute accumulation, and ion balance of
680 barley plants. *Photosynthetica* **47**: 79–86.
- 681 **Yousefi M, Naseri L, Zaare-Nahandi F. 2019.** Nitric oxide ameliorates salinity tolerance in
682 Pyrodwarf pear (*Pyrus communis*) rootstocks by regulating polyamine content. *Iranian Journal*
683 *of Plant Physiology* **10**: 3023–3033.
- 684 **Zarafshar M, Akbarinia M, Askari H, Hosseini SM, Rahaie M, Struve D, Striker GG. 2014.**
685 Morphological, physiological and biochemical responses to soil water deficit in seedlings of
686 three populations of wild pear (*Pyrus boissieriana*). *Biotechnology Agronomy Society and*
687 *Environment* **18**: 353–366.

Figure 1

Figure 1 Principal Component Analysis biplot showing variable loadings and cultivar-treatment groups on first two principal components in pear.

Lines radiating from the centre reflect relative contribution and directionality. PBC- Punjab Beauty control, PBS- Punjab Beauty sodic, PathC- Patharnakh control, PathS- Patharnakh sodic, Abbreviations: TH- tree height, TCSA- trunk cross sectional area, CV- canopy volume, TC- total leaf chlorophyll, MDA- malondialdehyde, HP- hydrogen peroxide (H_2O_2), APX- ascorbate peroxidase, POX- peroxidase, CAT- catalase, SOD- superoxide dismutase, Na- leaf Na^+ , K- leaf K^+ , Na.K- leaf Na^+/K^+ ratio.




Figure 2

Figure 2 Principal Component Analysis biplot showing variable loadings and cultivar-treatment groups on first two principal components in peach.

Lines radiating from the centre reflect relative contribution and directionality. PtpC- Partap control, PtpS- Partap sodic, SPC- Shan-e-Punjab control, SPS- Shan-e-Punjab sodic, Abbreviations: TH- tree height, TCSA- trunk cross sectional area, CV- canopy volume, TC- total leaf chlorophyll, MDA- malondialdehyde, HP- hydrogen peroxide (H_2O_2), APX- ascorbate peroxidase, POX- peroxidase, CAT- catalase, SOD- superoxide dismutase, Na- leaf Na^+ , K- leaf K^+ , Na.K- leaf Na^+/K^+ ratio.




Figure 3

Figure 3 Correlation plot showing Pearson's bivariate correlations between the measured traits in pear.

Ellipse size and color reflect the strength and direction (positive or negative) of the correlation. Individual cells marked with cross (X) denote non-significant correlations.

Abbreviations: TH- tree height, TCSA- trunk cross sectional area, CV- canopy volume, TC- total leaf chlorophyll, MDA- malondialdehyde, HP- hydrogen peroxide (H_2O_2), APX- ascorbate peroxidase, POX- peroxidase, CAT- catalase, SOD- superoxide dismutase, Na- leaf Na^+ , K- leaf K^+ , Na.K- leaf Na^+/K^+ ratio.




Figure 4

Figure 4. Correlation plot showing Pearson's bivariate correlations between the measured traits in peach.

Ellipse size and color reflect the strength and direction (positive or negative) of the correlation. Individual cells marked with cross (X) denote non-significant correlations.

Abbreviations: TH- tree height, TCSA- trunk cross sectional area, CV- canopy volume, TC- total leaf chlorophyll, MDA- malondialdehyde, HP- hydrogen peroxide (H_2O_2), APX- ascorbate peroxidase, POX- peroxidase, CAT- catalase, SOD- superoxide dismutase, Na- leaf Na^+ , K- leaf K^+ , Na.K- leaf Na^+/K^+ ratio.




Table 1(on next page)

Table 1 Analysis of Variance (ANOVA) for different traits in pear and peach.

Table 1 Analysis of Variance (ANOVA) for different traits in pear and peach.

Trait	Source	F	p	ω^2	F	p	ω^2
		Pear			Peach		
Tree height (m)	Treatment (T)	83.55	<0.001	0.354	218.76	<0.001	0.902
	Cultivar (C)	134.00	<0.001	0.571	5.95	0.031	0.021
	T x C	2.35	0.151*	0.006	3.62	0.081*	0.011
Trunk cross sectional area (cm ²)	Treatment (T)	50.98	<0.001	0.203	468.82	<0.001	0.379
	Cultivar (C)	179.72	<0.001	0.724	533.37	<0.001	0.431
	T x C	2.98	0.110*	0.008	219.17	<0.001	0.177
Canopy volume (m ³)	Treatment (T)	38.93	<0.001	0.606	411.53	<0.001	0.846
	Cultivar (C)	10.20	0.008	0.147	53.81	<0.001	0.109
	T x C	0.50	0.495*	0.000	7.15	0.020	0.013
Total leaf chlrophyll (mg/g FW)	Treatment (T)	36.52	<0.001	0.509	46.78	<0.001	0.366
	Cultivar (C)	18.39	0.001	0.249	55.54	<0.001	0.436
	T x C	1.86	0.197*	0.012	9.69	0.009	0.070
Malondialdehyde (nmoles/g FW)	Treatment (T)	30.65	<0.001	0.376	81.82	<0.001	0.818
	Cultivar (C)	32.93	<0.001	0.405	3.96	0.070*	0.030
	T x C	2.20	0.164*	0.015	0.03	0.861*	0.000
Hydrogen peroxide (mmoles/g FW)	Treatment (T)	84.27	<0.001	0.219	101.26	<0.001	0.772
	Cultivar (C)	281.57	<0.001	0.739	15.27	0.002	0.110
	T x C	0.95	0.349*	0.000	0.32	0.580*	0.000
Proline (mg/g FW)	Treatment (T)	40.49	<0.001	0.703	85.00	<0.001	0.853
	Cultivar (C)	0.42	0.530*	0.000	0.44	0.518*	0.000
	T x C	2.28	0.157*	0.023	0.06	0.814*	0.000
Ascorbate peroxidase (units/g FW)	Treatment (T)	147.59	<0.001	0.906	28.11	<0.001	0.092
	Cultivar (C)	0.42	0.527*	0.000	252.51	<0.001	0.856
	T x C	0.85	0.374*	0.000	0.21	0.652*	0.000
Peroxidase (units/g FW)	Treatment (T)	167.45	<0.001	0.500	130.39	<0.001	0.072
	Cultivar (C)	129.46	<0.001	0.386	1618.96	<0.001	0.898
	T x C	23.24	<0.001	0.067	39.18	<0.001	0.021
Catalase (units/g FW)	Treatment (T)	161.13	<0.001	0.421	197.25	<0.001	0.116
	Cultivar (C)	205.79	<0.001	0.539	1456.33	<0.001	0.860
	T x C	1.040e ⁻⁴	0.992*	0.000	24.82	<0.001	0.014
Superoxide dismutase (units/g FW)	Treatment (T)	88.32	<0.001	0.542	107.21	<0.001	0.554
	Cultivar (C)	55.54	<0.001	0.338	71.48	<0.001	0.368
	T x C	4.35	0.059*	0.021	0.01	0.911*	0.000
Leaf Na ⁺ (mg/g DW)	Treatment (T)	159.87	<0.001	0.745	423.82	<0.001	0.730
	Cultivar (C)	38.69	<0.001	0.177	121.42	<0.001	0.208
	T x C	1.80	0.205*	0.004	21.05	<0.001	0.035
Leaf K ⁺ (mg/g DW)	Treatment (T)	109.07	<0.001	0.590	252.43	<0.001	0.144
	Cultivar (C)	58.23	<0.001	0.312	1407.86	<0.001	0.806
	T x C	2.96	0.111*	0.011	71.96	<0.001	0.041
Leaf Na ⁺ /K ⁺ ratio	Treatment (T)	201.17	<0.001	0.871	651.86	<0.001	0.620
	Cultivar (C)	0.38	0.549*	0.000	360.43	<0.001	0.342
	T x C	15.25	0.002	0.062	25.19	<0.001	0.023

*Non-significant effect (p> 0.05).

Table 2 (on next page)

Table 2 Mean comparisons for tree growth parameters in pear and peach cultivars.

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.

Table 2 Mean comparisons for tree growth parameters in pear and peach cultivars.

Cultivar	Treatment	Tree height (m)	Trunk cross sectional area (cm ²)	Canopy volume (m ³)
Pear				
Punjab Beauty	Control	3.86±0.20a	36.76±2.63a	1.91±0.21a
	Sodic	3.23±0.12b	30.70±2.64b	1.06±0.19bc
Patharnakh	Control	3.03±0.20b	23.68±1.87c	1.53±0.53ab
	Sodic	2.14±0.13c	13.76±1.62d	0.47±0.08d
Peach				
Partap	Control	4.25±0.31a	231.17±9.61b	28.96±2.90a
	Sodic	2.71±0.11b	59.45±2.62c	2.80±0.18b
Shan-e-Punjab	Control	4.31±0.14a	270.22±12.0a	34.28±3.09a
	Sodic	3.05±0.18b	237.96±10.57b	14.20±1.68b

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.

Table 3(on next page)

Table 3 Mean comparisons for leaf physiological traits in pear and peach cultivars.

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD. TC- total leaf chlorophyll, MDA- malondialdehyde, H₂O₂- hydrogen peroxide.

Table 3 Mean comparisons for leaf physiological traits in pear and peach cultivars.

Cultivar	Treatment	TC (mg g ⁻¹ FW)	MDA (nmol g ⁻¹ FW)	H ₂ O ₂ (mmol g ⁻¹ FW)	Proline (mg g ⁻¹ FW)
Pear					
Punjab Beauty	Control	1.41±0.10a	8.95±0.25b	130.41±1.99b	3.78±0.43b
	Sodic	1.03±0.10bc	9.81±0.35a	146.27±3.23a	5.06±0.25a
Patharnakh	Control	1.12±0.05a	7.43±0.22c	105.74±4.32d	3.92±0.28b
	Sodic	0.88±0.14c	8.91±0.69b	118.55±2.43c	4.71±0.31a
Peach					
Partap	Control	1.56±0.11a	7.74±0.36b	138.12±4.28c	4.06±0.11b
	Sodic	1.08±0.15b	10.58±0.94a	175.57±8.77a	5.31±0.18a
Shan-e-Punjab	Control	1.05±0.05bc	7.04±0.11b	153.89±9.19b	4.16±0.19b
	Sodic	0.86±0.05c	10.0±0.78a	187.33±4.32a	5.39±0.50a

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD. TC- total leaf chlorophyll, MDA- malondialdehyde, H₂O₂- hydrogen peroxide.

Table 4(on next page)

Table 4 Mean comparisons for leaf anti-oxidant enzymes (units g⁻¹ FW) in pear and peach cultivars.

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.

Table 4 Mean comparisons for leaf anti-oxidant enzymes (units g⁻¹ FW) in pear and peach cultivars.

Cultivar	Treatment	Ascorbate peroxidase	Peroxidase	Catalase	Superoxide dismutase
Pear					
Punjab Beauty	Control	14.10±0.53b	183.68±5.80c	6.20±0.31b	64.09±1.88b
	Sodic	18.11±0.75a	224.07±4.92b	7.76±0.24a	80.62±4.40a
Patharnakh	Control	13.54±0.68b	220.47±3.47b	4.42±0.31c	56.23±1.29c
	Sodic	18.21±0.86a	238.84±3.74a	6.01±0.25b	66.84±2.08b
Peach					
Partap	Control	22.15±1.38b	307.41±5.85b	3.93±0.17d	46.88±1.31c
	Sodic	25.14±1.46a	326.60±5.42a	5.22±0.43c	54.47±1.55b
Shan-e-Punjab	Control	12.07±1.01d	178.22±6.93d	6.93±0.37b	53.00±1.57b
	Sodic	15.62±1.01c	219.20±5.92c	7.51±0.41a	60.84±1.54a

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.

Table 5(on next page)

Table 5 Mean comparisons for leaf Na⁺ and K⁺ and Na⁺: K⁺ ratio in pear and peach cultivars.

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.

Table 5 Mean comparisons for leaf Na⁺ and K⁺ and Na⁺: K⁺ ratio in pear and peach cultivars.

Cultivar	Treatment	Na ⁺ (mg/g DW)	K ⁺ (mg/g DW)	Na ⁺ : K ⁺ ratio
Pear				
Punjab Beauty	Control	0.82±0.07c	9.74±0.52a	0.08±0.01b
	Sodic	1.18±0.09a	8.01±0.31b	0.15±0.01a
Patharnakh	Control	0.59±0.05d	8.54±0.63b	0.07±0.01b
	Sodic	1.03±0.07b	6.09±0.45c	0.17±0.02a
Peach				
Partap	Control	1.13±0.10c	14.42±0.59a	0.08±0.01d
	Sodic	1.94±0.10a	9.81±0.25b	0.20±0.01b
Shan-e-Punjab	Control	0.91±0.04d	5.70±0.37c	0.16±0.01c
	Sodic	1.44±0.09b	4.32±0.26d	0.33±0.02a

Control and sodic treatments denote significantly different ($p < 0.001$) pH_s levels of ~ 8.2 and 8.8, respectively. Each data value represents mean (n=4) ± SD.