

Long-term outcomes and predictors of patients with ST elevated versus non-ST elevated myocardial infarctions in non-obstructive coronary arteries: A retrospective study in Northern China

Lin Chen^{1,2}, Yinghong Fan^{1,2}, Zhen Fang³, Ning Liu^{Corresp. 3}

¹ Yangzhou University, Institute of Gastroenterology, Affiliated Hospital of Yangzhou University, Yangzhou, Jiangsu Province, China

² Yangzhou University, Pancreatic Center, Department of Gastroenterology, The Affiliated Hospital of Yangzhou University, Yangzhou, Jiangsu Province, China

³ Department of Cardiology, Northern Jiangsu People's Hospital, Yangzhou, Jiangsu Province, China

Corresponding Author: Ning Liu
Email address: 844465791@qq.com

Background. Myocardial infarction with non-obstructive coronary arteries (MINOCA) is a heterogeneous disease entity with diverse etiologies and no uniform treatment protocols. Patients with MINOCA can be clinically classified into two groups based on whether they have an ST-segment elevation (STE) or non-ST segment elevation (NSTEMI), based on electrocardiogram (ECG) results, whose clinical prognosis is unclear. This study aimed to compare the outcomes and predictors of patients with STE and NSTEMI in the MINOCA population. **Methods.** We collected the data for 196 patients with MINOCA (115 with STE and 81 with NSTEMI) in China. Clinical characteristics, prognoses, and predictors of major adverse cardiovascular events (MACE) were analyzed during the follow-up of all patients. **Results.** The proportion of patients with STE was greater than that with NSTEMI in the MINOCA population. Patients with NSTEMI were older and had a higher incidence of hypertension. No differences were observed in the outcomes between the STE and NSTEMI groups during a median follow-up period of 49(37,46) months. No significant differences were observed in those with MACE (24.35% vs 22.22%, $P = 0.73$) and those without MACE. The multivariable predictors of MACE in the NSTEMI groups were Killip grades ≥ 2 (HR 9.035, CI 95%:1.657-49.263, $P=0.011$), reduced use of β -blockers during hospitalization (HR 0.238, CI 95%:0.072-0.788, $P=0.019$), and higher levels of low-density lipoprotein cholesterol (LDL-C) (HR 2.267, CI 95%: 1.008-5.097, $P=0.048$); the reduced use of β -blockers during hospitalization was the only independent risk factor of MACE in the STE group. **Conclusions.** There were differences between the clinical characteristics of patients with STE and NSTEMI in the MINOCA population, even though outcomes during follow-up were similar. Independent risk factors for major adverse cardiac events were not identical

in the STE and NSTEMI groups, which could be attributable to the differences in disease pathogenesis.

Long-term outcomes and predictors of patients with ST elevated versus non-ST elevated myocardial infarctions in non-obstructive coronary arteries: A retrospective study in Northern China

Lin Chen, MD^{1,2}, Yinghong Fan^{1,2}, Zhen Fang, MD³, Ning Liu, MD^{3*}

1 Pancreatic Center, Department of Gastroenterology, The Affiliated Hospital of Yangzhou University, Yangzhou University, Yangzhou 225001, Jiangsu Province, China

2 Institute of Gastroenterology, Affiliated Hospital of Yangzhou University, Yangzhou University, Yangzhou 225001, Jiangsu Province, China

3 Department of Cardiology, Northern Jiangsu People's Hospital, Yangzhou 225001, Jiangsu Province, China

*Corresponding author: Ning Liu

E-mail: 844465791@qq.com

Telephone: +86-18051062118

Abstract

Background. Myocardial infarction with non-obstructive coronary arteries (MINOCA) is a heterogeneous disease entity with diverse etiologies and no uniform treatment protocols. Patients with MINOCA can be clinically classified into two groups based on whether they have an ST-segment elevation (STE) or non-ST segment elevation (NSTEMI), based on electrocardiogram (ECG) results, whose clinical prognosis is unclear. This study aimed to compare the outcomes and predictors of patients with STE and NSTEMI in the MINOCA population.

Methods. We collected the data for 196 patients with MINOCA (115 with STE and 81 with NSTEMI) in China. Clinical characteristics, prognoses, and predictors of major adverse cardiovascular events (MACE) were analyzed during the follow-up of all patients.

Results. The proportion of patients with STE was greater than that with NSTEMI in the MINOCA population. Patients with NSTEMI were older and had a higher incidence of hypertension. No differences were observed in the outcomes between the STE and NSTEMI groups during a median follow-up period of 49(37,46) months. No significant differences were observed in those with MACE (24.35% vs 22.22%, $P = 0.73$) and those without MACE. The multivariable predictors of MACE in the NSTEMI groups were Killip grades ≥ 2 (HR 9.035, CI 95%:1.657-49.263, $P=0.011$), reduced use of β -blockers during hospitalization (HR 0.238, CI 95%:0.072-0.788, $P=0.019$), and higher levels of low-density lipoprotein cholesterol (LDL-C) (HR 2.267, CI 95%; 1.008-5.097, $P=0.048$); the reduced use of β -blockers during hospitalization was the only independent risk factor of MACE in the STE group.

Conclusions. There were differences between the clinical characteristics of patients with STE and NSTEMI in the MINOCA population, even though outcomes during follow-up were similar. Independent risk factors for major adverse cardiac events were not identical in the STE and NSTEMI groups, which could be attributable to the differences in disease pathogenesis.

Introduction

Myocardial infarction with nonobstructive coronary arteries (MINOCA) is a heterogeneous group of diseases with different pathogeneses. It is characterized by acute myocardial infarction with normal coronary arteries or mild coronary artery stenosis (stenosis < 50%), and occurs commonly in young women (Tamis-Holland et al. 2019). The prevalence of MINOCA reportedly ranges between 1-15% in patients with acute myocardial infarction (AMI), according to different studies (Abdu et al. 2019), and its overall prevalence was 6% in a recent meta-analysis (Pasupathy et al. 2015). MINOCA is a group of syndromes with multiple causes. Individuals with MINOCA can be classified into multiple subgroups, such as those with plaque rupture, coronary dissection, coronary artery spasm, and clinically unrecognized myocarditis or Takotsubo cardiomyopathy; all of these have different underlying pathophysiological mechanisms (Agewall et al. 2017; Niccoli et al. 2015). Therefore, it is potentially challenging to effectively treat MINOCA patients for whom multiple pathogenic mechanisms have various underlying causes. The pathogenesis and prognosis of MINOCA patients need to be assessed further in future studies.

Previous studies have reported that patients with MINOCA had lower rates of major adverse cardiovascular events (MACE) and mortality during follow-up than patients with MI-CAD (Montenegro Sá et al. 2018; Pasupathy et al. 2015; Pizzi et al. 2016). Although patients with MINOCA appear to have a slightly better long-term prognosis, compared to MI-CAD (MI with obstructive coronary artery disease) patients, studies conducted in recent years have shown that MINOCA is not always benign (Barr et al. 2018; Raparelli et al. 2018). Notably, a Swedish study conducted over 4 years has shown that adverse cardiovascular events occurred in 23.9% of MINOCA patients during follow-up; among these, the mortality rate could be as high as 13.4% (Lindahl et al. 2017). Moreover, a Japanese study also showed that MINOCA patients had a higher mortality rate within 30 days of follow-up, as compared to MI-CAD patients (4.48% VS 3.46%) (Ishii et al. 2020).

However, the differences in clinical features and prognosis between patients with ST-segment elevated myocardial infarction (STEMI) and non-ST segment elevated myocardial infarction

(NSTEMI) remain controversial. The occurrence of NSTEMI is more common than STEMI in the MINOCA population (Pasupathy et al. 2015; Xu et al. 2020). Previous studies have reported that STEMI patients had a poorer short-term prognosis and a more favorable long-term prognosis (Borrayo-Sánchez et al. 2018; Polonski et al. 2011). A large-scale Swedish study of MINOCA patients reported that during the 2.6-year follow-up period, the mortality rate for STEMI patients was 8%, while the mortality rate for NSTEMI patients was lower at 5% (Nordenskjöld et al. 2018), which was inconsistent with the results reported by Li et al. (Li et al. 2022). Nevertheless, some studies suggest that there were no differences in prognosis between patients with STEMI and NSTEMI (Montalescot et al. 2007). Some studies have shown that the history of atrial fibrillation, Killip grade, age, and treatment strategy were significant independent risk factors for prognosis in MINOCA patients (Montalescot et al. 2007; Polonski et al. 2011), while the predictors of prognosis in STE and NSTEMI patients are still unclear. Although the differences in prognosis between STEMI and NSTEMI patients in the AMI population have been reported hitherto, the differences in prognosis and predictors of prognosis among MINOCA patients with STE and NSTEMI remain unclear. This study aimed to compare the clinical features, prognosis, and predictors of MACE during the follow-up period among MINOCA patients with STE and NSTEMI in Northern China.

Materials & Methods

Patients

We conducted a retrospective study of patients who had been admitted to the First Hospital at Jilin University due to AMI from January 2015 to July 2018 and had undergone coronary angiography during hospitalization. Patients were included in the study if: (1) they met the diagnostic criteria specified in the AMI guidelines (Thygesen et al. 2018); (2) no occlusion of any infarct-related coronary artery and <50% stenosis could be observed in all epicardial vessels; (3) the patient received no other alternative diagnosis during clinical presentation (e.g., non-ischemic causes such as sepsis, acute renal failure, pulmonary embolism, and myocarditis); and

(4) age >18 years. Patients were excluded if: (1) thrombolytic therapy had been performed prior to coronary angiography; (2) they had a previous myocardial infarction or coronary revascularization; (3) previously underwent cardiac surgery; (4) had malignant tumors.

This study has been conducted in accordance with the Declaration of Helsinki and was approved by the Ethical Review Board of the hospital (the First Hospital of Jilin University, Changchun, China). Patient informed consent was waived, as this study was retrospective.

Data collection

Most of the data were obtained from the medical records at the First Hospital of Jilin University that contained data on the baseline characteristics, biochemical markers, electrocardiogram (ECG) images, coronary angiography, and medications provided during hospitalization. Basic patient information (e.g., age, sex) and past medical history (e.g., smoking history, history of hypertension, hyperlipidemia, diabetes, arrhythmias) were recorded in detail. The history of arrhythmias including previous atrial arrhythmias or ventricular arrhythmias or heart block. We collected information regarding biochemical markers, including blood cardiac troponin-T(cTnT), creatine kinase-MB(CK-MB), brain natriuretic peptide (BNP), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), triglyceride (TG) and indicators of echocardiography, including LV (left ventricle) and LVEF (left ventricular ejection fraction) in 24 hours after hospitalization. We classified the patients into the STE and NSTEMI groups based on their ECG results. STE and NSTEMI were defined in accordance with the Fourth Universal Definition of Myocardial Infarction (Thygesen et al. 2018).

The primary clinical endpoint of our study was the occurrence of major adverse cardiovascular events (MACE), including rehospitalization for increased chest pain that did not meet the criteria of AMI, based on ECG results and myocardial injury marker levels, and occurrence of non-fatal MI, heart failure, stroke, heart valve replacement, and all-cause deaths, which included cardiovascular and non-cardiovascular deaths. A diagnosis of MI was made if patients exhibited

the dynamic development of cardiac troponin in conjunction with symptoms suggestive of myocardial ischemia. Cardiovascular death was defined as death because of acute coronary syndrome(ACS), cardiac rupture, severe arrhythmias, or refractory severe heart failure. A stroke was defined as an ischemic cerebral infarction caused by thrombotic or embolic occlusions in any major intracranial artery. A diagnosis of heart failure (HF) was established according to the current guidelines of the European Society of Cardiology(ESC) (Ponikowski et al. 2016).

Statistical analysis

Statistical analysis was performed using SPSS 25.0 software. Normally distributed continuous variables were presented as mean \pm standard deviation (SD) values. Non-normally distributed continuous variables were presented in terms of the median and inter-quartile range (IQR). Categorical variables were presented as counts and percentages. An independent sample t-test and the Mann-Whitney U test were used to perform a comparison of continuous variables between groups. Categorical variables were compared using the Chi-square and Fisher's exact tests. We used logistic regression analysis to evaluate the independent risk factors of outcomes in the STE and NSTEMI groups, while the adjusted OR for MACE was calculated via logistic regression analysis. All the tests performed were two-sided tests and values were identified to be statistically significant at a P-value < 0.05 .

Results

Baseline characteristics of patients

In our study, the median follow-up period was 49 (37,46) months. A total of 9696 patients were diagnosed with MI; among these, 196 patients (2.02%) satisfied the diagnostic criteria for MINOCA. Based on the ECG results, 115 patients (58.7%) were included in the STE group, while 81 patients (41.3%) were included in the NSTEMI group (Figure 1). A comparison of the baseline characteristics between patients with STE and NST among the MINOCA population has been shown in Table 1.

In comparison to NSTEMI patients, patients with STEMI were younger. Patients with NSTEMI had a higher incidence of hypertension, whereas no significant differences were observed in the incidence of other coronary risk factors (e.g., diabetes, hyperlipidemia, previous arrhythmia, smoking history). The medications administered at discharge have been shown in Table 1. There were no significant differences between the two groups except for the fact that the more frequent use of aspirin and lower use of ACEI/ARB at admission in the STEMI group. Thus, the proportions of patients using clopidogrel, β -blockers, and statins were similar in the two groups. The level of serum glucose on admission in the NSTEMI group was higher than that in the STEMI group, while the other laboratory parameters were not significantly different among the two groups.

Follow-up

During a median follow-up period of 49 months (interquartile range [IQR] 37-61), MACE occurred in 46 (23.47%) out of a total of 51 patients. In the STEMI and NSTEMI groups, we observed the occurrence of MACE in 28 (24.35%) and 18 (22.22%) patients, respectively. The data are shown in Table 2. Thirty-one cases of MACE were observed in 28 patients (24.35%) in the STEMI group, including in patients who had to undergo rehospitalization for chest pain (4 cases, 3.48%), non-fatal MI (3 cases, 2.61%), heart failure (14 cases, 12.17%), stroke (5 cases, 4.35%), and all-cause deaths (5 cases, 4.35%). In the NSTEMI group, 20 cases MACE (24.35%) were observed in 18 patients (22.22%); these included chest pain (4 patients, 4.49%), non-fatal MI (2 patients, 2.47%), heart failure (8 patients, 9.88%), stroke (3 patients, 3.70%), heart valve replacement (2 patients, 2.47%), and all-cause death (1 patient, 1.23%). There were no statistical differences in the prevalence of MACE between the NSTEMI and STEMI groups ($P=0.73$). In this study, 5 patients died of cardiogenic diseases. During the follow-up period, there was no significant difference in the incidence of chest pain, non-fatal MI, heart failure, stroke, heart valve replacement, and all-cause death between the STEMI and NSTEMI groups among the MINOCA population ($P>0.05$).

Predictive factors

Univariate analysis showed that older age, Killip grade ≥ 2 , longer hospitalization duration, being born male, lower use of β -blockers during hospitalization, and red blood cell counts were significant risk factors for MACE in the STE group (Table 3).

We conducted a multivariate analysis adjusted for age, Killip grades, hospitalization duration, sex, use of β -blockers during hospitalization, red blood cell counts, history of diabetes, and level of low-density lipoprotein cholesterol (LDL-C). The results showed that Killip grades ≥ 2 (HR 9.035, CI 95%: 1.657-49.263, $P=0.011$), lowered use of β -blockers during hospitalization (HR 0.238, CI 95%: 0.072-0.788, $P=0.019$) and higher LDL-C levels (HR 2.267, CI 95%: 1.008-5.097, $P=0.048$) were independent risk factors for MACE in patients with STE (Table 4).

Univariate analysis showed that older age and lowered use of β -blockers during hospitalization were associated with a higher extent of occurrence of MACE in the NSTEMI group (Table 3).

The age and extent of use of β -blockers and aspirin during hospitalization were adjusted via multivariate analysis. The results revealed that the lowered use of β -blockers during hospitalization was the only independent risk factor for MACE in patients with NSTEMI (HR 0.303, CI 95%: 0.093-0.991, $P=0.048$). Thus, the use of β -blockers could improve the prognosis of MINOCA patients with NSTEMI.

Discussion

The objective of this study was to compare the prognosis and predictors of MACE among MINOCA patients with STE and NSTEMI. Our major findings were as follows: (1) There were differences in clinical features between the STE and NSTEMI groups among MINOCA patients; (2) there was no statistical difference in the incidence of MACE between the STE and NSTEMI groups during follow-up; (3) the independent risk predictors of MACE in MINOCA patients with STE include a higher level of LDL-C, Killip grades ≥ 2 , and lowered use of β -blockers during hospitalization, whereas the lowered use of β -blockers during hospitalization was the only multivariable predictor of MACE in MINOCA patients with NSTEMI.

MINOCA has always been a confusing clinical entity that is characterized by myocardial infarctions with normal or near-normal coronary arteries of angiography (Scalone et al. 2019). Due to the difference in sample size and definition among various cohorts, the incidence of MINOCA in patients with acute myocardial infarction (AMI) is 1-15% (Kilic et al. 2020; Nordenskjöld et al. 2018; Xu et al. 2020), which is consistent with the findings of our study. Although the underlying causes of MINOCA are diverse, patients can be classified into the STEMI and NSTEMI groups based on their electrocardiogram (ECG) results. Among MINOCA patients, the proportion of patients with NSTEMI is higher than that of those with STEMI (Pasupathy et al. 2015; Xu et al. 2020), which was in contrast to the findings of our study. This result may be attributable to the fact that our study is a single-center study with a small sample size. Certain previous studies have reported that there were significant differences in the clinical features of MINOCA patients with STEMI and NSTEMI (Borrayo-Sánchez et al. 2018; Hanssen et al. 2012). Recently, a Chinese study on MINOCA reported that patients with NSTEMI were older, mostly female, and had a higher incidence of atrial fibrillation. Furthermore, patients with STEMI were more likely to have a history of smoking and a higher diastolic blood pressure, whereas there were no significant differences in the incidence of other risk factors for coronary problems (e.g., hypertension, diabetes) between the two groups (Xu et al. 2020). Our study found that patients with NSTEMI had a higher age and a higher proportion of the patients had hypertensive disease, compared to STEMI patients, which was consistent with the findings of the study by Johnston et al. (Johnston et al. 2015). Therefore, these differences may be associated with the different pathogeneses of the two groups; this needs to be confirmed in multi-center and prospective studies with a large sample size.

The prognostic differences between STEMI and NSTEMI patients in the MINOCA population remain controversial. Previous studies have reported higher short-term mortality in STEMI patients and higher long-term mortality in NSTEMI patients (Borrayo-Sánchez et al. 2018; Johnston et al. 2015; Park et al. 2013), which was also observed in the MINOCA population (Nordenskjöld et al. 2018). Johnston et al. reported that all-cause mortality was significantly

higher in MINOCA patients with STEMI than in NSTEMI patients and that their long-term prognosis was poorer (Johnston et al. 2015). A recent study demonstrated that the mortality of patients with MINOCA presenting with STEMI was relatively high at 4.5% at year 1 (Gue et al. 2019). This might be related to the occurrence of congestive heart failure because of highly extensive and severe myocardial damage. However, no statistically significant differences in mortality were observed between STEMI and NSTEMI patients in this study. Our findings were similar to those of Xu (Xu et al. 2020) because we found that there was no statistical difference in the incidence of MACE (rehospitalization for chest pain, non-fatal MI, heart failure, stroke, heart valve replacement and all-cause death, etc.) in the follow-up period between the STEMI and NSTEMI groups, which may be related to the similar drug therapies administered to patients from different sub-groups.

There are several inconsistencies regarding the predictors of MACE in STEMI and NSTEMI patients in previous studies. One study reported that STEMI and NSTEMI patients differed significantly with regard to predictors of early and late-term mortality (Park et al. 2013; Polonski et al. 2011). In addition, the study conducted by Xu demonstrated that the predictors of MACE in MINOCA patients with STE and NSTE were different; the independent predictors of MACE in the NSTEMI group were age, lower level of TC, hypertension, and smoking history, and the strongest predictors in the STEMI group were reduced LVEF levels and a history of diabetes mellitus (Xu et al. 2020). A large meta-analysis showed that a further reduction in LDL-C levels was effective in reducing the incidence of prognostic cardiovascular disease and stroke (Baigent et al. 2010), which was consistent with our findings, which showed that a higher LDL-C level was an independent risk factor for MACE in the STEMI group. The use of statins in patients with MINOCA for reducing the LDL-C levels and stabilizing and controlling coronary plaque progression has had a beneficial prognostic impact (Choo et al. 2019). Johnston et al. found that among STEMI patients, the all-cause mortality was significantly higher in females than in males (Johnston et al. 2015), while this difference in mortality between the sexes was not observed in

our research. We suggest that this complexity is reflective of the heterogeneous features of MINOCA in terms of STE and NSTE. Currently, there is no uniform treatment for the MINOCA population. We found that β -blocker medication was a protective factor for MACE during the follow-up period in the MINOCA population with NSTEMI and STEMI, which is consistent with the findings of Ciliberti et al.(Ciliberti et al. 2021). However, the findings of the study by Abdu showed that the treatment of MINOCA with statins and ACEI/ARB had long-term beneficial effects on the outcome, whereas β -blocker and DAPT treatment seemed to have no significant effect on the occurrence of MINOCA(Abdu et al. 2019). The administration of characteristic therapies is necessary for patients in whom the occurrence of MINOCA is attributable to different underlying mechanisms. All the above-mentioned studies suggest that the use of secondary preventative medications for cardiovascular disease may significantly improve the prognosis of the MINOCA population and should be advocated, but this needs to be confirmed in multicenter studies with longer follow-up periods.

Limitations

There are several limitations to our study. One of the major limitations is that our study was a single-center retrospective study with a small sample size and a short follow-up period, because of which our findings might lead to biased findings. Second, cardiac magnetic resonance (CMR) cloud not be performed for all patients due to medical insurance-related issues and the lack of CMR may influence the accuracy of our findings in MINOCA patients. Finally, information regarding medications to be administered in the follow-up period could not be obtained for all patients. Hence, we could not further analyze whether the long-term use of secondary preventative medications was beneficial for patients with MINOCA. A larger multi-center randomized controlled study is necessary to clarify the results of this study.

290

291 **Conclusions**

292 In conclusion, the clinical characteristics of the STE and NSTEMI groups differed in patients with
293 MINOCA, whereas the outcomes during the 49-month follow-up were similar. The predictors for
294 MACE in patients between the STE group and NSTEMI group were not thoroughly identical.

295

296 **Acknowledgements**

297

298 **Competing Interests**

299 The authors declare there are no competing interests.

300 Human Ethics

301 We declare that this study has been approved by the Ethics Committee of the First Hospital of Jilin
302 University.

303

304

305

306

307 **References**

- 308 Abdu FA, Liu L, Mohammed AQ, Luo Y, Xu S, Auckle R, Xu Y, and Che W. 2019. Myocardial
309 infarction with non-obstructive coronary arteries (MINOCA) in Chinese patients: Clinical
310 features, treatment and 1 year follow-up. *Int J Cardiol* 287:27-31. 10.1016/j.ijcard.2019.02.036
- 311 Agewall S, Beltrame JF, Reynolds HR, Niessner A, Rosano G, Caforio AL, De Caterina R, Zimarino M,
312 Roffi M, Kjeldsen K, Atar D, Kaski JC, Sechtem U, and Tornvall P. 2017. ESC working group
313 position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J*
314 38:143-153. 10.1093/eurheartj/ehw149

Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhala N, Peto R, Barnes EH, Keech A, Simes J, and Collins R. 2010. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 376:1670-1681. 10.1016/s0140-6736(10)61350-5

Barr PR, Harrison W, Smyth D, Flynn C, Lee M, and Kerr AJ. 2018. Myocardial Infarction Without Obstructive Coronary Artery Disease is Not a Benign Condition (ANZACS-QI 10). *Heart Lung Circ* 27:165-174. 10.1016/j.hlc.2017.02.023

Borrayo-Sánchez G, Rosas-Peralta M, Ramírez-Arias E, Saturno-Chiu G, Estrada-Gallegos J, Parra-Michel R, Hernandez-García HR, Ayala-López EA, Barraza-Felix R, García-Rincón A, Adalid-Arellano D, Careaga-Reyna G, Lázaro-Castillo JL, Betancourt-Hernández LE, Camacho-Casillas R, Hernández-Gonzalez M, Celis-Quintal G, Villegas-González B, Hernández-Carrillo M, Benítez Arechiga ZM, Flores-Morales A, and Sepúlveda-Vildosola AC. 2018. STEMI and NSTEMI: Real-world Study in Mexico (RENASCA). *Arch Med Res* 49:609-619. 10.1016/j.arcmed.2019.01.005

Choo EH, Chang K, Lee KY, Lee D, Kim JG, Ahn Y, Kim YJ, Chae SC, Cho MC, Kim CJ, Kim HS, and Jeong MH. 2019. Prognosis and Predictors of Mortality in Patients Suffering Myocardial Infarction With Non-Obstructive Coronary Arteries. *J Am Heart Assoc* 8:e011990. 10.1161/jaha.119.011990

Ciliberti G, Verdoia M, Merlo M, Zilio F, Vatrano M, Bianco F, Mancone M, Zaffalon D, Bonci A, Boscutti A, Infusino F, Coiro S, Stronati G, Tritto I, Gioscia R, Dello Russo A, Fedele F, Gallina S, Cassadonte F, Ambrosio G, Bonmassari R, De Luca G, Sinagra G, Capucci A, Kaski JC, and Guerra F. 2021. Pharmacological therapy for the prevention of cardiovascular events in patients with myocardial infarction with non-obstructed coronary arteries (MINOCA): Insights from a multicentre national registry. *Int J Cardiol* 327:9-14. 10.1016/j.ijcard.2020.11.040

Gue YX, Corballis N, Ryding A, Kaski JC, and Gorog DA. 2019. MINOCA presenting with STEMI: incidence, aetiology and outcome in a contemporaneous cohort. *J Thromb Thrombolysis* 48:533-538. 10.1007/s11239-019-01919-5

- Hanssen M, Cottin Y, Khalife K, Hammer L, Goldstein P, Puymirat E, Mulak G, Drouet E, Pace B, Schultz E, Bataille V, Ferrières J, Simon T, and Danchin N. 2012. French Registry on Acute ST-elevation and non ST-elevation Myocardial Infarction 2010. FAST-MI 2010. *Heart* 98:699-705. 10.1136/heartjnl-2012-301700
- Ishii M, Kaikita K, Sakamoto K, Seki T, Kawakami K, Nakai M, Sumita Y, Nishimura K, Miyamoto Y, Noguchi T, Yasuda S, Tsutsui H, Komuro I, Saito Y, Ogawa H, and Tsujita K. 2020. Characteristics and in-hospital mortality of patients with myocardial infarction in the absence of obstructive coronary artery disease in super-aging society. *Int J Cardiol* 301:108-113. 10.1016/j.ijcard.2019.09.037
- Johnston N, Jönellid B, Christersson C, Kero T, Renlund H, Schenck-Gustafsson K, and Lagerqvist B. 2015. Effect of Gender on Patients With ST-Elevation and Non-ST-Elevation Myocardial Infarction Without Obstructive Coronary Artery Disease. *Am J Cardiol* 115:1661-1666. 10.1016/j.amjcard.2015.03.006
- Kilic S, Aydın G, Çoner A, Doğan Y, Arican Özlük Ö, Çelik Y, Ungan I, Tascanov MB, Düz R, Polat V, Özkan H, Özyaşar M, Tülüce K, Kurt D, Çetin N, Gül M, Inci S, Yılmaz Coskun F, Arı H, Zoghi M, Ergene AO, and Türk U. 2020. Prevalence and clinical profile of patients with myocardial infarction with non-obstructive coronary arteries in Turkey (MINOCA-TR): A national multi-center, observational study. *Anatol J Cardiol* 23:176-182. 10.14744/AnatolJCardiol.2019.46805
- Li M, He Y, Cheang I, Zhang Z, Liu Y, Wang H, and Kong X. 2022. Clinical characteristics and outcome in patients with ST-segment and non-ST-segment elevation myocardial infarction without obstructive coronary artery: an observation study from Chinese population. *BMC Cardiovasc Disord* 22:21. 10.1186/s12872-021-02359-x
- Lindahl B, Baron T, Erlinge D, Hadziosmanovic N, Nordenskjöld A, Gard A, and Jernberg T. 2017. Medical Therapy for Secondary Prevention and Long-Term Outcome in Patients With Myocardial Infarction With Nonobstructive Coronary Artery Disease. *Circulation* 135:1481-1489. 10.1161/circulationaha.116.026336

Montalescot G, Dallongeville J, Van Belle E, Rouanet S, Baulac C, Degrandt A, and Vicaute E. 2007. STEMI and NSTEMI: are they so different? 1 year outcomes in acute myocardial infarction as defined by the ESC/ACC definition (the OPERA registry). *Eur Heart J* 28:1409-1417. 10.1093/eurheartj/ehm031

Montenegro Sá F, Ruivo C, Santos LG, Antunes A, Saraiva F, Soares F, and Morais J. 2018. Myocardial infarction with nonobstructive coronary arteries: a single-center retrospective study. *Coron Artery Dis* 29:511-515. 10.1097/mca.0000000000000619

Niccoli G, Scalone G, and Crea F. 2015. Acute myocardial infarction with no obstructive coronary atherosclerosis: mechanisms and management. *Eur Heart J* 36:475-481. 10.1093/eurheartj/ehu469

Nordenskjöld AM, Baron T, Eggers KM, Jernberg T, and Lindahl B. 2018. Predictors of adverse outcome in patients with myocardial infarction with non-obstructive coronary artery (MINOCA) disease. *Int J Cardiol* 261:18-23. 10.1016/j.ijcard.2018.03.056

Park HW, Yoon CH, Kang SH, Choi DJ, Kim HS, Cho MC, Kim YJ, Chae SC, Yoon JH, Gwon HC, Ahn YK, and Jeong MH. 2013. Early- and late-term clinical outcome and their predictors in patients with ST-segment elevation myocardial infarction and non-ST-segment elevation myocardial infarction. *Int J Cardiol* 169:254-261. 10.1016/j.ijcard.2013.08.132

Pasupathy S, Air T, Dreyer RP, Tavella R, and Beltrame JF. 2015. Systematic review of patients presenting with suspected myocardial infarction and nonobstructive coronary arteries. *Circulation* 131:861-870. 10.1161/circulationaha.114.011201

Pizzi C, Xhyheri B, Costa GM, Faustino M, Flacco ME, Gualano MR, Fragassi G, Grigioni F, and Manzoli L. 2016. Nonobstructive Versus Obstructive Coronary Artery Disease in Acute Coronary Syndrome: A Meta-Analysis. *J Am Heart Assoc* 5. 10.1161/jaha.116.004185

Polonski L, Gasior M, Gierlotka M, Osadnik T, Kalarus Z, Trusz-Gluza M, Zembala M, Wilczek K, Lekston A, Zdrojewski T, and Tendera M. 2011. A comparison of ST elevation versus non-ST elevation myocardial infarction outcomes in a large registry database: are non-ST myocardial

infarctions associated with worse long-term prognoses? *Int J Cardiol* 152:70-77.
 10.1016/j.ijcard.2010.07.008

Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, and van der Meer P. 2016. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 18:891-975. 10.1002/ehhf.592

Raparelli V, Elharram M, Shimony A, Eisenberg MJ, Cheema AN, and Pilote L. 2018. Myocardial Infarction With No Obstructive Coronary Artery Disease: Angiographic and Clinical Insights in Patients With Premature Presentation. *Can J Cardiol* 34:468-476. 10.1016/j.cjca.2018.01.004

Scalone G, Niccoli G, and Crea F. 2019. Editor's Choice- Pathophysiology, diagnosis and management of MINOCA: an update. *Eur Heart J Acute Cardiovasc Care* 8:54-62. 10.1177/2048872618782414

Tamis-Holland JE, Jneid H, Reynolds HR, Agewall S, Brilakis ES, Brown TM, Lerman A, Cushman M, Kumbhani DJ, Arslanian-Engoren C, Bolger AF, and Beltrame JF. 2019. Contemporary Diagnosis and Management of Patients With Myocardial Infarction in the Absence of Obstructive Coronary Artery Disease: A Scientific Statement From the American Heart Association. *Circulation* 139:e891-e908. 10.1161/cir.0000000000000670

Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, and White HD. 2018. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol* 72:2231-2264. 10.1016/j.jacc.2018.08.1038

Xu B, Abdu FA, Liu L, Mohammed AQ, Yin G, Luo Y, Xu S, Lv X, Fan R, and Che W. 2020. Clinical Outcomes and Predictors of ST-Elevation Versus Non-ST-Elevation Myocardial Infarction with Non-Obstructive Coronary Arteries. *Am J Med Sci* 360:686-692. 10.1016/j.amjms.2020.07.016

Figure legends

Fig. 1 Flow chart of patients included in this study.

421 Abbreviations: AMI, acute myocardial infarction, CAG, coronary angiography, STE, ST-
 422 segment elevation, NSTEMI, non-ST segment elevation, MINOCA, non-obstructive coronary
 423 arteries

424

425

426

Figure 1

Flow chart of patients included in this study

Flow chart of cases collection in this study.




Table 1(on next page)

Table 1 Comparision of the baseline characteristics between STEMI and NSTEM among MINOCA population

1

2 Table 1 Comparison of the baseline characteristics between STEMI and NSTEMI among MINOCA population

Variables	STEMI (n=115)	NSTEMI (n=81)	P
Demographics			
Age(years)	52.93±12.68	56.47±11.21	0.045
Male,n(%)	81 (70.43)	60 (74.07)	0.577
Coronary risk factors			
Diabetes, n (%)	11 (9.57)	12 (14.81)	0.261
Hypertension, n (%)	47 (40.87)	48 (59.26)	0.011
hyperlipidaemia, n (%)	33 (28.70)	23 (28.39)	0.927
previous arrhythmia, n (%)	9 (7.83)	7 (8.64)	0.837
Smoking history, n (%)	83 (72.13)	52 (64.20)	0.235
Killip grade, n (%)			
1 grade	103 (89.57)	64 (79.01)	0.040
≥2 grades	12 (10.43)	17 (20.99)	
hospitalization days(days)	6 (4,8)	7 (4,8)	0.305
Medications during hospitalization			
Aspirin, n (%)	112 (97.39)	73 (90.12)	0.018
Clopidogrel, n (%)	108 (93.91)	73 (90.12)	0.415
β-blocker, n (%)	51 (44.35)	41 (50.62)	0.468
Statins, n (%)	110 (95.65)	77 (95.06)	0.721
ACEI/ARB(%)	44(38.26)	45(55.56)	0.017
Laboratory indicators			
Myoglobin (ng/ml)	94(46.8,309.00)	101.5(53.08,178.75)	0.917
cTnT((ng/ml))	3.02(0.18,13.5)	1.34(0.22,5.87)	0.076
CK-MB	5.95(1.08,33.45)	3.80(1.00,12.81)	0.102
BNP	112(27.13,297.75)	61.90(20.40,186.00)	0.106
WBC count(×10 ¹² /L)	8.24(6.24,10.38)	7.52(6.01,9.28)	0.158
NE(%)	5.24(3.94, 8.04)	4.99(3.71,6.55)	0.231
RBC count(×10 ¹² /L)	4.61(4.27,4.97)	4.69(4.45,4.94)	0.457

PLT count($\times 10^{12}/L$)	221.5 (182.00,266.75)	210 (172.25,255.75)	0.262
TC (mmol/L)	4.20 (3.67,4.95)	4.51 (3.75, 5.25)	0.115
LDL-C (mmol/L)	2.44 (1.99,1.45)	2.45 (2.09,3.35)	0.109
HDL-C (mmol/L)	1.18 (0.99,1.45)	1.2 (1.05,1.48)	0.305
TG (mmol/L)	1.45 (1.06,2.37)	1.51 (1.11,2.21)	0.718
Serum glucose(mmol/L)	5.19 (4.66,5.95)	5.55 (4.87,6.44)	0.047
Echocardiography			
LV(mm)	49 (46, 52)	50 (46.75,52)	0.333
LVEF(%)	57 (54,60)	59 (55, 60)	0.128

3 Abbreviation:cTnT,blood cardiac troponin-T,CK-MB,creatine kinase-MB,BNP,brain natriuretic
4 peptide,RBC,Red blood cell,WBC,White blood cell,NE,
5 neutrophilicgranulocyte,PLT,Platelet,TC,total cholesterol,LDL-C,low-density lipoprotein
6 cholesterol,HDL-C,high-density lipoprotein cholesterol,TG,triglyceride, LV, left
7 ventricle,LVEF,left ventricular ejection fraction .

8
9
10
11
12
13

Table 2(on next page)

Table 2 Comparision of the rate of MACE in MINOCA during follow-up period.

Table 2 Comparision of the rate of MACE in MINOCA during follow-up period.

	STEMI (n=115)	NSTEMI (n=81)	<i>P</i>
MACE,n(%)	28 (24.35)	18 (22.22)	0.73
Chest pain rehospitalization, n (%)	4 (3.48)	4 (4.94)	0.72
nonfatal MI, n (%)	3 (2.61)	2 (2.47)	1
Heart failure, n (%)	14 (12.17)	8 (9.88)	0.654
Stroke, n (%)	5 (4.35)	3 (3.70)	1
Heart valve replacement, n (%)	-	2 (2.47)	0.17
All-cause deaths, n (%)	5 (4.35)	1 (1.23)	0.404

Table 3(on next page)

Table 3 Univariate analysis of MACE among STEMI and NSTEMI population

1

2

Table 3 Univariate analysis of MACE among STEMI and NSTEMI population

Factors	MACE _{STE} (n=28)	P _{STE}	MACE _{NSTE} (n=18)	P _{NSTE}
Age(years)	57.39±13.74	0.003	60.44±9.94	0.049
Male,n(%)	15 (53.57)	0.025	15 (83.33)	0.376
Diabetes, n (%)	5 (17.86)	0.086	5 (27.78)	0.126
Hypertension, n (%)	14 (50.00)	0.259	13 (72.22)	0.204
hyperlipidaemia, n (%)	6 (21.43)	0.266	3 (16.67)	0.245
previous arrhythmia, n (%)	2 (7.14)	1.000	3 (16.67)	0.339
Smoking history, n (%)	19 (67.85)	0.558	11 (61.11)	0.757
Killip grade, n (%)				0.145
1 grade	20 (71.43)	0.001	12 (66.67)	
≥2 grades	8 (28.57)		6 (33.33)	
Laboratory indicators				
Myoglobin (ng/ml)	97.25 (50.20,248.25)	0.539	105.00 (53.63,173.25)	0.934
cTnT(ng/ml)	2.65 (0.07,14.57)	0.661	0.54 (0.19,8.99)	0.578
CK-MB(ng/ml)	4.17 (1.00,25.55)	0.481	3.79 (1.00,15.12)	0.986
BNP	120.00 (32.60,848.00)	0.305	85.45 (31.55,249.45)	0.150
WBC count(×10 ¹² /L)	8.20 (5.97,11.65)	0.931	7.52 (5.59,8.79)	0.461
RBC count(×10 ¹² /L)	4.43 (4.15,4.66)	0.016	4.68 (4.43,4.93)	0.773
PLT count(×10 ¹² /L)	219.00 (186.00,255.00)	0.401	198.50 (154.75,250.50)	0.450
TC (mmol/L)	4.11 (3.35,4.94)	0.276	4.68 (3.52,5.37)	0.775
LDL-C (mmol/L)	2.32 (1.77,2.80)	0.085	2.63 (1.98,3.20)	0.579
HDL-C (mmol/L)	1.21 (1.02,1.60)	0.372	1.31 (1.17,1.59)	0.113
TG (mmol/L)	1.33 (0.90,1.73)	0.170	1.36 (0.95,2.00)	0.229
Serum glucose(mmol/L)	5.46 (4.73,6.72)	0.262	5.62 (4.84,6.74)	0.775
Echocardiography				
LV (mm)	49.00 (46.00,52.00)	0.989	50.50 (47.75,55.00)	0.191
LVEF (%)	58.00 (51.00,60.00)	0.949	58.00 (50.75,60.50)	0.330

hospitalization days(days)	7.00 (5.00,8.00)	0.037	6.50 (3.75,9.00)	0.991
Medications during hospitalization				
Aspirin, n (%)	28 (100.00)	1.000	14 (77.78)	0.086
Clopidogrel, n (%)	26 (92.86)	0.634	17 (94.45)	0.677
β-blocker, n (%)	8 (28.57)	0.048	5 (27.78)	0.028
Statins, n (%)	28 (100.00)	0.571	17 (94.45)	1.000
ACEI/ARB(%)	25 (89.28)	0.754	16(88.89)	0.534

3 Abbreviation: cTnT, blood cardiac troponin-T, CK-MB, creatine kinase-MB, BNP, brain
 4 natriuretic peptide, RBC, Red blood cell, WBC, White blood cell, PLT, Platelet, TC, total
 5 cholesterol, LDL-C, low-density lipoprotein cholesterol, HDL-C, high-density lipoprotein
 6 cholesterol, TG, triglyceride, LV, left ventricle, LVEF, left ventricular ejection fraction.

7
 8
 9

Table 4(on next page)

Table 4 Multivariable predictors of MACE in STEMI patients.

Table 4 Multivariable predictors of MACE in STEMI patients.

Factors	OR	95%CI	<i>P</i>
Killip grade	9.035	(1.657,49.263)	0.011
β-blocker	0.238	(0.072,0.788)	0.019
LDL-C	2.267	(1.008,5.097)	0.048

Abbreviation:LDL-C,low-density lipoprotein cholesterol.