

Reducing options of ammonia volatilization and improving nitrogen use efficiency via organic and inorganic amendments in wheat (*Triticum aestivum* L.)

Pooja L Ramalingappa¹, Manoj Shrivastava¹, Shiva Dhar², Kalikinkar Bandyopadhyay³, Shiv Prasad¹, Sapna Langyan⁴, Ritu Tomer¹, Ashish Khandelwal¹, Sibananda Darjee¹, Renu Singh^{Corresp. 1}

¹ Division of Environment Science, ICAR-Indian Agricultural Research Institute, Delhi, India

² Division of Agronomy, ICAR-Indian Agricultural Research Institute, Delhi, India

³ Division of Physics, ICAR-Indian Agricultural Research Institute, Delhi, India

⁴ Division of Germplasm Evaluation, ICAR- National Bureau of Plant Genetic Resources, Delhi, India

Corresponding Author: Renu Singh

Email address: renu_icar@yahoo.com

Background. This study focuses to investigate the effect of organic and inorganic supplements on the reduction of ammonia (NH₃) volatilization, improvement in Nitrogen Use Efficiency (NUE), and wheat yield. **Methods.** A field experiment was conducted following a Randomized Block Design with 10 treatments *i.e.*, T₁-without nitrogen (control), T₂-Recommended Dose of Nitrogen (RDN), T₃-(N-(n-butyl) thiophosphoric triamide (NBPT @ 0.5% w/w of RDN), T₄-Hydroquinone (HQ @ 0.3% w/w of RDN), T₅-Calcium carbide (CaC₂ @ 1% w/w of RDN), T₆-Vesicular Arbuscular Mycorrhiza (VAM @ 10 kg ha⁻¹), T₇-(Azotobacter @ 50 g kg⁻¹ seeds), T₈-(Garlic powder @ 0.8% w/w of RDN), T₉-(Linseed oil @ 0.06% w/w of RDN), T₁₀-(Pongamia oil @ 0.06% w/w of RDN). **Results.** The highest NH₃ volatilization losses were observed in T₂ at about 20.4 kg ha⁻¹ per season. Significant reduction in NH₃ volatilization losses were observed in T₃ by 40%, T₄ by 27%, and T₈ by 17% when compared to the control treatment. Soil urease activity was found to be decreased in plots receiving amendments, T₃, T₄, and T₅. The highest grain yield was observed in the T₇ treated plot with 5.09 t ha⁻¹, and straw yield of 9.44 t ha⁻¹ in T₄. **Conclusion.** The shifting towards organic amendments is a feasible option to reduce NH₃ volatilization from wheat cultivation and improves NUE.

Reducing options of ammonia volatilization and improving nitrogen use efficiency via organic and inorganic amendments in wheat (*Triticum aestivum* L.)

Pooja L Ramalingappa¹, Manoj Shrivastava¹, Shiva Dhar², Kalikinkar Bandyopadhyay³, Shiv Prasad¹, Sapna Langyan⁴, Ritu Tomer¹, Ashish Khandelwal¹, Sibananda Darjee¹, and Renu Singh¹

¹Division of Environment Science, ICAR-Indian Agricultural Research Institute, New Delhi, 110012, India

²Division of Agronomy, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India

³Division of Agricultural Physics, ICAR-Indian Agricultural Research Institute, New Delhi, 110012, India

⁴Division of Germplasm Evaluation, ICAR-National Bureau of Plant Genetic Resources, New Delhi 110012, India

Corresponding Author:

Renu Singh¹

Division of Environment Science, ICAR-Indian Agricultural Research Institute, New Delhi, 110012, India

Email address: renu_icar@yahoo.com

Abstract

Background. This study focused on investigating the effect of organic and inorganic supplements on the reduction of ammonia (NH_3) volatilization, improvement in Nitrogen Use Efficiency (NUE), and wheat yield.

Methods. A field experiment was conducted following a Randomised Block Design with 10 treatments, *i.e.*, T₁-without nitrogen (control), T₂-Recommended Dose of Nitrogen (RDN), T₃-(N-(n-butyl) thiophosphoric triamide (NBPT @ 0.5% w/w of RDN), T₄-Hydroquinone (HQ @ 0.3% w/w of RDN), T₅-Calcium carbide (CaC_2 @ 1% w/w of RDN), T₆-Vesicular Arbuscular Mycorrhiza (VAM @ 10 kg ha⁻¹), T₇-(Azotobacter @ 50 g kg⁻¹ seeds), T₈-(Garlic powder @ 0.8% w/w of RDN), T₉-(Linseed oil @ 0.06% w/w of RDN), T₁₀-(Pongamia oil @ 0.06% w/w of RDN).

Results. The highest NH_3 volatilization losses were observed in T₂ at about 20.4 kg ha⁻¹ per season. Significant reductions in NH_3 volatilization losses were observed in T₃ by 40%, T₄ by 27%, and T₈ by 17% compared to the control treatment. Soil urease activity was found to be decreased in treatment T₃, T₄, and T₅ amended plots. The highest grain yield was observed in treatment T₇ with 5.09 t ha⁻¹, and straw yield was the maximum in treatment T₄ with 9.44 t ha⁻¹.

Conclusion. Based on the experimental data obtained, it can be concluded that organic amendments would be a feasible option to reduce NH_3 volatilization from wheat cultivation and improves NUE.

Introduction

Nitrogen (N) in the atmosphere is the principal source of all soil nitrogen. It naturally enters the soil through dead animal and plant residues, biological nitrogen fixation, and chemical N fertilizer applications. Nitrogen fertilizers have become essential to increase crop yield and enhance food quality (Leghari *et al.*, 2016). Food grain crops account for more than 69% of India's total N fertilizer intake, where wheat alone has a share of 24%. In the crop year 2020-21, India's wheat production increased from 109 Mt to 118 Mt to meet the growing

population's demand. However, in future, the wheat yield has to be increased by 1.5% per year to satisfy the growing population's demand (*Grain and Feed Annual report, 2021*). Ammonia volatilization is a significant cause of nitrogen depletion in agricultural soil worldwide, contributing to low N fertilizer usage, crop production, and indirect nitrous oxide (N₂O) emissions. Globally, the average NH₃ volatilization risks range from 0.9 to 64 % of the applied N (a mean of 17.6 %) (*Pan et al., 2016*). Ammonia volatilization into the atmosphere negatively affects agriculture, ecosystems, and human health.

Further, it also increases the loss of nitrogen for plant growth, thus increasing the cost of cultivation (*Brink & Van Grinsven, 2011*). Secondary particulate matter (PM₁₀) is formed when NH₃ reacts with other air contaminants such as sulfuric acid and nitric acid. It flies long distances and remains in the air for several days, and causes respiratory diseases in humans (*Bittman et al., 2014*).

A study conducted by Cao et al. (2013) found that the significant loss of N from applied fertilizer was through NH₃ volatilization, which was estimated to be 10.0-19.5% of total N loss. Further, they concluded that NH₃ volatilization might be the dominant pathway of N loss. Ammonia volatilization favourably occurs in N fertilizers like urea and organic manure. However, ammonia emission is undesirable as it removes N from the soil/plant system and releases it into the atmosphere. Volatilized NH₃ is deposited back to the earth's surface mainly through two processes which are i) Wet deposition through precipitation and ii) Dry deposition when combined with particulate matter. This seriously impacts the environment as it causes acidification of soil and water bodies and eutrophication of the natural ecosystem. It also acts as an indirect source of N₂O, a potent greenhouse gas. In some instances, it also directly affects plants under high concentrations and low temperatures. There are many other ways to control N loss, like avoiding the application of urea under high-risk conditions, deep placement, and using controlled-release N-fertilizers. But the more effective way to control is by using organic and inorganic amendments; some of them are chemical inhibitors like N-(n-butyl) thiophosphoric triamide (NBPT), Cyclohexyl phosphoric triamide (CHPT), Ammonium thiosulphate (ATS), Hydroquinone (HQ), Calcium carbide (CaC₂). Among these, NBPT is used globally, being the most effective in a market that has expanded at 16 % per year over the last ten years. NBPT-treated urea reduces NH₃ loss by 53% (*Cantarella et al., 2018*). Application of 12 kg N ha⁻¹ HQ on an Alluvial soil, in conjunction

with 120 kg urea-N ha⁻¹, decreased N₂O emission by 5% in rice and 7% in wheat systems when compared to the crops grown solely in the presence of 120 kg N ha⁻¹ urea (*Modolo et al., 2018*).

The slow-N-release coating technology is also suitable for consistent N supply to the plants and reducing loss and contamination effects. We have explored various organic and inorganic amendments to minimize NH₃ volatilization losses and enhance NUE and wheat crop productivity. The application of N inhibitors along with urea has increased the average grain yield by 6.8% (*Školníková et al., 2022*). Nitrification and urease inhibitors are being suggested to decrease N losses and thus increase crop nitrogen usage efficiency (*Abalos et al., 2014*). Different mechanisms are involved in reducing NH₃ volatilization in that NBPT retards the activity of the urease enzyme by competitive inhibition (*Fan et al., 2018*). Hydroquinone and Calcium carbide is also effective in reducing urease activity. Applying HQ to the soil inhibited urease enzyme activity and inhibited or enhanced the activity of other enzymes like polyphenol oxidase, dehydrogenase, protease and phosphatase (*Wolejko et al., 2020*). The CaC₂ has a negative impact on ammonia-oxidizing bacteria, which are present in the soil, and also reduces the activity of dehydrogenase and nitrate reductase enzymes, thereby reducing the N losses (*Mahmood et al., 2014*). Pongamia oil (karanjin) works as a nitrification inhibitor by reducing the *Nitrosomonas* activity without affecting *Nitrobacter spp.* Activity. The treatment with garlic extract, which is rich in the compound thiosulfinate, works on the mechanism of competitive inhibition of urease enzyme as it contains an organosulphur group (-S(O)-S) which is similar to urea which helps in inhibiting the hydrolysis process (*Mathialagan et al., 2017*).

The world's population will surpass 9.7 billion by 2050, posing a significant obstacle to achieving food sustainability. The projected increase in the world's population demands at least a 70% increase in agriculture production in developed countries and 100% in developing countries (*Mahmud et al., 2021*). Food security of the country and N-based global warming and environmental degradation are interlinked (*Bilal & Aziz, 2022*). The use of urease inhibitors in agricultural activities has already been investigated as one of the best methods for ensuring adequate nutritional security (*Modolo et al., 2018*). N-(n-butyl) thiophosphoric triamide treated urea lowers NH₃ loss by approximately 53%. The yield benefits from NBPT application on an average of 6.0 % and ranges from 0.8 to 10.2 % depending on crop types

(Cantarella et al., 2018). The effectiveness of NBPT in reducing NH_3 loss is well known, but there is still room for progress to increase the amount of inhibition and hence the retention of NBPT-treated urea in the field. However, these inorganic amendments are not economically viable, and most of them restricted their usage in the research areas due to high costs. Upadhyay, Tewari & Patra (2011) investigated that these chemicals are harmful and inhibit the growth of specific beneficial soil microorganisms that indirectly affects crop growth and development. Therefore, the best option that stands before us is using organic amendments to reduce NH_3 losses. Organic amendments are eco-friendly, economically viable, and, most importantly, biodegradable. Therefore, they act as promising N inhibitors. There are many plant-derived inhibitors like Neem (*Azadirachta indica*) oil, Pongamia (*Pongamia glabra*) oil, Linseed (*Linum usitatissimum*) oil, Garlic extract, and Mint (*Mentha spicata*) as N amendments in inhibiting N losses. Thiosulfinates (TS) present in fresh garlic extract act as a bio-inhibitor of urease enzyme activity and can be used as a potential urease inhibitor in agriculture. Thiosulfinates have the potential to inhibit the urease enzyme, and it takes three hours and thirty minutes to start acting on the urease enzyme after its' application (Ramli et al., 2014). Applying coated urea fertilizers such as neem-coated urea and pine oleoresin-coated urea in a vertisol reduced the NH_3 volatilization by 27.5% and 41.1%, respectively (Jadon et al., 2018). But these organic amendments lack commercialization and efficient utilization. So this study concentrates mainly on stressing the importance and comparative study of these organic and inorganic amendments to decrease ammonia volatilization loss and improve increasing NUE and productivity of Wheat (*Triticum aestivum* L.) crops.

Materials & Methods

Details of the experimental site

A field study was conducted at the research farm of the ICAR-Indian Agricultural Research Institute, New Delhi, located at 28° 40' N and 77° 12' E, at an altitude of 228.16 m above mean sea level (sub-tropical and semi-arid region), during November 2020 to April 2021. The Southwest monsoon contributed about 80% of rainfall, an average of 650 mm annually. The pH of the soil was 8 (± 0.1), with a sandy clay loam texture. The initial soil nutrients status, when analyzed before sowing of seeds, were found as follows, the available N was low

(157±0.5 kg ha⁻¹), medium level of available P (13.5±0.2 kg ha⁻¹), and the available K (196±0.8 kg ha⁻¹).

Experimental design and management

The experiment was conducted in a randomized block design (RBD) with 3 replications, and each plot area was 12 m² (4 m×3 m). In all treatments recommended dose of nitrogen (RDN) was applied except T₁ (control), where no RDN was used. The research was carried out through growing wheat variety (HD 2967) with 10 treatments in respective plots, namely labelled as T₁ (control), T₂ (RDN @ 150 kg ha⁻¹), T₃ (RDN + NBPT (N-(n-butyl) thiophosphoric triamide) @ 0.5% w/w RDN), T₄ (RDN +HQ (Hydroquinone) @ 0.3% w/w of RDN), T₅ (RDN + Calcium Carbide @ 1% w/w of RDN), T₆ (RDN + VAM (Vesicular Arbuscular Mycorrhiza @ 10 kg ha⁻¹), T₇ (RDN + Azotobacter @ 50 g kg⁻¹ seeds) T₈ (RDN + Garlic powder @ 0.8% w/w of RDN), T₉ (RDN + Linseed oil @ 0.06% w/w of RDN), and T₁₀ (RDN + Pongamia oil @ 0.06% w/w of RDN). Each dose was finalized based on the current usage of these amendments by firms, which were applied in 2 splits (One at the time of sowing and the second split application 30 days after sowing).

In order to study the effect of organic and inorganic amendments together, three inorganic amendments, two bio-fertilizers, and three organic amendments were chosen based on popularity and literature review. N-(n-butyl) thiophosphoric triamide (*Zanin et al., 2015*), Hydroquinone (*Modolo et al., 2018*), Calcium Carbide (*Sakariyawo et al., 2020*), Garlic powder (*Ramli et al., 2014*), Linseed oil, and Pongamia oil (*Majumdar, 2008*) were mixed with urea (RDN) as per the dosage. Wheat seeds were treated with *Azotobacter spp.* (as per ICAR RABI Agro-Advisory for Farmers) and used for sowing in T₇. Vesicular Arbuscular Mycorrhiza (as per ICAR RABI Agro-Advisory for Farmers) was applied directly into the soil during sowing with an RDN. The recommended dose of fertilizers (RDF) for wheat was applied as N: P₂O₅:K₂O (150:60:40 kg ha⁻¹). Urea, Single Super Phosphate (SSP), and Muriate of Potash (MOP) were used as a source of N, P₂O₅, and K₂O, respectively.

Method of Soil sample collection and analysis

Fresh soil samples were obtained from the 0-15 cm layer of soil at three separate sites from each treatment using an 8 cm tube auger. Three soil samples were obtained from each

treatment during tillering, flowering, grain filling, and physiological maturity of the crop. The total fresh soil samples were 30 in number and air-dried for 7 days, sieved through a 2 mm screen, mixed, and placed in plastic bags for further analysis.

Collection and analysis of NH₃

The ammonia volatilization was monitored after fertilizer application for up to 10 days using a forced air draft system method (*Bhaskar et al., 2022; Stumpe et al., 1984; Bremner, 1965*).

The closed chambers measuring 20 cm × 20 cm × 50 cm size made of 6 mm acrylic sheets were placed in the field. The volatilized NH₃ gas from the soil surface under different treatments was collected in a 2 % Boric acid solution containing a mixed indicator (methyl red and bromocresol green). The air inside the chamber was collected into boric acid traps using a vacuum pump having a flow rate of 3 L min⁻¹. The boric acid traps were changed every 24 hr. The volatilized NH₃ was determined by the titration of boric acid solution with 0.02 N sulphuric acid, and further calculations were done using the formula given below (Eq.1).

$$\text{Volatilized ammonia (mg per m sq. per day)} = \frac{A * 0.00028 * 1000}{L * B} \dots\dots\dots \text{Eq.1}$$

Total Nitrogen and NH₄⁺-N and NO₃⁻-N analysis

Total nitrogen content in soil was determined by the Kjeldahl method (*Kjeldahl, 1883*) during the initial and after harvesting of crop growth. The NH₄⁺-N and NO₃⁻-N were analyzed through steam distillation (*Keeney and Bremner, 1965*) during all four crop growth stages (Tillering, Flowering, Grain-filling, and Physiological maturity). The extract was prepared by taking 10 g soils with 0.25 g activated charcoal and 50 mL KCl solution and kept for shaking (30 min), then filtered with Whatman filter paper 1. From the same KCl extract, 10 mL each was taken in two different distillation flasks, and 100 mL of distilled water was added to each flask. In addition, 1g Devardas alloy was added for the case of NH₄⁺-N estimation and NO₃⁻-N estimation 1 g magnesium oxide (MgO) was added and distilled separately, and these ions were captured in 20 mL of 2% Boric acid and titrated against 0.02 N sulfuric acid. Further calculations were done using equations 2 and 3 for NH₄⁺-N estimation and NO₃⁻-N estimation, respectively.

$$\text{Exchangeable ammonical N \% in soil} = \frac{(V_s - V_b) * S * 0.014 * 100}{W} = Z1 \dots\dots\dots \text{Eq.2}$$

$$\text{Exchangeable NH}_4^+\text{-N (ppm)} = Z1 * 10^4$$

$$\text{Exchangeable nitrate N \% in soil} = \frac{(V_s - V_b) * S * 0.014 * 100}{W} = Z2 \dots \text{Eq. 3}$$

$$\text{Exchangeable NO}_3^- \text{-N (ppm)} = Z2 * 10^4$$

Where,

V_s denotes the volume of H₂SO₄ available for sample titration.

V_b denotes the volume of H₂SO₄ needed for blank titration.

S = H₂SO₄ power,

W = Weight of oven-dried soil used for analysis

223

224 Available Phosphorus (P) was analyzed using Olsens' estimation method (*Olsen et al., 1954*).

225 First, available P from the soil sample was extracted using 0.5 N NaHCO₃ solution buffer at

226 pH 8.5. Then available P in the extract was measured by an ascorbic acid method using a

227 spectrophotometer. Next, available potassium (K) in the soil was measured using an

228 ammonium acetate method (*Hanway & Heidel, 1952*), where available K was extracted by

229 shaking with neutral normal ammonium acetate for 5 min, and the K was determined using a

230 flame photometer. Finally, soil Organic Carbon was measured using Walkley and Black's

231 rapid titration method (*Walkley & Black, 1934*).

232

233 **Analysis of Nitrous oxide (N₂O) and other significant parameters**

234 Nitrous oxide flux was analyzed using the closed chamber method (*Herr et al., 2020*). In this

235 method, dark PVC boxes were installed, and the samples were drawn every 24 hrs in the

236 morning using syringes, evacuated into plastic vials, and analyzed chromatographically.

237 Denitrification losses were estimated by the denitrification enzyme assay method described by

238 Smith & Tiedje (*1979*).

239 Soil urease activity was analyzed at the 50% flowering stage, calorimetrically, by Bremner &

240 Douglas's (*1971*) method. Normalized Difference Vegetation Index (NDVI) was measured

241 using a green seeker (handheld crop sensor by Trimble) at the 50% flowering stage. Infrared

242 Gas Analyzer (LI-COR Model LI-6400X7 portable photosynthetic system) (IRGA) was used

243 to measure the photosynthetic rate and stomatal conductance. Soil Microbial Biomass Carbon

244 (MBC) and Soil Microbial Biomass Nitrogen (MBN) were determined by the chloroform

245 fumigation–extraction method described by Vance et al. (*1987*) and Brookes et al. (*1985*),

respectively. The N content in grains and straws was also measured using the Kjeldahl method (Kjeldahl, 1883). After harvesting the crop, yield attributes were calculated from each plot.

Calculation of Nitrogen Use Efficiency in wheat

The Nitrogen Use Efficiency can be defined as the ratio of outputs to inputs of nitrogen (*i.e.*, $NUE = N \text{ yield}/N \text{ input}$).

$$\text{Agronomic Efficiency of N (AEN) (kg ha}^{-1}\text{)} = \frac{GYF - GYN}{AFN} \dots\dots\dots \text{Eq. 4}$$

$$\text{Production Efficiency of N (PEN) (kg kg}^{-1}\text{)} = \frac{GYF - GYN}{TUN - CUN} \dots\dots\dots \text{Eq. 5}$$

$$\text{Apparent N Recovery (ANR) (\%)} = \frac{TUN - CUN}{AFN} \times 100 \dots\dots\dots \text{Eq. 6}$$

TUN = Total N uptake from the fertilized plots (kg ha⁻¹)

CUN = Total N uptake from unfertilized / control plots (kg ha⁻¹)

AFN = Amount of applied fertilizer N (kg ha⁻¹)

GYF = Grain yield in the fertilized plots (kg ha⁻¹)

GYN = Grain yield in unfertilized/control plots (kg ha⁻¹)

2.7 Statistical analysis

The measurements obtained from the experimental work were analyzed by using OPSTAT.01 Software (O.P. Sheoran, Hisar, India) was used to calculate ANOVA, and means were separated using Duncan Multiple Range Test (DMRT) at $\alpha=0.05$. The data for each variable was evaluated using variance protocol analysis for a Randomized block design, which was checked using the "F" test for statistical significance (Gomez & Gomez, 1984). The Standard Error of means (SEm) and Critical Difference (CD) parameters were calculated at a 5% significance level.

Results

Ammonia volatilization losses during the wheat growth period

The addition of organic and inorganic amendments in wheat crop have shown effective controlled of ammonia volatilization losses during entire cropping season as shown in **Figure 1**. Results revealed that significant losses of N through NH₃ volatilization were observed in the initial 5-6 days after N fertilizer application. The highest N volatilization losses were

observed in T₂ (20.4 kg ha⁻¹ season⁻¹) and the lowest losses were found in T₁ (6.4 kg ha⁻¹ season⁻¹). T₃ performed well in reducing the NH₃ losses by 40% as compared to T₂. NH₃ losses in T₃, T₄ and T₈ were recorded 12.4, 14.6, and 16.6 kg ha⁻¹ season⁻¹, respectively (Figure 2). Duncan Multiple Range Test revealed that T₃ (NBPT) and T₄ (HQ) effectively reduced the NH₃ losses. T₅ (CaC₂) and T₈ (Garlic powder) showed no significant difference among their means. However, among plant-based amendments, T₈ with garlic powder-treated urea showed better results by reducing NH₃ volatilization losses by 17% compared to only RDF-treated urea (T₂). Other treatments were not so effective in reducing the NH₃ losses.

Initial and final soil nutrient status

Available nitrogen content in soil after wheat crop harvest was found highest in treatment T₇ (*i.e.*, Azotobacter treated plot) which was 195 kg ha⁻¹. Detailed observations on Available N content for different treatments is given in Table 1. The highest SOC was observed in T₇, treated with Azotobacter spp., *i.e.*, 0.63%. It significantly increased the SOC content compared to the initial SOC (0.42%) before sowing the wheat crop, as shown in Figure 3. The highest level of available P was observed in T₆ treated with VAM, *i.e.*, 30.4 kg ha⁻¹, which is more than double the initial value of available P, as shown in Table 1. The initial level of available K was 196 kg ha⁻¹, while in the final analysis after harvesting the crop, it increased to 229 kg ha⁻¹ in T₁₀, as depicted in Table 1.

Ammonical N (NH₄⁺-N) and Nitrate (NO₃⁻-N) in the soil

The available nitrogen forms, mostly NH₄⁺-N and NO₃⁻-N in the soil, decreased from tillering to the grain filling stage (Figure 4.). The mean quantities of NH₄⁺-N in the soil at various crop growth stages (tillering, flowering, grain filling, and physiological maturity stage) were 78.6 kg ha⁻¹, 65.6 kg ha⁻¹, 57.9 kg ha⁻¹, and 53.5 kg ha⁻¹, respectively. The mean of NO₃⁻-N in the soil at all four growth stages was 70 kg ha⁻¹, 63.4 kg ha⁻¹, 56.5 kg ha⁻¹, and 51.1 kg ha⁻¹, respectively. During the tillering stage, the highest NH₄⁺-N (108 kg ha⁻¹) was observed in T₃, *i.e.*, treatment with NBPT amended urea, and the highest NO₃⁻-N (58.5 kg ha⁻¹) was found in RDN treated plot (T₂). In all the stages of plant growth, T₃ showed significantly higher values of NH₄⁺-N, which was 47% higher than T₂. Duncan Multiple Range Test also showed that the mean of treatments T₄, T₇, and T₈ was also significantly at par for the available NH₄⁺-N

content in the soil, and in the case of NO_3^- -N availability, the mean of treatments T_4 , T_7 , and T_5 also showed significant difference during all the stages of the wheat crop growth.

Nitrous oxide flux and denitrification losses from the soil

Nitrous oxide emission is directly related to the amount of N fertilizer available in the soil. The highest cumulative N_2O flux was observed in T_2 (RDF) plots with $6.52 \text{ kg N}_2\text{O-N ha}^{-1}$ season⁻¹. There were no significant differences between T_3 , T_4 , and T_5 in the case of N_2O emission reduction (Figure 3). But all of them were significant when compared with T_2 . Among plant-based inhibitor treatments, T_8 , having urea amended with garlic powder, showed effective results. On the other hand, the highest denitrification losses were observed in T_2 ($3.66 \text{ kg-N ha}^{-1}$) (Figure 5). N-(n-butyl) thiophosphoric triamide treated urea was comparatively effective by reducing the denitrification losses by 35 %.

Soil microbial biomass carbon and nitrogen in the soil

Soil biomass carbon and soil biomass nitrogen (found within living organisms like fungi and bacteria) were observed at the 50% flowering stage of the crop. The highest MBN content was reported in T_7 , i.e., *Azotobacter* spp. treated plot with 37 mg kg^{-1} , followed by T_6 , having VAM treated plot. The highest MBC (136 mg kg^{-1}) was observed in T_7 , *Azotobacter* spp. treated plot was 63.9% more than T_2 (Figure 6).

Soil urease enzyme activity in the soil

As shown in Figure 7, under the 50% flowering stage in T_2 , i.e., only RDN treated plot showed the highest urease activity ($18.9 \text{ mg urea g}^{-1} \text{ soil h}^{-1}$) and the lowest urease activity ($7.71 \text{ mg urea g}^{-1} \text{ soil h}^{-1}$) was observed in T_3 treatment containing NBPT which was indicated by the effectiveness of urease inhibitor.

Physiological parameters of wheat crop

Adding organic and inorganic amendments to the soil has significantly influenced the physiological parameters. The range of photosynthesis rate varied from $15.7 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ to $23.8 \mu \text{ mol m}^{-2} \text{ s}^{-1}$, which was recorded during the 50% flowering stage. The highest photosynthetic rate was observed in T_7 with $23.8 \mu \text{ mol m}^{-2} \text{ s}^{-1}$, followed by T_3 and T_6 ($23.3 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ and $22.9 \mu \text{ mol m}^{-2} \text{ s}^{-1}$).

The range of chlorophyll content varied from 0.66 to 0.77, recorded during the 50% flowering stage. The highest chlorophyll content was observed in both T₄ and T₇, with 0.77, followed by T₆ and T₁₀, showing 0.76. The highest LAI was observed in T₇ with 4.59, followed by T₃ and T₈, which recorded the same LAI, i.e., 4.16. The lowest reading was recorded in treatment T₁, i.e., 1.85. The highest stomatal conductance was observed in T₇ with 0.48 m mol m⁻² s⁻¹, followed by T₃ and T₈ (0.47 m mol m⁻² s⁻¹ and 0.43 m mol m⁻² s⁻¹), respectively.

Nitrogen uptake by wheat

The data about the nitrogen content analyzed in wheat grain and straw samples after harvest depicted the positive effect of both organic and inorganic amendments. The *Azotobacter spp.* treated plot (T₇) showed 38.7% higher nitrogen content in grains when compared to T₂ treatments, as shown in Table 1. The highest nitrogen content in straw was observed in T₃ (NBPT), almost double the straw nitrogen content of the T₁ (control) plot.

Wheat yield and nitrogen use efficiency

The data obtained in grain and straw yield showed significant improvement in yields due to the addition of organic and inorganic amendments. The highest grain yield was observed in T₃, i.e., treatment with NBPT amended urea with 5.09 t ha⁻¹, and straw yield was in T₄, i.e., treatment with hydroquinone amended urea (Table 2). However, in the case of straw yield, all the treatment means were not significantly different at $p=0.05$. The other yield attributes like biological yield, number of tillers per m⁻², number of spikes per m⁻², number of grains per spike, and harvest index were not significantly influenced by amendments addition ($p=0.05$). However, the test weight of the wheat grain of treatment T₈ (39.8 g) was significant among all the treatments.

As nitrogen content in the grains has increased on amending urea with organic and inorganic amendments, NUE was calculated in terms of Agronomic Efficiency of N (AEN), Production Efficiency of N (PEN), and Apparent N Recovery (ANR) using Equations 4,5, and 6. The highest AEN was found in T₃, with 12.9 kg ha⁻¹, and the lowest was in T₂, with 9.05 kg ha⁻¹ (Table 1). The highest PEN was observed in T₂ with 25.8 kg kg⁻¹, and the lowest was in T₇ (15.4 kg kg⁻¹). The average value of PEN was 17.9 kg kg⁻¹. The highest ARN (81 %) was seen in T₇, and the lowest was in T₂ (35 %). The mean value of ANR was observed to be 49.6%.

Discussion

Our experimental findings in the case of NH_3 volatilization losses depicted that T_3 outperformed all other treatments because of competitive inhibition of urease enzyme activity by NBPT in the soil, which has a significant role in urea hydrolysis process where urea molecule is converted into ammonium ion. Next to NBPT, the other inorganic amendments significantly decreased the NH_3 volatilization losses. The allicin (thiosulfinate) naturally present in garlic is a bio-inhibitor of urease activity, as it contains an organosulfur functional group like that of the ureas'. This might help allicin to decrease urease activity. This is in agreement with the research study conducted by Mathialagan et al. (2017). These amendments have practical implications in reducing the quantity of urea applied to the field, thereby reducing the N losses, even though many studies have been taken on the case of NBPT. There is still a need to improve the duration of inhibition of urease activity, the shelf life of urea fertilizer coated with NBPT, and the economic viability of these compounds. Agriculture alone contributes 80-90% of NH_3 emissions globally, mainly through volatilization from livestock and synthetic nitrogen fertilizer (Xu et al., 2019). It acts as a pollutant influencing the biosphere through haze formation and soil acidification. N_2O , a potent greenhouse gas, is also of great concern, contributes to global warming, and affects human and environmental health. Li et al. (2015) reported that the Limus® (a new urease inhibitor consisting of 75% NBPT and 25% N-(n-propyl) thiophosphoric triamide (NPPT)), showed an average 83% decrease in NH_3 losses during winter wheat season in China. Only the addition of NBPT can reduce NH_3 volatilization losses by 61% to 74%, as reported by Lasisi, Akinremi & Kumaragamage (2019). Affendi, Mansor & Samiri (2019) conducted a study by adding various chemical and natural urease inhibitors to reduce ammonia and nitrous oxide losses from soil. They reported similar findings in a combination of thiosulfate with urea. However, they found NBPT was more effective than the combined use of thiosulfate with urea to reduce NH_3 volatilization losses. Similar results were reported by Eduardo et al. (2016), reported a decrease of 31.6% N, which used to be lost if not applied with thiosulfinates, and concluded it was an upcoming natural urease inhibitor.

The final analysis of TNC, available N, P, and K, reported an increase in all these nutrient contents to the initial analysis, which helps the next sown crop in nutrition. T₇ treated with *Azotobacter spp.* performed well in increasing N and K content in the soil, and T₆ was treated with VAM to increase P content in the soil. An increase in the N content of the soil could be due to adding these bio-fertilizers as amendments that significantly increased residual N and reduced the N fertilizer application for the next crop, which will be sown in that plots. Vesicular Arbuscular Mycorrhiza improves P mobilization; hence uptake by wheat crops and P content in soil increases. The addition of bio-fertilizers as an amendments source to reduce N losses has improved the soil SOC.

The highest SOC was observed in T₇, which was treated with *Azotobacter spp.* i.e., 0.63%. It has doubled the SOC content compared to the initial SOC before sowing the wheat crop, i.e., 0.42%. A similar effect of these bio-fertilizers in increasing total nitrogen content was observed in a study on Bio-fertilizer affecting structural dynamics, function, and network patterns of the sugarcane rhizospheric microbiota by Liu et al. (2021). Kader et al. (2002) also found similar results while working on the effect of *Azotobacter spp.* Inoculants on wheat yield and nitrogen uptake. This result is supported by Suri et al. (2011) while working on the influence of VAM and applied P on root colonization in wheat.

The available N forms, mostly NH₄⁺-N and NO₃⁻-N, decreased from the tillering stage to the grain-filling stage due to uptake and losses. During the tillering stage, the highest NH₄⁺-N and NO₃⁻-N was observed in T₃, i.e., treatment with NBPT amended urea due to a slowdown of urease activity, thereby enhancing the accumulation of NH₄⁺ and NO₃⁻ ions, and this increased the plant uptake. The urease activity was slowed down when NBPT-amended urea was applied to the soil, thus, enhancing the accumulation of NH₄⁺ ions and increasing the plant uptake of NH₄⁺ and NO₃⁻ ions. During N mineralization from the applied N fertilizers, most of the N remains as NH₄⁺ ions, then rapidly converting into NO₃⁻ due to the inhibitory effect of NBPT. That also helps in reducing NO₃⁻ leaching. A similar result was found by Dhakar et al. (2015) while working on the impact of nitrification inhibitors and various nitrogen sources on soil nitrogen distribution in Kinnow orchards.

As NH₄⁺ availability was initially low for the conversion into N₂O, the losses were low in the initial 7 days due to the application of these inhibitors. The application of urease inhibitors as

amendments and urea regulates the concentration of NH_4^+ and NO_3^- in the soil, thereby commanding NH_3 and N_2O emissions (Ding et al., 2015). Denitrification losses depend on soil moisture, soil temperature, and N-ions availability. Even though NBPT has no direct effect on lowering denitrification losses, it was observed that it had influenced the reduction of the emission of N_2O (Cassim et al., 2021). The reduction in the availability of NO_3^- has controlled the processes of denitrification and N_2O emission, mainly in the summer. The primary controller of denitrification processes in winter wheat is soil aeration and temperature (Aulakh et al., 2001).

Amendments like bio-fertilisers *Azotobacter spp.* and VAM positively affect soil biomass carbon and soil biomass nitrogen. These microbes help decompose organic matter and release the essential nutrients for plant uptake, increasing the MBC and MBN of the soil. Similar results were observed by Faujdar (2011) while working on the effect of FYM, bio-fertilizers, and zinc on nutrient transformations, soil properties, and yield of maize, and their residual effect on wheat.

The soil urease enzyme was suppressed by N inhibitors by a competitive inhibition mechanism, thereby reducing or slowing down the enzyme's activity. Rapid losses in N are mainly due to urea hydrolysis by the urease enzyme, which drastically increases pH and NH_4^+ ion concentration in the soil (Liu et al., 2018). So, this rise in pH and ion concentration increases NH_3 volatilization losses. In the entire growth period of wheat, the urease activity in the soil was always at a peak, especially during tillering stage; as it grew to maturity, its' activity dropped. Similar results were reported by Fu et al. (2019) while working on the effects of urease and nitrification inhibitors on the soil.

The physiological characteristics of wheat analyzed (photosynthesis rate, chlorophyll content, LAI, and stomatal conductance) were also significantly influenced by organic and inorganic amendments. In this study, photosynthesis rate, chlorophyll content, LAI, and stomatal conductance were slightly increased in T_7 treatment with azotobacter because of an increase in nitrogen availability, which positively affects these physiological parameters. Furthermore, followed by Azotobacter treated plot, NBPT treated plot showed significant

improvement in all these parameters.

The increase in the N content in wheat straw and grain might be due to improved nitrogen availability due to a reduced N loss and better NUE. As nitrogen is the most limiting nutrient in the growth and development of the wheat crop, the better uptake of the nutrient N will affect plant growth which will help to attain global food security by increasing the wheat yield by 1.5% per year. These results were similar to those obtained by Singh et al. (2018) while working on certain microorganisms like *Bacillus* spp. and *Azotobacter* spp. Mukhtar, Bashir & Nawaz (2018) found that microorganisms enhance crop growth by making nutrients available by fixing nitrogen and phosphate solubilization. In addition, *azotobacter* spp. can produce a few metabolites like phytohormones and exopolysaccharides, which help crops absorb nutrients and develop roots (Hindersah et al., 2020). Both are inorganic amendments that have effectively improved the plant's nutrient availability, thereby increasing yield. These findings were similar to those of Kumar et al. (2015), who reported an increase in grain and biological yield by 22.6% and 17.4%, respectively. It was also reported by Galindo et al. (2020) that NBPT has some effect on metabolic pathways in decreasing urease enzyme activity hence increasing the NUE.

Furthermore, slowing down nutrient release has improved the period of availability of N and increased the crop's N uptake (Liu et al., 2020). These positive effects, in turn, enhanced wheat's growth, development, and yield. However, in reducing NH₃ volatilization NBPT, HQ amendments performed much better than other amendments, but these amendments have some adverse effects on soil microorganism activity as per the studies by Upadhyay, Tewari & Patra (2011).

Conclusions

The current study concludes that both organic and inorganic amendments significantly reduce NH₃ losses, thus increasing available soil N and enhancing the NUE of the crops. Treatment containing NBPT reduced the losses by 40% compared to only RDN without any amendments. These results may be due to strong inhibition of urease activity by NBPT in the soil. Among plant-based amendments, garlic powder-treated urea showed better results in

decreasing NH₃ volatilization losses by 17% compared to only RDN-treated urea. These positive effects, in turn, enhanced the growth, development, and yields of wheat. Even though inorganic amendments performed much better than organic amendments in reducing NH₃ volatilization, opting for organic amendments gives way to sustainable agriculture. Thus, the focus should be more on organic than inorganic amendments to reduce NH₃ emissions from agricultural fields.

References

- Abalos D, Jeffery S, Sanz-Cobena A, Guardia G, Vallejo A. 2014. Meta-analysis of the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency. *Agriculture, Ecosystems & Environment* 189: 136-144.
- Affendi NMN, Mansor N, Samiri SS. 2020. Addition of chemical and natural urease inhibitors reduces ammonia and nitrous oxide losses. *Journal of Soil Science and Plant Nutrition* 20(1): 253-258.
- Aulakh MS, Khera TS, Doran JW, Bronson KF. 2001. Denitrification, N₂O and CO₂ fluxes in rice-wheat cropping system as affected by crop residues, fertilizer N and legume green manure. *Biology and Fertility of Soils* 34(6): 375-389.
- Bhaskar KA, Al-Hashimi A, Meena M, Meena VS, Langyan S, Shrivastava M, Sayyed RZ, El-Enshasy HA, Almunqedhi BMA, Singh R. 2022. Conservation agricultural practices for minimizing ammonia volatilization and maximizing wheat productivity. *Environmental Science and Pollution Research* 29: 9792–9804 DOI: <https://doi.org/10.1007/s11356-021-16370-4>.
- Bilal HM, Aziz T. 2022. Nitrogen use efficiency in crop production: issues and challenges in South Asia. *Nitrogen Assessment*. 127-148.
- Bittman S, Dedina M, Howard CM, Oenema O, Sutton MA. 2014. Options for ammonia mitigation. Guidance from the UNECE Task Force on Reactive Nitrogen, chapter 8, Centre for Ecology and Hydrology, Edinburgh, UK.
- Bremner JT. 1965. Inorganic forms of nitrogen. *Methods of Soil Analysis, Part 2 Chemical and Microbiological Properties* 9: 1179-1237.

Bremner JM, Douglas LA. 1971. Inhibition of urease activity in soils. *Soil Biology & Biochemistry* 3(4): 297-307.

Bremner JM, Keeney DR. 1965. Steam distillation methods for determination of ammonium, nitrate and nitrite. *Analytica Chimica Acta* 32: 485-495.

Brink C, Van Grinsven H. 2011. Cost and benefits of nitrogen in the environment. The European Nitrogen Assessment, chapter 22, Cambridge University Press.

Brookes PC, Landman A, Pruden G, Jenkinson DS. 1985. Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biology & Biochemistry* 17(6): 837-842.

Cameron KC, Di H J, Moir JL. 2013. Nitrogen losses from the soil/plant system, a review. *Annals of Applied Biology* 162(2): 145-173.

Cantarella H, Otto R, Soares JR, de Brito Silva AG. 2018. Agronomic efficiency of NBPT as a urease inhibitor, A review. *Journal of Advanced Research* 13:19-27.

Cao Y, Tian Y, Yin B, Zhu Z. 2013. Assessment of ammonia volatilization from paddy fields under crop management practices aimed to increase grain yield and N efficiency. *Field Crops Research* 147: 23-31.

Cassim BMAR, Kachinski WD, Besen MR, Coneglian CF, Macon CR, Paschoeto GF, Batista MA. 2021. Duromide increase NBPT efficiency in reducing ammonia volatilization loss from urea. *Revista Brasileira de Ciencia Avicola* 45.

Dhakar MK, Singh AK, Patel VB, Singh SK, Datta SP, Kumar R, Khanna M. 2015. Effect of different nitrogen sources and nitrification inhibitors on soil nitrogen distribution in Kinnow orchard. *Indian Journal of Horticulture* 72(2): 178-182.

Ding WX, Chen ZM, Yu HY, Luo JF, Yoo GY, Xiang J, Yuan JJ. 2015. Nitrous oxide emission and nitrogen use efficiency in response to nitrophosphate, N-(n-butyl) thiophosphoric triamide and dicyandiamide of a wheat cultivated soil under sub-humid monsoon conditions. *Biogeosciences* 12(3), 803-815.

Eduardo LC, Silva DRG, Faquin V, de Almeida Gonçalves B, Cancellier LL, Spehar CR. 2016. Ammonia volatilization from enhance defficiency urea on no-till maize in Brazilian cerrado with improved soil fertility. *Ciencia e Agrotecnologia* 40(2):133–144.

Fan X, Yin C, Yan G, Cui P, Shen Q, Wang Q, Chen H, Zhang N, Ye M, Zhao Y and Li T. 2018. The contrasting effects of N-(n-butyl) thiophosphoric triamide (NBPT) on N₂O

emissions in arable soils differing in pH are underlain by complex microbial mechanisms. *Science of the Total Environment* 642:155-167.

Faujdar RS. 2011. Effect of FYM, biofertilizers and zinc on nutrient transformations, soil properties and yield of maize (*Zea mays* L.) and their residual effect on wheat (*Triticum aestivum* L.) on typic haplustept, MPUAT, Udaipur.

Fu Q, Abadie M, Bland A, Carswell A, Misselbrook TH, Clark IM, Hirsch PR. 2020. Effects of urease and nitrification inhibitors on soil N, nitrifier abundance and activity in a sandy loam soil. *Biology and Fertility of Soils* 56(2): 185-194.

Galindo FS, Teixeira Filho MCM, Buzetti S, Pagliari PH, Santini JMK. 2020. Can NBPT urease inhibitor in combination with Azospirillum brasilense inoculation improve wheat development? *Nutrient Cycling in Agroecosystems* 1-13.

Gomez K, Gomez AA. 1984. Statistical procedure for agriculture research. Wiley and Sons.

Grain and Feed Annual report. 2021. United States Department of Agriculture. 2-3. DOI: <https://apps.fas.usda.gov>.

Hanway JJ, Heidel H. 1952. Soil analysis methods as used in Iowa state college soil testing laboratory. *Iowa Agriculture*. 57: 1-31.

Herr C, Mannheim T, Müller T, Ruser R. 2020. Effect of nitrification inhibitors on N₂O emissions after cattle slurry application. *Agronomy* 10(8): 1174.

Hindersah R, Setiawati MR, Asmiran P, Fitriatin BN. 2020. Formulation of Bacillus and Azotobacter consortia in liquid cultures, preliminary research on microbes-coated urea. *International Journal of Agriculture System* 8(1): 1-10.

ICAR RABI Agro-Advisory for Farmers-2021-22:<https://icar.org.in/content/icar-rabi-agro-advisory-farmers>

Jain N, Singh GP, Yadav R, Pandey R, Ramya P, Shine MB, Prabhu KV. 2014. Root trait characteristics and genotypic response in wheat under different water regimes. *Cereal Research Communications* 42(3): 426-438.

Jadon P, Selladurai R, Yadav SS, Coumar MV, Dotaniya ML, Singh AK & Kundu S. 2018. Volatilization and leaching losses of nitrogen from different coated urea fertilizers. *Journal of soil science and plant nutrition* 18(4): 1036-1047.

Kader MA, Mian MH, Hoque MS. 2002. Effects of Azotobacter inoculant on the yield and nitrogen uptake by wheat. *Journal of Biological Sciences* 2(4): 259-261.

Kjeldahl C. 1883. A new method for the determination of nitrogen in organic matter. *Analytical Chemistry* 22: 366.

Kumar B, Dhar S, Vyas AK, Paramesh V. 2015. Impact of irrigation schedules and nutrient management on growth, yield and root traits of wheat (*Triticum aestivum*) varieties. *Indian Journal of Agronomy* 60(1): 87-91.

Lasisi AA, Akinremi OO, Kumaragamage D. 2019. Efficacy of a new N-(n-butyl) thiophosphoric triamide formulation in reducing ammonia volatilization from urea-based fertilizers. *Canadian Journal of Soil Science* 99(4): 395-405.

Leghari SJ, Wahocho NA, Laghari GM, Hafeez Laghari A, MustafaBhabhan G, HussainTalpur K, Lashari AA. 2016. Role of nitrogen for plant growth and development, A review. *Advances in Environmental Biology* 10(9): 209-219.

Li Q, Yang A, Wang Z, Roelcke M, Chen X, Zhang F, Liu X. 2015. Effect of a new urease inhibitor on ammonia volatilization and nitrogen utilization in wheat in north and northwest China. *Field Crops Research* 175: 96-105.

Liu G, Yang Z, Du J, He A, Yang H, Xue G, Yu C, Zhang Y. 2020. Adding NBPT to urea increases N use efficiency of maize and decreases the abundance of N-cycling soil microbes under reduced fertilizer-N rate on the North China Plain, *Plos one*, 15(10): pp. e0240925

Liu Q, Pang Z, Yang Z, Nyumah F, Hu C, Lin W, Yuan Z. 2021. Bio-fertilizer Affects Structural Dynamics, Function, and Network Patterns of the Sugarcane Rhizospheric Microbiota. *Microbial Ecology* 1-17.

Liu Y, Dai Q, Jin X, Dong X, Peng J, Wu M, Liang N, Pan B, Xing B. 2018. Negative impacts of biochars on urease activity: high pH, heavy metals, polycyclic aromatic hydrocarbons, or free radicals? *Environmental Science & Technology* 52(21): 12740-12747.

Mahmood R, Ali A, Yaseen M, Ahmad M. 2014. Role of encapsulated calcium carbide in nitrogen economy, growth and yield of crops: a review. *Journal of Animal and Plant Sciences* 24(2).

Mahmud K, Panday D, Mergoum A, Missaoui A. 2021. Nitrogen Losses and Potential Mitigation Strategies for a Sustainable Agroecosystem. *Sustainability* 13:2400. DOI: <https://doi.org/10.3390/su13042400>

Majumdar D. 2008. Unexploited botanical nitrification inhibitors prepared from Karanja plant. *Indian Journal of Natural Products and Resources* 7: 58-67.

Mathialagan R, Mansor N, Al-Khateeb B, Mohamad MH, Shamsuddin MR. 2017. Evaluation of allicin as soil urease inhibitor. *Procedia Engineering*. 184: 449-459.

Modolo LV., da-Silva CJ, Brandão DS, Chaves IS. 2018. A minireview on what we have learned about urease inhibitors of agricultural interest since mid-2000s. *Journal of Advanced Research* 13: 29-37.

Mukhtar H, Bashir H, Nawaz A. 2018. Optimization of growth conditions for *Azotobacter* species and their use as biofertilizer. *Journal of Bacteriology and Mycology* 6(5): 274-278.

Olsen SR. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate (No. 939). US Department of Agriculture.

Pan B, Lam SK, Mosier A, Luo Y, Chen D. 2016. Ammonia volatilization from synthetic fertilizers and its mitigation strategies: A global synthesis. *Agriculture, Ecosystems & Environment* 232: 283-289.

Ramli NK, Majeed Z, Shuib A, Mansor N, Man Z. 2014. The potential of thiosulfates in garlic extract as urease bioinhibitor. *In Applied Mechanics and Materials* 625:796-799. Trans Tech Publications Ltd.

Sakariyawo OS, Oyeledun KO, Adeyemi NO, Atayese MO. 2020. Nitrogen use efficiency and performance of maize (*Zea mays* L.) cultivars as influenced by calcium carbide and inorganic nitrogen application rates in a derived savanna. *Journal of Plant Nutrition* 43(6): 784-797.

Sheoran OP, Tonk DS, Kaushik LS, Hasija RC and Pannu RS .1998. Statistical Software Package for Agricultural Research Workers. Recent Advances in information theory, Statistics & Computer Applications by D.S. Hooda & R.C. Hasija Department of Mathematics Statistics, CCS HAU, Hisar:139-143

Singh G, Kumar S, Sidhu GS, Kaur R. 2018. Effect of integrated nutrient management on yield of wheat (*Triticum aestivum* L.) under irrigated conditions. *International journal of chemical studies* 6:904-907.

Školníková M, Škarpa P, Ryant P, Kozáková Z, Antošovský J. 2022. Response of Winter Wheat (*Triticum aestivum* L.) to Fertilizers with Nitrogen-Transformation Inhibitors and Timing of Their Application under Field Conditions. *Agronomy* 12(1): 223.

Smith MS, JM Tiedje. 1979. Phases of denitrification following oxygen depletion in soil. *Soil Biology & Biochemistry* 2: 262-267.

Stumpe JM, Vlek PLG, Lindsay WL. 1984. Ammonia volatilization from urea and urea phosphates in calcareous soils. *Soil Science Society of America Journal* 48(4): 921-992.

Suri VK, Choudhary AK, Chander G, Verma TS. 2011. Influence of vesicular-arbuscular mycorrhizal fungi and applied phosphorus on root colonization in wheat and plant nutrient dynamics in a phosphorus-deficient acid alfisol of Western Himalayas. *Communications in Soil Science and Plant Analysis* 42(10): 1177-1186.

Upadhyay RK, Tewari SK, Patra DD. 2011. Natural nitrification inhibitors for higher nitrogen use efficiency, crop yield, and for curtailing global warming. *Journal of Tropical Agriculture* 49: 19-24

Vance ED, Brookes PC, Jenkinson DS. 1987. An extraction method for measuring soil microbial biomass. *Soil Biology & Biochemistry* 19(6): 703-707.

Walkley A, Black IA. 1934. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science* 37(1): 29-38.

Wołejko E, Jabłońska-Trypuć A, Wydro U, Butarewicz A, and Łozowicka B. 2020. Soil biological activity as an indicator of soil pollution with pesticides—a review. *Applied Soil Ecology* 147: 103356.

Xu R, Hanqin Tian, Shufen Pan, Stephen A, Yucheng Feng, William D, Jian Chen and Jia Yang. 2019. Global ammonia emissions from synthetic nitrogen fertilizer applications in agricultural systems: Empirical and process-based estimates and uncertainty. *Global Change Biology* 25:314–326, <https://doi.org/10.1111/gcb.14499>.

Zanin L, Tomasi N, Zamboni A, Varanini Z, Pinton R. 2015. The urease inhibitor NBPT negatively affects DUR3-mediated uptake and assimilation of urea in maize roots. *Frontiers in plant science* 6: 1007.

Table 1(on next page)

Effect of different amendments on available nitrogen, phosphorous, potassium and soil organic carbon in soil, N concentration in wheat grains and straw

Effect of different amendments on available nitrogen, phosphorous, potassium and soil organic carbon in soil, N concentration in wheat grains and straw (statistically significant at $p=0.05$. Means followed by common alphabets are not significantly different among themselves by DMRT)

Treatments	Available N (kg ha ⁻¹) after harvesting of the crop	Available P (kg ha ⁻¹) after harvesting of the crop	Available K (kg ha ⁻¹) after harvesting of the crop	Nitrogen content in grain (%)	Nitrogen content in straw (%)	Agronomic Efficiency of N (kg ha ⁻¹)	Production Efficiency of N (kg kg ⁻¹)	Apparent N Recovery (%)
T-1	141 ^g	17.9 ^g	214 ^c	1.5 ^g	0.32 ^h	0	0	0
T-2	173 ^{def}	19.7 ^f	221 ^b	1.91 ^f	0.39 ^g	9.1	25.8	35.2
T-3	172 ^{ef}	22.5 ^e	222 ^b	2.09 ^c	0.65 ^a	12.9	19.8	65.2
T-4	171 ^f	23.7 ^d	222 ^b	2.06 ^{cde}	0.62 ^b	11.6	18.5	62.8
T-5	177 ^{cde}	23.7 ^{de}	223 ^b	2.02 ^{de}	0.56 ^c	9.8	19.8	49.4
T-6	187 ^b	30.4 ^a	228 ^a	1.97 ^b	0.53 ^d	9.1	16.5	55.4
T-7	195 ^a	27.8 ^b	229 ^a	1.98 ^a	0.63 ^{ab}	12.5	15.4	80.8
T-8	180 ^c	26.9 ^{cb}	229 ^a	1.92 ^b	0.54 ^d	10.1	17.4	57.7
T-9	181 ^{cb}	27.58 ^{bc}	227 ^a	2.02 ^{cd}	0.41 ^f	9.6	21.5	44.8
T-10	178 ^{cd}	26.8 ^c	229 ^a	1.91 ^e	0.44 ^e	10.7	24	44.5
Statistical Significance (F test)	Significant	Significant	Significant	Significant	Significant	-	-	-
C.D.	5.55	1.58	3.30	0.10	0.02	-	-	-
SE(m)	1.85	0.53	1.10	0.03	0.01	-	-	-
SE(d)	2.62	0.75	1.56	0.05	0.01	-	-	-
C.V.	1.83	3.70	0.85	2.88	2.20	-	-	-

1

2 **Table 1. Effect of different amendments on available nitrogen, phosphorous, potassium and soil organic carbon in soil, N**
3 **concentration in wheat grains and straw (statistically significant at p=0.05. Means followed by common alphabets are not**
4 **significantly different among themselves by DMRT)**

Table 2 (on next page)

Effect of different amendments on yield attributes and yield of wheat. Mean of straw yield, biological yield, No. of tillers, spikes, grains per spike and harvest index

Effect of different amendments on yield attributes and yield of wheat. Mean of straw yield, biological yield, No. of tillers, spikes, grains per spike and harvest index (NS - Non-significant ($P = 0.05$) and mean of grain yield and test weight were found S-Significant ($P=0.05$) when compared to F-table value. Duncan's Multiple Range Test (DMRT) for comparison of mean yields and test weight of different amendment treatments ($\alpha=0.05$) was conducted. Means of grain yield and test weight of different treatments followed by different alphabets are significantly different among themselves and Means with the same letter are not significantly different.)

Treatments	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Biological yield (t ha ⁻¹)	No. of tillers	No. of spikes	No. of grains per spike	Harvest Index	Test weight (g)
T-1	3.14 ^c	7.95	11.09	340.67	302.67	36.00	28.34	34.82 ^b
T-2	4.50 ^{ab}	8.85	13.35	448.67	423.67	40.00	33.72	39.35 ^a
T-3	5.09 ^{ab}	7.82	12.90	410.67	374.00	40.00	39.41	39.53 ^a
T-4	4.89 ^a	9.44	14.33	483.33	456.00	39.33	34.12	39.07 ^a
T-5	4.61 ^{ab}	8.16	12.77	432.33	402.33	40.67	36.09	39.40 ^a
T-6	4.51 ^b	8.30	12.81	403.67	370.67	40.67	35.21	39.58 ^a
T-7	5.01 ^{ab}	8.40	13.42	448.00	419.00	41.33	37.37	39.53 ^a
T-8	4.65 ^{ab}	8.42	13.07	448.00	416.00	38.67	35.59	39.76 ^a
T-9	4.59 ^{ab}	8.42	13.01	459.00	428.33	40.00	35.29	39.53 ^a
T-10	4.74 ^{ab}	8.23	12.97	466.00	430.00	40.00	36.57	39.16 ^a
Statistical Significance	S	NS	NS	NS	NS	NS	NS	S
C.D.	0.76	N/A	N/A	N/A	N/A	N/A	N/A	1.34
SE(m)	0.25	0.99	1.05	3.25	33.98	33.40	1.03	0.54
SE(d)	0.36	1.40	1.49	4.59	48.05	47.24	1.45	0.77
C.V.	9.59	20.41	14.08	15.86	13.56	14.22	4.49	2.41

1

2 **Table 2. Effect of different amendments on yield attributes and yield of wheat. Mean of straw yield, biological yield, No. of**
3 **tillers, spikes, grains per spike and harvest index**

4 (NS - Non-significant ($P = 0.05$) and mean of grain yield and test weight were found S-Significant ($P=0.05$) when compared to F-table
 5 value. Duncan's Multiple Range Test (DMRT) for comparison of mean yields and test weight of different amendment treatments
 6 ($\alpha=0.05$) was conducted. Means of grain yield and test weight of different treatments followed by different alphabets are significantly
 7 different among themselves and Means with the same letter are not significantly different.)

8

9

10

11

Figure 1

Figure 1. Temporal graph of ammonia flux and effect of organic and inorganic amendments on ammonia flux.




Figure 2

Figure 2. Effect of various organic and inorganic amendments on ammonia volatilization losses from the soil




Figure 3

Figure.3 Effect of various organic and inorganic amendments on soil organic carbon (SOC) in the soil after the harvesting of wheat




Figure 4

Figure 4. Effect of various organic and inorganic amendments on NH_4^+ -N and NO_3^- -N concentration in soil




Figure 5

Figure 5. Effect of various organic and inorganic amendments on nitrous oxide flux and denitrification losses in soil




Figure 6

Figure 6. Effect of various organic and inorganic amendments on soil microbial biomass nitrogen and carbon (MBN and MBC) in soil




Figure 7

Figure 7. Effect of various organic and inorganic amendments on soil urease activity in wheat field

