

Does the combination of exercise and cognitive training improve working memory in older adults? A systematic review and meta-analysis

Yiqing Wu¹, Ming Zang², Biye Wang^{1,3}, Wei Guo^{Corresp. 1,3}

¹ College of Physical Education, Yangzhou University, Yangzhou, China

² College of Electrical Engineering, Chuzhou Polytechnic, Chuzhou, China

³ Institute of Sports, Exercise and Brain, Yangzhou University, Yangzhou, China

Corresponding Author: Wei Guo

Email address: guowei@yzu.edu.cn

Background. Cognitive functioning is dependent on working memory and a decline in working memory is the main cause of cognitive aging. Many studies have suggested that physical exercise or cognitive intervention can effectively improve working memory in the elderly. However, it is still unknown whether a combination of exercise and cognitive training (CECT) is more effective than either intervention alone. The present systematic review and meta-analysis were undertaken to evaluate the effect of CECT on working memory in the elderly.

Methods. The review was registered in the International Prospective Systematic Review (PROSPERO, CRD42021290138). Systematic searches were conducted on Web of Science, Elsevier Science, PubMed and Google Scholar. The data were extracted according to PICOS framework. Comprehensive Meta-Analysis (CMA) software was used to perform the meta-analysis, moderator analysis and publication bias testing.

Results. The current meta-analysis included 21 randomized controlled trials (RCT). Results showed that CECT had a significantly greater impact on working memory in older adults compared to no intervention groups (SMD = 0.29, 95%CI [0.14, 0.44], $p < 0.01$), with no significant difference between CECT and exercise (SMD = 0.16, 95%CI [-0.04, 0.35], $p = 0.12$) or cognitive intervention alone (SMD = 0.08, 95%CI [-0.13, 0.30], $p = 0.44$). Furthermore, the positive effect of CECT was moderated by intervention frequency and cognitive state.

Conclusions. The CECT can effectively improve working memory of older adults, but the effect of CECT compared to single intervention needs to be further explored.

Does the combination of exercise and cognitive training improve working memory in older adults? A systematic review and meta-analysis

Yiqing Wu¹, Ming Zang², Biye Wang^{1,3}, Wei Guo^{1,3}

¹ College of Physical Education, Yangzhou University, Yangzhou, China

² College of Electrical Engineering, Chuzhou Polytechnic, Chuzhou, China

³ Institute of Sports, Exercise and Brain, Yangzhou University, Yangzhou, China

Corresponding Author:

Wei Guo^{1,3}

Yangzhou University, No.196 Huayang West Road, Hangjiang District, Jiangsu, P.R. China

Email address: guowei@yzu.edu.cn

Abstract

Background. Cognitive functioning is dependent on working memory and a decline in working memory is the main cause of cognitive aging. Many studies have suggested that physical exercise or cognitive intervention can effectively improve working memory in the elderly. However, it is still unknown whether a combination of exercise and cognitive training (CECT) is more effective than either intervention alone. The present systematic review and meta-analysis were undertaken to evaluate the effect of CECT on working memory in the elderly.

Methods. The review was registered in the International Prospective Systematic Review (PROSPERO, CRD42021290138). Systematic searches were conducted on Web of Science, Elsevier Science, PubMed and Google Scholar. The data were extracted according to PICOS framework. Comprehensive Meta-Analysis (CMA) software was used to perform the meta-analysis, moderator analysis and publication bias testing.

Results. The current meta-analysis included 21 randomized controlled trials (RCT). Results showed that CECT had a significantly greater impact on working memory in older adults compared to no intervention groups (SMD = 0.29, 95%CI [0.14, 0.44], $p < 0.01$), with no significant difference between CECT and exercise (SMD = 0.16, 95%CI [-0.04, 0.35], $p = 0.12$) or cognitive intervention alone (SMD = 0.08, 95%CI [-0.13, 0.30], $p = 0.44$). Furthermore, the positive effect of CECT was moderated by intervention frequency and cognitive state.

Conclusions. The CECT can effectively improve working memory of older adults, but the effect of CECT compared to single intervention needs to be further explored.

Keywords: combined intervention, physical exercise, cognitive training, working memory, older adults

Introduction

Working Memory (WM), a system of temporarily storing and efficiently manipulating information, is essential to effective cognitive functioning (Baddeley, 2012). A decline in working memory is the main cause of cognitive aging, and seriously affects daily life of the elderly (Goh & Park, 2009; Salthouse, 2012). However, it has recently been shown that the aging brain maintains plasticity (Passow, Thurm & LiC, 2014). Therefore, interventions that improve working memory in the elderly may delay cognitive aging (Lampit, Hallock & Valenzuela, 2014; Logan, 2014; Bonnechère, Langley & Sahakian, 2020; Cox & Lautenschlager, 2020).

For elderly with dementia or other cognitive disorders, non-pharmaceutical methods can reduce medical expenses and improve the quality of the life. Physical exercise is a promising non-pharmacological behavioral intervention for cognitive aging, inhibiting cognitive decline related to age and neurodegenerative diseases (Bherer, Erickson & Liu-Ambrose, 2013; Lin et al., 2019; Jia et al., 2019). Previous meta-analyses have also demonstrated the efficacy of physical exercise training on working memory in the elderly with or without dementia or mild cognitive impairment (MCI), which is a transitional state between normal aging and dementia (Petersen, 2004; Law et al., 2020; Zhidong et al., 2021). One possible mechanism for this effect

is the increase in brain-derived neurotrophic factor (BDNF) associated with exercise, as higher expression levels of BDNF correlate with slower cognitive decline (Buchman et al., 2016).

In addition to physical exercise, cognitive training intervention has been demonstrated to effectively delay cognitive aging (Nguyen, Murphy & Andrews, 2019). Cognitive training may enhance dopaminergic neurotransmission, leading to the consolidation and enhancement of cognition performance in older adults (Passow, Thurm & LiC, 2017). Prior studies have shown that cognitive training-based interventions have a positive impact on cognitive functioning in the elderly at a variety of cognitive states (Giuli et al., 2016). Furthermore, multi domain cognitive training has proven effective in improving working memory in healthy older people (Hong et al., 2021).

Over the last decade, research has found that the combination of cognitive training and exercise (CECT) can improve cognitive function to a larger extent than exercise or cognitive interventions alone (Benzing & Schmidt, 2017; Niederer et al., 2019). The combined intervention is a multimodal intervention strategy, combining physical exercise with cognitively challenging activities. However, the impact of the combination of the two interventions in an older population remains controversial. Some research has found a superior effect of CECT in healthy older adults compared to physical exercise or cognitive training alone (Shatil, 2013; Zhu et al., 2016; Joubert & Chainay, 2018). Similar results were also found in the elderly with MCI, i.e., the combined intervention also have significant cognitive benefits in older adults with MCI (Köbe et al., 2016; Law et al., 2014; Shimada et al., 2018; Yang et al., 2020). While some studies have yielded inconsistent results, showing that combined intervention have no advantage compared to single exercise or cognitive training (Fiatarone Singh et al., 2014; Hackney et al., 2015; Chainay, Joubert & Massol, 2021). Thus, no definitive conclusions have been drawn about the superiority of combined exercise and cognitive intervention. In addition, combined intervention can be divided into simultaneous or sequential modes (Tait et al., 2017). Sequential mode provides intervention modalities in separate sessions, usually during the same period (Ngandu et al., 2015; Damirchi, Hosseini & Babaei, 2018; Fabre et al., 2002; Legault et al., 2011). Simultaneous mode means that cognitive training and exercise are to be conducted at the same time (e.g., dual task, exergames and video dancing) (Eggenberger et al., 2016; Ordnung et al., 2017; Norouzi et al., 2019; Adcock et al., 2020). However, it is still unclear which combination modes is most beneficial for cognitive function in older adults. Therefore, more investigations are needed to demonstrate the benefit of combined exercise and cognitive training.

Previous research on combined interventions has concentrated on the effects of CECT on overall cognition (Joubert & Chainay, 2018; Karssemeijer et al., 2017; Lauenroth et al., 2016; Zhu et al., 2016) or executive functioning (Guo et al., 2020; Wollesen et al., 2020) and few studies focused on working memory. To our knowledge, there have been no previous meta-analyses performed to study the effect of CECT on working memory in older adults. Thus, the main aim of this meta-analysis was to investigate whether CECT is effective in delaying the decline of working memory in older adults.

Materials & Methods

This meta-analysis was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. The protocol of this review was registered in the International Prospective Systematic Review (PROSPERO, CRD42021290138).

Search strategy

Electronic searches of online databases including Web of Science, Elsevier Science, PubMed and Google Scholar were conducted to collect relevant literature since the start of each database until April of 2021. The following keywords were used in the searches: exercise intervention terms (“aerobic exercise” OR “multidomain exercise training” OR “physical activity”) AND cognitive intervention terms (“mental training” OR “cognitive training” OR “working memory training”) AND combination of intervention terms (“combined” OR “simultaneous” OR “exergame” OR “dual-task” OR “video dancing” OR “cognitive-motor game”) AND relevant working memory terms (“cognitive function” OR “working memory” OR “executive function”) AND older adults terms (“old” OR “elderly” OR “aging”). Additional articles were obtained through a list of references from published reviews.

Inclusion Criteria

Inclusion and exclusion criteria were determined based on PICOS (population, intervention, comparison, outcome, and study design). The following criteria were used for study inclusion: (1) population: participants were elderly (aged 60 and over) and included healthy individuals, along with those with MCI and dementia; (2) intervention: combination of exercise and cognitive trainings; (3) comparison: at least one comparison group (combined intervention group with cognitive intervention alone or physical exercise intervention alone or a control group receiving no intervention); (4) outcome: measurements of working memory included at least one outcome that could be used to calculate an effect size ; (5) study design: randomized controlled trial (RCT) design; and (6) other: written in English. Exclusion criteria included: (1) population: participants aged less than 60 years old; (2) intervention: without combination of exercise and cognitive intervention; (3) comparison: studies that did not compare the combined intervention group with other intervention groups alone; (4) outcome: studies without relevant data on working memory pre and post intervention; (5) study design: no RCT design; and (6) others: noninterventional studies, reviews and theoretical articles, case and protocol articles, unpublished studies and papers, studies that were not written in English.

Data extraction and assessment of study quality

In this systematic review, working memory tasks were included based on following criteria: (1) participants are required to store and manipulate information during the task, such as Digit Span Backward test, N-back task, and Corsi block-tapping task, (2) the authors declared the tasks were used to measure working memory (You et al., 2009; Damirchi, Hosseini & Babaei, 2018). Tasks that simply stored information with little manipulation were excluded, such as Short-term Memory task, Immediate Recall, Forward Digit Span, the Word List test (Linde & Alfermann, 2014; Mrakic-Sposta et al., 2018; Romera-Liebana et al., 2018; Jardim et al., 2021).

Data referring to working memory tasks were extracted from combined intervention versus exercise intervention alone, or cognitive intervention alone, and no intervention using a spreadsheet. The mean, standard deviation (SD), and the number of participants (N) in each group at pretest and posttest were extracted. If the means and SDs were not mentioned, the change values of the means and SDs after intervention or mean difference of the 95% confidence interval were extracted.

To assess study quality, two independent researchers evaluated the risk of bias for individual studies using the PEDro scale (Maher et al., 2003). Each criterion met on the scale was scored a point, for a maximum score of 11. Articles scoring 8 or higher were considered high quality while those scoring below 8 which were considered low quality.

The moderator analysis was conducted to assess the effects of combined intervention on working memory in the elderly. The following moderators were analyzed: the mode of combination (separate versus simultaneous), cognitive status (healthy versus MCI versus dementia), intervention length (short versus medium versus long), frequency (< 3 sessions per week versus ≥ 3 sessions per week), session length (≤ 60 min versus > 60 min), no intervention group (active versus passive), study quality (high quality versus low quality).

Data analysis

The statistical data was quantified using Comprehensive Meta-Analysis (CMA) software and calculated the effect size. The effect of the intervention was measured by the standardized mean difference (SMD) of changes after the combined intervention group versus the no intervention group, single exercise intervention group and single cognitive intervention group (Borenstein et al., 2009). The calculation equation of SMD are as follows:

$$SMD = \frac{\bar{x}_1 - \bar{x}_2}{S_{within}}$$

$$S_{within} = \sqrt{\frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}}$$

If one study had multiple task that measured working memory, outcomes were combined into an average effect size to avoid influencing results. The combined effect size with 95% confidence interval (CI) was calculated to determine the efficacy of CECT on working memory. The I^2 index was calculated to analyze the heterogeneity of the included studies. 0% indicated that no heterogeneity has been observed and the higher of the I^2 value reflected on more significant of the heterogeneity. The Egger's regression intercepts and funnel charts were performed to estimate publication bias. Funnel charts was symmetrical which indicated no risk of bias and studies were outside the funnel sharp reflecting on high risk of bias. If $I^2 < 50\%$, $p \geq 0.05$, representing the studies had no statistical heterogeneity, so the fixed effect model was used for analysis and if $I^2 \geq 50\%$, $p \geq 0.05$, indicating had statistical heterogeneity between the studies, so the random effect model was used for analysis.

Results

Identification of studies

Preliminarily, a total of 1,379 articles were obtained from database searches, of these, 275 articles were removed for duplication and 1104 were excluded on the basis of titles or abstracts. Of the remaining 55 articles, 38 were excluded based on information found in the full text. Finally, 21 eligible articles were included in this meta-analysis, 4 of which were included from previous reviews. The specific process of article selection, according to the recommended PRISMA flow diagram is shown in Figure 1.

(insert Figure 1 about here)

Characteristics of included studies

Twenty-one RCT articles were included in the current meta-analysis. The characteristics of included studies are shown in Table 1. Among the included studies, 16 were conducted on healthy older adults (Fabre et al., 2002; You et al., 2009; Legault et al., 2011; Mailliot, Perrot & Hartley, 2012; Shatil, 2013; Gschwind et al., 2015; Nishiguchi et al., 2015; Rahe et al., 2015; Eggenberger et al., 2016; Schättin et al., 2016; Ordnung et al., 2017; Kalbe et al., 2018; Norouzi et al., 2019; Adcock et al., 2020; Takeuchi et al., 2020; Dana, 2019), 4 focused on older adults with MCI (Suzuki et al., 2013; Combourieu Donnezan et al., 2018; Damirchi, Hosseini & Babaei, 2018; Bae et al., 2019), and 1 focused on older adults with dementia (Karssemeijer et al., 2019). Sample size ranged from 13 to 153. The mean age range for older adults was 65.8 to 79.2. Of all 21 studies, 6 included four comparison groups, 3 included three comparison groups, and 12 included two comparison groups. Of the 21 studies, 7 studies used separate interventions, while 14 studies adopted a simultaneous combination of exercise and cognitive intervention, (3 of the 14 used the dual task and 7 of the 14 used the exergame). The length of intervention ranged from 4 to 24 weeks and the frequency was one to six sessions each week. Five of the studies included active control groups (leisure activity and health education) and the rest included passive control groups (adhered to normal life). Types of exercise interventions included aerobic training, flexibility training, muscle strength training, balance training, endurance training, and resistance training. The types of cognitive interventions included multidomain cognitive training, working memory training, and cognitive games. Only 5 articles (Mailliot, Perrot & Hartley, 2012; Suzuki et al., 2013; Ordnung et al., 2017; Damirchi, Hosseini & Babaei, 2018; Norouzi et al., 2019) explicitly reported the gender distribution of participants, of which there were a total of 134 male and 101 female participants. The other included articles did not clearly report the gender distribution of participants. The quality of studies ranged from 6 to 11 on the PEDro scale and the specific scoring details are shown in Table S1.

(insert Table 1 about here)

Combined intervention versus no intervention group

Fig. S1 presented the funnel plot of included studies and the result showed no significant asymmetry (Egger's regression intercept = 1.50, $p > 0.05$).

A total of 15 articles (Adcock et al., 2020; Bae et al., 2019; Combourieu Donnezan et al., 2018; Damirchi, Hosseini & Babaei, 2018; Dana, 2019; Fabre et al., 2002; Gschwind et al., 2015; Karssemeijer et al., 2019; Legault et al., 2011; Maillot, Perrot & Hartley, 2012; Nishiguchi et al., 2015; Norouzi et al., 2019; Ordnung et al., 2017; Shatil, 2013; Suzuki et al., 2013) reported on the effects of CECT compared to no intervention on working memory in older adults. A fixed effect model for meta-analysis was used, as the heterogeneity test showed $Q_{(14)} = 17.31$, $p = 0.24$, $i^2 = 19.10$. As shown in Fig. 2, the combined intervention group showed significant improvement in working memory compared to the no intervention group (SMD = 0.29, 95%CI [0.14, 0.44], $p < 0.01$).

(insert Fig. 2 about here)

Combined intervention versus single exercise intervention

The funnel plot revealed that one study had a disproportionately large effect size and Egger's test was significant (Egger's regression intercept = 2.71, $p < 0.05$), so one study was removed from further analysis (Norouzi et al., 2019). The funnel plot after removal of one outlier was presented in fig. S2 and had no significant asymmetry.

Eleven studies (Combourieu Donnezan et al., 2018; Damirchi, Hosseini & Babaei, 2018; Dana, 2019; Eggenberger et al., 2016; Fabre et al., 2002; Karssemeijer et al., 2019; Legault et al., 2011; Schättin et al., 2016; Shatil, 2013; Takeuchi et al., 2020; You et al., 2009) compared the effect of CECT to single physical exercise. The results of our analysis revealed no significant difference on working memory in the effect of between CECT and single physical exercise (SMD = 0.16, 95%CI [-0.04, 0.35], $p = 0.12$) (Fig. 3). The heterogeneity test was not significant ($Q_{(10)} = 8.46$, $i^2 = 0$, $p = 0.58$), so a fixed effect model was used.

(insert Fig. 3 about here)

Combined intervention versus cognitive intervention alone

The funnel plot showed that the result had no significant asymmetry (fig. S3). Nine articles (Fabre et al., 2002; Legault et al., 2011; Shatil, 2013; Rahe et al., 2015; Combourieu Donnezan et al., 2018; Damirchi, Hosseini & Babaei, 2018; Kalbe et al., 2018; Dana, 2019; Takeuchi et al., 2020) compared the effect of combined intervention to cognitive intervention alone. The results (Fig. 4) of our meta-analysis revealed no significant difference in the effects of combined intervention versus cognitive intervention alone on working memory (SMD = 0.08, 95%CI [-0.13, 0.30], $p = 0.44$). The heterogeneity test was not significant ($Q_{(8)} = 2.13$, $i^2 = 0$, $p = 0.98$), so a fixed effect model was used.

(insert Fig. 4 about here)

Moderator analysis

The results of the moderator analysis are shown in Table 2. A heterogeneity test revealed a significant influence of cognitive status on the effects of CECT on working memory ($Q_{(2)} = 7.67, p = 0.02$). The criterion used to categorize participants' cognitive status was based on explicit declarations made by the authors, identifying individuals as either having mild cognitive impairment (MCI) or dementia. Combined intervention had a significant influence among healthy participants ($SMD = 0.24, 95\%CI [0.06, 0.42], p = 0.01$) and those with MCI ($SMD = 0.58, 95\%CI [0.29, 0.87], p < 0.01$) but no significant effect among those with dementia ($SMD = -0.17, 95\%CI [-0.64, 0.31], p = 0.49$).

A heterogeneity test revealed a significant influence of intervention frequency on the effects of CECT on working memory ($Q_{(1)} = 5.53, p = 0.02$). CECT was significantly effective in low frequency intervention ($SMD = 0.52, 95\%CI [0.28, 0.76], p < 0.01$), but no significant effect in high frequency intervention ($SMD = 0.15, 95\%CI [-0.03, 0.34], p = 0.11$).

In terms of mode of combination, simultaneous ($SMD = 0.31, 95\%CI [0.14, 0.47], p < 0.01$) training were proven effective, but no significance was found for sequential ($SMD = 0.23, 95\%CI [-0.09, 0.55], p = 0.16$) training. CECT was significantly effective regardless of intervention time: Short ($SMD = 0.48, 95\%CI [0.13, 0.83], p = 0.01$); medium ($SMD = 0.27, 95\%CI [0.01, 0.53], p = 0.05$); and long intervention length ($SMD = 0.23, 95\%CI [0.03, 0.44], p = 0.03$). Combined intervention was significantly effective in low quality ($SMD = 0.44, 95\%CI [0.21, 0.67], p < 0.01$) studies, and had no significance in high quality ($SMD = 0.18, 95\%CI [-0.01, 0.38], p > 0.05$) studies. Compared with no intervention, long combined intervention sessions ($SMD = 0.31, 95\%CI [0.07, 0.55], p = 0.01$) and short combined sessions ($SMD = 0.28, 95\%CI [0.09, 0.46], p < 0.01$) were both significantly beneficial for working memory. In terms of the no intervention group, CECT had a positive effect in passive ($SMD = 0.33, 95\%CI [0.14, 0.51], p < 0.01$) but no significant effect in active control group ($SMD = 0.22, 95\%CI [-0.03, 0.47], p = 0.08$).

(insert Table 2 about here)

Discussion

The current meta-analysis included 21 articles and explored the effect of CECT on working memory in older adults. The results support efficacy of CECT on working memory as compared to those receiving no intervention and the effect is modulated by the frequency of intervention and cognitive status.

The results indicate that there is a significant benefit of CECT on working memory in the elderly compared to no intervention. However, the effects of combined intervention are not significantly different than either physical exercise or cognitive intervention alone. These results disagree with previous meta-analysis and there are potential reasons for this discrepancy. First, other studies have explored the effect of CECT on cognitive function (Zhu

et al., 2016; Gheysen et al., 2018; Yang et al., 2020; Meng et al., 2021), while the current study specifically explored working memory. Second, there are a limited number of studies directly comparing efficacy of CECT with single exercise or cognitive intervention. Third, prior studies compared the efficacy of CECT on cognitive function in healthy older adults (Zhu et al., 2016), while the current meta-analysis extended findings to those with MCI and dementia. Lastly, different types of exercise intervention (e.g., aerobic exercise, flexibility, balance, coordination, and endurance training) may affect cognitive functioning differently. A combination of systematic exercise training and stimulation of cognitive tasks not only promotes new connections between nerve cells in the brain, but also promotes the formation of new neurons and enhances synapses and plasticity (Bennett et al., 1964). The CECT is effective for improved cognitive function and delayed cognitive decline, with positive effects for older people to participate in activities of daily living and improved quality of life.

The current meta-analysis showed that the effects of CECT on working memory in the elderly is moderated by the frequency of intervention and cognitive state. CECT showed significant positive improvements in working memory in both healthy individuals and those with MCI. These results are in line with previous studies showing that CECT improves overall cognitive function and executive function (Gavelin et al., 2020; Guo et al., 2020). In the current study, low frequency interventions were significantly beneficial to working memory comparing to high frequency interventions. While there is general consensus that greater training frequency generates greater benefits, the results of the current meta-analysis found that low frequency intervention is more than three times as effective as high frequency intervention. Some previous studies have also indicated that high-frequency interventions are less effective than low-frequency interventions (Lampit, Hallock & Valenzuela, 2014; Guo et al., 2020) suggesting that high-frequency interventions may cause cognitive fatigue in older people (Holtzer et al., 2011), negatively affecting cognitive functioning. Moreover, although several studies have indicated that CECT can have a positive impact on the cognitive function of older adults, there is still controversy surrounding which combination mode is more effective. Some meta-analyses have reported that both simultaneous and sequential CECT can improve the cognitive function of older adults (Zhu et al., 2016; Guo et al., 2020). However, a different meta-analysis has found a significant effect only for simultaneous CECT, with no notable effect for sequential CECT (Gheysen et al., 2018). This outcome is consistent with the results of the present meta-analysis. One possible explanation for this finding is the temporary nature of the increase in brain-derived neurotrophic factor (BDNF). Previous research has demonstrated that higher levels of BDNF can enhance working memory performance (Erickson et al., 2013); however, BDNF levels typically return to baseline within 10-60 minutes after physical activity (Knaepen et al., 2012). Thus, simultaneous CECT may offer the greatest neurotrophic benefits compared to sequential CECT.

It is important to note that there was no significant difference between CECT and single physical exercise or cognitive intervention. These results may be affected by the low number of studies that have compared CECT with single physical exercise or cognitive intervention, as majority of studies compare combined intervention to no intervention. This is a notable

limitation of the current study. Briefly, the efficacy of CECT was demonstrated, but whether CECT is more beneficial than either physical exercise or cognitive intervention alone is not clear. In the future, it is advisable that studies include multiple intervention groups for comparison. In addition, the majority of current studies on the efficacy of CECT have used some cognitive tasks, such as Digit Span task, Trail Making Test, Stroop task and so on. Future studies are needed to investigate whether biological risk factors (e.g., vascular risk factors and APOE genotype) would influence the efficacy of CECT, which could increase the level of evidence. Furthermore, given the substantial number of studies included in the current meta-analysis utilizing exergame interventions, it may be worthwhile for future studies to investigate the impact of such interventions on working memory in older adults.

The current research has some other limitations that should be noted when considering the results. First, the current study did not investigate post-intervention follow-up and therefore, the maintenance of improved working memory as a result of intervention cannot be assessed. Second, working memory is a complex cognitive domain and was measured in a variety of ways in the included studies, possibly introducing variability within the results. Finally, other potential moderator variables, such as baseline levels of exercise or cognitive health, were not analyzed.

Conclusions

The presented meta-analysis indicates that CECT is a promising way to improve working memory in older adults and that the effect is influenced by the frequency of intervention and cognitive status. However, there is no evidence that combined intervention is superior to exercise or cognitive intervention alone.

Grant Disclosures

This work was supported by the National Social Science Foundation of China (Grant No. 18CTY014).

Competing Interests

The authors declare that they have no competing interests.

References

- Adcock M, Fankhauser M, Post J, Lutz K, Zizlsperger L, Luft AR, Guimarães V, Schättin A, de Bruin ED. 2020. Effects of an In-home Multicomponent Exergame Training on Physical Functions, Cognition, and Brain Volume of Older Adults: A Randomized Controlled Trial. *Frontiers in Medicine* 6. DOI: 10.3389/fmed.2019.00321.
- Baddeley A. 2012. Working memory: Theories, models, and controversies. *Annual Review of Psychology* 63:1–29. DOI: 10.1146/annurev-psych-120710-100422.
- Bae S, Lee S, Lee S, Jung S, Makino K, Harada K, Harada K, Shinkai Y, Chiba I, Shimada H. 2019. The effect of a multicomponent intervention to promote community activity on cognitive function in

- 371 older adults with mild cognitive impairment: A randomized controlled trial. *Complementary*
372 *Therapies in Medicine* 42:164–169. DOI: 10.1016/j.ctim.2018.11.011.
- 373 Bennett EL, Diamond MC, Krech D, Rosenzweig MR. 1964. Chemical and anatomical plasticity of brain.
374 *Science* 146:610–619. DOI: 10.1126/science.146.3644.610.
- 375 Benzing V, Schmidt M. 2017. Cognitively and physically demanding exergaming to improve executive
376 functions of children with attention deficit hyperactivity disorder: A randomised clinical trial. *BMC*
377 *Pediatrics* 17. DOI: 10.1186/s12887-016-0757-9.
- 378 Bherer L, Erickson KI, Liu-Ambrose T. 2013. A review of the effects of physical activity and exercise on
379 cognitive and brain functions in older adults. *Journal of Aging Research* 2013. DOI:
380 10.1155/2013/657508.
- 381 Bonnechère B, Langley C, Sahakian BJ. 2020. The use of commercial computerised cognitive games in
382 older adults: a meta-analysis. *Scientific Reports* 10:1–14. DOI: 10.1038/s41598-020-72281-3.
- 383 Borenstein M, Hedges L v., Higgins JPT, Rothstein H. 2009. *Introduction to meta-analysis*. West Sussex:
384 John Wiley & Sons.
- 385 Buchman AS, Yu L, Boyle PA, Schneider JA, de Jager PL, Bennett DA. 2016. Higher brain BDNF gene
386 expression is associated with slower cognitive decline in older adults. *Neurology* 86:735–741. DOI:
387 10.1212/WNL.0000000000002387.
- 388 Chainay H, Joubert C, Massol S. 2021. Behavioural and ERP Effects of Cognitive and Combined
389 Cognitive and Physical Training on Working Memory and Executive Function in Healthy Older
390 Adults. *Advances in cognitive psychology* 17:58–69. DOI: 10.5709/ACP-0317-Y.
- 391 Combourieu Donnezan L, Perrot A, Belleville S, Bloch F, Kemoun G. 2018. Effects of simultaneous
392 aerobic and cognitive training on executive functions, cardiovascular fitness and functional abilities
393 in older adults with mild cognitive impairment. *Mental Health and Physical Activity* 15:78–87. DOI:
394 10.1016/j.mhpa.2018.06.001.
- 395 Cox KL, Lautenschlager NT. 2020. The evidence for the effects of physical exercise training on cognitive
396 function for those with mild cognitive impairment. *Alzheimer's & Dementia* 16:40752. DOI:
397 10.1002/alz.040752.
- 398 Damirchi A, Hosseini F, Babaei P. 2018. Mental Training Enhances Cognitive Function and BDNF More
399 Than Either Physical or Combined Training in Elderly Women With MCI: A Small-Scale Study.
400 *American Journal of Alzheimer's Disease and other Dementias* 33:20–29. DOI:
401 10.1177/1533317517727068.
- 402 Dana A. 2019. Comparing the Effects of Physical, Cognitive and Combined Rehabilitation on the
403 Improvement of Working Memory and Cognitive Flexibility of the Elderly. *Iranian Journal of*
404 *Learning and Memory* 2:67-77. DOI: 10.22034/IEPA.2019.91052.
- 405 Erickson KI, Banducci SE, Weinstein AM, MacDonald AW, Ferrell RE, Halder I, Flory JD, Manuck SB.
406 2013. The Brain-Derived Neurotrophic Factor Val66Met Polymorphism Moderates an Effect of
407 Physical Activity on Working Memory Performance. *Psychological Science* 24:1770–1779. DOI:
408 10.1177/0956797613480367.
- 409 Eggenberger P, Wolf M, Schumann M, de Bruin ED. 2016. Exergame and balance training modulate
410 prefrontal brain activity during walking and enhance executive function in older adults. *Frontiers in*
411 *Aging Neuroscience* 8. DOI: 10.3389/fnagi.2016.00066.
- 412 Fabre C, Chamari K, Mucci P, Massé-Biron J, Préfaut C. 2002. Improvement of cognitive function by
413 mental and/or individualized aerobic training in healthy elderly subjects. *International Journal of*
414 *Sports Medicine* 23:415–421. DOI: 10.1055/s-2002-33735.

- 415 Fiatarone Singh MA, Gates N, Saigal N, Wilson GC, Meiklejohn J, Brodaty H, Wen W, Singh N, Baune
416 BT, Suo C, Baker MK, Foroughi N, Wang Y, Sachdev PS, Valenzuela M. 2014. The Study of
417 Mental and Resistance Training (SMART) study—resistance training and/or cognitive training in
418 mild cognitive impairment: a randomized, double-blind, double-sham controlled trial. *Journal of the*
419 *American Medical Directors Association* 15:873–880. DOI: 10.1016/J.JAMDA.2014.09.010.
- 420 Gavelin HM, Dong C, Minkov R, Bahar-Fuchs A, Ellis KA, Lautenschlager NT, Mellow ML, Wade AT,
421 Smith AE, Finke C, Krohn S, Lampit A. 2020. Combined physical and cognitive training for older
422 adults with and without cognitive impairment: A systematic review and network meta-analysis of
423 randomized controlled trials. *Ageing Research Reviews*. DOI: 10.1101/2020.08.08.20170654.
- 424 Gheysen F, Poppe L, DeSmet A, Swinnen S, Cardon G, de Bourdeaudhuij I, Chastin S, Fias W. 2018.
425 Physical activity to improve cognition in older adults: Can physical activity programs enriched with
426 cognitive challenges enhance the effects? A systematic review and meta-analysis. *International*
427 *Journal of Behavioral Nutrition and Physical Activity* 15:1–13. DOI: 10.1186/s12966-018-0697-x.
- 428 Giuli C, Papa R, Lattanzio F, Postacchini D. 2016. The Effects of Cognitive Training for Elderly: Results
429 from My Mind Project. *Rejuvenation Research* 19:485–494. DOI: 10.1089/rej.2015.1791.
- 430 Goh JO, Park DC. 2009. Neuroplasticity and cognitive aging: The scaffolding theory of aging and
431 cognition. *Restorative Neurology and Neuroscience* 27:391–403. DOI: 10.3233/RNN-2009-0493.
- 432 Gschwind YJ, Eichberg S, Ejupi A, de Rosario H, Kroll M, Marston HR, Drobits M, Annegarn J,
433 Wieching R, Lord SR, Aal K, Vaziri D, Woodbury A, Fink D, Delbaere K. 2015. ICT-based system
434 to predict and prevent falls (iStoppFalls): Results from an international multicenter randomized
435 controlled trial. *European Review of Aging and Physical Activity* 12:1–11. DOI: 10.1186/s11556-
436 015-0155-6.
- 437 Guo W, Zang M, Klich S, Kawczyński A, Smoter M, Wang B. 2020. Effect of combined physical and
438 cognitive interventions on executive functions in older adults: A meta-analysis of outcomes.
439 *International Journal of Environmental Research and Public Health* 17:1–19. DOI:
440 10.3390/ijerph17176166.
- 441 Hackney ME, Byers C, Butler G, Sweeney M, Rossbach L, Bozzorg A. 2015. Adapted Tango Improves
442 Mobility, Motor–Cognitive Function, and Gait but Not Cognition in Older Adults in Independent
443 Living. *Journal of the American Geriatrics Society* 63:2105–2113. DOI: 10.1111/JGS.13650.
- 444 Holtzer R, Shuman M, Mahoney JR, Lipton R, Verghese J. 2011. Cognitive Fatigue defined in the
445 context of attention networks. *Aging, Neuropsychology, and Cognition* 18:108–128. DOI:
446 10.1080/13825585.2010.517826.
- 447 Hong X, Chen Y, Wang J, Shen Y, Li Q, Zhao B, Guo X, Feng W, Wu W, Li C. 2021. Effects of multi-
448 domain cognitive training on working memory retrieval in older adults: behavioral and ERP
449 evidence from a Chinese community study. *Scientific Reports* 11:1–13. DOI: 10.1038/s41598-020-
450 79784-z.
- 451 Jardim NYV, Bento-Torres NVO, Costa VO, Carvalho JPR, Pontes HTS, Tomás AM, Sosthenes MCK,
452 Erickson KI, Bento-Torres J, Diniz CWP. 2021. Dual-Task Exercise to Improve Cognition and
453 Functional Capacity of Healthy Older Adults. *Frontiers in Aging Neuroscience* 13:1–16. DOI:
454 10.3389/fnagi.2021.589299.
- 455 Jia RX, Liang JH, Xu Y, Wang YQ. 2019. Effects of physical activity and exercise on the cognitive
456 function of patients with Alzheimer disease: a meta-analysis. *BMC geriatrics* 19. DOI:
457 10.1186/S12877-019-1175-2.

- Joubert C, Chainay H. 2018. Aging brain: The effect of combined cognitive and physical training on cognition as compared to cognitive and physical training alone – A systematic review. *Clinical Interventions in Aging* 13:1267–1301. DOI: 10.2147/CIA.S165399.
- Kalbe E, Roheger M, Paluszak K, Meyer J, Becker J, Fink GR, Kukolja J, Rahn A, Szabados F, Wirth B, Kessler J. 2018. Effects of a Cognitive Training With and Without Additional Physical Activity in Healthy Older Adults: A Follow-Up 1 Year After a Randomized Controlled Trial. *Frontiers in Aging Neuroscience* 10:1–14. DOI: 10.3389/fnagi.2018.00407.
- Karssemeijer EGA, Aaronson JA, Bossers WJR, Donders R, Olde Rikkert MGM, Kessels RPC. 2019. The quest for synergy between physical exercise and cognitive stimulation via exergaming in people with dementia: A randomized controlled trial 17 Psychology and Cognitive Sciences 1701 Psychology 11 Medical and Health Sciences 1103 Clinical Sciences. *Alzheimer's Research and Therapy* 11:1–13. DOI: 10.1186/s13195-018-0454-z.
- Karssemeijer EGA (Esther), Aaronson JA (Justine), Bossers WJ (Willem), Smits T (Tara), Olde Rikkert MGM (Marcel), Kessels RPC (Roy). 2017. Positive effects of combined cognitive and physical exercise training on cognitive function in older adults with mild cognitive impairment or dementia: A meta-analysis. *Ageing Research Reviews* 40:75–83. DOI: 10.1016/j.arr.2017.09.003.
- Köbe T, Witte AV, Schnelle A, Lesemann A, Fabian S, Tesky VA, Pantel J, Flöel A. 2016. Combined omega-3 fatty acids, aerobic exercise and cognitive stimulation prevents decline in gray matter volume of the frontal, parietal and cingulate cortex in patients with mild cognitive impairment. *NeuroImage* 131:226–238. DOI: 10.1016/J.NEUROIMAGE.2015.09.050.
- Knaepen K, Goekint M, Heyman EM, Meeusen R. 2012. Neuroplasticity — Exercise-Induced Response of Peripheral Brain-Derived Neurotrophic Factor. *Sports Medicine* 2010 40:9 40:765–801. DOI: 10.2165/11534530-000000000-00000.
- Lampit A, Hallock H, Valenzuela M. 2014. Computerized Cognitive Training in Cognitively Healthy Older Adults: A Systematic Review and Meta-Analysis of Effect Modifiers. *PLoS Medicine* 11. DOI: 10.1371/journal.pmed.1001756.
- Lauenroth A, Ioannidis AE, Teichmann B. 2016. Influence of combined physical and cognitive training on cognition: A systematic review. *BMC Geriatrics* 16:21–23. DOI: 10.1186/s12877-016-0315-1.
- Law LLF, Barnett F, Yau MK, Gray MA. 2014. Effects of combined cognitive and exercise interventions on cognition in older adults with and without cognitive impairment: a systematic review. *Ageing research reviews* 15:61–75. DOI: 10.1016/J.ARR.2014.02.008.
- Law CK, Lam FM, Chung RC, Pang MY. 2020. Physical exercise attenuates cognitive decline and reduces behavioural problems in people with mild cognitive impairment and dementia: a systematic review. *Journal of Physiotherapy* 66:9–18. DOI: 10.1016/j.jphys.2019.11.014.
- Legault C, Jennings JM, Katula JA, Dagenbach D, Gaussoin SA, Sink KM, Rapp SR, Rejeski WJ, Shumaker SA, Espeland MA. 2011. Designing clinical trials for assessing the effects of cognitive training and physical activity interventions on cognitive outcomes: The Seniors Health and Activity Research Program Pilot (SHARP-P) Study, a randomized controlled trial. *BMC Geriatrics* 11. DOI: 10.1186/1471-2318-11-27.
- Lin S, Yang Y, Qi Q, Wei L, Jing N, Jie Z, Xia L, Shifu X. 2019. The beneficial effect of physical exercise on cognitive function in a non-dementia aging Chinese population. *Frontiers in Aging Neuroscience* 11:1–8. DOI: 10.3389/fnagi.2019.00238.

- 500 Linde K, Alfermann D. 2014. Single versus combined cognitive and physical activity effects on fluid
501 cognitive abilities of healthy older adults: A 4-month randomized controlled trial with follow-up.
502 *Journal of Aging and Physical Activity* 22:302–313. DOI: 10.1123/JAPA.2012-0149.
- 503 Logan PGB and GD. 2014. Extended Practice and Aerobic Exercise Interventions Benefit Untrained
504 Cognitive Outcomes in Older Adults: A Meta-Analysis. *Bone* 23:1–7. DOI: 10.1111/j.1532-
505 5415.2011.03761.x.Extended.
- 506 Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. 2003. Reliability of the PEDro scale for
507 rating quality of randomized controlled trials. *Physical Therapy* 83:713–721. DOI:
508 10.1093/ptj/83.8.713.
- 509 Maillot P, Perrot A, Hartley A. 2012. Effects of interactive physical-activity video-game training on
510 physical and cognitive function in older adults. *Psychology and Aging* 27:589–600. DOI:
511 10.1037/a0026268.
- 512 Meng Q, Yin H, Wang S, Shang B, Meng X, Yan M, Li G, Chu J, Chen L. 2021. The effect of combined
513 cognitive intervention and physical exercise on cognitive function in older adults with mild
514 cognitive impairment: a meta-analysis of randomized controlled trials. *Aging Clinical and*
515 *Experimental Research*. DOI: 10.1007/s40520-021-01877-0.
- 516 Mrakic-Spota S, di Santo SG, Franchini F, Arlati S, Zangiacomi A, Greci L, Moretti S, Jesuthasan N,
517 Marzorati M, Rizzo G, Sacco M, Vezzoli A. 2018. Effects of combined physical and cognitive
518 virtual reality-based training on cognitive impairment and oxidative stress in MCI patients: A pilot
519 study. *Frontiers in Aging Neuroscience* 10:1–11. DOI: 10.3389/fnagi.2018.00282.
- 520 Nguyen L, Murphy K, Andrews G. 2019. Cognitive and neural plasticity in old age: A systematic review
521 of evidence from executive functions cognitive training. *Ageing Research Reviews* 53:100912. DOI:
522 10.1016/j.arr.2019.100912.
- 523 Niederer D, Plaumann U, Seitz T, Wallner F, Wilke J, Engeroff T, Giesche F, Vogt L, Banzer W. 2019.
524 How does a 4-week motor–cognitive training affect choice reaction, dynamic balance and cognitive
525 performance ability? A randomized controlled trial in well-trained, young, healthy participants.
526 *SAGE Open Medicine* 7:205031211987002. DOI: 10.1177/2050312119870020.
- 527 Nishiguchi S, Yamada M, Tanigawa T, Sekiyama K, Kawagoe T, Suzuki M, Yoshikawa S, Abe N,
528 Otsuka Y, Nakai R, Aoyama T, Tsuboyama T. 2015. A 12-week physical and cognitive exercise
529 program can improve cognitive function and neural efficiency in community-dwelling older adults:
530 A randomized controlled trial. *Journal of the American Geriatrics Society* 63:1355–1363. DOI:
531 10.1111/jgs.13481.
- 532 Norouzi E, Vaezmosavi M, Gerber M, Pühse U, Brand S. 2019. Dual-task training on cognition and
533 resistance training improved both balance and working memory in older people. *Physician and*
534 *Sportsmedicine* 47:471–478. DOI: 10.1080/00913847.2019.1623996.
- 535 Ordnung M, Hoff M, Kaminski E, Villringer A, Ragert P. 2017. No overt effects of a 6-week exergame
536 training on sensorimotor and cognitive function in older adults. A preliminary investigation.
537 *Frontiers in Human Neuroscience* 11. DOI: 10.3389/fnhum.2017.00160.
- 538 Passow S, Thurm F, LiC SC. 2014. Plasticity of the aging brain: New directions in cognitive
539 neuroscience. *Science* 346:579–582. DOI: 10.1126/science.1254604.
- 540 Passow S, Thurm F, LiC SC. 2017. Activating developmental reserve capacity via cognitive training or
541 non-invasive brain stimulation: Potentials for promoting fronto-parietal and hippocampal-striatal
542 network functions in old age. *Frontiers in Aging Neuroscience* 9:1–20. DOI:
543 10.3389/fnagi.2017.00033.

- Petersen RC. 2004. Mild cognitive impairment as a diagnostic entity. In: *Journal of Internal Medicine*. 183–194. DOI: 10.1111/j.1365-2796.2004.01388.x.
- Rahe J, Becker J, Fink GR, Kessler J, Kukolja J, Rahn A, Rosen JB, Szabados F, Wirth B, Kalbe E. 2015. Cognitive training with and without additional physical activity in healthy older adults: Cognitive effects, neurobiological mechanisms, and prediction of training success. *Frontiers in Aging Neuroscience* 7:1–15. DOI: 10.3389/fnagi.2015.00187.
- Romera-Liebana L, Orfila F, Segura JM, Real J, Fabra ML, Möller M, Lancho S, Ramirez A, Marti N, Cullell M, Bastida N, Martinez D, Giné M, Cendrós P, Bistuer A, Perez E, Fabregat MA, Foz G. 2018. Effects of a primary care-based multifactorial intervention on physical and cognitive function in frail, elderly individuals: A randomized controlled trial. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* 73:1668–1674. DOI: 10.1093/gerona/glx259.
- Salthouse T. 2012. Consequences of age-related cognitive declines. *Annual Review of Psychology* 63:201–226. DOI: 10.1146/annurev-psych-120710-100328.
- Schättin A, Arner R, Gennaro F, de Bruin ED. 2016. Adaptations of prefrontal brain activity, executive functions, and gait in healthy elderly following exergame and balance training: A randomized-controlled study. *Frontiers in Aging Neuroscience* 8. DOI: 10.3389/fnagi.2016.00278.
- Shatil E. 2013. Does combined cognitive training and physical activity training enhance cognitive abilities more than either alone? A four-condition randomized controlled trial among healthy older adults. *Frontiers in Aging Neuroscience* 5:1–12. DOI: 10.3389/fnagi.2013.00008.
- Shimada H, Makizako H, Doi T, Park H, Tsutsumimoto K, Verghese J, Suzuki T. 2018. Effects of Combined Physical and Cognitive Exercises on Cognition and Mobility in Patients With Mild Cognitive Impairment: A Randomized Clinical Trial. *Journal of the American Medical Directors Association* 19:584–591. DOI: 10.1016/J.JAMDA.2017.09.019.
- Suzuki T, Shimada H, Makizako H, Doi T, Yoshida D, Ito K, Shimokata H, Washimi Y, Endo H, Kato T. 2013. A Randomized Controlled Trial of Multicomponent Exercise in Older Adults with Mild Cognitive Impairment. *PLoS ONE* 8. DOI: 10.1371/journal.pone.0061483.
- Tait JL, Duckham RL, Milte CM, Main LC, Daly RM. 2017. Influence of Sequential vs. Simultaneous Dual-Task Exercise Training on Cognitive Function in Older Adults. *Frontiers in aging neuroscience* 9. DOI: 10.3389/FNAGI.2017.00368.
- Takeuchi H, Magistro D, Kotozaki Y, Motoki K, Nejad KK, Nouchi R, Jeong H, Sato C, Sessa S, Nagatomi R, Zecca M, Takanishi A, Kawashima R. 2020. Effects of Simultaneously Performed Dual-Task Training with Aerobic Exercise and Working Memory Training on Cognitive Functions and Neural Systems in the Elderly. *Neural Plasticity* 2020. DOI: 10.1155/2020/3859824.
- Wollesen B, Wildbrecht A, van Schooten KS, Lim ML, Delbaere K. 2020. The effects of cognitive-motor training interventions on executive functions in older people: A systematic review and meta-analysis. *European Review of Aging and Physical Activity* 17. DOI: 10.1186/s11556-020-00240-y.
- Yang C, Moore A, Mpofu E, Dorstyn D, Li Q, Yin C. 2020. Effectiveness of Combined Cognitive and Physical Interventions to Enhance Functioning in Older Adults with Mild Cognitive Impairment: A Systematic Review of Randomized Controlled Trials. *Gerontologist* 60:E633–E642. DOI: 10.1093/geront/gnz149.
- You JH, Shetty A, Jones T, Shields K, Belay Y, Brown D. 2009. Effects of dual-task cognitive-gait intervention on memory and gait dynamics in older adults with a history of falls: A preliminary investigation. *NeuroRehabilitation* 24:193–198. DOI: 10.3233/NRE-2009-0468.

587 Zhidong C, Wang X, Yin J, Song D, Chen Z. 2021. Effects of physical exercise on working memory in
 588 older adults: a systematic and meta-analytic review. *European Review of Aging and Physical*
 589 *Activity* 18. DOI: 10.1186/s11556-021-00272-y.

590 Zhu X, Yin S, Lang M, He R, Li J. 2016. The more the better? A meta-analysis on effects of combined
 591 cognitive and physical intervention on cognition in healthy older adults. *Ageing Research Reviews*
 592 31:67–79. DOI: 10.1016/j.arr.2016.07.003.

593

Figure 1

PRISMA flow chart of study selection process




Figure 2

Forest plot for the effect sizes of the combined interventions compared to the no intervention




Figure 3

Forest plot for the effect sizes of the combined interventions compared to the exercise intervention




Figure 4

Forest plot for the effect sizes of the combined interventions compared to the cognitive intervention




Table 1 (on next page)

Characteristics of Included Studies

1 Table 1. Characteristics of Included Studies.

Study	Characteristics						Intervention Methods		WM Measures Task	No intervention Group Activities	Study Quality
	Sample Size	Age (Mean)	Cognitive Status	Comparison	Cognitive Intervention	Exercise Intervention	Combination Mode	Intervention Plan			
Fabre 2002	32	60-76 (65.9)	Health	CECT:CT:ET :CG	Mental training	Aerobic training	Separate	60-90 min/session, 3 sessions/week, 8 weeks	WMS-LM I	Leisure activities	7
You 2009	13	64-84 (68.3)	Health	CECT:ET	Memory recall tasks	Walking	Simultaneous (dual-task)	30 min/session, 5 sessions/week, 6 weeks	Memory recall task	No	6
Legault 2011	67	70-85 (76.4)	Health	CECT:CT:ET :CG	Memory training	Aerobic, flexibility training	Separate	50-150 min/session, 3.5 sessions/week, 16 weeks	The 1-Back, 2-Back Tests	Health Education	8
Maillot 2012	30	65-78 (73.5)	Health	CECT:CG	Nintendo Wii games	Nintendo Wii games	Simultaneous (exergame)	60 min/session, 2 sessions/week, 12 weeks	Spatial Span task	No	7
Suzuki 2012	50	65-92 (76.0)	MCI	CECT:CG	Cognitive stimulation	A multicomponent exercise	Simultaneous	90 min/session, 2 sessions/week, 12 weeks	WMS-LM I	Health Education	7
Shatil 2013	122	65-93 (76.8)	Health	CECT:CT:ET :CG	Multidomain cognitive training	Multidomain exercise training	Separate	40-45 min/session, 6 sessions/week, 16 weeks	Auditory working memory test	No	6
Nishiguchi 2015	48	≥60 (73.3)	Health	CECT:CG	Cognitive-motor training	Multidomain exercise training	Simultaneous	90 min/session, 1 sessions/week, 12 weeks	WMS-LM I ,1-back	No	8
Rahe 2015	68	50-85 (68.4)	Health	CECT:CT	Multidomain cognitive training	Strength, endurance, flexibility, coordination	Separate	90 min/session, 2 sessions/week, 7 weeks	WAIS-II(DSB)	No	8
Gschwind 2015	153	≥65 (74.7)	Health	CECT:CG	iStoppFalls exergame with additional cognitive tasks	Balance, strength	Simultaneous (exergame)	60 min/session, 3 sessions/week, 16 weeks	Digit Span Backward (DSB)	Education booklet (passive)	8
Eggenberger 2016	33	>65 74.9	Health	CECT:ET	Cognitive-motor training	Cognitive-motor training	Simultaneous (exergame)	30 min/session, 3 sessions/week, 8 weeks	working memory test	No	8
Schättin 2016	27	≥65 79.2	Health	CECT:ET	Cognitive-motor training	Cognitive-motor training	Simultaneous (exergame)	30 min/session, 3 sessions/week, 8-10 weeks	Working memory test	No	8
Ordnung 2017	30	69.2	Health	CECT:CG	The Microsoft X box 360™	The Microsoft X box 360™	Simultaneous (exergame)	60 min/session, 2 sessions/week, 6 weeks	n-back task	No	7

Damirchi 2017	44	60-85 (68.4)	MCI	CECT:CT:ET :CG	“Modified My Better Mind” program	Walking	Separate	45 min/session, 3 sessions/week, 8 weeks	Forward Digit Span test	No	7
Kalbe 2018	55	50-85 (68.1)	Health	CECT:CT	Multidomain cognitive training	Strength, flexibility, coordination, and endurance	Separate	90 min/session, 2 sessions/week, 7 weeks	Digit Span Backward (DSB)	No	8
Donnezan 2018	69	76.7	MCI	CECT:CT:ET :CG	Multidomain cognitive training	Aerobic training on bikes	Simultaneous	60 min/session, 2 sessions/week, 12 weeks	the Digit Span Backward test	No	7
Bae 2019	83	≥65 (75.9)	MCI	CECT:CG	“KENKOJISEIC HI”system	“KENKOJISEICHI ”system	Simultaneous	90 min/session, 2 sessions/week, 24 weeks	Corsi block-tapping task	Health Education	8
Norouzi 2019	60	≥65 (68.3)	Health	CECT:ET:CG	12 cognitive tasks	Resistance training	Simultaneous (dual-task)	60-80 min/session, 3 sessions/week, 4 weeks	n-back	No	7
Karssemeijer 2019	115	>60 (79.2)	Dementia	CECT:ET:CG	Multidomain cognitive training	Aerobic bicycle training	Simultaneous (exergame)	30-50 min/session, 3 sessions/week, 12 weeks	WAIS-III Digit Span, WMS-III Spatial Span	Relaxation and flexibility exercises	8
Dana 2019	48	60-75 (65.8)	Health	CECT:CT:ET :CG	Multidomain cognitive training	Aerobic treadmill exercises	Separate	60 min/session, 3 sessions/week, 8 weeks	Working memory test	in the waiting list	7
Adcock 2020	31	65-70 (73.9)	Health	CECT:CG	The Active@Home exergame	The Active@Home exergame	Simultaneous (exergame)	30-40 min/session, 3 sessions/week, 16 weeks	The digit span backward task	No	8
Takeuchi 2020	93	65-75 (65.9)	Health	CECT:CT:ET	Working memory training	Aerobic exercise training	Simultaneous (dual-task)	60 min/session, 3 sessions/week, 12 weeks	A digit span task, WMS-LM I, 2 back	No	8

2 MCI: mild cognitive impairment. CECT: combination of exercise and cognitive training. ET: exercise training. CT: cognitive training.
3 CG: control group (no intervention group). WMS-LM I: Wechsler Memory Scale-Logical memory immediate recall. DSB: Digit Span
4 Backward. WAIS-III: Wechsler Adult Intelligence Scale-III.

5

Table 2(on next page)

Moderator Analysis for the Combined Intervention Group vs. the No Intervention Group

1 Table 2. Moderator Analysis for the Combined Intervention Group vs. the No Intervention Group.

Moderator	Level	No. of studies	SMD	95% Confidence Interval	I ²	Homogeneity Test		
						Q	df	p
Mode of combination	Separate	5	0.23	-0.09 to 0.55	4.25	0.17	1	0.68
	Simultaneous	10	0.31**	0.14 to 0.47	30.56			
Cognitive status	Healthy	10	0.24*	0.06 to 0.42	0.00	7.67	2	0.02
	MCI	4	0.58**	0.29 to 0.87	0.00			
	Dementia	1	-0.17	-0.64 to 0.31	0.00			
Intervention length	Short (<12 weeks)	5	0.48*	0.13 to 0.83	0.00	1.47	2	0.48
	Medium (≥12 to <16 weeks)	5	0.27	0.01 to 0.53	42.92			
	Long (≥16 weeks)	5	0.23*	0.03 to 0.44	30.95			
Frequency	Low (<3 sessions/week)	6	0.52**	0.28 to 0.76	0.00	5.53	1	0.02
	High (≥3 sessions/week)	9	0.15	-0.03 to 0.34	4.19			
Session length	Short (≤60min)	9	0.28**	0.09 to 0.46	34.94	0.06	1	0.81
	Long (>60min)	6	0.31*	0.07 to 0.55	0.00			
No intervention group	Active	5	0.22	-0.03 to 0.47	48.93	0.44	1	0.51
	Passive	10	0.33**	0.14 to 0.51	0.38			
Study quality	Low quality (<8 scores)	9	0.44**	0.21 to 0.67	0.00	2.80	1	0.09
	High quality (≥8 scores)	6	0.18	-0.01 to 0.38	39.36			

*p<0.05, **p<0.01