

Association between statin use and risk of gallstone disease and cholecystectomy: A meta-analysis of 590,086 patients

Yu Chang¹ Equal first author, Hong-Min Lin² Equal first author, Kuan-Yu Chi³, Wan-Ying Lin³, Tsung-Ching Chou¹ Corresp.

¹ National Cheng Kung University Hospital, Tainan, Taiwan, [ORCID](#)

² Chi-Mei Medical Center, Tainan, Taiwan, [ORCID](#)

³ Taipei Medical University, Taipei, Taiwan

Corresponding Author: Tsung-Ching Chou

Email address: chou5182@gmail.com

Background Statins have been reported to reduce the risk of gallstone disease. However, the impacts of different durations of statin use on gallstone disease have not been clarified. The aim of this study is to perform a systematic review with meta-analysis to update and to elucidate the association between statin use and the risk of gallstone disease and cholecystectomy. **Methods** Medline, Embase and Cochrane Library were searched from the inception until August 2022 for relevant articles investigating the difference in the risk of gallstone disease between statin users and non-users (PROSPERO, ID: CRD42020182445). Meta-analyses were conducted using odds ratios (ORs) with corresponding 95% confidence intervals (CIs) to compare the risk of gallstone disease and cholecystectomy between statin user and nonusers.. **Results** Eight studies enrolling 590,086 patients were included. Overall, the use of statins was associated with a marginally significant lower risk of gallstone disease than nonusers (OR, 0.91; 95% CI, 0.82 to 1.00). Further subgroup analysis showed that short-term users, medium-term users, and long-term users were associated with a significantly higher risk (OR, 1.18; 95% CI, 1.11 to 1.25), comparable risk (OR, 0.93; 95% CI, 0.83 to 1.04), and significantly lower risk of gallstone diseases (OR, 0.78; 95% CI, 0.68 to 0.90) respectively, compared to nonusers. **Conclusions** Patients with medium-term or long-term use of statins without discontinuation are at a lower risk of gallstone disease or cholecystectomy.

Association between statin use and risk of gallstone disease and cholecystectomy: A meta-analysis of 590,086 patients

Yu Chang^{1†}, M.D, Hong-Min, Lin ^{2†}, M.D, Kuan-Yu Chi^{3,4}, M.D, Wan-Ying, Lin⁵, M.D, Tsung-Ching Chou^{1*}, M.D

¹Departement of Surgery, National Cheng Kung University Hospital, National Cheng Kung University, Tainan, Taiwan

² Department of Family Medicine, Chi-Mei Medical Center, Tainan, Taiwan

³Departement of Education, Center for Evidence-Based Medicine, Taipei Medical University Hospital, Taipei, Taiwan

⁴Department of Internal Medicine, Taipei Medical University Hospital, Taipei, Taiwan

⁵Department of Family Medicine, Taipei Medical University Hospital, Taipei, Taiwan

†Authors equally contributed to the manuscript

*Corresponding author

Tsung-Ching Chou

Attending Doctor, Department of Surgery, National Cheng Kung University Hospital

Tel: 886-6-2353535, Ext 3440

E-Mail: chou5182@gmail.com

24

25

26 **Abstract**

27 **Background**

28 Statins have been reported to reduce the risk of gallstone disease. However, the impacts of
29 different durations of statin use on gallstone disease have not been clarified. The aim of this
30 study is to perform a systematic review with meta-analysis to update and to elucidate the
31 association between statin use and the risk of gallstone disease and cholecystectomy.

32 **Methods**

33 Medline, Embase and Cochrane Library were searched from the inception until August 2022 for
34 relevant articles investigating the difference in the risk of gallstone disease between statin users
35 and non-users (PROSPERO, ID: CRD42020182445). Meta-analyses were conducted using odds
36 ratios (ORs) with corresponding 95% confidence intervals (CIs) to compare the risk of gallstone
37 disease and cholecystectomy between statin user and nonusers..

38 **Results**

39 Eight studies enrolling 590,086 patients were included. Overall, the use of statins was associated
40 with a marginally significant lower risk of gallstone disease than nonusers (OR, 0.91; 95% CI,
41 0.82 to 1.00). Further subgroup analysis showed that short-term users, medium-term users, and
42 long-term users were associated with a significantly higher risk (OR, 1.18; 95% CI, 1.11 to
43 1.25), comparable risk (OR, 0.93; 95% CI, 0.83 to 1.04), and significantly lower risk of gallstone
44 diseases (OR, 0.78; 95% CI, 0.68 to 0.90) respectively, compared to nonusers.

45 **Conclusions**

46 Patients with medium-term or long-term use of statins without discontinuation are at a lower risk

of gallstone disease or cholecystectomy.

Introduction

Gallstone is one of the most common indications for hospitalization in developed countries¹. The prevalence of gallstone in the western population is between 10-30%². Regarding the treatment of symptomatic cholelithiasis, cholecystectomy is an optimal management with its surgical volume reaching up to 70,000 in the United States in 2008^{1,3}. Gallstones primarily fall into two compositions, cholesterol and pigment stones, with the former accounting for 80-90% of all cases². Nowadays, 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors, namely statins, are widely used for the treatment and prevention of cardiovascular disease due to their well-established therapeutic effects on low-density lipoprotein cholesterol (LDL-C), which is one of the major risk factors for atherosclerosis and coronary heart disease⁴. In addition to the effects on cardiovascular system, statins significantly decrease cholesterol biosynthesis, which may potentially reduce cholesterol concentration in the biliary system⁵. Although it has been proven in animal studies that statin may alter biliary lipid composition, thereby reducing the risk of gallstone development^{6,7}, in human studies it differed widely in the outcomes. Some studies ascertained statins' efficacy of decreasing cholesterol concentration in bile or dissolving gallstones⁸⁻¹¹, whereas others found little effect of statins on bile composition¹²⁻¹⁴. The latest meta-analysis¹⁵ in 2015 concluded that statin users have lower risk of gallstone disease. However, considerable heterogeneity has remained unexplored, and the impacts of different durations and discontinuation of statin use on gallstone varied in the literature. For example, while Erichsen et al.¹⁶ reported lower risk of gallstone disease among long-term user, more

recently, Biétry et al.¹⁷ reported comparable risk of gallstone disease among long-term user and nonusers. Moreover, the definition of statin current user or former users was not consistent across published studies. Hence, we sought to perform a meta-analysis to integrate the available and relevant literature and determine the association between statin use and risk of gallstone disease and cholecystectomy.

Methods

We conducted the present systematic review and meta-analysis following Cochrane Handbook for Systematic Reviews and Interventions¹⁸ and reported the results based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses and Meta-analysis Of Observational Studies in Epidemiology guidelines (**Supplemental Method 1 and 2**). Electronic databases of Medline, Embase, and Cochrane library were searched from the inception until August 2022, encompassing all languages. This study was registered on PROSPERO (ID: CRD42020182445). Before the registration, we completed the formal screening of search results against eligibility criteria, since PROSPERO accepts registration for reviews that have not started data extraction. Two investigators (Y.C and K.Y.C) independently conducted the search to determine relevant studies to be included, and any discrepancy was addressed by reaching a consensus or by consulting a senior reviewer (T.C.C). Search details are presented in the **Supplemental Method 3**.

Eligibility criteria

Articles with the following criteria were included: (1) Randomized controlled trials, prospective studies, retrospective studies and case-control studies; (2) studies involving human adults without history of gallstone disease or cholecystectomy as participants; (3) studies reporting

clinical outcome as with/without diagnosis of gallstone disease or record of cholecystectomy. Studies not investigating the use of statins were excluded. In case of duplicate studies with an accumulated number of patients or increased durations of follow-up, only the most complete reports were included

Data extraction

Two investigators (Y.C and K.Y.C) independently extracted relevant information from tables or results of eligible articles. Extracted data included first author name, publication year, country where the study was conducted, follow-up durations, number of participants, sex, age range, underlying diseases, and method used to identify cases. Additionally, we obtained adjusted odds ratio (OR) and standard error (SE) from each study. For studies that reported outcomes by relative risk (RR), since the extracted RR and SE were adjusted, the formula¹⁹ for the conversion between RR and OR may be inappropriate. Therefore, we regarded the adjusted RR as OR since most of the included studies have incidence of outcome less than 10% and all ORs converted from RRs were between 0.5 and 2.5, correction may not be desirable²⁰.

Quality assessment

Two investigators (Y.C and H.M.L) independently completed a critical appraisal of included literature by using the Risk Of Bias In Non-randomized Studies - of Interventions (ROBINS-I) tool²¹ for retrospective cohorts and National Heart, Lung, and Blood Institute tool for case-control studies. Any item on which assessors did not reach a consensus was addressed through discussion with third investigator (T.C.C).

Statistical analysis

Meta-analysis was conducted using R software with the “metafor” package²² (**Supplemental**

Method 4). Regarding the risk of gallstone disease, we used the inverse variance (IV) method to pool the odds ratio obtained from each study regarding risk of gallstone disease among statin users and non-users. The restricted maximum likelihood (REML)²³ method was exploited as a heterogeneity estimator for conducting random-effects meta-analyses given that the between-trial variance was inevitable. P values <0.05 were considered statistically significant. Heterogeneity was assessed using I^2 statistics proposed by Higgins and Thompson²⁴, with estimated values of $I^2 < 25\%$, $25\% < I^2 < 50\%$, and $I^2 \geq 50\%$ indicating low, moderate, and high heterogeneity, respectively. We aimed to explore the statistical heterogeneity of our meta-analysis by performing subgroup analysis based on the durations of statin use (long-term, medium-term and short-term) and current or former statin users.

Results

Study selection

Our search strategy identified 1007 references from Medline, Embase, and Cochrane library. After title and abstract screening, we excluded duplicates (n= 49) and irrelevant references (n= 942) and retrieved 16 full texts for further review. Eventually, 6 case-control studies^{16-16,17,26-29} and 2 retrospective cohort studies^{25,30} met the eligibility criteria for qualitative and quantitative synthesis. **Figure 1** shows the flowchart of study selection.

Characteristic of included studies and patients (Table 1)

Among 8 studies^{16,17,25-30} enrolling 590,086 participants from 1994 to 2012, 5 studies^{17,25,26,228,29} recruited patients who underwent cholecystectomy due to gallstone diseases, and other 3 studies^{16,27,30} included patients with the diagnosis of gallstone diseases. Of note, 5 studies^{16,17,26-28} specifically categorized patients into statin current, past, or nonusers, and the other 3 studies^{25,29,30} referred to statin users as any statin prescription prior to the index date, defined as

the first-time diagnosis of gallstone diseases or the date of cholecystectomy in patients without the record of cholelithiasis. Detailed patient characteristics were summarized in **Supplemental Table 1 and 2**. Regarding the durations of statin use, 3 of the included studies^{16,17,28} classified the durations of statin use based on the numbers of prescription prior to the index date (**Supplemental Table 3 and Table 4**).

Quality assessment of included studies

Supplemental Table 5 demonstrated the summary of quality assessment for case-control studies using the National Heart, Lung, and Blood Institute tool, and the overall quality was good in 3 studies and the other 3 were fair. **Supplemental Table 6** demonstrated the risk of bias assessment using ROBINS-I tool for two retrospective cohort studies, which were evaluated as moderate risk of bias resulted from bias due to confounding.

Risk of cholelithiasis and cholecystectomy

Overall, the pooled result of 8 studies^{16,17,25-30} demonstrated that the use of statins was associated with a marginally significant lower risk of gallstone disease than nonusers, however, with substantial heterogeneity (OR, 0.91; 95% CI, 0.82 to 1.00, $I^2=86\%$; **Figure 2**).

Durations of statin use

To explore the source of heterogeneity, we performed subgroup analysis according to the durations of statin use in three studies^{16,17,28} that exclusively classified the durations based on the number of prescriptions prior to the index date (short-term, 1-4; medium, 5-19; long-term, over 20). Notably, our subgroup analysis seemed to show a dose-response tendency (**Figure 3**). Short-term use of statins is associated with higher risk of gallstone disease in both current (OR, 1.16;

95% CI, 1.07 to 1.26, $I^2 = 0\%$; with gallstone disease: 804 patients; without gallstone disease: 4430 patients) and former users (OR, 1.20; 95% CI, 1.10 to 1.30, $I^2 = 35\%$; with gallstone disease: 717 patients; without gallstone disease: 3469 patients). Medium-term current statin user had significant lower risk of gallstone disease compared to nonusers (OR, 0.86; 95% CI, 0.81 to 0.91, $I^2 = 15\%$; with gallstone disease: 1737 patients; without gallstone disease: 10923 patients) and former users had comparable risk of gallstone disease compared to nonusers (OR, 1.05; 95% CI, 0.91 to 1.21, $I^2 = 50\%$; with gallstone disease: 587 patients; without gallstone disease: 3259 patients). Long-term use of statin significantly reduced the risk of gallstone disease in current users (OR, 0.73; 95% CI, 0.62 to 0.87, $I^2 = 82\%$; with gallstone disease: 1451 patients; without gallstone disease: 9096 patients) but former users had comparable risk of gallstone disease compared to nonusers (OR, 0.87; 95% CI, 0.72 to 1.05, $I^2 = 0\%$; with gallstone disease: 204 patients; without gallstone disease: 1327 patients). Additionally, the pooled analysis of 4 studies^{16,25,27,30} involving statin-users over 1-year period showed a significantly lower risk of gallstone diseases (OR, 0.83; 95% CI, 0.77 to 0.89; $I^2 = 4\%$; *Figure 4*).

Discussion

Risk of cholecystectomy and cholelithiasis

Statins are one of the most common prescribed regimens in the treatment and prevention of cardiovascular disease³¹. In addition to lowering the risk of cardiovascular disease, some studies have demonstrated that statins may reduce the risk of urolithiasis^{32,33}. Notably, our study revealed that statin users had marginally significant reduced risk of gallstone disease, compared to non-users, however, with high statistical heterogeneity ($I^2=71\%$).

One of the sources of heterogeneity would arise from the differences in demographics among included studies. For example, there were wide variation in prevalence of diabetes, ischemic heart diseases, and cerebrovascular diseases among included studies. Moreover, baseline characteristics that may be associated with gallstone disease such as hyperlipidemia, estrogen exposure or hormone therapy were only reported in two studies. On the other hand, we acquired adjusted ORs from included studies and pooled in meta-analysis, however, the variables adjusted for were not consistent in each study, which would contribute to the statistical heterogeneity. Most of the included studies defined case as “cholecystectomy required”. The use of statin may not only impact the formation of gallstone but the symptom severity of gallstone disease. For instance, Suuronen et al. found a decreased number of all cholecystectomies but increased number of laparoscopic cholecystectomy (LC) with statin use in their population-based study³⁴. They described one of the reasons may be an increased incidence of mild symptomatic gallstones leading to LC. The utilization of different approaches may reflect the severity of the disease, however, none of included studies investigated the difference between open and laparoscopic cholecystectomy.

Gallstone disease risk of statin users with different durations of use

Our results demonstrated that there was a marginally significant reduction in the risk of developing gallstone disease among statin-ever users. The subgroup analysis showed a dose-response relationship between the number of prescription and the risk of gallstone diseases, and, of note, the protective effect of stain was only observed in medium and long-term current users. However, this result had to be interpreted with caution since the durations of statin use was solely based on the prescription of medication in medical records. Paucity of detailed dosage of statin prescription and information for drug compliance may hindered the accuracy of statin

exposure. Moreover, the cut-off time selected for the definition of current and former users varied from 90 days to 180 days representing an obvious source of conceptual heterogeneity. As a matter of fact, long-term statin users only accounted for 15% of all statin users in our analysis. We postulated that the underlying reason for such small proportion may be associated with either discontinuation of statins or low adherence to medications. Although statins are one of the most effective medications for the secondary prevention of cardiovascular diseases, non-compliance and discontinuation with this medication are common problems³⁵⁻³⁷. Although patient-perceived side effect is the most common reason for statin discontinuation³⁸, the incidence of statins' most common side effect, namely myopathy, is only estimated to be around 0.01%³⁹. Of note, those not adhering to statins, mostly due to self-perceived stable disease condition⁴⁰, may experience relapsing dyslipidemia following statin discontinuation leading to worsening cardiovascular and survival outcomes^{41,42}. Consequently, we propose that the recurrence of dyslipidemia may be responsible for neutral and detrimental effects on gallstone formation in medium- and short-term users, respectively.

Compare to previous synthesis

In our study, we included three more studies^{17,27,30} compared with the previous one. In accordance with the study by Kan et al.¹⁵, our results demonstrated patients with medium-term use and long-term use of statin users conferred to lower risk of gallstone disease. However, we addressed several concerns that were not well-elucidated in previous study. First, we clarified the impacts of statin discontinuation on gallstone disease. Second, we excluded cross-sectional study in our meta-analysis since this type of study is estimating the prevalence but incidence⁴³. The cross-sectional study by Caroli-Bosc⁴⁴ in the previous meta-analysis might not contribute to statistical heterogeneity, however, conceptually, it brings between-study variance. Third,

methodologically, we performed our meta-analysis through a more robust heterogeneity estimator based on the latest simulation study⁴⁵.

Limitation

The findings of our meta-analysis must be interpreted within the context of the study limitations. First, we included case-control studies and retrospective cohort studies, which induced a potential bias due to the confounding that was generally not well-controlled. Second, we investigated the protective effect of statin on gallstone development, however, the definition of statin use, and gallstone disease were not consistent in each study. Third, baseline characteristics that may be associated with gallstone formation such as hyperlipidemia, estrogen exposure or hormone therapy were not reported in several studies, which hindered the accuracy of our findings. Last but not least, healthy user and adherer effect are two easily overlooked sources of bias, especially in observational studies⁴⁶. Patients who are willing to take and adhere to statins are more likely to be engaged with healthy behaviors, contributing to the protective effects of statins. Unfortunately, It is difficult to adjust these healthier behaviors through observational design, leading to unmeasured confounding bias.

Conclusions

Based on our meta-analysis, long-term and medium-term use of statins without discontinuation lower the risk of gallstone disease or cholecystectomy. Nonetheless, given the observational nature of the included studies, we propose that further prospective studies should be undertaken to overcome the aforementioned limitations.

Acknowledgments

248 Not applicable

249 Reference

- 250 1. Beckingham IJ. ABC of diseases of liver, pancreas, and biliary system. Gallstone disease.
251 Bmj. 2001;322(7278):91-94.
- 252 2. Marschall HU, Einarsson C. Gallstone disease. J Intern Med. 2007;261(6):529-542.
- 253 3. Lammert F, Miquel JF. Gallstone disease: from genes to evidence-based therapy. J
254 Hepatol. 2008;48 Suppl 1:S124-135.
- 255 4. Awad K, Serban MC, Penson P, et al. Effects of morning vs evening statin administration
256 on lipid profile: A systematic review and meta-analysis. J Clin Lipidol. 2017;11(4):972-
257 985.e979.
- 258 5. Kallien G, Lange K, Stange EF, Scheibner J. The pravastatin-induced decrease of biliary
259 cholesterol secretion is not directly related to an inhibition of cholesterol synthesis in
260 humans. Hepatology. 1999;30(1):14-20.
- 261 6. Davis KG, Wertin TM, Schriver JP. The Use of Simvastatin for the Prevention of
262 Gallstones in the Lithogenic Prairie Dog Model. Obesity Surgery. 2003;13(6):865-868.
- 263 7. Abedin MZ, Narins SC, Park EH, Smith PR, Kirkwood KS. Lovastatin alters biliary lipid
264 composition and dissolves gallstones: A long-term study in prairie dogs. Digestive Diseases
265 and Sciences. 2002;47(10):2192-2210.
- 266 8. Porsch-Ozçürümez M, Hardt PD, Schnell-Kretschmer H, et al. Effects of fluvastatin on
267 biliary lipids in subjects with an elevated cholesterol saturation index. European journal of
268 clinical pharmacology. 2001;56(12):873-879.
- 269 9. Smith JL, Roach PD, Wittenberg LN, et al. Effects of simvastatin on hepatic cholesterol
270 metabolism, bile lithogenicity and bile acid hydrophobicity in patients with gallstones.
271 Journal of Gastroenterology and Hepatology (Australia). 2000;15(8):871-879.
- 272 10. Chapman BA, Burt MJ, Chisholm RJ, Allan RB, Yeo KHJ, Ross AG. Dissolution of
273 gallstones with simvastatin, an HMG CoA reductase inhibitor. Digestive Diseases and
274 Sciences. 1998;43(2):349-353.
- 275 11. Wilson IR, Hurrell MA, Pattinson NR, Chapman BA. The effect of simvastatin and
276 bezafibrate on bile composition and gall-bladder emptying in female non-insulin-dependent
277 diabetics. Journal of Gastroenterology and Hepatology (Australia). 1994;9(5):447-451.
- 278 12. Miettinen TE, Kiviluoto T, Taavitsainen M, Vuoristo M, Miettinen TA. Cholesterol
279 metabolism and serum and biliary noncholesterol sterols in gallstone patients during
280 simvastatin and ursodeoxycholic acid treatments. Hepatology. 1998;27(3):649-655.
- 281 13. Smit JW, van Erpecum KJ, Renooij W, et al. The effects of the 3-hydroxy, 3-
282 methylglutaryl coenzyme A reductase inhibitor pravastatin on bile composition and
283 nucleation of cholesterol crystals in cholesterol gallstone disease. Hepatology.
284 1995;21(6):1523-1529.

14. Sharma BC, Agarwal DK, Baijal SS, Saraswat VA. Pravastatin has no effect on bile lipid composition, nucleation time, and gallbladder motility in persons with normal levels of cholesterol. *Journal of Clinical Gastroenterology*. 1997;25(2):433-436.
15. Kan HP, Guo Wb, Tan YF, Zhou J, Liu CD, Huang YQ. Statin use and risk of gallstone disease: A meta-analysis. *Hepatology Research*. 2015;45(9):942-948.
16. Erichsen R, Frøslev T, Lash TL, Pedersen L, Sørensen HT. Long-term statin use and the risk of gallstone disease: A population-based case-control study. *American Journal of Epidemiology*. 2011;173(2):162-170.
17. Biétry FA, Reich O, Schwenkglenks M, Meier CR. Statin use and risk of cholecystectomy—A case-control analysis using Swiss claims data. *Expert Opinion on Drug Safety*. 2016;15(12):1577-1582.
18. Julian Higgins JT, Jacqueline Chandler, Miranda Cumpston, Tianjing Li, Matthew Page, Vivian Welch. *Cochrane Handbook for Systematic Reviews of Interventions* version 6.0. In:2019.
19. Grant RL. Converting an odds ratio to a range of plausible relative risks for better communication of research findings. *Bmj*. 2014;348:f7450.
20. Zhang J, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *Jama*. 1998;280(19):1690-1691.
21. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919-i4919.
22. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software*. 2010;36(3).
23. Harville DA. Maximum Likelihood Approaches to Variance Component Estimation and to Related Problems. *Journal of the American Statistical Association*. 1977;72(358).
24. Higgins JPT. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557-560.
25. Tsai C, Leitzmann MF, Willett WC, Giovannucci EL. Statin Use and the Risk of Cholecystectomy in Women. *Gastroenterology*. 2009;136(5):1593-1600.
26. Merzon E, Weiss NS, Lustman AJ, Elhayani A, Dresner J, Vinker S. Statin administration and risk of cholecystectomy: A population-based case-control study. *Expert Opinion on Drug Safety*. 2010;9(4):539-543.
27. González-Pérez A, García Rodríguez LA. Gallbladder disease in the general population: Association with cardiovascular morbidity and therapy. *Pharmacoepidemiology and Drug Safety*. 2007;16(5):524-531.
28. Bodmer M, Brauchli YB, Krähenbühl S, Jick SS, Meier CR. Statin use and risk of gallstone disease followed by cholecystectomy. *JAMA - Journal of the American Medical Association*. 2009;302(18):2001-2007.
29. Chiu HF, Chen CC, Kuo HW, Lee IM, Wu TN, Yang CY. Statin use and the risk of gallstone disease: A population-based casecontrol study. *Expert Opinion on Drug Safety*. 2012;11(3):369-374.

30. Martin D, Schmidt R, Mortensen EM, Mansi I. Association of Statin Therapy and Risks of Cholelithiasis, Biliary Tract Diseases, and Gallbladder Procedures: Retrospective Cohort Analysis of a US Population. *Annals of Pharmacotherapy*. 2016;50(3):161-171.
31. Chou R, Dana T, Blazina I, Daeges M, Jeanne TL. Statins for Prevention of Cardiovascular Disease in Adults: Evidence Report and Systematic Review for the US Preventive Services Task Force. *Jama*. 2016;316(19):2008-2024.
32. Cohen AJ, Adamsky MA, Nottingham CU, et al. Impact of Statin Intake on Kidney Stone Formation. *Urology*. 2019;124:57-61.
33. Sur RL, Masterson JH, Palazzi KL, et al. Impact of statins on nephrolithiasis in hyperlipidemic patients: a 10-year review of an equal access health care system. *Clin Nephrol*. 2013;79(5):351-355.
34. Suuronen S, Niskanen L, Paajanen P, Paajanen H. Declining cholecystectomy rate during the era of statin use in Finland: a population-based cohort study between 1995 and 2009. *Scandinavian journal of surgery : SJS : official organ for the Finnish Surgical Society and the Scandinavian Surgical Society*. 2013;102(3):158-163.
35. Navar AM, Roe MT, White JA, et al. Medication discontinuation in the IMPROVE IT trial. *Circ Cardiovasc Qual Outcomes* 2019;12(1): e005041. 6.
36. Toth PP, Granowitz C, Hull M, Anderson A, Philip S. Long-term statin persistence is poor among high-risk patients with dyslipidemia: a real-world administrative claims analysis. *Lipids Health Dis* 2019;18(1):175.
37. Wei L, Wang J, Tompson P, Struthers S, MacDonald TM. Adherence to statin treatment and readmission of patients after myocardial infarction: a six year follow up study. *Heart*. 2002;88(3):229-33
38. Bradley CK, Wang TY, Li S, Robinson JG, Roger VL, Goldberg AC, Virani SS, Louie MJ, Lee LV, Peterson ED, Navar AM. Patient-Reported Reasons for Declining or Discontinuing Statin Therapy: Insights From the PALM Registry. *J Am Heart Assoc*. 2019 Apr 2;8(7)
39. Thompson PD, Clarkson P, Karas RH. Statin-associated myopathy. *JAMA*. 2003 Apr 2;289(13):1681-90. doi: 10.1001/jama.289.13.1681.
40. Kamal-Bahl SJ, Burke T, Watson D, Wentworth C. Discontinuation of lipid modifying drugs among commercially insured United States patients in recent clinical practice. *Am J Cardiol*. 2007 Feb 15;99(4):530-4.
41. Daskalopoulou SS, Delaney JA, Filion KB, et al. Discontinuation of statin therapy following an acute myocardial infarction: a population-based study. *Eur Heart J* 2008; Jul 29
42. Rea F, Biffi A, Ronco R, Franchi M, Cammarota S, Citarella A, Conti V, Filippelli A, Sellitto C, Corrao G. Cardiovascular Outcomes and Mortality Associated With Discontinuing Statins in Older Patients Receiving Polypharmacy. *JAMA Netw Open*. 2021 Jun 1;4(6)
43. Sedgwick P. Cross sectional studies: advantages and disadvantages *BMJ* 2014; 348 :g2276
44. Caroli-Bosc FX, Le Gall P, Pugliese P, et al. Role of fibrates and HMG-CoA reductase inhibitors in gallstone formation: epidemiological study in an unselected population. *Digestive Diseases & Sciences*.46(3):540-544.

45. Langan D, Higgins JPT, Jackson D, et al. A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses. *Res Synth Methods*. 2019;10(1):83-98.

46. Frank WH, Patrick AR, Brookhart MA. Healthy user and related biases in observational studies of preventive interventions: a primer for physicians. *J Gen Intern Med*. 2011 May;26(5):546-50.

Table 1 (on next page)

Characteristics of included studies

Characteristics of included studies

Table 1 Characteristics of included studies

Study, year	Study type	Country	Inclusion period	Match method	Exclusion criteria	Case definition	Statin use
Biétry 2016 ²³	CC	Switzerland	2008~2014	1:4 matched on age, sex, and index date	Cancer or HIV	Cholecystectomy only	Prior to index date†, Current user: last prescription <180 days Former user: last prescription >180 days
Bodmer 2009 ¹⁹	CC	UK	1994~2008	1:4 matched on age, sex, and index date	Alcoholism, drug abuse, cancer or HIV	First time diagnosed gallstone disease followed by cholecystectomy in 2 years; or cholecystectomy only	Prior to index date, Current user: last prescription <90 days Former user: last prescription >90 days
Chiu 2012 ²¹	CC	Taiwan	1996~2009	1:1 matched on age, sex, and index date	Cancer	First time diagnosed gallstone disease followed by cholecystectomy in 2 years; or cholecystectomy only	Any prescription prior to index date
Erichsen 2010 ¹⁷	CC	Denmark	1996~2008	1:10 matched on age, sex	Preexisting gallstone disease or liver, bile duct or pancreatic cancer	Diagnosis of gallstone, cholecystitis or a record gallbladder surgery (cholecystectomy or drainage)	Prior to index date, Current user: last prescription <90 days; Former user: last prescription >90 days
González-Pérez 2007 ¹⁸	CC	UK	1996~1996	1:4 matched on age and sex	Preexisting gallstone disease or cancer	Symptomatic gallstone	Prior to index date, Current user: last prescription <30 days;

							Former user: last prescription >30 days
Merzon 2010 ¹⁶	CC	Israel	2003~2006	1:4 matched on age and sex	N/A	Cholecystectomy due to gallstone disease	Prior to index date, Last prescription <180 days prior to index date
Martin 2015 ²⁴	RC	US	2003~2012	1:1 propensity matched	History of burn, trauma, statin use <90 days, or starting statin use after baseline period	Diagnosis of cholelithiasis	Statin use lasting over 90 days prior to index date
Tsai 2009 ¹⁵	RC and PC	US	1994~2000	N/A	Prior cholecystectomy or gallstone disease or cancer	Self-reported cholecystectomy	Prior to index date, Current user: self-reported current use from inception to 2000, Former user: no present use in 2000

CC, Case-control study, RC, retrospective cohort study, NA, not applicable

2
3
4
5
6
7
8
9
10

Figure 1

PRISMA flow diagram

The PRISMA flow diagram demonstrates a total of 1007 potential references were extracted initially and meta-analysis included 8 studies meeting the eligibility criteria.



PRISMA 2009 Flow Diagram




Figure 2

Risk of gallstone disease among statin users and nonusers

Forest plot for risk of gallstone disease among statin users versus nonusers. The size of squares is proportional to the weight of each study with the error bar showing the 95% CI. Red diamond represents the pooled estimate of OR with the 95% CI. CI, confidence interval; IV, inverse variance method; OR, odds ratio; SE, standard error of logOR; TE, logOR





Figure 3


Risk of gallstone disease based on different duration of statin use

Forest plot for risk of gallstone disease among short-term, medium-term and long-term statin users compared to nonusers. Subgroup analysis was performed based on current and former users. The size of squares is proportional to the weight of each study with the error bar showing the 95% CI. Red diamond represents the pooled estimate of OR with the 95% CI. CI, confidence interval; IV, inverse variance method; OR, odds ratio; SE, standard error of logOR; TE, logOR

Short-term user



Medium-term user



Long-term user




Figure 4

Risk of gallstone disease among long-term statin users (over one year)

Forest plot for risk of gallstone disease among long-term statin users versus nonusers.

Subgroup analysis was performed based on current and former users. The size of squares is proportional to the weight of each study with the error bar showing the 95% CI. Red diamond represents the pooled estimate of OR with the 95% CI. CI, confidence interval; IV, inverse variance method; OR, odds ratio; SE, standard error of logOR; TE, logOR

