Metabolic shift and the effect of mitochondrial respiration in osteogenic differentiation of dental pulp stem cells (#79009)

First submission

Guidance from your Editor

Please submit by 18 Dec 2022 for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the <u>materials page</u>.

Custom checks

5 Figure file(s)

5 Raw data file(s)

Human participant/human tissue checks

- Have you checked the authors ethical approval statement?
- Does the study meet our <u>article requirements</u>?
- Has identifiable info been removed from all files?
- Were the experiments necessary and ethical?

Structure and Criteria



Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING
- 2. EXPERIMENTAL DESIGN
- 3. VALIDITY OF THE FINDINGS
- 4. General comments
- 5. Confidential notes to the editor
- You can also annotate this PDF and upload it as part of your review

When ready submit online.

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your guidance page.

BASIC REPORTING

- Clear, unambiguous, professional English language used throughout.
- Intro & background to show context.
 Literature well referenced & relevant.
- Structure conforms to <u>PeerJ standards</u>, discipline norm, or improved for clarity.
- Figures are relevant, high quality, well labelled & described.
- Raw data supplied (see <u>PeerJ policy</u>).

EXPERIMENTAL DESIGN

- Original primary research within Scope of the journal.
- Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
- Rigorous investigation performed to a high technical & ethical standard.
- Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

- Impact and novelty not assessed.

 Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
- All underlying data have been provided; they are robust, statistically sound, & controlled.



Conclusions are well stated, linked to original research question & limited to supporting results.



Standout reviewing tips



The best reviewers use these techniques

п	n
	v

Support criticisms with evidence from the text or from other sources

Give specific suggestions on how to improve the manuscript

Comment on language and grammar issues

Organize by importance of the issues, and number your points

Please provide constructive criticism, and avoid personal opinions

Comment on strengths (as well as weaknesses) of the manuscript

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Your introduction needs more detail. I suggest that you improve the description at lines 57-86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

- 1. Your most important issue
- 2. The next most important item
- 3. ...
- 4. The least important points

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.



Metabolic shift and the effect of mitochondrial respiration in osteogenic differentiation of dental pulp stem cells

Lingyun Wan Equal first author, 1, Linyan Wang Equal first author, 2, Ran Cheng 1, Li Cheng Corresp., 1, Tao Hu

Corresponding Authors: Li Cheng, Tao Hu

Email address: chengli@scu.edu.cn, hutao@scu.edu.cn

Background.

Metabolism shifts from glycolysis to mitochondrial oxidative phosphorylation is important during differentiation of stem cells. Mitochondria function as directors in cell differentiation rather than bystanders. However, metabolic shift and the effect of mitochondria regulating differentiation of human dental pulp stem cells (hDPSCs) remain unclear.

Methods.

Human dental pulp stem cells were collected from 5 healthy donors. Cell differentiation was induced by osteogenic mineralized medium. The activities of alkaline phosphatase, hexokinase, pyruvate kinase, and lactate dehydrogenase were measured by enzyme activity kits. The extracellular acidification rate and the mitochondrial oxygen consumption rate were detected. The mRNA level of COL-1, ALP, TFAM and NRF1 was analyzed. The protein level of p-AMPK and AMPK were measured by western blot.

Results.

Glycolysis decreased after a slight increase, while mitochondrial oxidative phosphorylation kept on rising when cells growing in osteogenic mineralized medium. Thereby, the metabolism of differentiating cells switched to mitochondrial respiration. Next, inhibition of mitochondrial respiration with carbonyl cyanidem-chlorophenylhydrazone, a mitochondrial uncoupler, depressed hDPSCs differentiation with less ALP activity and decreased ALP and COL-1 mRNA expression. Furthermore, mitochondrial uncoupler leaded to AMPK activation. 5-aminoimidazole-4-carboxamideribonucleotide, an AMPK activator simulated the effect of mitochondrial uncoupler by inhibiting cell differentiation, mitochondrial biogenesis and mitochondrial morphology. Mitochondrial uncoupler and activation of AMPK depressed mitochondrial oxidative phosphorylation and inhibited differentiation, suggesting that they may serve as a regulator to guit osteogenic differentiation from impaired mitochondrial oxidative phosphorylation.

¹ West China Hospital of Stomatology, Sichuan University, Chengdu, Sichuan, China

² Chengdu second People's Hospital, Chengdu, Sichuan, China



1 Metabolic shift and the effect of mitochondrial respiration in

2 osteogenic differentiation of dental pulp stem cells

- 3 Lingyun Wan^{1#}, LinyanWang^{2#}, Ran Cheng¹, Cheng Li¹, Tao Hu¹
- 4 ¹, State Key Laboratory of Oral Diseases, Frontier Innovation Center for Dental Medicine Plus,
- 5 National Clinical Research Center for Oral Diseases, Dept. of Preventive Dentistry, West China
- 6 Hospital of Stomatology, Sichuan University, Chengdu, Sichuan, China.
- ⁷ Pepartment of Stomotology, Chengdu second People's Hospital, Chengdu, Sichuan, China
- 8 Corresponding authors:
- 9 Tao Hu¹,Cheng Li¹
- 10 State Key Laboratory of Oral Diseases, Frontier Innovation Center for Dental Medicine Plus,
- 11 National Clinical Research Center for Oral Diseases, Dept. of Preventive Dentistry, West China
- 12 Hospital of Stomatology, Sichuan University, Chengdu, Sichuan, China.
- $13 \quad \hbox{E-mail address:} hut ao@scu.edu.cn \ , chengli@scu.edu.cn$
- 14 Tel: 86-28-85503486
- 15 *, Contributed to this work equally.

16

17



18 Abstract

- 19 Background.
- 20 Metabolism shifts from glycolysis to mitochondrial oxidative phosphorylation is important
- 21 during differentiation of stem cells. Mitochondria function as directors in cell differentiation
- 22 rather than bystanders. However, metabolic shift and the effect of mitochondria regulating
- 23 differentiation of human dental pulp stem cells (hDPSCs) remain unclear.
- 24 Methods.
- 25 Human dental pulp stem cells were collected from 5 healthy donors. Cell differentiation was
- 26 induced by osteogenic mineralized medium. The activities of alkaline phosphatase, hexokinase,
- 27 pyruvate kinase, and lactate dehydrogenase were measured by enzyme activity kits. The
- 28 extracellular acidification rate and the mitochondrial oxygen consumption rate were detected.
- 29 The mRNA level of COL-1, ALP, TFAM and NRF1 was analyzed. The protein level of p-AMPK
- and AMPK were measured by western blot.
- 31 Results.
- 32 Glycolysis decreased after a slight increase, while mitochondrial oxidative phosphorylation kept
- on rising when cells growing in osteogenic mineralized medium. Thereby, the metabolism of
- 34 differentiating cells switched to mitochondrial respiration. Next, inhibition of mitochondrial
- 35 respiration with carbonyl cyanidem-chlorophenylhydrazone, a mitochondrial uncoupler,
- 36 depressed hDPSCs differentiation with less ALP activity and decreased ALP and COL-1 mRNA
- 37 expression. Furthermore, mitochondrial uncoupler leaded to AMPK activation.5-
- 38 aminoimidazole-4-carboxamideribonucleotide, an AMPK activator simulated the effect of





39	mitochondrial uncoupler by inhibiting cell differentiation, mitochondrial biogenesis and
40	mitochondrial morphology. Mitochondrial uncoupler and activation of AMPK depressed
41	mitochondrial oxidative phosphorylation and inhibited differentiation, suggesting that they may
12	serve as a regulator to quit osteogenic differentiation from impaired mitochondrial oxidative
13	phosphorylation.
14	Keywords: dental pulp stem cells; osteogenic differentiation; glycolysis; mitochondrial
45	oxidative phosphorylation,
46	
17	



49

Introduction

the ability to differentiate into multiple types of cell types, including but not limited to 50 odontoblast/osteoblast, chondrocyte, adipocytes, myogenic and neural cells(Sui et al., 2020; 51 Tsutsui, 2020). Due to their easily isolation and great potential in tissue engineering and 52 regenerative medicine, DPSCs are widely used in various fields. Numerous studies are 53 dedicating to uncover the detailed mechanisms involved in their self-renewal ability and multi-54 lineage differentiation potential. The capacity of dental pulp stem cells to differentiate is 55 important for dental pulp repair and dentin rebuilding (Sui et al., 2020). 56 Energy metabolism plays the critical role in guiding the proliferation, differentiation and 57 many life processes of stem cells. Energy production by aerobic oxidative phosphorylation is the 58 59 main function of mitochondria (Rossmann et al., 2021). In stem cells, mitochondria produce the energy not only for the maintenance of homeostasis, but also for differentiation(Tsutsui, 60 61 2020). The metabolism predilection shifts between mitochondrial oxidative phosphorylation 62 (OXPHOS) and glycolysis change along cell status and different levels of mitochondrial maturation, which depend on the pluripotency stage (Khacho & Slack, 2018; Wanet et al., 2015). 63 For instance, fully differentiated cells rely more on OXPHOS, while undifferentiated stem cells 64 rely more on glycolysis(Ly, Lynch & Ryall, 2020). Some evidence also shows a metabolic 65 switch from anaerobic glycolysis toward OXPHOS upon differentiation of hematopoietic stem 66 67 cells (HSCs). Quiescent HSCs rely on glycolysis, while the active HSCs change to OXPHOS metabolism when cells undergo differentiation (Morganti, Cabezas-Wallscheid & Ito, 2022). 68

Human dental pulp stem cells (hDPSCs), as a source of adult multipotent stem cells, present

dot should be removed before literature citing



Moreover, the dynamic distribution, subcellular content and structural of mitochondria, 69 have been shown to exhibit peculiar functions in the processes of cellular differentiation and 70 reprogramming (Khacho & Slack, 2017). Mitochondrial-mediated shift accompanied by 71 mitochondrial remodeling and dynamics are vital to the differentiation and the fate of neural 72 stem cells (Coelho et al., 2022). Our previous study demonstrated that at the initial stage of 73 hDPSCs differentiation, both the OXPHOS and glycolysis were upregulated (Wang et al., 2016). 74 It should be emphasized whether the mitochondrial and glycolysis change exist during hDPSCs 75 continuous differentiation. Other questions that need to be addressed are the role of mitochondria 76 upon differentiation of hDPSCs. 77 Some studies are underway to unveil the bidirectional crosstalk between mitochondria and 78 differentiation, such as reactive oxygen species production, energy-sensing pathways, hypoxia-79 80 inducible factor pathway. Adenosine monophosphate (AMP)—activated protein kinase (AMPK), as a central metabolic sensor sensitive to the AMP/ATP ratio, can be activated by various types 81 of cellular stress(Aslam & Ladilov, 2022; Li et al., 2013). AMPK is involved in the regulation 82 the fate of stem cell, and serves as a key regulator of cell differentiation (Liu et al., 2021; Sun et 83 al., 2017; Yang et al., 2016) Previous reports showed that AMPK was required for immune cell 84 differentiation and adipogenic differentiation (Rambold & Pearce, 2018; Son et al., 2019; Yang 85 et al., 2016). AMPK activation established a metabolic barrier to impede the cellular 86 reprogramming into iPSCs(Vazquez-Martin et al., 2013; Vazquez-Martin et al., 2012). AMPK 87 influenced osteogenic differentiation of human dental pulp mesenchymal stem cells (Pantovic et 88 al., 2013). The study also detected the change of AMPK when OXPHOS was intervened.



Metabolism is an important regulator for stem cells and fate decision. Anaerobic glycolysis
may be an adaption to the low oxygen niche for stem cells and maintain their stemness. The shift
from glycolysis to OXPHOS is required for stem cells to differentiate. The reverse transition
from OXPHOS to glycolysis is required for the induction of pluripotency from somatic
cells(Zhang ,Menzies & Auwerx, 2018). Metabolism regulating of stem cells would be a
potential target for tissue engineer or regeneration medicine.

In all, the aim of the study is to study the change of mitochondrial OXPHOS and glycolysis during hDPSCs differentiation and its possible regulating factors.

Materials & Methods

Reagents and antibodies

Hexokinase (HK), pyruvate kinase (PK), lactate dehydrogenase (LDH), and alkaline phosphatase(ALP) enzyme activity kits were obtained from Jiancheng (Nanjing, China). MitoTracker Red CMXRos (M7512) and Alexa Fluor 488 Phalloidin (A12379) were obtained from Invitrogen (Carlsbad, CA, USA). XF Cell Energy Phenotype Test Kit, XFp extracellular flux analyzer, XFp culture microplates and bicarbonate-free DMEM were obtained from Seahorse Bioscience—Agilent Technologies (North Billerica, MA, USA). Hoechst 33342 (B2261), Carbonyl cyanide 3-chlorophenylhydrazone (CCCP, C2759) was purchased from Sigma-Aldrich (St Louis, MO, USA). The following antibodies were used: AMPK (1:1000, Abcam, ab80039), p-AMPK (1:1000, Abcam, ab133448), anti-GAPDH (1:1000, Abcam, ab9485), and HRP-conjugated secondary antibodies (Bio-Rad Laboratories, Hercules, CA, USA),



113

114

115

116

117

118

119

120

121

122

123

124

125

127

128

129

130

131

Alexa Fluor 594-conjugated secondary antibody (Invitrogen). 111

Cell culture and treatment

After receiving the informed consents from patients and their parents, human dental pulp tissues were collected from caries- and periodontitis-free third molars extracted for orthodontic purposes from healthy donors (n=5). The experimental procedures were conducted according to the Declaration of Helsinki. The protocol was approved by the Institutional Ethics Committee of West China Hospital of Stomatology (WCHSIRB-D-2017-052; WCHSIRB-D-2020-075-R1). Briefly, after washing with sterile PBS, dental pulp tissues were cut into fragments, digested with 3 mg/ml type I collagenase at 37°C for 30 min, placed on 25 cm² culture dishes, and maintained in standard medium which was low glucose Dulbecco modified Eagle medium (DMEM) containing 10% fetal bovine serum (FBS) (Gibco, Carlsbad, CA, USA), penicillin (100 units/mL), and streptomycin (100 mg/mL) at 37°C in 5% CO₂. Cells were sub-cultured when cells reached 80% confluence, and used at passage 3 in this study. Differentiation was induced by osteogenic mineralized medium (MM), which was the standard medium supplemented with 10 mM b-glycerophosphate, 50 mg/mL ascorbic acid and 100 nM dexamethasone (Sigma-Aldrich, St. Louis, MO, USA). All media were replaced every 2 days. Flow cytometry was used to 126 analyze cell surface marker expressions and the cells were also incubated in MM for further alizarin red staining (Supplementary Fig. S1). To inhibit OXPHOS activity, cells were cultured in a mineralized medium with a mitochondrial uncoupler, carbonyl cyanide 3-chlorophenylhydrazone (CCCP, 2μ M). In some experiments, cells were treated with 5-aminoimidazole-4-carboxamideribonucleotide (AICAR,



132 $500\mu M$)

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

Enzyme activities assay

Cells were collected and ultrasonically decomposed on ice. After centrifugation, the supernatant was to measure the activities of ALP, HK, PK and LDH with corresponding enzyme activity kits. Enzyme activities were calculated from optical density value and total protein.

Cell energy phenotype analysis

Our previous results showed that hDPSCs initiated to differentiate on day 3 induced in MM. At this critical point, cells possessed increased glycolytic and mitochondrial function. With the further differentiation, extracellular flux rates were measured in this study. Cells (4×10^3) cells/well) were plated on XFp cell culture microplates and divided into experimental group and control group. The experimental group was cultured in MM, while the control was grown in standard medium at 37 °C and 5% CO₂. On day 1, 3, 5 and 7, the extracellular acidification rate (ECAR) as an indicator of the glycolysis potential, and the mitochondrial oxygen consumption rate (OCR) as parameter of mitochondrial respiration were analyzed on a Seahorse XFp Bioanalyzer, using the Seahorse XF Cell Energy Phenotype Test Kit according to the manufacturer's instructions. One hour before beginning the measurement, cells were switched to Seahorse bicarbonate-free DMEM, adjusted to pH 7.4 with NaOH, were subsequently incubated in a non-CO₂ incubator at 37°C for 1 h. Then, OCR and ECAR tests were performed under baseline and stressed conditions, after Oligomycin (1 µM), an ATP synthase inhibitor, and CCCP, 2µM, a mitochondrial uncoupling agent, were injected into the microplates. Extracellular flux rates were analyzed by using Seahorse XF software after normalized to the total protein units,



and the ratio of OCR/ECAR were calculated.

Real-time quantitative PCR (qPCR)

Total RNA was extracted from hDPSCs treated in different conditions using TRIzol 155 (Invitrogen) and cDNA was reverse transcribed with PrimeScript RT reagent Kit (TakaKa, 156 Osaka, Japan). A negative control without the reverse transcriptase was also implemented. Real-157 time PCR was performed using a SYBR Green PCR kit (Takara) and the Roche LightCycler480 158 Real-Time PCR System (Roche, Basel, Switzerland). GAPDH was used as the internal reference 159 gene, and levels were relativized to the control group. The sequences of specific primers were: 160 5'-5'-AAGGAGAAAGAGGAGCCAAAGG-3'(forward) and 161 5'-AGCACCAGGGAAACCAGTCATAC-3'(reverse) for *COL-1*; 162 5'-CAGATGAAGTGGGAGTGCTTGT-3'(forward) 163 and 164 CTGATGTGGAGTATGAGAGTGACG-3'(reverse) for ALP; 5'-ATGGCGTTTCTCCGAAGCAT-3'(forward) 5'-TCCGCCCTATAAGCATCTTGAand 165 5'-AGGAACACGGAGTGACCCAA-3'(forward) 3'(reverse) for TFAM; 5'-166 TATGCTCGGTGTAAGTAGCCA-3'(reverse) 5'for NRF1; 167 CTTTGGTATCGTGGAAGGACTC-3'(forward) and5'-GTAGAGGCAGGGATGATGTTCT-3' 168 (reverse) for *GAPDH*. 169

Fluorescence microscopy

170

171 Cells were seeded on the coverslips at 1×10⁴ cells/well, and incubated with MitoTracker 172 Red CMXRos (MRC) in the FBS-free DMEM at a concentration of 100 nM at 37°C for 30 min. 173 After washing, cells were fixed in 4% paraformaldehyde for 10 min, and permeabilized with



174 0.05% Triton X-100 in PBS for 10 min at room temperature. Then, cells were incubated with
175 Alexa Fluor 488 Phalloidin to show the cytoskeleton for 1 hour at room temperature. Nuclei
176 were stained with Hoechst 33342 for 5 min.

For the immunofluorescence experiment, cells were first fixed in 4%paraformaldehyde for 10 min, and were permeabilized with 0.05% Triton X-100 in PBS for 10 min. After washing, cells were blocked with 5% BSA in PBS for 30 min and incubated with primary antibody against p-AMPK overnight at 4°C. Following further washing, cells were incubated with Alexa Fluor 594-conjugated secondary antibody. Nuclei were also visualized by Hoechst 33342 for 5 min. Fluorescence signals were obtained using an Olympus microscope (Olympus, Munster, Germany).

Immunoblotting

Total proteins from hDPSCs were extracted using total protein extraction kit (KeyGEN BioTECh, Nanjing, China) according to the manufacturer's recommendation. SDS/PAGE was performed on Bio-Tris 5-10% gradient polyacrylamide gels acrylamide. Proteins were transferred to PVDF membranes (Bio-Rad), and membranes were blocked with 5% BSA in TRIS-buffered saline, 0.1% Tween (TBS-T) for 60 min at room temperature. The membranes were incubated overnight with appropriate primary antibodies. Detection was subsequently performed with HRP-conjugated secondary antibody for 1 hour at room temperature, and the membranes were scanned using the GelDoc XR+ System (Bio-Rad Laboratories, Hercules, CA, USA). The density of Western blot band signals was monitored using Quantity One software.

Statistical analysis



Every experiment was independently replicated a minimum of three times, ANOVA and Student's t-test were performed to determine statistical significance between control group and experimental group using SPSS version 17.0 software. P-values were considered to be significant when p < 0.05 (*p < 0.05).

Results

During osteogenic differentiation, glycolysis function declined after an initial slight increase

In order to assess the status of osteogenic differentiation, the mRNA expression of cell differentiation makers (ALP and COL-I) and ALP activity were detected on day 0, 1, 3, 5, and 7 cultured in control and mineralized medium (MM). By q-PCR, the expression of ALP mRNA increased at 3 days after differentiation induction (p=0.005), and COL-I started to upregulate 5 days after treatment (p=0.044), compared with the control (Fig. 1A and B). To further understand the hDPSC differentiation, ALP activity was also monitored. As is reported in Fig. 1C, the increase of ALP activity occurred from day 3, and kept on rising during the process of differentiation (p=0.000).

The activity of several important enzymes in glycolysis was measured. The activities of HK (p=0.023), PK (p=0.013) and LDH (p=0.005) had a slight increase at day 3 when hDPSC started to differentiate, then gradually decreased to a much lower level than that of control at day 7 during differentiation of hDPSC (Fig. 1D, E, and F).

Mitochondrial OXPHOS predominated in energy production of the differentiating hDPSCs.

The data showed that mitochondrial transcription factor A (TFAM, controlling mtDNA



expression and mitochondrial biogenesis) expressions were elevated at day 3, 5 and 7 compared to the control (Fig. 2A, p=0.016, p=0.002, p=0.001). The expressions of TFAM in MM were also gradually upregulated since day 3 (Fig. 2A, p=0.000). However, the expression level of Nuclear Respiratory Factors 1 (NRFI, controlling nuclear encoded respiratory chain components expression) was unchanged compared to control cells (Fig. 2B).

Later, the mitochondrial OCR(mitochondrial respiration indicator) and ECAR(glycolysis indicator) during hDPSCs differentiation in baseline and stressed condition were analyzed by Seahorse XF Cell Energy Phenotype Test Kit. Two stressor compounds, oligomycin that causes an increase of glycolysis and carbonyl cyanide p-trifluoro methoxyphenylhydrazone that drives OCR rates higher, were used as to induce a stressed condition. Under the basal condition, consistent with upregulated TFAM mRNA expression, OCR of differentiating cell was greater than that of control at day 3 (Fig. 2C), but the ratio of OCR/ECAR had no difference (Fig. 2D). At the days 5 and 7, ECAR was down-regulated, and OCR was still rising. Moreover, the cells in MM exhibited higher OCR/ECAR compared to the control at day 5 and 7(Fig. 2D, p=0.004, p=0.001). On the metabolic stressed condition, a similar pattern of specific dynamic changes in mitochondrial OXPHOS and glycolysis could be observed (Fig. 2E).

Mitochondrial uncoupling interfered with hDPSCs' differentiation

A mitochondrial uncoupler, CCCP, was used to attenuate mitochondrial respiration(Kane et al., 2018). The cells were cultured in osteogenic mineralized medium with or without CCCP. Upon differentiation, both the basal and stressed mitochondrial OCR depressed after CCCP treatment, compared to cells cultured in MM (Fig. 3A). The ECAR of cells growing in presence

spaces should be left



of CCCP showed a decrease compared to the control at day 3 in baseline conditon (*p*=0.002), while there was no difference in other days (Fig. 3B). With the MitoTracker Red CMXRos staining, the mitochondrial morphology showed to be punctate, and failed to form functional network at the initial stage of differentiation (yellow arrow, Fig. 3C). Importantly, this suppression of mitochondrial respiration attenuated the differentiation of hDPSCs, manifested as the inhibited ALP activity and the restrained *ALP* and *COL-1* mRNA expression, compared to cells growing in absence of CCCP (Fig. 3D).

The activation of AMPK impaired OXPHOS-driven differentiation

The activation of AMPK was detected in cells cultured in MM with or without CCCP (2μ M) for 1-7 days. The ratio of AMPK phosphorylation/AMPK protein were increased at day 3,5,7 by using CCCP (Fig. 4A). Would AMPK influence hDPSCs differentiation? As shown on Fig. 4B, the activation of AMPK was induced significantly by AICAR treatment. Crucially, consistent with CCCP treatment, AICAR suppressed hDPSCs differentiation (Fig. 4C). Furthermore, the mRNA expression of *TFAM* was suppressed, and the mitochondrial fragmentation was observed, compared to the control since day 3 after treatment of AICAR (Fig. 4D and 4E). Collectively, these results indicated that the activation of AMPK contributed to hDPSCs differential inhibition and impaired OXPHOS.

Discussion

In this study, it was demonstrated that the dynamic shift of glycolysis and mitochondrial OXPHOS took place during differentiation of hDPSCs. Attenuating mitochondrial respiration

Conditon- condition



261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

with uncoupler CCCP suppressed hDPSCs differentiation and activated the central metabolicsensor, AMPK.

Metabolic shift accompanied hDPSCs' osteogenic differentiation

Recent studies have displayed that stem cells depend mostly on anaerobic glycolysis for ATP supply, while the differentiating cells depend on mitochondrial aerobic respiration. The metabolic shift and mitochondrial resetting into mature bioenergetic states are considered a hallmark of stem cell differentiation (Wanet et al., 2015). When hDPSCs differentiated, cells presented distinct energy phenotypes. At day 3, the level of ECAR showed that glycolysis increased rather than decreased. Studies have shown that glycolysis supplies rapid energy generation and substrates for the biosynthesis. Increased glycolysis can provide a quick production of ATP and sufficient substrates for the biosynthesis, which meet the anabolic demands of initial differentiation (Palmer et al., 2015). This implies that differentiation requires the supply of enough energy and substrates. The joint effects of glycolysis and mitochondrial OXPHOS may support the tremendous energy demand on day 3 of differention. Simutanously, higher level of OCR and increased ratio of OCR/ECAR herald a shift toward a characteristic of mitochondrial OXPHOS. The results showed that mitochondrial OXPHOS predominated in energy production of the differentiating hDPSCs at day 5 and day 7, when metabolic shift of hDPSCs were observed. The finding was similar to other reported stem cells (Coelho et al., 2022; Folmes et al., 2011; Prigione et al., 2010). Some studies suggested that mitochondrial content, such as mitochondrial DNA (mtDNA)

copy number, mitochondrial mass and biogenesis, tightly adapted to the change of cell



metabolism during pathological or physical activity(Clark & Parikh, 2020; Procaccio et al., 2014). In order to assay the mitochondrial content-related modulating factors during hDPSCs differentiation, the mRNA expression of *TFAM* and *NRF1* were detected in the study(Kang,Chu & Kaufman, 2018; Quan et al., 2020) As hDPSCs differentiation went onto day 5 and 7, upregulated *TFAM* expression were observed, which might induced increased mtDNA content and increased mitochondrial biogenesis(Tsutsui, 2020; Vega-Naredo et al., 2014). Nevertheless, it seemed that *NRF1* was not involved in the process of hDPSCs differentiation. The reason might be related to the difference of cell types.

AMPK participated in differential inhibition when mitochondrial activity attenuated

Metabolic shift plays an essential role in regulating cellular function, particularly in stem cell self-renewal, pluripotency, and plasticity(Andre et al., 2019). Mitochondrial biogenesis and metabolic shift toward OXPHOS are deemed as early events in differentiation process of multiple stem cells. Furthermore, Inoue S et al. suggested that mitochondrial respiration was more important in mouse hematopoietic stem cell commitment and differentiation than in reaching the final phase of cell differentiation(Inoue et al., 2010). Consistent with previous reports, our data showed that inhibiting mitochondrial function by CCCP from the beginning prevented differentiation of hDPSCs. It turned out that mitochondrial OXPHOS was required for differentiation. CCCP-treated hDPSCs exhibited an apparent fragmentation of mitochondrial tubules, likely due to a block in mitochondrial fusion.

The energy sensor AMPK was activated by the decreased intracellular ATP due to impaired OXPHOS. It was speculated that AMPK may be the direct regulator of the mitochondrial fission



and fusion machinery to mediate subsequent events. As an intracellular energy sensor, the vital role of AMPK was to restore the energy balance. Once activated, AMPK induces catabolic pathways to produce energy and prevents anabolic pathways, such as lipid and protein synthesis, to save energy(Wanet et al., 2015). On the other hand, considerable evidence shows that AMPK is involved in regulating cell fate, including controlling stem cell pluripotency and differentiation(Afinanisa ,Cho & Seong, 2021; Fernandez-Veledo et al., 2013; Liu et al., 2021). In the study, CCCP-treated hDPSCs exhibited an elevated level of p-AMPK, showing that the energy sensor AMPK was involved in the process of metabolism that influenced differentiation. AMPK activation was a potential modulator when the differentiation of hDPSCs had switched on.

To further verify the effect of AMPK activation. AICAR, a direct AMPK activator was applied. AICAR resulted in mitochondrial fragmentation and differential inhibition, which were similar to that induced by CCCP. For cell differentiation, numerous anabolic pathways, including the biosynthesis of proteins and lipids, are required to support specific function of differentiated cells. The mitochondrial biogenesis and stimulation of metabolism are also needed to produce adequate energy for the processes. CCCP-induced or AICAR-induced AMPK activation may serve as a regulator for the cells to quit differentiation from impaired OXPHOS, which might be useful for keeping stemness or to maintain homeostasis. Mitochondrial dysfunction as a regulator may be a viable therapeutic target for stem-cell-based therapies and interventions for cognitive defects(Khacho, Harris & Slack, 2019).

However, the detailed mechanism between mitochondria and cell differentiation requires



342

321	further investigation. Whether metabolic shift is prerequisite for cell differentiation and whether
322	mitochondrial OPHOXS can regulate cell differentiation in vivo remains to be further studied.
323	
324	Conclusion
325	It was demonstrated that increased mitochondrial function was indispensable for hDPSCs
326	differentiation, while the glycolysis gradually decreased in the stage of energy supply. It was also
327	unveiled that mitochondrial uncoupler CCCP depressed mitochondrial OXPHOS and inhibited
328	hDPSCs differentiation. Activation of AMPK also interfere with the mitochondrial morphology,
329	mitochondrial OXPHOS and cell differentiation. The findings helped to reveal the relationship
330	among glycolysis, mitochondrial OXPHOS and cell differentiation. CCCP or AMPK activation
331	may be a potential regulator to quit differentiation from impaired OXPHOS or to maintain
332	homeostasis.
333	
334	Acknowledgments
335	Not applicable.
336	
337	References
338	Afinanisa Q, Cho MK, Seong HA. 2021. AMPK localization: A key to differential energy
339	regulation. International Journal of Molecular Sciences 22(20) DOI
340	10.3390/ijms222010921
341	Andre E, De Pauw A, Verdoy R, Brusa D, Bouzin C, Timmermans A, Bertrand L,

Balligand JL. 2019. Changes of metabolic phenotype of cardiac progenitor cells during



343	differentiation. Neutral effect of stimulation of AMP-activated protein kinase. Stem Cens
344	and Development 28(22) :1498-1513 <u>DOI 10.1089/scd.2019.0129</u>
345	Aslam M, Ladilov Y. 2022. Emerging role of cAMP/AMPK signaling. Cells 11(2) DOI
346	10.3390/cells11020308
347	Clark AJ, Parikh SM. 2020. Mitochondrial metabolism in acute kidney injury. Seminars in
348	Nephrology 40(2) :101-113 <u>DOI 10.1016/j.semnephrol.2020.01.002</u>
349	Coelho P, Fao L, Mota S, Rego AC. 2022. Mitochondrial function and dynamics in neural stem
350	cells and neurogenesis: Implications for neurodegenerative diseases. Ageing Research
351	Reviews 80 :101667 DOI 10.1016/j.arr.2022.101667
352	Fernandez-Veledo S, Vazquez-Carballo A, Vila-Bedmar R, Ceperuelo-Mallafre V,
353	Vendrell J. 2013. Role of energy- and nutrient-sensing kinases AMP-activated protein
354	kinase (AMPK) and mammalian target of rapamycin (mTOR) in adipocyte differentiation.
355	IUBMB Life 65(7):572-583 DOI 10.1002/iub.1170
356	Folmes CD, Nelson TJ, Martinez-Fernandez A, Arrell DK, Lindor JZ, Dzeja PP, Ikeda Y,
357	Perez-Terzic C, Terzic A. 2011. Somatic oxidative bioenergetics transitions into
358	pluripotency-dependent glycolysis to facilitate nuclear reprogramming. Cell Metabolism
359	14(2):264-271 DOI 10.1016/j.cmet.2011.06.011
360	Inoue S, Noda S, Kashima K, Nakada K, Hayashi J, Miyoshi H. 2010. Mitochondrial
361	respiration defects modulate differentiation but not proliferation of hematopoietic stem
362	and progenitor cells. FEBS Letters 584(15):3402-3409 DOI
363	10.1016/j.febslet.2010.06.036
364	Kane MS, Paris A, Codron P, Cassereau J, Procaccio V, Lenaers G, Reynier P, Chevrollier
365	A. 2018. Current mechanistic insights into the CCCP-induced cell survival response.
366	Biochemical Pharmacology 148 :100-110 DOI 10.1016/j.bcp.2017.12.018
367	Kang I, Chu CT, Kaufman BA. 2018. The mitochondrial transcription factor TFAM in
368	neurodegeneration: Emerging evidence and mechanisms. FEBS Letters 592(5):793-811
369	DOI 10.1002/1873-3468.12989



370	Khacho M, Harris R, Slack RS. 2019. Mitochondria as central regulators of neural stem cell
371	fate and cognitive function. Nature Reviews: Neuroscience 20(1):34-48 DOI
372	<u>10.1038/s41583-018-0091-3</u>
373	Khacho M, Slack RS. 2017. Mitochondrial activity in the regulation of stem cell self-renewal
374	and differentiation. Current Opinion in Cell Biology 49:1-8 DOI
375	10.1016/j.ceb.2017.11.003
376	Khacho M, Slack RS. 2018. Mitochondrial and reactive oxygen species signaling coordinate
377	stem cell fate decisions and life long maintenance. Antioxidants and Redox Signaling
378	28(11):1090-1101 DOI 10.1089/ars.2017.7228
379	Li YY, Yu LF, Zhang LN, Qiu BY, Su MB, Wu F, Chen DK, Pang T, Gu M, Zhang W, Ma
380	WP, Jiang HW, Li JY, Nan FJ, Li J. 2013. Novel small-molecule AMPK activator
381	orally exerts beneficial effects on diabetic db/db mice. Toxicology and Applied
382	Pharmacology 273(2) :325-334 <u>DOI 10.1016/j.taap.2013.09.006</u>
383	Liu Y, Yamane J, Tanaka A, Fujibuchi W, Yamashita JK. 2021. AMPK activation reverts
384	mouse epiblast stem cells to naive state. iScience 24(7):102783 DOI
385	10.1016/j.isci.2021.102783
386	Ly CH, Lynch GS, Ryall JG. 2020. A metabolic roadmap for somatic stem cell fate. Cell
387	Metabolism 31(6):1052-1067 DOI 10.1016/j.cmet.2020.04.022
388	Morganti C, Cabezas-Wallscheid N, Ito K. 2022. Metabolic regulation of hematopoietic stem
389	cells. <i>Hemasphere</i> 6(7) :e740 <u>DOI 10.1097/HS9.0000000000000740</u>
390	Palmer CS, Ostrowski M, Balderson B, Christian N, Crowe SM. 2015. Glucose metabolism
391	regulates T cell activation, differentiation, and functions. Frontiers in Immunology 6:1
392	DOI 10.3389/fimmu.2015.00001
393	Pantovic A, Krstic A, Janjetovic K, Kocic J, Harhaji-Trajkovic L, Bugarski D, Trajkovic V.
394	2013. Coordinated time-dependent modulation of AMPK/akt/mTOR signaling and
395	autophagy controls osteogenic differentiation of human mesenchymal stem cells. Bone
396	52(1) :524-531 DOI 10.1016/j.bone.2012.10.024



397	rigione A, rauler B, Lurz K, Lenrach H, Aujaye J. 2010. The senescence-related
398	mitochondrial/oxidative stress pathway is repressed in human induced pluripotent stem
399	cells. Stem Cells 28(4):721-733 DOI 10.1002/stem.404
400	Procaccio V, Bris C, Chao de la Barca JM, Oca F, Chevrollier A, Amati-Bonneau P,
401	Bonneau D, Reynier P. 2014. Perspectives of drug-based neuroprotection targeting
402	mitochondria. Revue Neurologique 170(5):390-400 DOI 10.1016/j.neurol.2014.03.005
403	Quan Y, Xin Y, Tian G, Zhou J, Liu X. 2020. Mitochondrial ROS-modulated mtDNA: A
404	potential target for cardiac aging. Oxidative Medicine and Cellular Longevity
405	2020 :9423593 <u>DOI 10.1155/2020/9423593</u>
406	Rambold AS, Pearce EL. 2018. Mitochondrial dynamics at the interface of immune cell
407	metabolism and function. Trends in Immunology 39(1):6-18 DOI
408	10.1016/j.it.2017.08.006
409	Rossmann MP, Dubois SM, Agarwal S, Zon LI. 2021. Mitochondrial function in development
410	and disease. Disease Models & Mechanisms 14(6) DOI 10.1242/dmm.048912
411	Son J, Cho YW, Woo YJ, Baek YA, Kim EJ, Cho Y, Kim JY, Kim BS, Song JJ, Ha SJ.
412	2019. Metabolic reprogramming by the excessive AMPK activation exacerbates antigen-
413	specific memory CD8(+) T cell differentiation after acute lymphocytic choriomeningitis
414	virus infection. Immune Network 19(2):e11 DOI 10.4110/in.2019.19.e11
415	Sui B, Wu D, Xiang L, Fu Y, Kou X, Shi S. 2020. Dental pulp stem cells: From discovery to
416	clinical application. <i>Journal of Endodontics</i> 46(9S) :S46-S55 <u>DOI</u>
417	10.1016/j.joen.2020.06.027
418	Sun X, Yang Q, Rogers CJ, Du M, Zhu MJ. 2017. AMPK improves gut epithelial
419	differentiation and barrier function via regulating Cdx2 expression. Cell Death and
420	Differentiation 24(5) :819-831 <u>DOI 10.1038/cdd.2017.14</u>
421	Tsutsui TW. 2020. Dental pulp stem cells: Advances to applications. Stem Cells and Cloning:
422	Advances and Applications 13:33-42 DOI 10.2147/SCCAA.S166759
423	Vazquez-Martin A, Corominas-Faja B, Cufi S, Vellon L, Oliveras-Ferraros C, Menendez




424	OJ, Joven J, Lupu K, Wienendez JA. 2013. The initochondrial $H(\tau)$ -ATP symmase and
425	the lipogenic switch: New core components of metabolic reprogramming in induced
426	pluripotent stem (iPS) cells. Cell Cycle 12(2):207-218 DOI 10.4161/cc.23352
427	Vazquez-Martin A, Vellon L, Quiros PM, Cufi S, Ruiz de Galarreta E, Oliveras-Ferraros C,
428	Martin AG, Martin-Castillo B, Lopez-Otin C, Menendez JA. 2012. Activation of
429	AMP-activated protein kinase (AMPK) provides a metabolic barrier to reprogramming
430	somatic cells into stem cells. <i>Cell Cycle</i> 11(5) :974-989 <u>DOI 10.4161/cc.11.5.19450</u>
431	Vega-Naredo I, Loureiro R, Mesquita KA, Barbosa IA, Tavares LC, Branco AF, Erickson
432	JR, Holy J, Perkins EL, Carvalho RA, Oliveira PJ. 2014. Mitochondrial metabolism
433	directs stemness and differentiation in P19 embryonal carcinoma stem cells. Cell Death
434	and Differentiation 21(10) :1560-1574 DOI 10.1038/cdd.2014.66
435	Wanet A, Arnould T, Najimi M, Renard P. 2015. Connecting mitochondria, metabolism, and
436	stem cell fate. Stem Cells and Development 24(17):1957-1971 DOI
437	10.1089/scd.2015.0117
438	Wang L, Cheng L, Wang H, Pan H, Yang H, Shao M, Hu T. 2016. Glycometabolic
439	reprogramming associated with the initiation of human dental pulp stem cell
440	differentiation. Cell Biology International 40(3):308-317 DOI 10.1002/cbin.10568
441	Yang Q, Liang X, Sun X, Zhang L, Fu X, Rogers CJ, Berim A, Zhang S, Wang S, Wang B,
442	Foretz M, Viollet B, Gang DR, Rodgers BD, Zhu MJ, Du M. 2016. AMPK/alpha-
443	ketoglutarate axis dynamically mediates DNA demethylation in the Prdm16 promoter and
444	brown adipogenesis. <i>Cell Metabolism</i> 24(4) :542-554 <u>DOI 10.1016/j.cmet.2016.08.010</u>
445	Zhang H, Menzies KJ, Auwerx J. 2018. The role of mitochondria in stem cell fate and aging.
446	Development 145(8) DOI 10.1242/dev.143420
147	



Glycolysis increased first and then decreased during hDPSC differentiation.


Cells were cultured in control and mineralized medium for 0, 1, 3, 5, and 7 days, respectively. (A) mRNA expression of ALP was upregulated from day 3, (B) COL-1 was upregulated from day 5. (C) HK, a key glycolytic enzyme, increased at day 3 when cells initiated to differentiate, and then decreased from day 5. (D, E) The same trend was detected in PK and LDH. Data were mean \pm SD, *p<0.05, **p<0.01, ***p<0.001.



Cell energy phenotype shifted to mitochondrial OXPHOS as hDPSCs differentiated.

(A) TFAM expression, which is involved in mitochondrial biogenesis, upregulated from the initial stage of hDPSCs differentiation, (B) while NRF1 was not greatly changed upon differentiation. (C) On day 1, 3, 5, and 7, the OCR and ECAR of cells growing in basal condition were firstly measured, and then those of stressed condition in the absence of Oligomycin and FCCP were assayed to show the max metabolic potential. The ECAR was on the horizontal axis, and the OCR was on the vertical axis. The hollow rhombus was for basal condition, while the solid was for stressed condition. (D) For the differentiating cells, the ratio of basal OCR/ECAR, which was calculated to show the major energy phenotype, increased from day 5. (E) The stressed OCR/ECAR ratio had the similar tendency. Data were mean \pm SD, *p<0.05, **p<0.01, ***p<0.001.








As hDPSC differentiation, cell energy phenotype shifted to mitochondrial OXPHOS.

Cells were cultured in the MM with or without CCCP (2 μ M) for 1-7 days. (A) The change of OCR was monitored over time. With the CCCP treatment, cells9 OCR was depressed, compared to the control. (B) Except for the day 3, the ECAR of CCCP group had no significant change. (C) At day 3, fluorescence was used to detect the morphology of mitochondria: mitochondria - MitoTracker Red CMXRos (MRC), red; F-actin-Phalloidine, green; nucleus - Hoechst 33342, blue. The local magnification showed CCCP inhibited the form of mitochondria functional network. (D) The osteogenic differential makers, the mRNA expression of ALP and COL-1 and the activity of ALP were suppressed by CCCP. Scale bars:20 μ m. Data were mean \pm SD, *p<0.05, **p<0.01, ***p<0.001.



The activation of AMPK contributed to differential inhibition.

(A) mmunoblot analysis for AMPK and phosphorylated AMPK in cell lysates from hDPSCs treated with CCCP (2 μ M) or MM only for 1, 3, 5, and 7 days. GAPDH served as the standard. (B) Cells were treated with AMPK agonist AICAR (500 μ M) for 3 days, the p-AMPK was observed using immunofluorescence: p-AMPK, red; nucleus, blue. Scale bar: 50 μ M. (C) The ALP, COL-1 expression and ALP activity were detected to unveil the effect of AICAR in cell differentiation. (D) qPCR analysis of mitochondrial factor TFAM in AICAR-treated (500 μ M) cells. (E) Mitochondrial morphology in AICAR-treated and control cells were evaluated by fluorescence staining. Mitochondria-MitoTracker Red CMXRos (MRC), red; F-actin -Phalloidine, green; nucleus - Hoechst 33342, blue. Scale bar: 20 μ M. Data were mean±SD, * p <0.05, *** p <0.01, **** p <0.001.



