# Risk factors of self-reported physical child abuse during the COVID-19 pandemic in Japan: Work-related changes in men and fear of COVID-19 in women (#77873)

First revision

### Guidance from your Editor

Please submit by 22 Mar 2023 for the benefit of the authors .



### **Structure and Criteria**

Please read the 'Structure and Criteria' page for general guidance.



### **Custom checks**

Make sure you include the custom checks shown below, in your review.



### Raw data check

Review the raw data.



### Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

#### **Files**

Download and review all files from the <u>materials page</u>.

- 1 Tracked changes manuscript(s)
- 1 Rebuttal letter(s)
- 1 Figure file(s)
- 3 Table file(s)
- 1 Raw data file(s)

### Custom checks

### Human participant/human tissue checks

- Have you checked the authors ethical approval statement?
- Does the study meet our <u>article requirements</u>?
- Has identifiable info been removed from all files?
- Were the experiments necessary and ethical?

## Structure and Criteria



### Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING
- 2. EXPERIMENTAL DESIGN
- 3. VALIDITY OF THE FINDINGS
- 4. General comments
- 5. Confidential notes to the editor
- You can also annotate this PDF and upload it as part of your review

When ready submit online.

### **Editorial Criteria**

Use these criteria points to structure your review. The full detailed editorial criteria is on your guidance page.

#### **BASIC REPORTING**

- Clear, unambiguous, professional English language used throughout.
- Intro & background to show context.
  Literature well referenced & relevant.
- Structure conforms to <u>PeerJ standards</u>, discipline norm, or improved for clarity.
- Figures are relevant, high quality, well labelled & described.
- Raw data supplied (see <u>PeerJ policy</u>).

#### **EXPERIMENTAL DESIGN**

- Original primary research within Scope of the journal.
- Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
- Rigorous investigation performed to a high technical & ethical standard.
- Methods described with sufficient detail & information to replicate.

### **VALIDITY OF THE FINDINGS**

- Impact and novelty not assessed.

  Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
- All underlying data have been provided; they are robust, statistically sound, & controlled.



Conclusions are well stated, linked to original research question & limited to supporting results.



# Standout reviewing tips



The best reviewers use these techniques

Τ	p

# Support criticisms with evidence from the text or from other sources

### Give specific suggestions on how to improve the manuscript

### Comment on language and grammar issues

### Organize by importance of the issues, and number your points

# Please provide constructive criticism, and avoid personal opinions

Comment on strengths (as well as weaknesses) of the manuscript

### **Example**

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Your introduction needs more detail. I suggest that you improve the description at lines 57-86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

- 1. Your most important issue
- 2. The next most important item
- 3. ...
- 4. The least important points

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.



# Risk factors of self-reported physical child abuse during the COVID-19 pandemic in Japan: Work-related changes in men and fear of COVID-19 in women

Haruaki Naito Equal first author, 1, 2, Katsuya Nitta Equal first author, 1, Yasuhiro Kakiuchi Corresp. 1, 2

Corresponding Author: Yasuhiro Kakiuchi Email address: kakiuchi@yokohama-cu.ac.jp

Background: There is no global consensus on whether pandemic-related factors are related to child abuse. How the pandemic reinforces the risk factors of child abuse might depend largely on individuals' current and past lifestyles in each country. Some changes of lifestyles continue after the pandemic, and it is important to understand which factors is strongly associated with child abuse. We analyzed the pandemic-related characteristics of offenders and non-offenders of self-reported child physical abuse from Internet survey data in Japan and discussed how the pandemic affected physical child abuse and what caused the difference by gender.

Methods: We conducted a cross-sectional study on physical child abuse by caregivers based on the Internet survey from September to October 2021. We divided the participants who were living with their child aged less than 1 year to 14 years into offenders and non-offenders based on the answer to the question about physical child abuse. The population distribution of the sample was compared to that of caregivers under the same conditions in a large Japanese dataset. The association between their characteristics and physical child abuse was analyzed by univariate and multivariate analysis.

Results: The caregivers analyzed in the cohort had similar population distributions to those in the large Japanese dataset. As risk factors of male offenders, "work from home 4-7 days/week," "decreased work," "normal relationships with household members (compared to good relationships)," "COVID-19 infected, both themselves and household members, within a year," "unwillingness to receive COVID-19 vaccination because the license process of the vaccine is doubtful," "high levels of benevolent sexism," and "history of child abuse" were observed. As risk factors of female offenders, "bad relationships with household members (compared to good relationships)," "fear of COVID-19," "COVID-19 infected, both themselves and household members, within a year," "feelings of discrimination related to COVID-19 in the past two months," and "history of child verbal abuse" were observed.

Conclusions: Among male offenders, a significant relationship was observed regarding work-related changes, which may have been reinforced by the pandemic. Furthermore, the extent of the influence and fear of losing jobs caused by these changes may have varied according to the strength of gender roles and financial support in each country. Among female offenders, a significant relationship was observed regarding fear of infection itself, which is consistent with the findings of other studies. In terms of factors related to dissatisfaction with families, in some countries with prominent stereotyped gender roles, men are thought to experience difficulties adapting to work-related changes induced by crises, while women are thought to experience intense fear of the infection itself.

<sup>1</sup> Department of Forensic Medicine, School of Medicine, Kindai University, Sayama, Osaka, Japan

<sup>&</sup>lt;sup>2</sup> Department of Forensic Medicine, School of Medicine, Tokai University, Isehara, Kanagawa, Japan



- Risk factors of self-reported physical child abuse during the
- 2 COVID-19 pandemic in Japan: Work-related changes in men
- and fear of COVID-19 in women

4	Haruaki Naito <sup>#,1,2</sup> , Katsuya Nitta <sup>#,1</sup> , Yasuhiro Kakiuchi*, <sup>1,2</sup>
5	
6	<sup>1</sup> Department of Forensic Medicine, School of Medicine, Kindai University, Sayama, Osaka, Japan
7	
8	<sup>2</sup> Department of Forensic Medicine, School of Medicine, Tokai University, Isehara, Kanagawa,
9	Japan
10	
11	#Contributed equally
12	
13	*Corresponding author
14	E-mail: kakiuchi@yokohama-cu.ac.jp
15	
16	
17	
18	
19	
20	
21	
22	



### Abstract

24	Background: There is no global consensus on whether pandemic-related factors are related to
25	child abuse. How the pandemic reinforces the risk factors of child abuse might depend largely on
26	individuals' current and past lifestyles in each country. Some changes of lifestyles continue after
27	the pandemic, and it is important to understand which factors is strongly associated with child
28	abuse. We analyzed the pandemic-related characteristics of offenders and non-offenders of self-
29	reported child physical abuse from Internet survey data in Japan and discussed how the pandemic
30	affected physical child abuse and what caused the difference by gender.
31	Methods: We conducted a cross-sectional study on physical child abuse by caregivers based on
32	the Internet survey from September to October 2021. We divided the participants who were
33	living with their child aged less than 1 year to 14 years into offenders and non-offenders based
34	on the answer to the question about physical child abuse. The population distribution of the
35	sample was compared to that of caregivers under the same conditions in a large Japanese dataset.
36	The association between their characteristics and physical child abuse was analyzed by
37	univariate and multivariate analysis.
88	Results: The caregivers analyzed in the cohort had similar population distributions to those in the
39	large Japanese dataset. As risk factors of male offenders, "work from home 4-7 days/week,"
10	"decreased work," "normal relationships with household members (compared to good
11	relationships)," "COVID-19 infected, both themselves and household members, within a year,"
12	"unwillingness to receive COVID-19 vaccination because the license process of the vaccine is
13	doubtful," "high levels of benevolent sexism," and "history of child abuse" were observed. As
14	risk factors of female offenders, "bad relationships with household members (compared to good



45 relationships)," "fear of COVID-19," "COVID-19 infected, both themselves and household

46 members, within a year," "feelings of discrimination related to COVID-19 in the past two

47 months," and "history of child verbal abuse" were observed.

48 Conclusions: Among male offenders, a significant relationship was observed regarding work-

related changes, which may have been reinforced by the pandemic. Furthermore, the extent of

50 the influence and fear of losing jobs caused by these changes may have varied according to the

strength of gender roles and financial support in each country. Among female offenders, a

significant relationship was observed regarding fear of infection itself, which is consistent with

the findings of other studies. In terms of factors related to dissatisfaction with families, in some

countries with prominent stereotyped gender roles, men are thought to experience difficulties

adapting to work-related changes induced by crises, while women are thought to experience

56 intense fear of the infection itself.

57

58

61

64

65

49

51

52

53

54

55

### Background

59 The negative impacts of child abuse are maintained throughout the victims' lifespan, and result

60 in various mood disorders, anxiety disorders, suicidal behavior, and the intergenerational cycle

of child abuse (Chapman et al., 2004; Fujiwara & Kawakami, 2011; Stickley et al., 2020;

62 Horikawa et al., 2016; Berlin, Appleyard & Dodge, 2011). Many studies have concluded that

63 more attention to child abuse is necessary during the COVID-19 pandemic because some risk

factors such as mental stress, economic stress, social isolation, and a decrease in the chances of

detecting child abuse must be reinforced by the pandemic (Romanou & Belton, 2020; Pereda &

66 Díaz-Faes, 2020; Brown et al., 2020; Amin & Parveen, 2022).



-	$\neg$
n	-/

However, there is a fundamental question of whether the pandemic equally reinforces the risk factors of child abuse in every country. A systematic review that analyzed 12 articles reported that the number of areas where abuse increased and decreased from pre-COVID-19 pandemic was approximately the same (Rapp et al., 2021). The impact of the COVID-19 pandemic on child abuse might largely depend on each country's past and current conditions, including the extent of fear of COVID-19, the degree of telework penetration, and attitudes toward gender roles.

Hence, we used Internet survey data and performed statistical analysis by including various factors directly and indirectly related to the COVID-19 pandemic. There are several drawbacks to using official reports or self-reports of child abuse concerning the quality and number of child abuse cases (Gilbert et al., 2009). However, it is plausible to estimate that self-reports were closer to the true number of child abuse cases during the pandemic when many child protection agencies and schools—major resources of official reports of child abuse—reduced their activities. This study is the first in Japan to report that some pandemic-related characteristics in offenders were significantly associated with physical child abuse; notable differences by gender were also observed. To the best of our knowledge, no study in Japan has identified a significant relationship between physical child abuse and pandemic-related factors of offenders.

### Methods

### 88 Data collection

89 Data was obtained from the Japan COVID-19 and Society Internet Survey (JACSIS) 2021. The





90 survey focused on public health issues regarding COVID-19 pandemic and was conducted by an Internet research company (Rakuten Insight, Inc.), which had 2.3 million registered panelists. 91 Before the survey, the target number of responses was set in each group, stratified by age, sex, 92 and prefecture, based on the population distribution. The population distribution was calculated 93 using the Comprehensive Survey of Living Conditions (CSLC) in Japan. From September 27 to 94 95 October 5, 2021, questionnaires with pre-defined options were distributed through the Internet to the 33,081 respondents of JACSIS 2020 and JASTIS 2015-2020 aged 16-81 years, in seven 96 waves. From October 23 to 28, the target numbers were not reached in some stratified groups; 97 98 therefore, questionnaires were distributed to the new 26,138 panelists by random sampling, based on the difference between the target and collected numbers in each group. In the second 99 wave, all groups reached the target numbers and the survey was completed. Hence, 100 101 questionnaires were distributed to 59,219 panelists, and the respondents represented the Japanese population regarding age, gender, and residential prefecture using a simple random sampling 102 process (the Japan COVID-19 and Society Internet Survey, 2022). 103 104 Invalid responses were measured using three questions, Q18, Q26, and Q68, which were similar 105 106 to the exclusion criteria used in previous JACSIS studies (Yamada et al., 2021; Minoura et al., 2021). In Q18, "Please choose the second option from the bottom," people who did not choose 107 the correct option were excluded. In Q26 about 20 types of diseases, people who answered 108 109 "currently" to all nine types of diseases (hypertension, diabetes, asthma, atopic dermatitis, allergic rhinitis, myocardial infarction or angina, stroke, cancer, and chronic pain over three 110 111 months) were excluded. In Q68, regarding the use of alcohol and drugs, people who answered 112 "almost every day" or "sometimes" to all the options i.e., alcohol, sleeping pills, anxiolytic,



illegal opioids, organic solvent, dangerous drugs, cannabis, stimulant drugs, cocaine, and heroin, 113 were also excluded. 114 Outcome Variables 115 We focused on the question about child physical abuse: 116 "Have you used violence on your child in the past two months?" 117 118 The question was only asked to caregivers living with child aged less than 1 year to 14 years. Those who answered "Yes" were classified as an experimental group, and those who answered 119 "No" were classified as a control group. Those who answered "not sure" or "not want to answer" 120 were excluded from the analysis. 121 **Explanatory Variables** 122 The following sociodemographic characteristics of the subjects were obtained: "age (20-29, 30-123 39, 40-49, and 50-59 years)," "work from home (none, 1-3 days/month, 1-3 days/week, 4-7 124 days/week, and unemployed)," "change of job status within a year (did not change, work 125 decreased, leave of absence or furlough, resigned, and others)," "relationships with household 126 members (good, normal, and bad)," "benevolent sexism scale (<=24 and >24)," "experienced or 127 witnessed physical abuse by parents before age of 18 (no and yes)," and "experienced verbal 128 129 abuse by parents before the age of 18 (no and yes)." 130 131 The following pandemic-related characteristics of the subjects were also obtained: "fear of 132 COVID-19 (not strong and strong)," "infection with COVID-19 within a year (neither myself nor household member, only myself, only household member, and both myself and household 133 member)," "feelings of discrimination related to COVID-19 (never, before two months ago, and 134 135 in the past two months)," and "willingness to receive COVID-19 vaccination (yes, no because



the license process of the vaccine is doubtful, and no due to other reasons)."

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

136

Regarding "fear of COVID-19," those who answered "very applicable" to any one of the questions regarding physically or mentally serious reactions to simply considering COVID-19 were defined as having a strong fear of COVID-19. Among the all reasons of "unwillingness to receive COVID-19 vaccination," "the license process of the vaccine is doubtful" is the most skeptical reason. Other reasons indicate that people did not want to receive the vaccination because they were concerned with the side effects, doubted its effectiveness, had low risk of serious complications, were previously infected, or did not have time to get the vaccine. "Benevolent sexism scale" was set to see whether the attitudes toward gender role are related to child abuse. Glick et al. proposed that sexism is divided into hostile and benevolent one in modern times, and that benevolent individuals have a more negative impact on women's career advancement than hostile ones (Glick & Fiske, 1996; Dardenne, Dumont & Bollier, 2007; King et al., 2012; Jones et al., 2014). The original sexism scale by Glick and Fiske consists of 11 questions each for hostile and benevolent individuals. In 2018, to make the scale more practical to measure benevolent sexism in Japan, a new benevolent sexism scale with eight questions corresponding to question numbers 1, 4, 8, 14, 15, 16, 18, and 19 in the original scale, was developed (Morinaga et al., 2018). The JACSIS 2019 fully used the scale in Q38. Victims of dating violence or child abuse had reportedly higher scores for benevolent sexism (Cuadrado-Gordillo, Fernández-Antelo & Martín-Mora Parra, 2020; Vives-Cases et al., 2021; Melissa Abi Rached, Ahmed Hankir & Rashid Zaman, 2021). To our knowledge, common criteria for the benevolent sexism scale do not exist; we set 24 out of 48 as a cut-off value for the explanatory variable.



159	Statistical analysis
160	First, our sample was compared to that of caregivers in the CSLC data in 2019 under the same
161	conditions, to show the representativeness of the valid respondents. Second, Pearson's chi-square
162	$(\chi^2)$ test was performed to determine the significance of the overall variables. Third, a
163	multivariate logistic regression analysis was performed to estimate the strength of associated risk
164	factors to physical child abuse. At 5% significance level, these gender-stratified analyses were
165	conducted by EZR ver. 1.54.
166	Ethics approval
167	Ethical approval was obtained from the Research Ethics Committee of the Osaka International
168	Cancer Institute (approval no. 20084). Before answering the distributed questionnaire, in the first
169	page of the questionnaire, all the participants read the purpose of the study, how personal
170	information would be handled, and how to decline participation in the study. Those who
171	answered the questionnaire were considered to have consented to participate in the study.
172	
173	Results
174	Figure 1 shows the flowchart of participant selection. Of the 2,296 male caregivers with children
175	aged less than 1 year to 14 years, 134 answered "Yes," 2,034 answered "No," and 128 answered
176	"Unknown/Do not want to answer" to the question of physical child abuse over the past 2
177	months. Of the 2,329 female caregivers with children aged less than 1 year to 14 years, 187
178	answered "Yes," 2,038 answered "No," and 104 answered "Unknown/Do not want to answer" to
179	the same question.

PeerJ reviewing PDF | (2022:09:77873:1:0:NEW 27 Feb 2023)





182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

Table 1 shows the comparison between our sample and that of caregivers from the CSLC data under the same conditions, aged 20–59 years with children aged less than 1 year to 14 years. In our study cohort, of both mele and female caregivers, 15.6% answered "YES" or "No" to the question. Of both male and female caregivers, 0.7–0.9% answered "Unknown" or "Do not want to answer." In the CSLC data, the caregivers under the same conditions were 17.1-18.6% men aged 16–81 years. Regarding age, the study cohort caregivers had a similar distribution to that of the caregivers from CSLC, compared with the non-analyzed caregivers. Regardless of the responses, the respondents of JACSIS had a higher prevalence of International Standard Classification of Education (ISCED)  $\geq$ 6 (men: 67.2–69.3%, women: 43.6–44.2%) than those from the CSLC data (men: 36.7%, women: 21.1%). Table S1 shows the numbers of male offenders and non-offenders according to their characteristics and the results of Pearson's  $\chi^2$  test and multivariate logistic regression analysis. Among men, self-reported physical child abuse was significantly associated with "work from home 4-7 days/week (Odds Ratios [ORs]=1.81, P value [P]=0.028)," "decrease of work (ORs=1.66, P=0.025)," "normal relationships with household members (compared to good relationships, ORs=1.57, P=0.039)," "COVID-19 infected, both themselves and household members, within a year (ORs=4.10, P=0.007)," "unwillingness to receive COVID-19 vaccination because the license process of the vaccine is doubtful (ORs=2.85, P=0.006)," "high levels of benevolent sexism (ORs=1.65, P=0.018)," "history of physical child abuse (ORs=2.96, P<0.001)," and "history of verbal child abuse (ORs=2.24, P=0.004)." All VIF values were less than 1.5.

203





Table S2 shows the number of female offenders and non-offenders according to their characteristics and the results of Pearson's  $\chi^2$  test and multivariate logistic regression analysis. Among women, self-reported physical child abuse was significantly associated with "bad relationships with household members (compared to good relationships, ORs=3.51, P<0.001)," "strong fear of COVID-19 (ORs=1.44, P=0.028)," "COVID-19 infected, both themselves and household members, within a year (ORs=3.22, P=0.038)," "feelings of discrimination related to COVID-19 in the past two months (ORs=2.08, P=0.044)," and "history of verbal child abuse (ORs=2.66, P<0.001)." It was significantly and negatively associated with "aged 50-59 (compared to aged 20-29, ORs=0.29, P=0.029)" and "unwillingness to receive COVID-19 vaccination for other reasons than the license process of the vaccine is doubtful (ORs=0.55, P=0.037)." All VIF values were less than 2.2.

### **Discussion**

### The importance of self-reported child abuse

Compared to officially reported child abuse data, self-reported data may include a greater number of milder cases. However, society is busy with specific responses to address previously reported severe cases. The number of officially reported child abuse cases is increasing across developed countries, while the number of cases identified or suggested as child abuse has not changed significantly (Ministry of Health, Labour and Welfare, 2022; Department for Education, 2019; Children's Bureau, 2020; Australian Institute of Health and Welfare, 2020). This may indicate that people are more conscious of abuse or that mild cases are actually increasing, but society has a limited capacity to respond to all officially reported cases. Although mild violence is not necessarily consistent with severe violence, it is a risk factor for difficult cases in



229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

227 previously reported data.

### The validity of our sample

As Table 1 shows, the caregivers analyzed in our study cohort had a similar population distribution with regard to sex and age, compared with those from the CSLC data, while the nonanalyzed caregivers did not. This may have been caused by a trend of younger people being more likely to respond "Unknown/Do not want to answer" to the question about physical child abuse. While we cannot determine how many offenders answered "Unknown/Do not want to answer," these options are factors that support the quality of submitted data. In this web-based survey, panelists had to fill in all the questions to submit the data, and these additional options prevented them from giving contradictory answers, and enabled us to set the exclusion criteria. In addition, web-based surveys are reported to have better quality responses to delicate questions than do face-to-face interactions (Hasegawa, 2007). Our sample was biased toward people with higher educational backgrounds, similar to a previous report comparing the CSLC data and Internet surveys (Taniguchi & Omori, 2022). Considering the relationship between low educational background and child abuse, our results may be adaptable to a group in which child abuse is unlikely. Common risk factors for both genders The common risk factors of self-reported physical child abuse in offenders of both genders are "COVID-19 infected, both themselves and household members, within a year," "poor relationships with household members," and "history of child abuse." The intergenerational cycle of child abuse is known worldwide as a major risk factor (Berlin, Appleyard & Dodge, 2011; Horikawa et al., 2016; Assink et al., 2018; Anderson et al., 2018). In this pandemic,

"COVID-19 infected, both themselves and household members, within a year" had a high odds



251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

ratio, exceeding that of "history of child abuse." Considering that "only themselves being infected" did not show a high prevalence of self-reported physical child abuse, depression symptoms due to COVID-19 or Long COVID would not significantly change their responses. Therefore, physical child abuse is very likely to occur in a household where the offender and someone else were infected within a year. Considering the high odds ratios exceeding those of "history of child abuse," abuse is more likely to occur during periods of physical, mental, or social stress, when household members are frustrated with each other. For offenders, this stress may differ between "only offender infected (with COVID-19)" and "Offender and another household member infected." For example, it is difficult to determine the source of the infection in most cases. When several family members are simultaneously infected, they may often assume that others might be responsible for the infection. Some strong risk factors were found like intergenerational chain of child abuse and low socioeconomic status, but practical intervention is not possible. The negative impacts of COVID-19 have been widely studied, and this result might be one of the risks for increased stress, especially toward their families in particular situations (Amin & Parveen, 2022).

### Male offenders

Risk factors specific to male offenders were "work from home 4-7 days/week," "decreased work," "unwillingness to receive COVID-19 vaccination because the license process of the vaccine is doubtful," and "high benevolent sexism scale." Benevolent sexism measures the attitude toward stereotyped gender roles; men should work, and women should do household chores. We estimate that high levels of benevolent sexism of men reflect an emphasis on the importance of an occupation, leading to the strong stress caused by "work from home 4-7 days/week" and "decreased work." The strength of gender roles is inferred from the gender gap in the labor force



274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

participation. Substantial gaps in labor force participation prevalence between married men and women (25-30%) were observed in countries such as Japan, South Korea, Italy, and Brazil, compared to the gaps (less than 15%) in countries such as France, Sweden, Finland, England, and Denmark (ILOSTAT, 2022). In Japan, half of the employed women guit their jobs after childbirth (Ministry of Health, Labour and Welfare in Japan, 2021). As Japan has very established gender roles, it is unclear whether the work-related risk factors in our results are also found in countries with gender roles as prominent, such as in Italy and Brazil. Notably, law telework penetration is consistent with high gender gap in the labor force participation to some extent. It is reported that telework penetration before the pandemic was low in Japan, Italy, and Brazil and high in France, Sweden, Finland, England, and Denmark (OECD, 2021). The pandemic has increased telework globally, but it was a rapid change for the country with prominent stereotyped gender roles, such as Japan. Moreover, "work from home 4-7 days/week" and "decreased work" are associated with increased time spent at home and anxiety about potential job loss. For men with strong stereotyped gender roles, the sudden increase in time with their children is largely inconsistent with their previous lifestyles. This may be a primary cause of dissatisfaction with their children and spouses because they were previously uninvolved in childcare. In addition, the situation where the husband is the only working member can increase their dissatisfaction with their family when faced with the uncertainty of unemployment. It is reported that "working remotely" and "becoming unemployed during the COVID-19 pandemic" were not associated with anxiety symptoms in Finland (Savolainen et al., 2021). The telework system introduced during the pandemic will likely continue in many companies. Globally, as there was no precedent for the sudden increase in time at home for men, it is difficult to speculate how long the estimated risk factors due to men's familial dissatisfaction will continue in countries with strong gender roles.

1	Λ	1
$\mathcal{L}$	ч	n

Anxiety over decreasing or losing jobs is also influenced by financial support in each country. Many developed countries have job retention schemes and support the income of workers who are temporarily unemployed or whose work hours have been decreased. Japan, England, and Germany showed the lowest unemployment rates before and during the pandemic (OECD, 2020). On the other hand, in Japan, wages supported by job retention schemes for workers with 100% reductions in work hours are much higher than the amounts provided by unemployment benefits. In countries with large differences, such as in Japan, Portugal, Denmark, and Poland, anxiety surrounding job loss might be high (OECD, 2020).

"Unwillingness to receive COVID-19 vaccination because the license process of the vaccine is doubtful" was a risk factor for male offenders. Considering that "fear of COVID-19" and "feelings of discrimination related to COVID-19 infection" were risk factors associated with female offenders, men with no vaccination owing to skepticism might be subject to domestic discord due to their uncooperative attitude toward infection prevention.

### Female offenders

Risk factors specific to female offenders were "bad relationships with household members (compared to good relationships)," "strong fear of COVID-19," and "feelings of discrimination related to COVID-19 in the past two months." Unlike the discussion on the work-related risk factors in men, the infection-related risk factors are not always explained by gender roles. In Northern European countries, women showed a higher prevalence of anxiety and depression symptoms due to the COVID-19 pandemic (Savolainen et al., 2021; Johansson et al., 2013). Importantly, infection-related factors are gradually reduced with the abatement of the pandemic.



320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

This study had some limitations. First, we found that self-reported physical child abuse was likely to occur in the household where the offender and someone were infected within a year, with a high odds ratio similar to "history of child abuse." However, we could not identify who the someone is, and a particular combination might influence the relationship with child abuse. For example, while the combination of infected offender and child might indicate a simple mechanism of child abuse, the combination of infected offender and spouse might indicate a complex mechanism of child abuse regarding childcare, partly explained by the strength of gender roles. Second, we could not determine whether the offenders were biological parents or parents-in-law because there was no definition of parents in JACSIS. It has been reported that parents-in-law abuse children more than biological parents, and different types of abuse are observed (Turner et al., 2013; Daly & Wilson, 1994; Baba et al., 2020). Third, this was a crosssectional study for which causality was not identified. Fourth, although the analyzed respondents were representative of the population in terms of age and sex, they may have biased the characteristics and influenced the results. Our sample included many people with high educational backgrounds, similar to a previous study (Taniguchi & Omori, 2022). Respondents may have had milder personalities than the average for the population, which may have in turn resulted in a mild and low prevalence of self-reported child abuse.

337

338

339

340

341

### Conclusion

Physical child abuse was very likely to be reported in the "COVID-19 infected, both themselves and household members, within a year" households. Among male offenders, a significant relationship was observed in the change in work situations, such as "work from home 4–7"



343

344

345

346

347

348

days/week" and "decreased work." Difficulty adapting to work-related changes and the fear of
losing jobs may be high in countries with established gender roles and insufficient financial
support. Among female offenders, a significant relationship was observed regarding fear of
infection, which is consistent with the findings of previous studies on other pandemics (Savolainen
et al., 2021; Johansson et al., 2013). When emergencies, such as pandemics, keep people at home
or change past lifestyles, the parental stress might be more prevalent in countries with strong
stereotyped gender roles.

349

350

### Acknowledgements

- We thank all members of JACSIS Survey Team. We would like to thank Honyaku Center Inc.
- 352 for English language editing.

353

354

### References

- Amin UA, Parveen AP. 2022. Impact of COVID-19 on children. *Middle East Current Psychiatry*
- 356 29:94. DOI: 10.1186/s43045-022-00256-3.
- 357 Australian Institute of Health and Welfare. 2020. Child protection Australia 2018–19. Available at
- 358 https://www.aihw.gov.au/reports/child-protection/child-protection-australia-2018-
- 359 *19/summary* (accessed February 17, 2023).
- Baba S, Eshak ES, Shirai K, Fujiwara T, Yamaoka Y, Iso H. 2020. Factors Associated With Family
- Member's Spanking of 3.5-year-old Children in Japan. *Journal of Epidemiology* 30:464–
- 362 473. DOI: 10.2188/jea.JE20190160.
- 363 Berlin LJ, Appleyard K, Dodge KA. 2011. Intergenerational Continuity in Child Maltreatment:
- Mediating Mechanisms and Implications for Prevention. *Child development* 82:162–176.



365	DOI: 10.1111/j.1467-8624.2010.01547.x.
366	Brown SM, Doom JR, Lechuga-Peña S, Watamura SE, Koppels T. 2020. Stress and parenting
367	during the global COVID-19 pandemic. Child Abuse & Neglect 110:104699. DOI:
368	10.1016/j.chiabu.2020.104699.
369	Chapman DP, Whitfield CL, Felitti VJ, Dube SR, Edwards VJ, Anda RF. 2004. Adverse childhood
370	experiences and the risk of depressive disorders in adulthood. Journal of Affective
371	Disorders 82:217–225. DOI: 10.1016/j.jad.2003.12.013.
372	Children's Bureau. 2020.Child Maltreatment 2018. Available at
373	https://www.acf.hhs.gov/cb/report/child-maltreatment-2018 (accessed February 17, 2023).
374	Cuadrado-Gordillo I, Fernández-Antelo I, Martín-Mora Parra G. 2020. Search for the Profile of
375	the Victim of Adolescent Dating Violence: An Intersection of Cognitive, Emotional, and
376	Behavioral Variables. International Journal of Environmental Research and Public Health
377	17:8004. DOI: 10.3390/ijerph17218004.
378	Daly M, Wilson MI. 1994. Some differential attributes of lethal assaults on small children by
379	stepfathers versus genetic fathers. Ethology and Sociobiology 15:207–217. DOI:
380	10.1016/0162-3095(94)90014-0.
381	Dardenne B, Dumont M, Bollier T. 2007. Insidious dangers of benevolent sexism: Consequences
382	for women's performance. Journal of Personality and Social Psychology 93. DOI:
383	10.1037/0022-3514.93.5.764.
384	Department for Education. 2019. Characteristics of children in need: 2018 to 2019. Available at
385	https://www.gov.uk/government/statistics/characteristics-of-children-in-need-2018-to-
386	2019 (accessed February 17, 2023).
387	Fujiwara T, Kawakami N. 2011. Association of childhood adversities with the first onset of mental



388	disorders in Japan: Results from the World Mental Health Japan, 2002–2004. <i>Journal of</i>
389	Psychiatric Research 45:481–487. DOI: 10.1016/j.jpsychires.2010.08.002.
390	Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. 2009. Burden and
391	consequences of child maltreatment in high-income countries. The Lancet 373:68-81.
392	DOI: 10.1016/S0140-6736(08)61706-7.
393	Glick P, Fiske ST. 1996. The Ambivalent Sexism Inventory: Differentiating hostile and benevolent
394	sexism. Journal of Personality and Social Psychology 70:491-512. DOI: 10.1037/0022-
395	3514.70.3.491.
396	Hasegawa K. 2007. "You do not have to answer the question?" [Translated from Japanese].
397	Advanced social research:49–64.
398	Horikawa H, Suguimoto SP, Musumari PM, Techasrivichien T, Ono-Kihara M, Kihara M. 2016.
399	Development of a prediction model for child maltreatment recurrence in Japan: A historical
400	cohort study using data from a Child Guidance Center. Child Abuse & Neglect 59:55-65.
401	DOI: 10.1016/j.chiabu.2016.07.008.
402	ILOSTAT. 2022.Labour force participation rate by sex, age and marital status (%). Available at
403	https://ilostat.ilo.org/topics/women/ (accessed November 22, 2022).
404	Johansson R, Carlbring P, Heedman Å, Paxling B, Andersson G. 2013. Depression, anxiety and
405	their comorbidity in the Swedish general population: point prevalence and the effect on
406	health-related quality of life. <i>PeerJ</i> 1:e98. DOI: 10.7717/peerj.98.
407	Jones K, Stewart K, King E, Morgan WB, Gilrane V, Hylton K. 2014. Negative consequence of
408	benevolent sexism on efficacy and performance. Gender in Management: An International
409	Journal 29:171–189. DOI: 10.1108/GM-07-2013-0086.
410	King EB, Botsford W, Hebl MR, Kazama S, Dawson JF, Perkins A. 2012. Benevolent Sexism at



111	work. Gender Differences in the Distribution of Chantenging Developmental Experiences.
112	Journal of Management 38:1835–1866. DOI: 10.1177/0149206310365902.
113	Melissa Abi Rached, Ahmed Hankir, Rashid Zaman. 2021. Patriarchal Upbringing in the 21st
114	Century: The Impact Emotional Abuse Related to Parental Styles Has on the Genesis of
115	Gender Inequality and the Development of Unresolved Trauma in Children. Psychiatria
116	Danubina 33:121–132.
117	Ministry of Health, Labour and Welfare. 2022. Number of cases of child abuse consultation in
118	Japan, 1992-2021 (only in Japanese). Available at
119	https://www.mhlw.go.jp/content/11900000/000987725.pdf
120	Ministry of Health, Labour and Welfare in Japan. 2021. Overview of the 11th longitudinal survey
121	of children born in 2010 and the mothers (only in Japanese). Available at
122	https://www.mhlw.go.jp/toukei/saikin/hw/syusseiji/20/index.html (accessed November 28,
123	2022).
124	Minoura A, Ishimaru T, Kokaze A, Tabuchi T. 2021. Increased Work from Home and Low Back
125	Pain among Japanese Desk Workers during the Coronavirus Disease 2019 Pandemic: A
126	Cross-Sectional Study. International Journal of Environmental Research and Public
127	Health 18:12363. DOI: 10.3390/ijerph182312363.
128	Morinaga Y, Sakata K, Kitakaji Y, Oike M, Fukudome K. 2018. Benevolent Sexism Scale for
129	Japanese (only in Japanese). The Proceedings of the Annual Convention of the Japanese
130	Psychological Association 82:1AM – 140. DOI: 10.4992/pacjpa.82.0_1AM-140.
131	OECD. 2020. Job retention schemes during the COVID-19 lockdown and beyond. Available at
132	https://www.oecd.org/coronavirus/policy-responses/job-retention-schemes-during-the-
133	covid-19-lockdown-and-beyond-0853ba1d/ (accessed February 16, 2023).



134	OECD. 2021. Teleworking in the COVID-19 pandemic: Trends and prospects. Available at
135	https://www.oecd.org/coronavirus/policy-responses/teleworking-in-the-covid-19-
136	pandemic-trends-and-prospects-72a416b6/ (accessed November 22, 2022).
37	Pereda N, Díaz-Faes DA. 2020. Family violence against children in the wake of COVID-19
38	pandemic: a review of current perspectives and risk factors. Child and Adolescent
39	Psychiatry and Mental Health 14:40. DOI: 10.1186/s13034-020-00347-1.
40	Rapp A, Fall G, Radomsky AC, Santarossa S. 2021. Child Maltreatment During the COVID-19
141	Pandemic. Pediatric Clinics of North America 68:991–1009. DOI:
42	10.1016/j.pcl.2021.05.006.
43	Romanou E, Belton E. 2020. The impact of COVID-19 on child abuse in the UK. Available at
144	https://learning.nspcc.org.uk/research-resources/2020/social-isolation-risk-child-abuse-
145	during-and-after-coronavirus-pandemic/(accessed January 30, 2023).
146	Savolainen I, Oksa R, Savela N, Celuch M, Oksanen A. 2021. COVID-19 Anxiety—A
47	Longitudinal Survey Study of Psychological and Situational Risks among Finnish
48	Workers. International Journal of Environmental Research and Public Health 18. DOI:
49	10.3390/ijerph18020794.
50	Stickley A, Waldman K, Ueda M, Koyanagi A, Sumiyoshi T, Narita Z, Inoue Y, DeVylder JE, Oh
51	H. 2020. Childhood neglect and suicidal behavior: Findings from the National Comorbidity
152	Survey Replication. Child Abuse & Neglect 103:104400. DOI:
53	10.1016/j.chiabu.2020.104400.
54	Taniguchi M, Omori S. 2022.Bias in Internet-Based Surveys- A Comparative Study Using the
55	Census and an Interview-Based Survey Available at
56	https://english.nira.or.jp/papers/research_reports/2022/03/bias-in-internet-based-






157	surveysa-comparative-study-using-the-census-and-an-interview-based-survey-
158	.html?_ga=2.159685438.2030227239.1676684935-
159	1829064643.1676684935&_gl=1*1ehpmp2*_ga*MTgyOTA2NDY0My4xNjc2Njg0OTM1
160	*_ga_RE51W1NQ6C*MTY3NjY4NDkzNS4xLjEuMTY3NjY4NTc0NS40NS4wLjA.
161	the Japan COVID-19 and Society Internet Survey. 2022. Homepage of JACSIS study (only in
162	Japanese). Available at https://jacsis-study.jp/ (accessed April 20, 2022).
163	Turner HA, Finkelhor D, Hamby SL, Shattuck A. 2013. Family structure, victimization, and child
164	mental health in a nationally representative sample. Social Science & Medicine 87:39-51.
165	DOI: 10.1016/j.socscimed.2013.02.034.
166	Vives-Cases C, Sanz-Barbero B, Ayala A, Pérez-Martínez V, Sánchez-SanSegundo M, Jaskulska
167	S, Antunes das Neves AS, Forjaz MJ, Pyżalski J, Bowes N, Costa D, Waszyńska K,
168	Jankowiak B, Mocanu V, Davó-Blanes MC. 2021. Dating Violence Victimization among
169	Adolescents in Europe: Baseline Results from the Lights4Violence Project. International
170	Journal of Environmental Research and Public Health 18:1414. DOI:
171	10.3390/ijerph18041414.
172	Yamada K, Wakaizumi K, Kubota Y, Murayama H, Tabuchi T. 2021. Loneliness, social isolation,
173	and pain following the COVID-19 outbreak: data from a nationwide internet survey in
174	Japan. Scientific Reports 11:18643. DOI: 10.1038/s41598-021-97136-3.
175	



### Figure 1

Flowchart of the analyzed participants selection.







### Table 1(on next page)

Epidemiological characteristics of caregivers from the Japan COVID-19 and Society Internet Survey (JACSIS) 2021 and the Comprehensive Survey of Living Conditions (CSLC) 2019.



- **Table 1:**
- 2 Epidemiological characteristics of caregivers from the Japan COVID-19 and Society
- 3 Internet Survey (JACSIS) 2021 and the Comprehensive Survey of Living Conditions (CSLC)
- 4 **2019.**

		Analyzed	Non-analyzed	Caregivers
	n (%)	caregivers	caregivers	from CSLC
		2168/13870	128/13870	35819/206437
Men	Number/Total	(15.6)	(0.9)	(17.4)
	Age			
	20-29	105 (4.8)	13 (10.0)	1807 (5.0)
	30-39	820 (37.8)	55 (43.0)	12887 (36.0)
	40-49	1004 (46.3)	53 (41.4)	17033 (47.6)
	50-59	239 (11.0)	7 (5.5)	4092 (11.4)
	ISCED >=6	1503 (69.3)	86 (67.2)	13129 (36.7)
		2225/14305	104/14305	40464/217960
Woman	Number/Total	(15.6)	(0.7)	(18.6)
	Age			
	20-29	222 (10.0)	16 (15.4)	2949 (7.3)
	30-39	952 (42.8)	53 (51.0)	16821 (41.6)
	40-49	930 (41.8)	30 (28.8)	18083 (44.7)
	50-59	121 (5.4)	5 (4.8)	2611 (6.5)
	ISCED >=6	969 (43.6)	46 (44.2)	8549 (21.1)

<sup>5</sup> ISCED: International Standard Classification of Education