

Contraceptive use among reproductive-age females with disabilities in central Sidama National Regional State, Ethiopia: a multilevel analysis

Zelalem Tenaw^{Corresp., 1}, Taye Gari², Achamyesh Gebretsadik³

¹ Midwifery, Hawassa University, Hawassa, Sidama, Ethiopia

² Public health, Hawassa University, Hawassa, Sidama, Ethiopia

³ Public health, Hawassa University, Hawassa, Please select:, Ethiopia

Corresponding Author: Zelalem Tenaw
Email address: abigiatenaw@gmail.com

Background: Contraceptive use is an important and cost-effective intervention to prevent unwanted pregnancies. People with disabilities face discrimination when it comes to using contraception and are doubly burdened by unwanted pregnancies. However, the status of contraceptive use and associated factors among reproductive-aged females with disabilities was not adequately determined in Ethiopia. **Objective:** This study aimed to assess contraceptive use and associated factors among reproductive-age females with disabilities in Dale and Wonsho districts and Yirgalem city administration of central Sidama National Regional State, Ethiopia. **Methods:** A community-based cross-sectional study was conducted among randomly selected 620 reproductive-age females with disabilities living in the selected districts from June 20 to July 15, 2022. The data were collected through face-to-face interviewing techniques using a structured questionnaire. A multilevel logistic regression analysis model was employed to analyze the data. The adjusted odds ratio (AOR) with a 95% confidence interval (CI) was used to report the measures of associations. **Results:** In this study, 27.3 % (95% CI: 23.8 %, 31.0 %) of the reproductive-age females with disabilities were current contraceptive users. Regarding the methods, 82 (48.5%) of the reproductive-age females with disabilities used implants. Having good contraceptive knowledge (AOR = 9.03; 95% CI: 4.39, 18.6), transport accessibility to health facilities (AOR=2.28; 95% CI: 1.32, 3.94), being an adult (25 to 34 years old) (AOR = 3.04; 95% CI: 1.53, 6.04), having a hearing disability (AOR = 0.38; 95% CI: 0.18, 0.79), having paralysis of the extremities (AOR = 0.06; 95% CI: 0.03, 0.12), and wheel-chaired disability (AOR = 0.10; 95% CI: 0.05, 0.22) were factors associated with contraceptive use. **Conclusion:** Contraceptive use among reproductive-age females with disabilities is low. Transport accessibility, contraceptive knowledge, being in the age groups of 25 to 34 years, and the types of disability determine their contraceptive use. Therefore, designing appropriate

strategies to provide contraceptive education and information and provide contraceptive services in their homes is important to enhance contraceptive use.

Contraceptive use among reproductive-age females with disabilities in central Sidama National
Regional State, Ethiopia: a multilevel analysis

Zelalem Tenaw (ZT)^{1*}, Taye Gari (TG)² Achamyesh Gebretsadik (AG)²

^{1*}Department of Midwifery, College of Medicine and Health Sciences, Hawassa University,
Ethiopia

²School of Public health, College of Medicine and Health Sciences, Hawassa University, Ethiopia

Corresponding author (Zelalem Tenaw, Hawassa University, Ethiopia: Email:
abigiatenaw@gmail.com)

Email addresses:

ZT: abigiatenaw@gmail.com

TG: tayegari@gmail.com

AG: agtsadik@gmail.com

25

26 Abstract

27 **Background:** Contraceptive use is an important and cost-effective intervention to prevent
28 unwanted pregnancies. People with disabilities face discrimination when it comes to using
29 contraception and are doubly burdened by unwanted pregnancies. However, the status of
30 contraceptive use and associated factors among reproductive-aged females with disabilities was
31 not adequately determined in Ethiopia.

32 **Objective:** This study aimed to assess contraceptive use and associated factors among
33 reproductive-age females with disabilities in Dale and Wonsho districts and Yirgalem city
34 administration of central Sidama National Regional State, Ethiopia.

35 **Methods:** A community-based cross-sectional study was conducted among randomly selected 620
36 reproductive-age females with disabilities living in the selected districts from June 20 to July 15,
37 2022. The data were collected through face-to-face interviewing techniques using a structured
38 questionnaire. A multilevel logistic regression analysis model was employed to analyze the data.
39 The adjusted odds ratio (AOR) with a 95% confidence interval (CI) was used to report the
40 measures of associations.

41 **Results:** In this study, 27.3 % (95% CI: 23.8 %, 31.0 %) of the reproductive-age females with
42 disabilities were current contraceptive users. Regarding the methods, 82 (48.5%) of the
43 reproductive-age females with disabilities used implants. Having good contraceptive knowledge
44 (AOR = 9.03; 95% CI: 4.39, 18.6), transport accessibility to health facilities (AOR=2.28; 95% CI:
45 1.32, 3.94), being an adult (25 to 34 years old) (AOR = 3.04; 95% CI: 1.53, 6.04), having a hearing
46 disability (AOR = 0.38; 95% CI: 0.18, 0.79), having paralysis of the extremities (AOR = 0.06;
47 95% CI: 0.03, 0.12), and wheel-chaired disability (AOR = 0.10; 95% CI: 0.05, 0.22) were factors
48 associated with contraceptive use.

49 **Conclusion:** Contraceptive use among reproductive-age females with disabilities is low. Transport
50 accessibility, contraceptive knowledge, being in the age groups of 25 to 34 years, and the types of
51 disability determine their contraceptive use. Therefore, designing appropriate strategies to provide
52 contraceptive education and information and provide contraceptive services in their homes is
53 important to enhance contraceptive use.

Introduction

More than one billion people in the world are estimated to have disabilities. The majority of them are from developing countries (Zziwa Swaibu, Babikako Harriet et al. 2019). People with disabilities are the most discriminated against and marginalized group in many countries, particularly in developing countries, including Ethiopia (MacKay Don 2006, Hosseinpoor, Stewart Williams et al. 2013). In Sidama National Regional State, Ethiopia, we have observed that many females with disabilities are sacrificing themselves with unwanted pregnancies and unwanted children. Due to these problems, women with disabilities have been facing many challenges and have spent their lives begging around the roads and in churches with their unwanted pregnancies and children.

Contraceptive methods are chemicals, drugs, and surgical procedures used to prevent unwanted pregnancy (Jain and Muralidhar 2011). Although people with disabilities have a reproductive right to access and use contraception, coverage of contraceptive use in developing countries, including Ethiopia, is low when compared to developed countries. This is evidenced by the fact that 70.1% (Haynes, Boulet et al. 2018) of disabled women in the United States of America, 67.4% (Aslan, Yılmaz et al. 2021) in Turkey, 34% (Olajide, Omisore et al. 2014) in Nigeria, 26.9% (Trani, Browne et al. 2011) in Sierra Leone, 26.1% (Ayiga and Kigozi 2016) in Uganda, 17% (Kumi-Kyereme 2021) in Ghana, 16% (Odhiambo 2012) in Kenya, and 18% (Beyene, Munea et al. 2019) to 34% (Yesgat, Gebremeskel et al. 2020) in Ethiopia have access to contraceptives.

In Ethiopia, various factors associated with contraceptive use among females with disabilities were identified. Of the reported factors, marital status, age, types of disabilities, knowledge and attitude towards family planning methods, the presence of nearby health facilities providing family

planning services, keeping confidentiality and privacy in the health facility, having a good self-perception, and educational and economic status were the most common (Tsegay, Gebremariam et al. 2017, Beyene, Muneza et al. 2019, Yimer Awol Seid 2019, Mekonnen Alemayehu Gonie, Bayleyegn Alebachew Demelash et al. 2020, Yesgat, Gebremeskel et al. 2020). Not only do the factors listed above influence contraceptive use, but disability, such as types of disabilities and contraceptive methods, also has an administrative impact on contraceptive use.

In Ethiopia, few studies were conducted to determine the prevalence of contraceptive use and associated factors among reproductive-age females with disabilities from 2013 to 2019 (Tsegay, Gebremariam et al. 2017, Beyene, Muneza et al. 2019, Yimer Awol Seid 2019, Mekonnen Alemayehu Gonie, Bayleyegn Alebachew Demelash et al. 2020, Yesgat, Gebremeskel et al. 2020). These studies considered only urban female residents, deaf and blind females, and females enrolled in supporting organizations and considered only individual-level factors. Contraceptive coverage in these populations is also inconsistent, ranging from 18% (Beyene, Muneza et al. 2019) to 34% (Yesgat, Gebremeskel et al. 2020).

Therefore, this study aimed to determine the prevalence of contraceptive use and its associated factors among reproductive-age females with disabilities by considering rural and urban residency, all types of disability (except mental disability), and individual and community-level factors.

101

102

103

104 **Methods and materials**

105 **Study design and setting**

106 A community-based cross-sectional study was conducted from June 20 to July 15, 2022, to
 107 determine the prevalence and factors associated with contraceptive use among reproductive-age
 108 females with disabilities in Sidama National Regional State, Ethiopia. The study was conducted in
 109 the Dale and Wonsho districts and in the Yirgalem city administration. According to the Sidama
 110 National Regional State Report (2021), the total population of Dale and Wonsho districts and
 111 Yirgalem city administration was 469,455 (Sidama Region Health Bureau 2021, Sidama Region
 112 Health Bureau 2022). The two districts are the health and demographic surveillance sites of
 113 Hawassa University. Both districts are known for their coffee production and highly dense
 114 populations. In the districts and city administration, there are 56 rural and 10 urban kebeles (the
 115 lowest political administrative units in Ethiopia). The districts and city administration have one
 116 hospital, 16 health centers, and 54 health posts.

117 **Population**

118 Reproductive-age females with disabilities in Dale and Wonsho districts and Yirgalem city
 119 administration in Sidama National Regional State were the source population. Reproductive-age
 120 females with disabilities who lived in the selected kebeles for at least six months were the study
 121 population, except for those who have dual disabilities (i.e., cannot see and hear) and are seriously
 122 ill during the data collection time.

123 **Sample size and sampling procedure**

124 The sample size for the first objective (prevalence) was determined by using Epi Info version 7
 125 software with the assumptions of a 95% confidence interval with 33.7% contraceptive use among

reproductive-age women with disability (Yibeltal Mesfin Yesgat, Feleke Gebremeskel et al. 2020), a level of significance (α) of 0.05, a 5% margin of error ($d = 0.05$), and a design effect of 1.64. The sample size for factors associated with contraceptive use was also computed using Epi-Info version 7 with the assumptions of a two-sided confidence level of 95%, a power of 80, a ratio of one (unexposed: exposed), and a percent outcome in the unexposed group (14.5) versus a percent outcome in the exposed group (24.5). Accordingly, the maximum (530) sample size was determined by marital status (Beyene, Muneza et al. 2019). The sample size from the prevalence of 563 was larger than the associated factors' maximum sample size of 530. After adjusting for an anticipated 10% nonresponse rate, the final sample size was 620.

The sample size was proportionally allocated to the randomly selected 30 kebeles (20 rural and 10 urban) based on the number of reproductive-age females with disabilities. Before conducting this study, a house-to-house census was done to determine the number and identify reproductive-age females with disabilities in each kebele. Reproductive-age females with disabilities were registered during the census using the tracing form. The registration form was used to select study participants using a simple random sampling technique.

Variables

The outcome variable was contraceptive use. Whereas the independent variables were marital status, age, types of disability, educational status, knowledge about family planning, income, self-perception, attitude toward family planning, health care providers' attitudes, the presence of family planning provision at a nearby health facility, and the keeping of confidentiality and privacy by the health facility.

Data collection procedures and quality assurance

The questionnaires (data collection tools) were developed by reviewing different existing literature, like EDHS 2016 (Central Statistical Agency (CSA) [Ethiopia] and ICF 2016, Mekonnen Alemayehu Goniye, Bayleyegn Alebachew Demelash et al. 2020, Yesgat, Gebremeskel et al. 2020), which consists of personal and socio-demographic characteristics and contraceptive use-related issues. After developing and pretesting the data collection tool, six data collectors and one

supervisor who are fluent speakers of Sidamu Afoo and who have data collection experience were employed. The data were collected through face-to-face interviewing techniques using structured questionnaires. Two of the data collectors were proficient in sign language and collected the data from reproductive-age females with hearing disabilities. The interview was conducted in a place where confidentiality and privacy are assured. To assure the quality of the data collection, a three-day data collector training was given. The data collection tool was first prepared in English and then translated into Afoo-Sidamu, a local language, and then back to English to check the consistency. The trained data collectors did a pre-test on 31 (5%) reproductive-age females with disabilities in *Lokie kebele* Hawassa city to check the tools, and corrections were made based on the feedback. The principal investigator (PI) monitors and controls the overall process of data collection and makes appropriate corrections for any issues raised during data collection. The PI also checked the completeness of the questionnaires daily.

Data management and analysis

The Kobo Collect version 2021.3.4 application was used to collect the data. Following collection, the data were imported into Stata version 16 for analysis using the "SSC install kobo2stata" command. The cleaning and organizing of the data were done in Stata. The types of variables were clarified, and the distribution was checked by running the frequency for categorical data and mean \pm SD (standard deviation) for continuous variables. A multilevel logistic regression analysis model was used to account for the kebele level. Before using the multilevel logistic analysis model, we checked the intraclass correlation coefficient (ICC) level with the chi-square significance level to determine whether using the multilevel logistic analysis model is justifiable. The ICC of 0.12 and its chi-square ($P = 0.001$) significance level showed that using a multilevel analysis model is reasonable. Then, bi-variable multilevel logistic regression was done to identify eligible variables ($P\text{-value} < 0.20$) for multivariable multilevel logistic regression analysis. The multivariable multilevel logistic regression was performed to check for the presence of an association between level one or level two variables and contraceptive use. To determine whether a significant

association existed and its strength, variables with adjusted odds ratios with a 95% confidence interval and a P-value <0.05 were considered.

Ethical considerations

The ethical clearance was obtained from the Institutional Review Board at the College of Medicine and Health Sciences of Hawassa University with an approval number of Ref.No: IRB/143/14. After approval, a support letter was written to the Sidama National Regional Public Health Institute. Then, after obtaining the support letter from Sidama National Regional Public Health Institute, the permission and cooperation letter were given to the woreda health offices. Finally, the woreda health offices wrote a permission letter to selected kebeles, asking them to cooperate and give consent to conduct the study. Written consent was obtained from the study participants to collect the data. There is no risk in participating in this survey. People with disabilities having different health problems were linked to nearby health facilities for possible support and follow-up.

205

206

207

208

209

210

211

212

213 **Results**

214 **Socio-demographic characteristics of study participants**

215 A total of 620 reproductive-age females with disabilities were included in this study. The mean
216 (SD) age of the study participants was 28.12 (8.54) years. Of the study participants, 55.32% had no
217 formal education (illiterate) and almost all (98.90%) were not employed. Most (83.71%) of the
218 reproductive-age females with disabilities had no occupation, and 54.20% were married (Table 1).

219 **Contraceptive knowledge and attitude**

220 Among the study participants, 382 (61.6%) had good knowledge about contraceptives. Regarding
221 attitude, 303 (48.9%) of reproductive-age females with disabilities had a positive attitude towards
222 contraceptive use.

223 **Contraceptive use prevalence**

224 In this study, the overall prevalence of current contraceptive use among reproductive-age females
225 with disabilities was 27.3% (95% CI: 23.8, 31.0), of which 19.19% (95% CI: 16.17, 22.52) were
226 from rural residents and 8.06 (95% CI: 6.04, 10.49) were from urban residents. From the overall
227 contraceptive use, 20.3% (95% CI: 17.2, 23.7) were married, and 7% (95% CI: 5.06, 9.22) were
228 unmarried.

229 **Types of contraceptive methods used**

Of the contraceptive method users, 82 (48.5%) of the reproductive-age females with disabilities used implants, followed by injectable (36%), oral contraceptive pills (12%), and intrauterine contraceptive devices and condoms (4%).

Reasons for not using contraceptives and their plan to use in the future

This study tried to identify the possible reasons for not using contraceptives among the 451 non-users of contraceptives of reproductive age females with disabilities. Of the respondents, the majority, 161 (36%) did not use it due to a lack of information about contraceptives, and 23 (5%) are due to the plan to have a baby in the near future (Figure 1). Regarding their future plans for contraceptive use, 147 (32.59%) had the plan to use, 209 (46.34%) had no plan to use, and 95 (21.06%) were not sure about their future plans.

Factors associated with contraceptive use

Random effect model

In the zero model (model I), 12 % of the variability in contraceptive use was at the community level (kebele level). This may be attributable to other unobserved community factors ($ICC = 0.12$), which were supported by the chi-square test ($P < 0.001$). This finding also showed that using a multilevel analysis model is reasonable.

Fixed effect model

In the bivariable logistic regression, marital status, education, occupation, self-perception, age, transport accessibility, contraceptive knowledge, types of disability, and residence were significantly associated with contraceptive use, but in the multivariable multilevel logistic regression analysis (after adjusting for the possible confounders), contraceptive knowledge, transport accessibility to the health facility, age, and types of disability were significantly associated with contraceptive use.

Reproductive-age females with a disability who knew about contraceptives had nine (AOR = 9.03; 95% CI: 4.39, 18.6) times higher odds of contraceptive use compared with those who had no

contraceptive knowledge. On the other hand, reproductive-age females with disabilities who had transport accessibility to health facilities had a twofold (AOR = 2.28; 95% CI: 1.32, 3.94) higher likelihood of contraceptive use compared with those who had no transport accessibility. Regarding age, reproductive-age females with disabilities who were 25 to 34 years old had three (AOR = 3.04; 95% CI: 1.53, 6.04) times higher odds of contraceptive use compared with those who were in the age group of 15 to 24 years old. Participants with hearing disabilities were 62% (AOR = 0.38; 95% CI: 0.18, 0.79), those with extremity paralysis were 94% (AOR = 0.06; 95% CI: 0.03, 0.12), and those with wheel-chair disabilities were 90% (AOR = 0.10; 95% CI: 0.05, 0.22) less likely to use contraceptives than their counterparts with vision disabilities (Table 2).

Discussion

The prevalence of contraceptive method use among reproductive-age females with disabilities was 27.3%. In the multivariable multilevel logistic regression analysis, contraceptive knowledge, transport accessibility to the health facility, being an adult (25 to 34 years old), and types of disability were significantly associated with contraceptive use.

This study revealed that the contraceptive method use prevalence of 27.3% was almost similar to the studies conducted in Uganda, 26.1% (Ayiga and Kigozi 2016), in Sierra Leone, 26.9% (Trani, Browne et al. 2011) and in Ethiopia, 24.5% (Mekonnen Alemayehu Gonie, Bayleyegn Alebachew Demelash et al. 2020) and 27.2% (Tsegay, Gebremariam et al. 2017). On the other hand, the prevalence of contraceptive use in the current study is higher than in the previous studies conducted in Kenya, 16% (Odhiambo 2012), Ghana, 17% (Kumi-Kyereme 2021), and in Ethiopia 18% (Beyene, Muneza et al. 2019). The possible reasons might be the long time interval between previous studies (2008) (Odhiambo 2012) and our study (2022). And, due to age group differences in the study population, school young people in the previous study (Kumi-Kyereme 2021) and

reproductive-age females with disabilities in the current study. And also due to information bias from the source of data, since the information was collected from caregivers (Odhiambo 2012). The other possible elucidation might be due to sample size differences. However, the prevalence of contraceptive use is lower than the studies conducted in Namibia, 32.7% (Loeb and Grut 2005), Nigeria, 34% (Olajide, Omisore et al. 2014) and in Ethiopia 34% (Yesgat, Gebremeskel et al. 2020). The possible justification might be that the Namibia study (Loeb and Grut 2005) was conducted among married women with disabilities, the chance of having unprotected sex increased among married people, and people with disabilities have a higher desire to prevent pregnancy (Casebolt M Tara, Singh Kavita et al. 2022).

In this study, contraceptive knowledge is found to be significantly associated with contraceptive use. Those participants who had contraceptive knowledge had a higher chance of using contraceptives compared with those who had no contraceptive knowledge. The finding is consistent with the studies conducted in Uganda (Ayiga and Kigozi 2016) and Nigeria (Olajide, Omisore et al. 2014). The possible justification might be due to the power of knowledge to create awareness and overcome some cultural and social constraints that may act as a barrier to the use of contraceptives (Beyene, Muneza et al. 2019).

Those who had transport availability to the health facility had a greater chance of contraceptive use when compared with those who had no transport accessibility to the health facility. As it is known, most people with disabilities face a physical challenge (Olajide, Omisore et al. 2014) in accessing health facilities. Due to this, transportation is very important for accessing health facilities and getting contraceptive methods. Being an adult (25 to 34 years old) increased the chance of contraceptive use when compared with the age group of 15 to 24 years old. The possible justification is that the chance of marriage and unprotected sexual intercourse will increase among 25 to 34-year-old females with disabilities, and the chance of using contraception will also increase due to the greater desire of people with disabilities to avoid pregnancy (Casebolt M Tara, Singh Kavita et al. 2022). Compared with vision impairment, the probability of using contraceptives by hearing-disability reproductive age females with disabilities had 62% lower odds of contraceptive use, 94% lower odds of contraceptive use by extremity paralysis disabilities, and 90% lower odds by wheel-chaired disabilities. This finding is inconsistent with studies conducted in Gondar, the Amhara region, Ethiopia, and Addis Ababa. The studies revealed that the probability of

contraceptive use increased among vision-impaired females with disabilities when compared with other types of disabilities (Beyene, Muneza et al. 2019). The possible reason for the difference might be that visually impaired females with disabilities had an increased chance of accessing information access through different social media, most commonly radio. Radio is one of the most accessible and effective channels of information transmission for people with disabilities in developing countries, including Ethiopia (Beyene, Muneza et al. 2019).

These findings may be important for different stakeholders who are concerned about reproductive-age females with disabilities and their reproductive health services, specifically contraceptive use. This study was conducted among all types of reproductive-age females with disabilities who reside in urban and rural areas. In the previous studies, rural residents with disabilities were excluded from the contraceptive use assessment studies. The other strength of this study was the use of multilevel analysis to check the effect of kebele-level variables on contraceptive use. However, due to the sensitivity and principles of contraceptive use, this study did not consider reproductive-age females with mental disabilities. Therefore, this study could be generalized to all reproductive-age females with disabilities except for mental disabilities.

Conclusion

Contraceptive use among reproductive-age females with disabilities, specifically among the unmarried, is noticeably low in Dale and Wonsho districts and Yirgalem city administration, Sidama National Regional State, Ethiopia. Contraceptive knowledge, accessible transportation to the health facility, being an adult (25 to 34 years old), and having specific types of disability were factors associated with contraceptive use among reproductive-age females with disabilities. Therefore, designing appropriate strategies to provide contraceptive education and information and provide contraceptive services in their homes is important to enhance contraceptive use.

Acknowledgements

The authors would like to thank the data collectors and the study participants for their valuable contributions to this study. We are also grateful to the Sidama National Regional State Public Health Institute and selected woredas and kebeles for their assistance and permission to undertake this research.

344
345
346
347
348
349
350
351
352
353
354
355
356
357

358 **References**

359 Aslan, E., B. Yılmaz and Z. Acar (2021). "Reproductive Health, Sexual Function and Satisfaction
360 Levels in Women with Physical, Hearing, and Visual Disabilities." Sexuality and Disability: 1-14.
361 Ayiga, N. and S. Kigozi (2016). "Access to and Uptake of Contraception by Women with
362 Disabilities."
363 Beyene, G. A., A. M. Muneza and G. A. Fekadu (2019). "Modern Contraceptive Use and Associated
364 Factors among Women with Disabilities in Gondar City, Amhara Region, North West Ethiopia: A
365 Cross Sectional Study." Afr J Reprod Health **23**(2): 101-109.
366 Beyene, G. A., A. M. Muneza and G. A. Fekadu (2019). "Modern contraceptive use and associated
367 factors among women with disabilities in Gondar City, Amhara Region, north west Ethiopia: a
368 cross sectional study." African journal of reproductive health **23**(2): 101-109.

Casebolt M Tara, Singh Kavita, Speizer Ilene S and Halpern Carolyn T (2022). "Use of modern contraceptives by women with disabilities in Rajasthan, India: An analysis of the annual health survey." Sexual & Reproductive Healthcare **31**: 100699.

Central Statistical Agency (CSA) [Ethiopia] and ICF (2016). Ethiopia Demographic and Health Survey.

Haynes, R. M., S. L. Boulet, M. H. Fox, D. D. Carroll, E. Courtney-Long and L. Warner (2018). "Contraceptive use at last intercourse among reproductive-aged women with disabilities: an analysis of population-based data from seven states." Contraception **97**(6): 538-545.

Hosseinpour, A. R., J. A. Stewart Williams, J. Gautam, A. Posarac, A. Officer, E. Verdes, N. Kostanjsek and S. Chatterji (2013). "Socioeconomic inequality in disability among adults: a multicountry study using the World Health Survey." American journal of public health **103**(7): 1278-1286.

Jain, R. and S. Muralidhar (2011). "Contraceptive methods: needs, options and utilization." The Journal of Obstetrics and Gynecology of India **61**(6): 626-634.

Kumi-Kyereme, A. (2021). "Sexual and reproductive health services utilisation amongst in-school young people with disabilities in Ghana." Afr J Disabil **10**: 671.

Kumi-Kyereme, A. J. A. J. o. D. (2021). "Sexual and reproductive health services utilisation amongst in-school young people with disabilities in Ghana." **10**.

Loeb, M. and L. Grut (2005). "Women with disabilities sharing knowledge: Education, Employment, Reproductive history." SINTEF Rapport.

MacKay Don (2006). "The United Nations Convention on the rights of persons with disabilities." Syracuse journal of international law and commerce **34**: 323.

Mekonnen Alemayehu Gonie, Bayleyegn Alebachew Demelash, Aynalem Yared Asmare, Adane Tigist Demssew, Muluneh Mikyasa Arega and Asefa Meaza (2020). "Level of knowledge, attitude, and practice of family planning and associated factors among disabled persons, north-Shewa zone, Amhara regional state, Ethiopia,." Contraception Reproductive Medicine **5**(1): 1-7.

Odhiambo, M. (2012). Determinants of contraceptive use: a comparative study of disabled and non-disabled women in Kenya, University of Nairobi, Kenya.

Olajide, F. O., A. G. Omisore, O. O. Arije, O. T. Afolabi and A. O. Olajide (2014). "Awareness and use of modern contraceptives among physically challenged in-school adolescents in Osun state, Nigeria." African Journal of Reproductive Health **18**(2): 87-96.

Sidama Region Health Bureau (2021). Health facility data with Catchment Population (Updated Meskerem 2014 E.C)

Sidama Region Health Bureau (2022). Health facility and population prifile. Hawassa, Ethiopia, Sidama Region Health Bureau.

Trani, J.-F., J. Browne, M. Kett, O. Bah, T. Morlai, N. Bailey and N. Groce (2011). "Access to health care, reproductive health and disability: a large scale survey in Sierra Leone." Social science & medicine **73**(10): 1477-1489.

Trani, J.-F., J. Browne, M. Kett, O. Bah, T. Morlai, N. Bailey, N. J. S. s. Groce and medicine (2011). "Access to health care, reproductive health and disability: a large scale survey in Sierra Leone." **73**(10): 1477-1489.

Tsegay, K., H. Gebremariam and F. Haile (2017). "Modern Contraceptive Methods Utilization and Associated Factors among Women with Disabilities in Mekelle Town, Tigray, Ethiopia." International Journal of Pharmaceutical and Biological Sciences Fundamentals.

Yesgat, Y. M., F. Gebremeskel, W. Estifanous, Y. Gizachew, S. Jemal, N. Atnafu and K. Nuriye (2020). "Utilization of family planning methods and associated factors among reproductive-age women with disability in Arba Minch Town, Southern Ethiopia." Open access journal of contraception **11**: 25.

Yibeltal Mesfin Yesgat, Feleke Gebremeskel, Wubshet Estifanous, Yordanos Gizachew, Seid Jemal, Natnael Atnafu and Keyredin Nuriye (2020). "Utilization of Family Planning Methods and Associated Factors Among Reproductive-Age Women with Disability in Arba Minch Town, Southern Ethiopia." Journal of contraception **11**: 25-35.

Yimer Awol Seid, M. L. M. (2019). "Modern contraceptive methods knowledge and practice among blind and deaf women in Ethiopia. A cross-sectional survey." BMC Womens Health **19**(1): 1-13.

Zziwa Swaibu, Babikako Harriet, Kwesiga Doris, Kobusingye Olive, Bentley Jacob A , Oporia Frederick, Nuwematsiko Rebecca, Bachani Abdulgafoor, Atuyambe Lynn M and Paichadze Nino (2019). "Prevalence and factors associated with utilization of rehabilitation services among people with physical disabilities in Kampala, Uganda. A descriptive cross-sectional study." BMC public health **19**(1): 11.

Figure 1

Reasons for not using contraceptives among reproductive-age females with disabilities in Sidama National Regional State, Ethiopia, 2022




Table 1(on next page)

Socio-demographic characteristics of reproductive-age females with disabilities in Sidama Regional Stata, Ethiopia, 2022 (N=620)

Table 1: Socio-demographic characteristics of reproductive-age females with disabilities in Sidama Regional Stata, Ethiopia, 2022 (N=620)

Variable		Number	Percent
Age in years mean(SD)	28.12 (8.54)		
Marital status of the participants	Married	336	54.2
	Never married	239	38.6
	Widowed	20	3.20
	Divorced	25	4.00
Residency	Rural	381	61.45
	Urban	239	38.55
Participants educational status	Illiterate	343	55.32
	Primary school	192	30.96
	Secondary school	78	12.60
	Vocational and technique	7	1.12
Employment status	Employed	6	1.10
	Unemployed	538	98.90
wealth index of household	Low	216	34.84
	Medium	196	31.61
	High	208	33.55

3

Table 2(on next page)

Multilevel logistic regression analysis for factors associated with contraceptive use among reproductive-aged females with disabilities in Sidama National Regional State, 2022

1 **Table 2:** Multilevel logistic regression analysis for factors associated with contraceptive use
 2 among reproductive-aged females with disabilities in Sidama National Regional State, 2022

Variables		Contraceptive		COR with 95% CI	AOR with 95% CI
		Use	Not use		
Marital status	Not married	48	238	1.00	1.00
	Married	121	215	2.79 (1.92,4.35)*	1.53 (0.87, 2.69)
Education	Illiterate	85	258	1.00	1.00
	Literate	84	198	1.30 (0.89,1.91)*	1.13 (0.68, 1.88)
Occupation	No occupation	130	389	1.00	1.00
	Have occupation	39	62	1.88 (1.02,2.68)*	1.17 (0.63, 2.16)
Self-perception	Bad	66	127	1.00	1.00
	Good	103	324	0.61 (0.41,0.92)*	0.99 (0.57, 1.70)
Age (Years)	15 to 24	28	193	1.00	1.00
	25 to 34	88	153	3.96 (2.61,7.22)*	3.04 (1.53, 6.04)**
	35 to 44	48	85	3.89 (1.89,6.02)*	2.20 (0.99, 4.85)
	45 to 49	5	20	1.70 (0.52,4.85)	2.09 (0.56, 7.83)
Transport accessibility	Yes	129	257	2.34 (1.52,3.59)*	2.28 (1.32, 3.94)**
	No	40	194	1.00	1.00
Distance to a health facility	≥30 minutes on foot	56	90	1.39 (0.84, 2.30)*	1.17 (0.64, 2.11)
	<30 minutes on foot	113	361	1.00	1.00
knowledge	Knowledgeable	143	239	4.85 (2.97,7.90)*	9.03 (4.39, 18.6)**
	Not knowledgeable	26	212	1.00	1.00
Wealth index	Rich	67	141	1.21(0.76,1.92)	1.41 (0.77, 2.57)
	Medium	46	150	0.99(0.61,1.63)	1.22 (0.66, 2.24)
	Poor	56	160	1.00	1.00
Disability type	Hearing	30	100	0.26 (0.14,0.48)*	0.38 (0.18, 0.79)**
	Extremity	36	185	0.18 (0.10,0.31)*	0.06 (0.03, 0.12)**
	Wheel-chaired	40	84	0.39 (0.22,0.71)*	0.10 (0.05, 0.22)**
	Vision	63	82	1.00	1.00
Residence	Urban	50	189	0.65 (0.34,1.21)*	0.84 (0.42, 1.71)

	Rural	119	262	1.00	1.00
--	-------	-----	-----	------	------

3
*: P-value < 0.2; **: P-value < 0.05; AOR: Adjusted odds ratio; CI: Confidence interval

4