

Research progress on the bulb expansion and starch enrichment in taro (*Colocasia esculenta* (L). Schott)

Erjin Zhang¹, Wenyan Shen¹, Weijie Jiang¹, Wenlong Li¹, Xiaping Wan¹, Xurun Yu¹, Fei Xiong^{Corresp. 1}

¹ Yangzhou University, Yangzhou, Jiangsu province, China

Corresponding Author: Fei Xiong
Email address: feixiong@yzu.edu.cn

Background. Taro is an important potato crop in the world, which can be used as food, vegetable, feed and industrial raw materials. The yield and quality of taro are mainly determined by the expansion degree of taro bulb and the filling condition of starch, while the expansion of taro bulb is a complex biological process. However, few information is reviewed on research progress on the bulb expansion and starch enrichment in taro.

Methodology. PubMed, Web of Science and China National Knowledge Infrastructure databases were searched for relevant articles. After removing duplicate articles and the articles with little relevance, 73 articles were selected for the review.

Results. This article introduces the formation and development of taro bulb for the workers engaged in taro research. The content includes the process of amyloplast formation at the cytological level and the changes in bulb expansion and starch enrichment at physiological levels, which involve endogenous hormones and key enzyme genes for starch synthesis. The effects of environment and cultivation methods on taro bulb expansion were also reviewed.

Conclusions. The future research directions and research focus about the development of taro bulb were prospected. Limited research has been conducted on the physiological mechanism and hormone regulation pathway of taro growth and development, taro bulb expansion, key gene expression, and starch enrichment. Therefore, the above research will become the key research direction in the future.

Research progress on the bulb expansion and starch enrichment in taro (*Colocasia esculenta* (L.) Schott)

Er-jin Zhang¹, Wen-yuan Shen¹, Wei-jie Jiang¹, Wen-long Li¹, Xia-ping Wan¹, Xu-run Yu¹, Fei Xiong¹

¹ Jiangsu Key Laboratory of Crop Genetics and Physiology, Yangzhou University, Yangzhou, China

Corresponding Author:

Fei Xiong¹

Yangzhou University, Yang Zhou, Jiang Su, 225009, China

Email address: feixiong@yzu.edu.cn

Abstract

Background. Taro is an important potato crop in the world, which can be used as food, vegetable, feed and industrial raw materials. The yield and quality of taro are mainly determined by the expansion degree of taro bulb and the filling condition of starch, while the expansion of taro bulb is a complex biological process. However, few information is reviewed on research progress on the bulb expansion and starch enrichment in taro.

Methodology. PubMed, Web of Science and China National Knowledge Infrastructure databases were searched for relevant articles. After removing duplicate articles and the articles with little relevance, 73 articles were selected for the review.

Results. This article introduces the formation and development of taro bulb for the workers engaged in taro research. The content includes the process of amyloplast formation at the cytological level and the changes in bulb expansion and starch enrichment at physiological levels, which involve endogenous hormones and key enzyme genes for starch synthesis. The effects of environment and cultivation methods on taro bulb expansion were also reviewed.

Conclusions. The future research directions and research focus about the development of taro bulb were prospected. Limited research has been conducted on the physiological mechanism and hormone regulation pathway of taro growth and development, taro bulb expansion, key gene expression, and starch enrichment. Therefore, the above research will become the key research direction in the future.

Introduction

Taro is an underground bulbous crop planted in the tropical and subtropical regions. It originated in China, India, Malaysia, and other regions, and is widely cultivated in Asia, Africa, and other regions. It has a cultivation history of more than 2,000 years in China. At present, taro can be classified using three methods. According the ecological type, it can be classified into two classification which we called aquatic taro planted in paddy field and dry taro planted in orchard soil. According the eating part, it can be classified into three classifications which we called

petiole taro whose mainly edible part is the petiole, flower taro whose mainly edible part is the flower, and bulb taro whose mainly edible part is the bulb. According the bulb tillering habit, it can be classified into three classifications which we called Kui taro, multi-cormels taro, and multi-head taro. (Zhang,1982; Wu et al., 2021). The three classifications of taros in agricultural production of China exhibit various characteristics (Figure 1). Kui taro has few sub taro and single mother-taro which is the main edible part. In multi-cormels taro, the sub taros are numerous and grow in groups which have strong tillering ability. In multi-head taro, there are few sub taros and the tillers of the mother taros grow in groups. The fundamental difference of taro bulb tiller is the difference of expansion between mother and sub taro. This phenomenon is caused by the difference in the number of chromosome and differential expression of genes (Zhu et al., 2018). Studies on taro chromosomes has proved that Kui taro is diploid ($2n=2x=28$), while the multi-head and multi-cormels taro are triploid ($2n=3x=42$) (Huang et al., 2014; Wu et al., 2021). The chromosome multiples are related to the geographical distribution. Diploid is more common in hot and humid areas with low altitudes, and triploid is more common in dry and cold areas with high altitudes.

The taro bulb, which is rich in starch and carbohydrates, is one of the staple food (Njintang et al., 2008). In comparison with potato, sweet potato, and cassava, taro starch granules are small and has a diameter of approximately 1.5 μm , making it easy to digest and providing therapeutic and health care functions. Taro starch particles are small and uniform and have good cold and hot stability, good whiteness, and good adhesion. Therefore, it can be used as a brightener in cosmetics (Qi et al., 2015). Taro bulbs are rich in many nutrients, such as starch, and the starch content of different varieties of taro could reach 10%–36% (Falade et al., 2013). Taro starch can be used as an industrial raw material. It could improve the stability of Pickering emulsion (Zhang et al., 2020), and it could be used for the treatment of wastewater generated in the printing and dyeing process (Zhou et al., 2018) and as an enhancer of starch film (Dai et al., 2015). Taro contains a large amount of dietary fiber and various essential amino acids, but its fat content is low (Sefa-Dedeh et al., 2002). It is also rich in vitamins and amino acid, making it is a food for people of all ages (Han et al., 2018). Taro leaf contain flavonoids, but its biosynthesis of flavonoids in taro bulb is still unknown (Iwashina et al., 1999). Taro is a food and medicine homologous crop, which is often used for the treatment of diarrhea, internal bleeding, asthma, and skin diseases (Prajapati et al., 2011). It could also lower blood sugar and cholesterol (Sebnem et al., 2012). Therefore, taro has good prospects for medicinal development.

According to the statistics of the Food and Agriculture Organization of United Nations, taro is the 14th largest vegetable crop, with a global planting area of 1.6 million hectares and an annual output of 11.7 million tons. The cultivation area in China ranks first in the world and has a large number of wild resources and local varieties, which are mainly distributed in southern regions such as Yunnan, Taiwan, and the Yangtze River Basin. In recent years, taro has become an important input and export trade industry. The total export trade volume and total value were higher than the imports (Chang et al., 2019). With the rapid development of the taro industry, the

demand for taro has increased. It has also introduced new and high requirements for taro production and basic theoretical research.

Survey methodology

PubMed, Web of Science and China National Knowledge Infrastructure databases were searched for relevant articles. A total of 2525 articles appeared in the PubMed database using “taro” as the search term and the date of publication from 1975/1/1 to 2022/5/1. After narrowing the search with keywords including “bulb of taro”, “bulb expansion in taro”, “bulb starch enrichment in taro”, “amyloplast enrichment process of taro bulbs”, “development of taro bulbs”, “regulation of hormones on bulb development”, “genes in starch synthesis”, and “regulation of hormones on bulb swelling”, 1532 studies were obtained. With “taro” as the search term, 2525 articles published between 1975–2022 appeared in the Web of Science database, of which 914 articles were selected as above. After removing duplicate articles and the articles with little relevance, 73 articles were selected for the review.

Taro expansion and development process

Taro is a perennial herbaceous plant, which belongs to the Araceae family, but it is generally cultivated as an annual crop in agricultural production. Lebot V identified the plants as *Colocasia esculenta* based on its morphological examination (Ivancic A, 1999). According to the growth and development characteristic of taro (Figure 2), its life cycle is divided into five periods, namely, embryonic stage, seedling stage, plant vigorous growth stage, taro expansion stage, and bulb dormancy stage (Sun et al., 2014). Taro bulbs undergo metamorphosis during evolution. According to the authoritative book (Huang et al., 2006), some terms such as “mother taro”, “grandson taro”, “great grandson taro” and so on are merely direct translation of the local Chinese terms. Here take multi-cormels taro and multi-head taro for example. The bulb of taro is composed of mother and sub taro, and some varieties include grandson and great grandson taro (Figure 3). The mother taro of taro bulb is developed by the top bud of the seed taro. The sub taro is developed from the lateral bud on the mother taro, and the grandson taro is developed from the lateral bud of the sub taro. An axillary bud is present on each node of mother taro, and it can develop into sub taro. The axillary bud grown in the leaf axil and the leaf have a cogrowth relationship. The growth site grows clockwise, and the angle between adjacent axillary buds is 144°. The positions of each axillary bud are roughly in a straight line, showing typical 2/5 phyllodes features (Xu et al., 2020). The sub taro of Kui taro grows first and then thickens, and the top swells to form a smaller sub taro; the sub taro of taro with multi-cormels elongates and thickens, and the whole bulb also swells to form a conical shape; three small-molecular-weight specific proteins during bulb development (CSP) in the sub taro, namely, CSP2, CSP3 and CSP4, have spatiotemporal specificity and are related to the occurrence and expansion of the sub taro. The relative differences in the contents of CSP1, CSP2, and CSP4 in the mother taro are related to the apical dominance of the terminal buds of the mother taro. The greater the difference is, the more obvious the apical dominance and the lower the developmental degree of the sub taro is (Zhu et al., 2018). However, Luiz found that the expression of the gene (TC1), which encodes a

corm globulin protein(G1d) related to curculin, during bulb development is spatiotemporal globulin protein (Castro et al., 1992). The gene that determines bulb expansion is present at the beginning of bulb differentiation. The role of TC1 gene in the process of bulb expansion remains to be further studied. The proportion of fresh weight of taro first increases and then decreases in the growth process. Similarly, the water content of the bulbs first increases and then decreases, while the dry matter content gradually increases, showing an S-shaped growth trend. The dry matter accumulation of the sub taro in the later stage is greater than that of mother taro (Xu et al., 2020).

During the growth and development process of mother taro, from the perspective of component content, the weight gradually increased and the water content also gradually decreased. At the same time, the total starch content gradually increased and the soluble sugar content first increased and then decreased. From the point of Cytology, the volume of parenchyma cells continued to increase, and the internal starch content gradually increased. The number and diameter of starch bodies showed an upward trend, and the number of starch granules in a single amyloplast also increases (Zhu et al., 2017). During taro bulb expansion, the number of annual rings on the surface of the bulb gradually increased. From the point of Cytology, the volume of the parenchyma cells in the bulb continued to increase. At the same time, the number and size of amyloplasts gradually increased so that the entire cell was enriched; with the expansion of the bulb, the vascular tissue gradually developed and perfected, and the area of the sieve tube increased in number and became irregularly arranged; the taro bulb was densely covered with mucous cavities, which spread out from the center of the bulb, but no mucus cavity was observed in the epidermal cells (Sheng et al., 2021).

Amyloplast enrichment process of taro bulbs

At present, the research on the development and proliferation of amyloplasts mainly focuses on the endosperm cells of grains. In the development of endosperm amyloplasts, the number of plastids continued to divide, and the number increased; then the number and volume of starch granules increased in the plastid; individual starch granules increased in size and then began to form small amyloplasts. As the development progressed, the envelope of the small amyloplasts was further extended and expanded. Finally, the newly expanded envelope is filled with starch granules, thereby forming the large amyloplasts. The amyloplasts of the large starch granules in the wheat endosperm formed small starch granules by budding, while the amyloplasts of the small starch granules proliferated by constriction or budding (Wei et al., 2002; Wei et al., 2008).

A few reports have focused on the proliferation mode of amyloplasts in underground rhizome crops. The starch bodies of the complex starch contained in sweet potato tubers exist in the form of "diplex", "triad", and "multiple"; for the growth mode of the amylosome, single grain starch directly expands and grows; the amyloplasts of multigranular starch first split into monomers, and then grow in the form of monomers; the amyloplasts of complex starch do not divide during the growth process and form large complex amyloplasts (Jing et al., 2013). According to the morphology of amyloids observed by scanning electron microscope, the cassava root amyloid membrane is either "constricted", "wrinkled", or "sprouted", forming

multiple differentiation centers of amyloplasts; with the continuous expansion of the amyloplast, the membrane structure was degraded and disappeared, and the irregular starch granules were released (Min et al., 2010).

Generally, the accumulation process of taro bulb starch is divided into three stages, namely, starch formation, rapid starch accumulation, and amyloplast enrichment. The specific performance is the formation of amyloplasts and the continuous increase in diameter, followed by the continuous increase in the number of starches, and finally the increase in the number of starches in the amyloplasts. The surface of taro amyloplast is mostly round and oval, belonging to complex starch, containing multiple polyhedral starch granules. Take Kui taro for example. in the early stage of development, amyloid was mostly distributed at the edge of the cell, and then gathered at the center of the cell. Later in development, the number and size of amyloplasts continued to increase (Figure 4). Transmission electron microscopy showed that the amyloplasts of taro bulbs were large, and the large amyloplasts split into several small starch granules. Some small starch granules (Figure 5A) were observed close to the free state at the edge of amyloplasts, and these granules were loosely arranged and smaller in size. With the development of the bulb, the starch granules with large diameters in the amyloplasts were mostly concentrated in one area, and their arrangement was relatively compact. The small starch granules were mostly distributed on the other side or around, and their arrangement was loose (Figure 5B).

Taro bulb starch originates from the precursor plastid, and starch granules are formed in the plastid and free in the cytoplasm of the membrane (Sheng, 2021). As the starch granules increase, the volume of the plastids increases, and the starch granules develop from the original irregular shape to a round shape. In the later stage, they gradually extruded each other into polygons. The amyloplast of taro bulbs were mostly complex starches. Amyloid proliferation is divided into two stages. In the first stage, the number of starch granules in the amyloplast changes; the starch granules gradually increase, fill up, squeeze, and deform each other. The large starch granules split into many small starch granules. The second stage involves the proliferation of amyloplasts. Taro amyloplasts can be split via membrane constriction proliferation and capsular vesicle proliferation. Capsule constriction and proliferation squeeze the starch granules in the amyloplast to both sides through the inward depression of the amylosome envelope. Further, it is constricted into multiple amyloplasts, and the encapsulated vesicles proliferate in which the original amyloplasts spit out vesicles from the envelope, and new amyloplasts are generated and proliferated in the vesicles.

Effects of environment and cultivation methods on the development of taro bulbs

Taro adapts to high temperature and humidity environment and is not resistant to low temperature and frost. The optimum temperature for germination is 12–15 °C. The optimum temperature for growth is generally 25–30 °C. If the temperature is very low, the growth of taro will slow down or stop. If the temperature is very high, the condition will not be conducive to the development and expansion of taro bulb. (Wang et al., 2007) Different varieties of taro have different requirements for temperature. Taro with Multi-cormels can be planted at low

temperatures, making it widely distributed in temperate zones. Kui taro has strict requirements for high temperature. Therefore, Kui taro is mostly produced in tropical and subtropical regions with high temperature and humidity (Chang et al., 2019). Taro requires sunlight, and the light saturation point is approximately 50,000 lux. Taro is shade tolerant and can grow under scattered light. The light intensity, composition, and light time remarkably affect the growth of taro, but strong light is conducive to the growth of taro and improves yield and quality. Under blue violet light, the leaves of taro are large and thick, and the petioles are thick and short, and this condition is conducive to the growth and development of bulbs. In red and yellow light, the leaves are small, and the petioles are slender, and this condition is not conducive to the growth and development of bulbs. In the early stage of taro development, a longer light time is required to promote the increase of leaf area and the accumulation of photosynthetic products. The later stage of taro development requires a shorter light time to facilitate the formation and expansion of bulbs (Chang et al., 2019). Taro requires dampness, but there are differences between classifications. Aquatic taro is grown in paddy fields, but dry taro cannot be flooded for a long time. Taro has different requirements for humidity before different growth stages. The field should be kept moist during the germination stage to induce the germination of taro. During the plant vigorous growth stage, the water demand is large, and the water supply needs to be guaranteed; the soil should be kept dry before harvesting to maintain good condition for the harvesting and storage of bulbs (Huang et al., 2016). Taro does not require very strict soil texture. Loose and fertile soil with deep soil layer and convenient irrigation and drainage is conducive to the growth of taro and the expansion of bulbs. Taro can grow normally in soil with pH of 4.1–9.1, but the optimum pH is 5.5–7.0. A highly acidic or highly alkaline soil is not conducive to the growth and development of taro (Huang et al., 2016).

Different cultivation methods remarkably affect the yield of taro. Film mulching can provide soil temperature in the early stage of taro growth, which is beneficial to the growth of taro and the expansion of bulbs. The growth and yield of taro bulbs in perforated film-covering cultivation was better than that in ridge film-covering cultivation. However, considering the inconvenience of cultivating a large amount of soil during the growth period of taro, ridge and perforation film-covering cultivation easily form green taro, thus affecting the quality of bulbs and the taste of eating (Wang et al., 2001). If no freezing damage is observed after emergence, early sowing is conducive to the development of taro root. The plant height of taro increases, the number of taro and taro increases, and the yield increases, but these changes only slightly affect the number of taro and the shape index of taro. If the planting is late, the life cycle of the taro will be shortened, which is not conducive to the growth and development of the bulb. This condition will lead to insufficient bulb expansion, thereby reducing the yield (Zheng, 2008). Nitrogen and potassium fertilizers have obvious effects on the yield and quality of taro, and potassium fertilizer has a greater effect than nitrogen fertilizer. A significant interaction effect was observed among nitrogen, potassium, and phosphorus fertilizer. Within the reasonable range of potassium and nitrogen fertilizer application, the yield gradually increases with the increase of the amount of fertilizer. Excessive fertilizer application will reduce the yield. Phosphate fertilizer

alone only slightly affects the development of taro bulbs, and no obvious rule has been established. Reasonable fertilization is beneficial to the growth, development, and yield increase of taro bulbs (Song et al., 2004).

Regulation of hormones on bulb development

Regulation of endogenous hormones on bulb expansion and starch enrichment

Hormones is an important endogenous substance that regulates plant growth and is a key factor in bulb formation (Durbak et al., 2012). Some genes and proteins related to bulb formation are also closely related to plant hormone signaling pathways (Aksenova et al., 2012). Different plant hormones have different functions in bulb expansion, and gibberellin (GA) can inhibit or delay tuber formation (Vreugdenhil et al., 1989). Absciscic acid (ABA) does not participate in the induced metamorphosis process of tubers, but it counteracts the antagonism of other hormones (Shu et al., 2017). Auxin (IAA) can promote the metamorphic development of tubers and promote plant root development, and its concentration affects tuberous root thickening (Wang et al., 2006). Although GA and ABA are not directly related to tuber formation, they are related to the ratio of GA3/ABA. The balance of “inducing substances” and “inhibiting substances” is a key factor for tuber formation (Liu, 2001). Cytokinins (CTK) are mainly involved in the formation of tubers. Matsuo and Mitsuzono (1988) reported that the content of zeatin riboside (ZR) is significantly positively correlated with the formation and thickening of sweet potato tubers (Matsuo et al., 1988). The overexpression of CTK synthesis gene *ipt* in potato could form more tubers (Tao et al., 2010). IAA-related genes such as auxin response factor (ARFs) and Aux/IAAs are specifically expressed in early tuber development (Kloosterman et al., 2008). IAA and GA3 are necessary for potato stolon elongation. ABA and jasmonic acid (JA) are positive regulators for inducing tuber formation. GA3 is a negative regulator (Liu et al., 2019). JA and methyl jasmonate (MeJA), as classes of plant growth regulators, play an important role in tuber and bulb formation (Sarkar et al., 2006).

Plant endogenous hormone regulation is closely related to starch anabolism (Kim et al., 2005). The enlargement of plant bulbs mainly depends on starch accumulation and cell division enlargement, and starch accumulation mainly depends on sucrose synthesis and transportation. Plant hormone signal transduction affects starch accumulation. ABA can induce the expression of starch synthesis genes and enhance the transduction of sugar signals to promote starch accumulation (Akihiro et al., 2005). The level of GA at the grain filling stage of wheat is positively correlated with the final grain yield and starch yield. GA plays an important role in starch accumulation in wheat grains. Changes in endogenous hormone levels may indirectly affect starch accumulation in grains by affecting regulatory enzymes and regulatory processes (Xie et al., 2003). Scientists added IAA to MS medium, and the potato tuber starch content and starch granule size increased by 15%–30% (Gukasyan et al., 2005). In the study of tulip bulbs, IAA and ZR indirectly promoted starch accumulation by increasing the activity of ADP-glucose pyrophosphorylase (AGPase), thus catalyzing the production of a large number of products. Endogenous hormones may promote starch accumulation by participating in the starch synthesis pathway (Miao et al., 2016). Hormones have multiple roles and interact to form a regulatory

network, thereby regulating tuber development (Jung et al., 2013). In the early stage of taro development (bulb is about 1cm in diameter), the content of endogenous hormones ABA, bulb endogenous zeatin (Z), and ZR showed an upward trend, while the content of IAA, GA3, and JA showed a downward trend. In the later stage of development (bulb is about 13cm in diameter), the content of endogenous hormones ABA, IAA, Z, and GA3 showed an upward trend. The content of ZR and JA showed a downward trend, but the contents of IAA, GA3, and JA hormones were generally high during the whole development process (Sheng, 2021). Other related studies on endogenous hormones on the growth and development of taro bulbs and starch enrichment have not been conducted.

Regulation of growth, development, and starch enrichment of bulbs by exogenous hormones

The effect of exogenous hormones on the rhizome expansion of potato, sweet potato, and other potato crops has been studied. Yang (2005) used four auxins to spray potatoes (Yang, 2005). The results showed that the four auxins increased plant height and stem diameter and prolonged the photosynthetic accumulation in the later stage. This condition allowed the tubers to accumulate more organic matter during the expansion stage, thereby substantially increasing the yield. The exogenous application of IAA can promote the formation of potato stolons and the development of tubers. It is achieved by accelerating starch accumulation and starch granule enlargement, which are beneficial to the formation and development of tubers (Gukasyan et al., 2005; Roumeliotis et al., 2012). GA also promotes the occurrence of stolons. Stolons appeared on the second day after the medium containing GA3 and IAA was added, and the occurrence continued throughout the tuber setting period (Lian et al., 2002). However, the addition of GA alone could inhibit or delay the formation of potato tubers, and inactivation of the active GA gene could promote potato tuber formation (Xu et al., 1998; Roumeliotis et al., 2012). Treatment with exogenous GAs inhibited Sucrose synthase (SS) and Soluble starch synthase (SSS) activity, thus decreasing the sucrose and starch contents in tubers (Vreugdenhil et al., 1999). ABA is a promoting factor for the formation of potato tubers, and timely spraying is beneficial to potato formation (Krauss et al., 1982; Garcia et al., 2014). Varying results have been obtained about the role of ABA in the development of tuber plants. GA3 inhibits the formation of potatoes in vitro, while ABA promotes its tuber formation (Hu et al., 2017). Exogenously applied ABA can promote tuber expansion (Xu et al., 2022). However, in vitro, ABA cannot make stolons metamorphose into tubers smoothly (Yang, 2005). ABA does not participate in the induced metamorphosis process of tubers, but its presence counteracts the respective physiological activities of other hormones (Xu et al., 2022). CTK can promote potato tuber development, regulate the balance between source and sink, and participate in the transport of nutrients to storage organs (Roitsch et al., 2000). When a certain concentration of CTK is applied exogenously, the biomass of tubers remarkably increases, and the transformation of stolons to tubers is accelerated (Romanov, 2009). In vitro, CTK inhibits sucrose invertase activity but activates phosphorylase and AGPase, thereby promoting starch accumulation (Zhu et al., 2016). Therefore, CTK is an important factor in inducing tuber formation (Quan et al., 2002). The

exogenous application of JA and its derivatives can induce the swelling of the stolon top, and the content of endogenous JA increases during this process (Abdala et al., 2002). After exogenous JA treatment, the intracellular sucrose accumulates, thus increasing the osmotic pressure of the cell wall, changing the structure of the cell wall, and increasing the cell ductility. More polysaccharides such as cellulose, hemicellulose, and pectin accumulate, indicating that JA controls the expansion of the cell by regulating the synthesis of intracellular sugar (Takahashi et al., 1995). This phenomenon induces the formation of the apical meristem of potato stolon and promotes tuber development (Cenzano et al., 2003).

The development of taro bulbs is remarkably affected by exogenous hormones. The diameter of taro bulbs that were irrigated with auxin increased significantly, the weight increased, and the filling degree of amyloplasts in parenchyma cells increased. Low concentration of 6-BA can promote the development of bulbs, but it is not conducive to the enrichment of amyloplasts, and high concentrations of 6-BA have a certain inhibitory effect on the development of bulbs. GA3 promotes the elongation of the petioles of taro plants, but it does not promote the expansion of the bulbs. High concentrations of GA3 (100–200 mg/L) have an inhibitory effect on the development of taro bulbs, but it promotes the development of taro and increases the number of taro (Sheng, 2021). Limited studies have focused on the effects of exogenous hormones on the development of taro bulbs and their enrichment of amyloplasts, and further research is needed.

Role of key enzyme genes in starch synthesis during starch enrichment

In crops mainly harvesting underground storage organs, the synthesis and accumulation of starch is a complex physiological and biochemical process, which is the result of the synergistic interaction of multiple enzymes. The key enzymes of starch synthesis in root crops, such as potato and lotus root, have been widely studied. The changes in AGPase and soluble starch synthase (SSS) activities have important effects on starch synthesis in potato tubers (Tang, 2015). However, SS and AGPase can remarkably promote the synthesis of starch in the process of lotus root rhizome expansion, and their activities affect the starch content of lotus root rhizomes at the mature stage (Li et al., 2006). Based on the study of substance accumulation and changes in related enzyme activities during the development of yam, sucrose phosphate synthase activity plays a key regulatory role in the development of yam tubers and is closely related to the main functional substances (Liang et al., 2011). Based on the study of taro bulbs, AGPase activity is positively correlated with total starch content (Zang et al., 2016). With the gradual deepening of the research on starch metabolism pathways, people have new understanding of the key enzyme gene sequences and related expression regulators in the pathway.

In sweet potato, the key enzyme genes of starch synthesis such as AGPase and SS have been cloned into the gene sequence. The expression and regulation of these genes have been studied, and these key enzyme genes play a key role in the sweet potato starch metabolism pathway (Tang et al., 2011). The genes controlling sweet potato starch synthesis include granule bound starch synthase (GBSS) gene I, SSS genes I and II, starch branching enzyme (SBE) genes

I and II, starch debranching enzyme (DBE) gene, AGPase gene A/B/C, sucrose synthase (SS) genes I and II, and isoamylase (ISA) gene (Kim et al., 2009).

GBSS I is a key enzyme that controls starch synthesis, and it catalyzes the synthesis of amylose. Otani et al. (2007) interfered with the expression of GBSS I by RNAi technology to make sweet potato taste more glutinous. SSSII can affect the structure of amylopectin and reduce the gelatinization temperature of starch (Otani et al., 2007). The reduction of starch gelatinization temperature is conducive to simplifying the starch hydrolysis process and reducing the production cost of starch fermentation (Takahata et al., 2010). AGPase improves the starch content of potato tubers (Song et al., 2005). However, the synergistic expression of starch synthesis related genes under exogenous sucrose treatment promotes the conversion of sucrose to starch (Ahn et al., 2010). Peak synthase has been widely studied, but no direct research has been conducted on taro starch synthase.

Conclusions and future direction

The development of taro bulbs is a complex biochemical process, including the accumulation of morphogenesis and assimilation products, involving gene expression, material metabolism, nutrient input, and the effect of external environmental conditions (Figure 6). Limited studies have been conducted in the world. Understanding the development process, expansion mechanism, and regulation mechanism of taro bulbs has a guiding role in the production of taro and is important to ensuring food security and responding to food crises. At present, the research on the development of taro mainly focuses on evolutionary classification, genotype and isozyme analysis, cultivation, production, processing, and utilization. Limited research has been conducted on the physiological mechanism and hormone regulation pathway of taro growth and development, taro bulb expansion, key gene expression, and starch enrichment.

Therefore, the following research will become the key research direction in the future.

Expansion of taro bulbs and the regularity of the development and spatial distribution of starch bodies

Starch is the main storage material of taro bulbs, and amyloplasts are the organelles that synthesize and accumulate starch. The development of amyloplasts determines the yield and quality of taro. Limited studies locally and abroad have focused on the development of taro corm and amyloplast, and they remain in the preliminary stage. The fine structure observation of amyloplast development, its proliferation mode, and the spatial distribution characteristics of taro corm amyloplast are not clear. In the future, the occurrence, division, proliferation, and enrichment of amyloplasts in parenchyma cells of different types of taro bulbs, the differences of physical and chemical properties of taro starch at different development stages, and the development and enrichment characteristics of amyloplasts in different spatial parts should be focused on.

Role of key enzyme genes in starch synthesis in starch enrichment

Starch is the main storage material of taro bulbs. The expansion process of taro is closely related to the synthesis of starch, and the genes related to starch synthesis are closely related to starch synthesis, which directly determine the starch content of taro. The research on starch

synthase gene has remarkably progressed in wheat, rice, potato and other crops, but limited research has been conducted on taro starch synthase gene. Therefore, the differences in the expression of key enzyme genes (e.g., AGPase, GBSS, SSS, and SBE) for taro starch synthesis, the roles of these genes in regulating the starch enrichment process, and the exploration of individual gene functions will become the focus of research.

Regulation of hormones on bulb swelling and starch enrichment

The expansion of taro bulbs mainly depends on the increase in the number and volume of parenchyma cells, and this process results from the synergistic action of various hormones, especially IAA, GA, CTK, and other hormones. The changes of hormones during bulb development, the relationship between hormones and bulb expansion, and the relationship between the expression of hormone synthesis genes and signal transduction related genes and taro starch enrichment need to be investigated.

Hormone-regulated pathways promoting bulb expansion and starch enrichment

Exogenous hormones and plant growth regulators have important regulatory effects on taro bulb swelling, starch enrichment, and yield increase. The effects of exogenous substances such as 6-BA, 2,4-D, GA₃, PP333, and 5-aminolevulinic acid (5-ALA) on the development, yield, and quality of taro corn, as well as the type and concentration of the best exogenous hormone to promote corm expansion and starch enrichment need to be studied, and plant growth regulators for increased yield and improved quality of taro should be developed to provide an important theoretical basis for taro production.

In a word, the development of taro bulbs still requires a lot of research. With the deepening of research and the solution of key problems, the production of taro will continue to improve, and the development and utilization of taro will be more efficient.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

E.J.Z., F.X. and W.Y.S. conceived the outline of the manuscript.

E.J.Z. and F.X. wrote the manuscript.

W.Y.S., W.J.J., W.L.L., X.R.Y., X.P.W. provided revisions.

All authors read and approved the final manuscript.

Data Avail

The following information was supplied regarding data availability:

There is no raw data or code in this literature review.

Ability Statement

All data were collected from the published research papers.

References

- Abdala G, Castro G, Miersch O, Pearce, D. 2002. Changes in jasmonate and gibberellin levels during development of potato plants (*Solahum tuberosum* L.). *Plant Growth Regulation*.36:121-126.
- Ahn YO, Kim S.H, Kim CY, Lee JS, Kwak S, Lee H. 2010. Exogenous sucrose utilization and starch biosynthesis among sweetpotato cultivars. *Carbohydrate Research*. 345(1): 55-60.
- Akihiro T, Mizuno K, Fujimura F. 2005. Gene expression of ADP-glucose pyrophosphorylase and starch contents in rice cultured cells are cooperatively regulated by sucrose and ABA. *Plant Cell Physiology*. 46(6):937-946.
- Aksenova NP, Konstantinova TN, Golyanovskaya SA, Sergeeva LI, Romanov GA. 2012. Hormonal regulation of tuber formation in potato plants. *Russian Journal of Plant Physiology*. 59(4):451-466.
- Castro LAD, Carneiro M, Neshich DDC, Paiva GRD. 1992. Spatial and temporal gene expression patterns occur during corm development. *Plant Cell*. 4:1549-1559.
- Cenzano A, Vigliocco A, Kraus T, Abdala G. 2003. Exogenously applied jasmonic acid induces changes in apical meristem morphology of potato stolons. *Annals of Botany*. 91:915-919.
- Chang L, Wang X. 2019. Overview of Development Status of Taro Industry in the World. *Modern Agricultural Science and Technology*. 2(3):57-59.
- Dai L, Qiu C, Xiong L, Sun Q. 2015. Characterisation of corn starch-based films reinforced with taro starch nanoparticles. *Food Chemistry*. 174:82-88.
- Durbak A, Yao H, McSteen P. 2012. Hormone signaling in plant development. *Current Opinion in Plant Biology*.15(1):92-96.
- Falade KO, Okafor CA. 2013. Physicochemical properties of five cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) starches. *Food Hydrocolloids*. 30(1):173-181.
- Garcia MNM, Stritzler M, Capiati DA. 2014. Heterologous expression of Arabidopsis ABF4 gene in potato enhances tuberization through ABA-GA crosstalk regulation. *Planta*. 239:615-631.
- Gukasyan IA, Golyanovskaya SA, Grishunina EV, KKonstantinova TN, Romanov GA. 2005. Effect of Rol Transgenes, IAA, and Kinetin on Starch Content and the Size of Starch Granules in Tubers of In Vitro Potato Plants. *Russian Journal of Plant Physiology*. 52(6):809-813.
- Han X, Zhang DX, Wang L, Li Q. 2018. Research Progress on the Nutrition Components and Processing and Utilization of Taro. *China Fruit Vegetable*. 38(3):9-13.
- Hu Y, Zhan S, Ku WZ, Wang R, Xiao L. 2017. The Roles of ABA and GA3 in the Joint Regulation of Potato Tuber Formation and Starch Accumulation in Vitro. *Mol. Plant Breeding*.15(10):4210-4214.
- Huang XF, Peng J, Ke WD, Liu YM, Sun YL, Dong HX, Li F, Liu YP, Zhu HL, Li SM, Ye YY, Huang LC, Li MH, Wang Y, Zhong L, Zhou K. 2014. Analysis on Quality Characters of 206 Taro [*Colocasia esculenta* (Linn.)Schott] Germplasm Resources. *Journal of Plant Genetic Resources*. 15(3):519-525.
- Huang XF, Ke WD, Sun Y. 2016. High yield and high efficiency cultivation of high quality taro, 1st ed.; China Agriculture Press, China, pp.22-30.
- Huang XF, Ke WD. 2006. Descriptors and Data Standard for Taro [*Colocasia esculenta* (L). *Schott*], 1st ed.; China Agriculture Press, China.
- Ivancic A, Lebot V. 1999. Descriptors for Taro [*Colocasia esculenta* (L). *Schott*], Rome, Italy: International Plant Genetic Resources Institute.
- Jing YP, Li DL, Liu DT, Yu XR, Hu ML, Gu YJ, Wang Z. 2013. Anatomical Structure of the Tuberous Root Growth and Its Amyloplast Development in Sweet Potato. *Acta Botanica Boreali-Occidentallia Sinica*.33(12):2415-2422.
- Jung JKH, McCouch S. 2013. Getting to the roots of it: genetic and hormonal control of root architecture. *Frontiers Plant Science*.4:186.
- Kim KJ, Kim KS. 2005. Changes of endogenous growth substances during bulb maturation after flowering in *Lilium* oriental hybrid ‘Casa Blanca’. *Acta Horticulturae*. 570:661–667.
- Kim T, Goo Y, Lee C, Lee B, Bae, J, Lee S. 2009. The sweetpotato ADP-glucose pyrophosphorylase gene (ibAGP1) promoter confers high-level expression of the GUS reporter gene in the potato tuber. *Comptes Rendus Biologies*. 332(10):876-885.
- Kloosterman B, Koeyer DD, Griffiths R, Flinn B, Bachem C. 2008. Genes driving potato tuber initiation and growth: identification based on transcriptional changes using the POCI array. *Functional and Integrative Genomics*. 8(4):329-340.

- Krauss A, Marschner H. 1982. Influence of nitrogen nutrition d aylength and temperature on contents of gibberellic and abscisic acid and on tuberization in potato plants. *Potato Research*. 25:13-21.
- Li LJ, Zhang XD, Pan EC, Sun L, Xie K, Li G, Cao BS. 2006. Relationship of Starch Synthesis with It's Related Enzymes' Activities During Rhizome Development of Lotus. *Scientia Agricultura Sinica*. 39(11):2307-2312.
- Liu J. 2001. Formation Mechanism of Potato Tuber in Vitro and Regulation of Tuber Formation. Doctor's Thesis, Huazhong Agric. University, Wuhan, Hubei, China.
- Liu D, Xu M, Hu Y, Wang R, Tong J, Xiao L. 2019. Dynamic Changes of KeyPlant Hormones during Potato Tuber Development. *Molecular Plant Breeding*. 17(6):1998-2003.
- Lian Y, Zou Y, Dong HR, Jin LP, Lin H. 2002. The Changes of Several Endogenous Phytohormones during Microtuber Formation in vitro in *Solanum tuberosum* L. *Acta Horticulturae Sinica*, 29(6):537-541.
- Liang RF, Li CZ, Zhang J, He LF, Wei BH, Gan XQ, He HY. 2011. Changes of Matter Accumulation and Relative Enzymatic Activity during Yam Tuber Development. *Acta Agronomica Sinica*. 37(5):903-910.
- Matsuo T, Mitsuzono H. 1988. Variation in the levels of major free cytokinins and free abscisic acid during tuber development of sweet potato. *Journal of Plant Growth Regulation*.7:249-258.
- Min Y, Yao Y, Wang J, Hu XW, Guo JC. 2010. Observation on the structure and development of amyloplast in early stage of cassava storage root with scanning electronic microscope. *Journal of Chinese Electron Microscopy Society*. 29(4):379-384.
- Miao YY, Zhu ZB, Guo QS, Yang XH, Liu L, Sun Y, Wang CL. 2016. Dynamic changes in carbohydrate metabolism and endogenous hormones during *Tulipa edulis* stolon development into a new bulb. *Journal of Plant Biology*. 59(2):121-132.
- Njintang YN, Scher J, Mbofung CMF. 2008. Physicochemical, thermal properties and microstructure of six varieties of taro (*Colocasia esculenta* L. Schott) flours and starches. *Journal of Food Engineering*. 86:294-305.
- Otani M, Hamada T, Katayama K, Kitahara K, Kim S, Takahata Y, Suganuma T, Shimada T. 2007. Inhibition of the gene expression for granule-bound starch synthase I by RNA interference in sweet potato plants. *Plant Cell Reports*. 26:1801-1807.
- Prajapati R, Kalariya M, Umbarkar R, Parmar S, Sheth N. 2011. *Colocasia esculenta*: A potent indigenous plant. *International Journal of Nutrition Pharmacol Neurological Diseases*. 1:90-96.
- Qi HL, Yin ZY, Zheng XX. 2015. Study Progress on Starches of Root and Tuber Plants. *Journal of Chongqing Technol Business University*. (Nat Sci Ed).32:68-71.
- Quan F, Zhang AX, Cao XW. 2002.The Role of Plant Hormones in the Formation and Development of Potato Tubers. *Chinese Potato Journal*. 1:29-32.
- Roumeliotis E, Visser RG, Bachem CW. 2012b. A crosstalk of auxin and GA during tuber development. *Plant Signaling Behavior*. 7:1360-1363.
- Roumeliotis E, Kloosterman B, Oortwijn M, Kohlen W, Bachem C. 2012a. The effects of auxin and strigolactones on tuber initiation and stolon architecture in potato. *Journal of Experimental Botany*.63:4539-4547.
- Roitsch T, Ehneß R. 2000. Regulation of source/sink relations by cytokinins. *Plant Growth Regulation*. 32:359-367.
- Romanov GA. 2009. How do cytokinins affect the cell. *Russian Journal of Plant Physiology*. 56:268-290.
- Sarkar D, Pandey SK, Sharma S. 2006. Cytokinins antagonize the jasmonates action on the regulation of potato (*Solanum tuberosum*) tuber formation in vitro. *Plant Cell, Tissue and Organ Culture*. 87(3):285-295.
- Sefa-Dedeh S, Agyir-Sackey EK. 2002. Starch structure and some properties of cocoyam (*Xanthosoma sagittifolium* and *Colocasia esculenta*) starch and raphides. *Food Chemistry*. 79(4):435-444.
- Sebnem S, Sedef NE. 2012. Production of resistant starch from taro (*Colocasia esculenta* L. Schott) corm and determination of its effects on health by in vitro methods. *Carbohydrate Polymers*. 90(3):1204-1209.
- Sheng JY. 2021. Corm Development of Taro and Its Response to Exogenous Hormones. Master's Thesis, Yangzhou University, China.
- Shu K, Zhou WG, Yang WY. 2017. APETALA 2-domain-containing transcription factors: focusing on abscisic acid and gibberellins antagonism. *New Phytologist*. 217(3):977-983.
- Song CF, Xu K. 2004. Absorption and distribution of nitrogen, phosphorus and potassium in taro. *Plant Nutrition and Fertilizing Scince*.10:403-406.
- Song BT, Xie CH, Liu J. 2005. Expression of potato sAGP gene and its effects on contents of starch and reducing sugar of transgenic potato tubers. *Entia Agricultura Sinica*. 38:1439-1446.
- Sun JD, Sun XM, Yuan ZZ. 2014. Taro Cultivation, 1st ed.; China Agricultural Press, China, pp.34-36.
- Tao GQ, Letham DS, Yong JWH, Zhang K, John PCL, Schwartz O, Wong C, Farquhar GD. 2010. Promotion of shoot development and tuberisation in potato by expression of a chimaeric cytokinin synthesis gene at normal and elevated CO₂ levels. *Functional Plant Biology*.37:43-54.

- Tang HL. 2015.** The Activity Changes of Key Potato Starch Synthase Effects on the Starch Content of Tuber. Northeast Agric. University. China.
- Tang W, Li Q, Zhang YG, Wang X, Hou M, Ma DF. 2011.** Advances on Structure and Function of Key Enzymes and Genes Involved in Starch Biosynthesis in Sweetpotato. *Mol. Plant Breeding*. 9:1479-1485.
- Takahata Y, Tanaka M, Otani M, Katayama K, Kitahara K, Nakayachi O, Nakayama H, Yoshinaga M. 2010.** Inhibition of the expression of the starch synthase II gene leads to lower pasting temperature in sweetpotato starch. *Plant Cell Reports*. 29(6):535-543.
- Takahashi K, Fujino K, Kikuta Y, Fujino K. 1995.** Involvement of the accumulation of sucrose and the synthesis of cell wall polysaccharides in the expansion of potato cells in response to jasmonic acid. *Plant Science*. 111:11-18.
- Vreugdenhil D, Struik PC. 1989.** An integrated view of the hormonal regulation of tuber formation in potato (*Solanum tuberosum*). *Physiologia plantarum*. 75(4):525-531.
- Vreugdenhil D, Sergeeva LI. 1999.** Gibberellins and tuberization in potato. *Potato Research*. 42:471-481.
- Wang WH, Li M, Song BC, Liu CL, Zhang XX. 2001.** Effect of Different Culture Way on Growth and Yield of *Colocasia esculenta* Schott. *Journal of Laiyang Agricultural College*. 18:12-14.
- Wang QM, Zhang LM, Guan YA, Wang ZL. 2006.** Endogenous Hormone Concentration in Developing Tuberous Roots of Different Sweet Potato Genotypes. *Agricultural Science China*. 5:919-927.
- Iwashina T, Konishi T, Takayama A, Fukada A, Ootani S. 1999.** Isolation and Identification of the Flavonoids in the Leaves of Taro. *Annals of the tsukuba botanical garden*. 18:71-74.
- Wei CX, Lan SY, Xu ZX. 2002.** Relation between envelope and proliferation of amyloplast in endosperm cell of rice. *Journal of Chinese Electron Microscopy Society*. 21(2):123-128.
- Wei CX, Zhang J, Zhou WD, Chen YF, Xu RG. 2008.** Ultrastructural Observation on the Development of Small Starch Granule (Compound Starch Granule) of Wheat Endosperm. *Journal of Triticeae Crops*. 25(5):804-810.
- Wu W, Chang QT, Wang A. 2021.** Research Progress of *Colocasia esculenta* Germplasm Resources in China. *Anhui Agricultural Science*. 49:4-7.
- Xu RC, Yang MY, Xu JP, Liu SD. 2020.** Preliminary Study on Dynamic Characteristics of Biological Characters in Growth and Development of Taro (*Colocasia esculenta*). *Acta Agricultural Jiangxi*. 32: 44-48.
- Xu X, Lammeren AM, Vermeer E, Vreugdenhil D. 1998.** The role of gibberellin, abscisic acid, and sucrose in the regulation of potato tuber formation in vitro. *Plant Physiology*. 117:575-584.
- Xu Z, Zhen LW, Zhang Y, Zhang SS, Guo DP. 2022.** Regulatory network of tuberization in potato (*Solanum tuberosum* L.). *Plant Physiology Journal*. 58(1):109-118.
- Xie ZJ, Jiang D, Cao WX, Dai TB, Jing Q. 2003.** Relationships of endogenous plant hormones to accumulation of grain protein and starch in winter wheat under different post-anthesis soil water statuses. *Plant Growth Regulation*. 41(2):117-127.
- Yang LJ. 2005.** Plant Growth Substances and Their Application in Potato. *Heilongjiang Agricultural Science*. 1:49-50.
- Zang YW, Jiang FL, Cheng YQ, Kong XY, Wu Z. 2016.** Study on Dynamic Changes of the Major Carbohydrate Content and the Related Enzyme Activities during the Microcorm Development of *Colocasia esculenta*. *Acta Botanica Boreali-Occidentalia Sinica*. 36(4):700-705.
- Zhang GM, Zhang DX. 1990.** The relationship between geographic distribution and ploidy level of taro. *Euphytica*. 47:25-27.
- Zhang LL, Zhu YZ, Li DP, Jiang Y, Dong B, Li F. 2020.** Effect of Heat-Modified Starch from Laiyang Taro on the Formation and Stability of Emulsions. *Food Science*. 41(6):51-57.
- Zhang Z. 1982.** Origin, evolution and classification of tar. *Jiangxi Agricultural Science & Technology*. 7:25-31.
- Zheng F. 2008.** Effects of Different Sowing Dates and Fertilization Conditions on the Growth and Yield of Red Bud Taro. *Shanghai Vegetables*. 5:69-71.
- Zhou J, Li DF, Xu J, Wang LL, Qiu YX. 2018.** Adsorption of Cationic Taro Starch for Removal of C. I. Direct Violet 1 and C. I. Disperse Blue 56 in Wastewater. *Natural Product Research and Development*. 30:1078-1084.
- Zhu XW, Cui WX, Zhang EJ, Wang LL, Yu XR, Xiong F. 2018.** A Comparative Study on Morphological and Physicochemical Properties of Starches Isolated from Mother and Filial Bulbs of 'Binglang' and 'Xiangsha'. *Acta Horticulturae Sinica*. 45(7):1314-1326.
- Zhu XW, Cui WX, Zhang EJ, Sheng JY, Xiong F. 2017.** Morphological and physicochemical properties of starches isolated from three taro bulbs. *Starch-Strke*. 70:1-2.
- Zhu YM, Luo XL, Fan WJ. 2016.** Effects of endogenous cytokinin content on starch accumulation in root tuber of cassava. *Journal of Southern Agriculture*. 4:1279-1284.
- Zou LQ, Wang CX, Kuang XJ, Li Y, Sun C. 2016.** Advance in flavonoids biosynthetic pathway and synthetic biology. *China Journal of Chinese Materia Medica*. 41(22): 4124-4128.

Figure 1

three classifications of taros in agricultural production of China

(A) Kui taro; (B) Multi-cormels taro; (C) Multi-head taro.

Zhang identified the three samples in 1982 (Zhang, 1982).




Figure 2

The development of taro according to the diameter

The taro bulb is developed by the bud and the diameter increases with the development process. From left to right, the age is 2 weeks,4 weeks,8 weeks,14 weeks and 16 weeks.



Figure 3

The taro bulb's mother taro, sub taro and grandson taro.

(A) Multi-head taro; (B) Multi-cormels taro. Mother taro: it is developed by the top bud of the seed taro. Sub taro: it is developed from the lateral bud on the mother taro. Grandson taro: it is developed from the lateral bud of the sub taro.




Figure 4

The development process of amyloplasts in the parenchyma cells of Kui taro bulbs.

(A,E) Bulb diameter 1 cm; (B,F) Bulb diameter 5 cm; (C,G) Bulb diameter 9 cm; (D,H) Bulb diameter 13 cm; (Am) amyloid; (Cy) cytoplasm.




Figure 5

Transmission electron microscope pictures of starch grains of Kui taro bulbs

(A) Bulb diameter 1 cm; The proliferation of amyloplasts, the red box shows the small starch granules after the overflow and shrinkage, and the shape has changed from irregular to spherical. (B) Bulb di-iameter 5 cm; The arrangement of small starch granules in large starch bodies, red asterisks represent large starch granules, blue asterisks represent small starch granules, two types of starch granules are distributed on both sides of starch bodies.




Figure 6

Effects of external environmental factors and internal factors on taro bulb expansion and starch enrichment.

