

Experimental modulation of external microbiome affects nestmate recognition in harvester ants (*Pogonomyrmex barbatus*)

Andy Dosmann, Nassim Bahet, Deborah M Gordon

Social insects use odors as cues for a variety of behavioral responses, including nestmate recognition. Past research on nestmate recognition indicates cuticular hydrocarbons are important nestmate discriminators for social insects, but other factors are likely to contribute to colony-specific odors. Here we experimentally tested whether external microbes contribute to nestmate recognition in red harvester ants (*Pogonomyrmex barbatus*). We changed the external microbiome of ants through topical application of either antibiotics or microbial cultures. We then observed behavior of nestmates when treated ants were returned to the nest. Ants whose external microbiome was augmented with microbial cultures were much more likely to be rejected than controls, but ants treated with antibiotics were not. This result is consistent with the possibility that external microbes are used for nestmate recognition.

1 Title Page: Experimental modulation of external microbiome affects nestmate recognition in
2 harvester ants (*Pogonomyrmex barbatus*)

3 Dr. Andy Dosmann*
4 Department of Natural Sciences
5 Minerva Schools at Keck Graduate Institute
6 San Francisco, CA

7 Nassim Bahet
8 Stanford University Online High School
9 Stanford, CA

10 Dr. Deborah M. Gordon
11 Department of Biology
12 Stanford University
13 Stanford, CA

14 * - Corresponding author information:
15 24 Seymour St., Middlebury, VT 05753, USA
16 dosmann1@gmail.com

17 Experimental modulation of external microbiome affects nestmate recognition in harvester ants
18 (*Pogonomyrmex barbatus*)

19 Dosmann, A, Bahet, N., and Gordon, D. M.

20 ABSTRACT

21 Social insects use odors as cues for a variety of behavioral responses, including nestmate
 22 recognition. Past research on nestmate recognition indicates cuticular hydrocarbons are important
 23 nestmate discriminators for social insects, but other factors are likely to contribute to colony-
 24 specific odors. Here we experimentally tested whether external microbes contribute to nestmate
 25 recognition in red harvester ants (*Pogonomyrmex barbatus*). We changed the external
 26 microbiome of ants through topical application of either antibiotics or microbial cultures. We then
 27 observed behavior of nestmates when treated ants were returned to the nest. Ants whose external
 28 microbiome was augmented with microbial cultures were much more likely to be rejected than
 29 controls, but ants treated with antibiotics were not. This result is consistent with the possibility
 30 that external microbes are used for nestmate recognition.

31 *Keywords: Nestmate recognition, microbiome, harvester ants*

32 INTRODUCTION

33 In ants, individuals discriminate between nestmates and non-nestmates, which is often equivalent
 34 to kin discrimination, by using chemical cues (Vander Meer and Morel 1998; van Zweden and
 35 d’Ettorre 2010; Sturgis and Gordon 2012). Cuticular hydrocarbons (CHCs) are chemical

compounds that serve a variety of functions in social insects but very often are implicated in nestmate recognition by both empirical and theoretical research (Wagner et al. 2000, Howard and Blomquist 2005). CHCs are synthesized from lipids, and the CHC profile of an ant, which consists of various classes of alkanes, alkenes and methyl-branched hydrocarbons in differing proportions, can be both task and colony specific (Sturgis and Gordon 2012). In *Cataglyphis niger*, when hydrocarbons are isolated from the postpharyngeal gland and applied to one of a pair of live ants, the untreated ant behaves in a way that indicates CHCs function as nestmate discriminators (Lahav et al. 1999). Esponda and Gordon (2015) recently proposed a model of distributed nestmate recognition in which each ant in a colony has a flexible decision boundary in multidimensional odor space, acquired through interactions, that divides nestmates from non-nestmates.

For good reason, research on nestmate recognition in social insects has focused on CHCs, but other chemicals are also likely to play a role in nestmate recognition (Sturgis and Gordon 2012). Davis et al. (2013) reviewed studies showing that volatile chemicals emitted by microbes affect insect behavior ranging from foraging to oviposition. Lizé et al. (2014) showed that bacteria influence kin recognition in *Drosophila melanogaster*. Although CHCs alone are sufficient for nestmate recognition in some species of ants (Lahav et al. 1999; Wagner et al. 2000), this does not preclude a role for microbes. Both internal and external microbial communities are colony-specific in ants (Anderson et al. 2013; Hu et al. 2014), and thus potential nestmate discriminators. Over ten years ago, Matsuura (2001) experimentally showed that bacteria mediate nestmate recognition in termites (*Reticulitermes speratus*). However, the role of microbes in nestmate recognition of ants has not been investigated, except for a study by Richard et al. (2007) showing that ants from one colony of fungus growing ants (*Acromyrmex* spp.), if reared on a second colony's fungus, experienced less aggression when introduced to the second colony.

Here we test the hypothesis that the microbial composition on an ant's cuticle affects whether it is treated as a nestmate. Both pathogenic and protective microbes inhabit the surface of insect cuticles (Douglas 2015), and they could produce chemicals that affect nestmate recognition (Davis et al. 2013). Previous work in honey bees (*Apis mellifera*) showed that internal injection of bacteria caused significant changes in CHC profiles, and in turn nestmate recognition (Richard et al. 2012), but here we isolate the effect of external microbes on nestmate recognition. To test the hypothesis that the external microbiome affects nestmate recognition in ants, we experimentally modulated the microbes on the cuticles of red harvester ants (*Pogonomyrmex barbatus*) and tested whether those ants were rejected by their nestmates. *P. barbatus* can distinguish between nestmates and non-nestmates using CHCs alone (Wagner et al. 2000), but microbes may still play a role. We predicted that if the external microbiome plays a role in nestmate recognition, individuals with an altered external microbiome would not be accepted as nestmates when returned to their nest.

METHODS

Study species

Experiments were conducted in 3 queenright, brood-producing laboratory colonies of red harvester ants (*Pogonomyrmex barbatus*) obtained from a field site near Rodeo, NM. Details of lab conditions are found in Gordon et al. (2005).

Assay of nestmate recognition

On the day of the experiment, we first removed a group of ants from their nest's foraging arena ~4 hours prior to behavioral testing. Group size (9-20 ants) depended on the number of ants active in the foraging arena during sampling. Each ant then received one of five experimental or control treatments, described below, and was painted with a color corresponding to its treatment for identification during behavioral observation (Gordon et al. 2005). We then returned the group of treated and marked ants to a location near the nest entrance, where they interacted with resident nestmates. We performed the test with groups of ants at the nest entrance, rather than paired individuals in a foreign arena, because recent models indicate that accurate nestmate recognition occurs at the colony level (Johnson et al. 2011; Esponda and Gordon 2015) and empirical work indicates that testing ants in groups provides results that are more consistent and indicative of natural responses than tests with individuals (Roulston et al. 2003). An observer blind to the treatment groups then scored whether each ant was rejected from the nest during the following 20 minutes. Rejection, which was often preceded by aggressive acts like biting, consisted of a resident ant seizing a treated ant and carrying it out of the nest or away from the nest entrance towards the midden pile at the far end of the foraging arena.

Experimental treatment

We manipulated the external microbiome of ants ($N = 111$) using five different treatments: 1) Antibiotic group: To decrease the external microbiome, we coated each ant's body surface with a 1% rifampin/sterile water solution. 2) Microbial group: To increase the external microbiome, we coated each ant's body surface with a solution of Lysogeny Broth (LB) broth (Becton Dickinson, Franklin Lakes, NJ) that had previously been cultured from nestmates. Because only a small proportion of the microbiome survives in culture (Rappe and Giovannoni 2003), this solution modified and augmented an ant's external microbiome, and we then confirmed this change (see

below). In the control groups, we treated ants with 3) Water: sterile water, 4) Broth: sterile LB broth, or 5) Nest: simply removed ants from the nest and marked them before return.

Validation of treatment

To validate that our treatment produced the intended effect on the ants' external microbiome, we used the method of Ren et al. (2007) to culture the external microbiome of additional ants (N = 35). Although only a small proportion of microbial species can be cultured (Rappe and Giovannoni 2003), we used differences in the amount of culturable microbes obtained, measured by counting the number of microbial colony forming units (CFUs) that grew on LB agar plates, to indicate effectiveness of the treatments. A greater number of CFUs indicates a greater amount of external microbes on an ant.

Effect of treatment on survival

To determine whether the treatments had any beneficial or adverse effects, we treated additional ants (N = 72) and kept them in Petri dishes with water for two weeks. We checked on individuals once daily and recorded whether the ant was alive or dead.

Statistical analysis

To control for non-independence produced by colony membership and returning ants in groups (i.e. block design), we used a generalized linear mixed-effects model fitting block and colony as random effects, with treatment (Microbial, Antibiotic, LB Broth, Water, Nest) as a categorical predictor and rejection as a binary dependent variable. We used the *lmer* function in the *lme4*

package in R 3.0.2 for linear mixed-effects models, and the *multcomp* package to make Tukey post hoc comparisons of treatment effects.

Because the counts of CFUs were nonparametrically distributed, we used Wilcoxon Rank Sum Tests to assess the treatment validation. We made planned comparisons that would indicate treatment efficacy (Antibiotic vs. Broth, Water, and Nest; Microbial vs. Broth, Water, and Nest), correcting alpha for the false discovery rate according to Benjamini and Hochberg (1995).

To determine whether an experimental treatment significantly increased or decreased survival, we used a Cox proportional hazards model with the *survival* package in R (Crawley 2007), and used the *multcomp* package to make Tukey post hoc comparisons of treatment effects.

RESULTS

The experimental treatment modulated ants' external microbiome, as measured by number of colony forming units. The topical application of the microbial culture significantly increased in the number of CFUs obtained from an ant's cuticle compared to each of the controls (Microbial vs. Broth: $W = 49$, $P = 0.002$, $N = 10$, vs. Water: $W = 40.5$, $P = 0.006$, $N = 9$, vs. Nest: $W = 49$, $P = 0.002$, $N = 10$). Antibiotic treatment significantly decreased the number of CFUs compared to controls, except for the water treatment (Antibiotic vs. Broth: $W = 7.5$, $P = 0.035$, $N = 10$; vs. Water: $W = 13.5$, $P = 0.317$, $N = 9$; vs. Nest: $W = 7$, $P = 0.026$, $N = 10$). With 5 of 6 comparisons $P < 0.05$, our alpha, controlled for false discovery rate, is 0.042.

Changes in an ant's external microbiome influenced whether it was rejected from its nest by nestmates (Figure 1). Ants with experimentally augmented external microbiomes were more

likely to be rejected than ants treated with broth alone, sterile water, or only marked (all post hoc comparisons $|Z| > 3.8$, $P < 0.001$). Ants treated with the antibiotic solution were not more likely to be rejected than ants treated with broth alone, sterile water, or only marked (all post hoc comparisons $|Z| < 1.1$, $P > 0.80$).

Ants in the Microbial group died significantly sooner than ants given the water treatment ($Z = -3.085$, $P = 0.017$). There were no significant differences in survival between the Microbial group and the other control groups (all post hoc comparisons $|Z| < 1.64$, $P > 0.46$), or between the Antibiotic group and the control groups (all post hoc comparisons $|Z| < 1.173$, $P > 0.76$).

DISCUSSION

We found that as we predicted, ants treated with additional microbes were rejected by resident colony members as if they were non-nestmates. This aligns with our predictions and supports our hypothesis that the external microbiome plays a role in nestmate recognition in *P. barbatus*. Contrary to predictions, ants treated with the antibiotic solution were not rejected by resident colony members. Because the antibiotic treatment was delivered in a sterile water solution, ants' CHC profiles were likely unaltered, in which case many familiar odors would persist. This is consistent with the possibility that the presence of a foreign odor is more likely to lead to the recognition of a non-nestmate than absence of a familiar odor (van Zweden and d'Ettorre 2010; Esponda and Gordon 2015), as well as experimental data showing that carpenter ants (*Camponotus herculeanus*) reject non-nestmates by recognizing novel CHCs (Guerrieri et al. 2009).

The rejection of ants with augmented external microbiomes could be interpreted as a form of social immunity (Cremer et al. 2007), in which ants with an unfamiliar or unusual microbiome are perceived as sick nestmates in need of quarantine. Social immunity may involve a microbially-mediated form of nestmate recognition, since a microbe that causes an individual to be placed on the ‘reject’ side of another ant’s boundary may or may not be a pathogen. In this way, social immunity could constitute an additional function of microbially-mediated nestmate recognition. We found a significant difference in survival between ants from the Microbial group and those treated with water, but not between ants from the Microbial group and those from the Broth and Nest groups. Thus, the extent to which the microbial treatment may have included pathogens is not resolved and we do not know the extent to which rejection of treated ants contributed to social immunity.

In addition to further research on social immunity, our results suggest several research directions that may provide insight on the role of microbes in chemical communication. First, it will be important to learn what microbial species are key to nestmate recognition and what chemicals they produce. In hyenas, fermentative bacteria appear to play a key role in chemical communication (Theis et al. 2013). Second, it will be important to establish how much microbial communities differ among colonies, and how consistent those communities are over time. In leafcutter ants (*Acromyrmex echinator*), the external microbiome is colony-specific and differences are maintained up to 10 years, even after colonies were kept in laboratory conditions (Anderson et al. 2013). Third, an ant’s microbiome may be task specific, as well as colony-specific. In *P. barbatus*, CHCs are task specific (Sturgis and Gordon 2013) and influence task allocation (Greene and Gordon 2003). It is likely that microbial communities of ants vary with task type, as foragers are exposed to different microbes from brood care or midden workers

(Cremer et al. 2007). Microbes may also inform task decisions alongside CHCs, or could directly affect the structure of CHCs.

In conclusion, our results support a role for microbes in nestmate recognition in harvester ants. They align with experimental data showing microbes influence nestmate/non-nestmate recognition in termites (Matsuura 2001) and leaf-cutting ants (Richard et al. 2007), and show that external microbes alone can impact nestmate recognition. This finding augments extensive evidence of CHC-mediated nestmate recognition in social insects (Vander Meer and Morel 1998; van Zweden and d'Ettorre 2010; Sturgis and Gordon 2012), and calls for further investigation of the role of microbes in chemical communication of social insects.


REFERENCES

- Anderson SB, Hansen LH, Sapountzis P, Sorensen SJ, Boomsma JJ (2013) Specificity and stability of the *Acromyrmex-Pseudonocardia* symbiosis. *Mol Ecol* 22:4307-4321
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc* 57:289-300
- Crawley MJ (2007) *The R Book*. John Wiley & Sons.
- Cremer S, Armitage SAO, Schmid-Hempel (2007) Social Immunity. *Curr Biol* 17:R693-R702
- Davis TS, Crippen TL, Hofstetter RW, Tomberlin JK (2013) Microbial Volatile Emissions as Insect Semiochemicals. *J Chem Ecol* 39:840-859
- Douglas AE (2015) Multiorganismal Insects: Diversity and Function of Resident Microorganisms. *Ann Rev Entomol* 60:17-34

- 204 Esponda F, Gordon DM (2015) Distributed nestmate recognition in ants. *Proc R Soc B* 282:1806
- 205 Gordon DM, Chu J, Lillie A, Tissot M, Pinter N (2005) Variation in the transition from inside to
- 206 outside work in the red harvester ant *Pogonomyrmex barbatus*. *Insect Soc* 52:212-217.
- 207 Greene MJ, Gordon DM (2003) Social insects: Cuticular hydrocarbons inform task decisions.
- 208 *Nature* 423:32.
- 209 Guerrieri FJ, Nehring V, Jorgensen CG, Nielsen J, Galizia CG, d'Ettorre P (2009) Ants recognize
- 210 foes not friends. *Proc R Soc B* 276:2461-2468.
- 211 Howard RW, Blomquist GJ (2005) Ecological, behavioral, and biochemical aspects of insect
- 212 hydrocarbons. *Ann Rev Entomol* 50:371-393
- 213 Hu Y, Lukasik P, Moreau CS, Russell JA (2014) Correlates of gut community composition across
- 214 an ant species (*Cephalotes varians*) elucidate causes and consequences of symbiotic
- 215 variability. *Mol Ecol* 23:1284-1300
- 216 Johnson BR, van Wilgenburg E, Tsutsui ND (2011). Nestmate recognition in social insects:
- 217 overcoming physiological constraints with collective decision making. *Behav Ecol*
- 218 *Sociobiol* 65:935-944.
- 219 Lahav S, Soroker V, Hefetz A, Vander Meer RK (1999) Direct Behavioral Evidence for
- 220 Hydrocarbons as Ant Recognition Discriminators. *Naturwissenschaften* 86:246-249
- 221 Lizé A, McKay R, Lewis Z (2014) Kin recognition in *Drosophila*: the importance of ecology and
- 222 gut microbiota. *ISME J* 8:469-477
- 223 Matsuura K (2001) Nestmate Recognition Mediated by Intestinal Bacteria in a Termite,
- 224 *Reticulitermes speratus*. *Oikos* 92:20-26
- 225 Rappe MS, Giovannoni SJ (2003) The uncultured microbial majority. *Ann Rev Microbiol*
- 226 57:369-394
- 227 Ren C, Webster P, Finkel SE, Tower J (2007) Increased Internal and External Bacterial Load
- 228 during *Drosophila* Aging without Life-Span Trade-Off. *Cell Metab* 6:144-152.

- 229 Richard FJ, Poulson M, Hefetz A, Errard C, Nash DR, Boomsma JJ (2007) The origin of the
230 chemical profiles of fungal symbionts and their significance for nestmate recognition in
231 *Acromyrmex* leaf-cutting ants. Behav Ecol Sociobiol 61:1637-1649.
- 232 Roulston TH, Buczkowski G, Silverman J (2003) Nestmate discrimination in ants: Effect of
233 bioassay on aggressive behavior. Insectes Sociaux 9:73-85.
- 234 Sturgis SJ, Gordon DM (2012) Nestmate recognition in ants (Hymenoptera: Formicidae): a
235 review. Myrmecological News 16:101-110
- 236 Sturgis SJ, Gordon DM (2013) Aggression is task dependent in the red harvester ant
237 (*Pogonomyrmex barbatus*). Behav Ecol 24:532-539
- 238 van Zweden JS, d'Ettorre P (2010) Nestmate recognition in social insects and the role of
239 hydrocarbons. Insect hydrocarbons: biology, biochemistry and chemical ecology 11:222-
240 243
- 241 Theis KR, Vankataraman A, Dycus JA, Koonter KD, Schmitt-Matzen EN, Wagner AP, Holekamp
242 KE, Schmidt TM (2013) Symbiotic bacteria appear to mediate hyena social odors. PNAS
243 110:19832-19837
- 244 Vander Meer RK, Morel L (1998) Nestmate Recognition in Ants. In Pheromone Communication
245 in Social Insects. Westview Press, Boulder, CO.
- 246 Wagner D, Tissot M, Cuevas W, Gordon DM (2000) Harvester ants utilize cuticular hydrocarbons
247 in nestmate recognition. J Chem Ecol 26:2245-2257

248 FIGURE



249 Figure 1 – Probability that an ant was rejected by its nestmates (i.e. not recognized as a nestmate)
 250 after manipulation of its external microbiome. Antibiotic = topical treatment with 1% rifampin
 251 solution. LB Broth = topical treatment with sterile LB broth. Microbial = topical treatment with
 252 microbes cultured from ants in LB broth. Nest = no topical treatment; individuals were only
 253 marked before return to nest. Water = topical treatment with sterile water.