

Investigating the relationship between inbreeding and life expectancy in dogs: mongrels live longer than pure breeds (#82902)

1

First revision

Guidance from your Editor

Please submit by **27 Apr 2023** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

1 Figure file(s)

5 Table file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Investigating the relationship between inbreeding and life expectancy in dogs: mongrels live longer than pure breeds

Fernando Mata ^{Corresp., 1}, Andreia Mata ²

¹ Center for Research and Development in Agrifood Systems and Sustainability, Instituto Politécnico de Viana do Castelo, Viana do Castelo, Portugal

² Department of Biosciences, Durham University, Durham, United Kingdom

Corresponding Author: Fernando Mata
Email address: fernandomata@ipvc.pt

This study aimed to investigate the establishment of a relationship between inbreeding and life expectancy in dogs. A dataset of $N = 30563$ dogs, containing information about their breed and longevity was subject to survival analysis. A Cox regression proportional hazards model was used to differentiate survivability in three groups of dogs (mongrel, cross-bred, and pure breed). The model was found significant ($p < 0.001$) with mongrel dogs surviving longer than cross-bred and both these surviving longer than pure breeds. A second Cox regression was also found significant ($p < 0.001$) differentiating the lifespan of different dog breeds, and correlating positively the hazard ratio and the Genetic Illness Severity Index for Dogs. The results show that survivability is higher in mongrel dogs followed by cross-bred with one of the ancestors only as a pure breed, and for the last pure breed dogs have the highest morbidity levels. Higher morbidity is associated with higher GISID scores, and therefore, higher inbreeding coefficients.

Investigating The Relationship Between Inbreeding And Life Expectancy In Dogs: Mongrels Live Longer Than Pure Breed

Fernando Mata¹, Andreia Mata²

¹CISAS – Center for Research in Agrifood Systems and Sustainability, Instituto Politécnico de Viana do Castelo, Viana do Castelo, Portugal

²Department of Biosciences, Durham University, Durham, United Kingdom

Corresponding Author:

Fernando Mata¹

Rua Escola Industrial e Comercial Nun'Álvares 34, Viana do Castelo, 4900-347, Portugal.

Email address: fernandomata@ipvc.pt

Abstract

This study aimed to investigate the establishment of a relationship between inbreeding and life expectancy in dogs. Using a dataset of N = 30563 dogs sourced from the VetCompass™ Program, UK, and made available by the Royal Veterinary College, University of London, containing information about their breed and longevity, was subject to survival analysis. A Cox regression proportional hazards model was used to differentiate survivability in three groups of dogs (mongrel, cross-bred, and pure breed). The model was found highly significant ($p < 0.001$) and we found that mongrel dogs had the highest life expectancy, followed by cross-bred dogs with only one purebred ancestor, and purebred dogs had the lowest life expectancy. A second Cox regression was also found highly significant ($p < 0.001$) differentiating the lifespan of different dog breeds and correlating positively the hazard ratio and the Genetic Illness Severity Index for Dogs (GISID). The results show that survivability is higher in mongrel dogs followed by cross-bred with one of the ancestors only as a pure breed, and pure breed dogs have the highest morbidity levels. Higher morbidity is associated with higher GISID scores, and therefore, higher inbreeding coefficients. These findings have important implications for dog breeders, owners, and animal welfare organizations seeking to promote healthier, longer-lived dogs.

Introduction

While dogs have been our faithful companions for thousands of years, their breeding and genetics have been the subject of growing concern in recent times. Dogs, the human's best friends, have been domesticated from the wolf during the late Pleistocene more than 15,000 years ago (Horard-Herbin et al., 2014). Since the beginning of this process, we have understood that dogs could be useful to human activities in many ways, such as hunting, guarding, herding, waste disposal, warfare, entertainment, pest control, transport, clothing, and even food (Janssens et al., 2018). Having so many potential utilitarian roles led to specialization, and humans started to shape dogs accordingly. More recently, in the past 200 to 300 years shape standards were envisaged, and dog's morphology was manipulated through closed population breeding (Axelsson et al., 2021). It was however, during the late 19th century, in Victorian Britain, that dog breeding standards started to be implemented, and the pedigree and pure blood concepts reached momentum (Worboys et al., 2018). The modern dog had been 'invented'. The modern dog has a multitude of shapes, sizes, colors, and hair types, but also behaviors and personalities, adapted to human needs. The closed population breeding of dogs is a modern practice aiming at the fixation of traits of interest (Axelsson et al., 2021). This bottlenecking of gene flow has also, however, undesirable consequences, as each individual carries deleterious genes with the potential to cause harm and affect fitness and health (Mabunda et al., 2022). Most of these genes are recessive and can affect the phenotype in homozygosity only. Inbreeding decreases the genetic load and creates in the descendants' genome long homozygous regions, increasing the potential for deleterious genes to express themselves (Bosse et al., 2019). The deleterious defects in pure-breed dogs have long ago been identified, and the number of problems identified is growing. Hodgman (1963) identified thirteen conditions, having highlighted the most important being hip dysplasia, patella luxation, entropion, retinal atrophy, and the elongated soft palate. More recently Asher et al. (2009) and Summers et al. (2010), in two companion papers, identified almost 400. At present, the Online Mendelian Inheritance in Animals (OMIA) database, (Nicholas & Tammen, 2023) identifies 856 trait disorders in dogs. The lifespan of dogs in relation to breeds and types has been presented and discussed (e.g. Patronek et al., 1997; Salvin et al., 2012) and very recently Teng et al. (2022) produced lifetables based on a large UK population above 30,000 dogs. The enormous diversity of dog breeds, ranging between the 1 kg Chihuahua and the 75 kg Saint Bernard, the 30 cm of the Dachshund and the 85 cm of the Great Dane, determines some variation in the different breeds' lifespan

(Fleming et al., 2011). Several factors of variation have been identified by the different epidemiological studies investigating lifespan in dogs: weight (Adams et al., 2010), neutering status (Moore et al., 2001), breed (Teng et al., 2022), and breed purity (Proschowsky et al., 2003). Asher et al. (2009) and Summers et al. (2010) in their two companion papers introduced the Generic Illness Severity Index for Dogs (GISID), where the severity of disorders is sensibly scored. In addition, the authors provide an estimate of these scores for the most common breeds in the UK. The authors classify the different disorders (inherited defects) as conformation related (C), conformation exacerbated (CD), and not previously linked to conformation (D). The present study used the lifespan data reported by Teng et al. (2022) to relate with the GISID scores obtained by Asher et al. (2009) and Summers et al. (2010). This study aims to investigate the establishment of a relationship between inbreeding and life expectancy in dogs. Understanding this relationship can be used in the management of breeding for gains in dogs' health and welfare. It is hypothesized that mongrel dogs may have a larger lifespan than crossbred and pure breeds based on a decrease in inbreeding coefficients, associated with carriers of deleterious genes. The findings of this study could help guide breeding practices and ultimately improve the health and welfare of dogs.

Materials & Methods

Data are open access (Creative Commons, CC BY 4.0) and were retrieved from the Royal Veterinary College, University of London, repository (O'Neill, 2022). The sample includes all dogs under primary veterinary care at clinics participating in the VetCompass™ Program, UK during 2016 (i.e., dogs with at least one clinical record in 2016) (Teng et al., 2022). The VetCompass™ Program, UK centralizes in a repository de-identified clinical records from veterinary practices in the UK (VetCompass™ Program, UK) and Australia (VetCompass™ Program, Australia). Data in this study includes dogs that had at least one clinical record in the year 2016, in The VetCompass™ Program, UK. The dog breeds recognized by any of the Kennel Club (KC), the American Kennel Club, and the Australian National Kennel Council were considered purebred, while all others were considered crosses. Crosses with ancestors not belonging to a recognized breed were considered mongrels. We have therefore obtained a lifespan data set of dogs classified as 'pure breed', cross with at least a pure breed ancestor ('cross bred'), and a cross without pure breed ancestors ('mongrel'). A total of N = 30563 entries

were found in the original dataset, which after data cleansing and elimination of outliers, resulted in a sample of N = 30470 dogs (n = 2406 mongrel, n = 3962 crossbred, and n = 24102 pure breed).

Outliers were identified after data transformation for a standard normal distribution (a normal distribution with mean with value zero and standard deviation with value one), and when their z score was above 3 or below -3 standard deviations. Data entered a Cox-regression proportional hazards model for survival analysis with 'Lifespan' as the 'time to event' variable, and 'Type of Dog' ('Mongrel', 'Cross Bred', and 'Pure Breed') as the factor to analyze. There were no censored entries, and the 'event' is 'Age at Death' (years). An ANOVA with an LSD test as post hoc was also used to differentiate means between 'Type of Dog'.

The different pure breed dogs were then compared with the inherited defects in pedigree dogs, namely the disorders related (Asher et al., 2009), and not related (Summers et al., 2010) to breed standards. At this stage, some breeds in the original dataset were eliminated from analysis once to compare with Asher et al. (2009) and Summers et al. (2010) we had to consider the breeds reported by these authors. Therefore breeds not entering the studies of Asher et al. (2009) and

Summers et al. (2010) were not considered. The new dataset comprises N = 19466 dogs distributed within breeds as shown in Table 1.

The number of identified disorders in the different breeds (inherited defects), were divided by the average scores obtained by the application of the GISID (Asher et al., 2009; Summers et al., 2010) (for C, CD, D, and Total), to obtain the variables 'Rate C', 'Rate D', 'Rate CD' and 'Rate Total'. These new variables giving an average score per disorder entered a Cox-regression proportional hazards model for survival analysis with 'Lifespan' as the 'time to event' variable, 'Breed' as a factor, and the 'Rate' variables as covariates. Again, there were no censored entries, and the event considered is 'Age at Death' (years).

The models were tested via the -2 Log likelihood test and their parameters via the Wald test. A cumulative survival plot was also produced for the first model.

Data were initially entered in a spreadsheet (Microsoft® Excel® for Microsoft 365 MSO, version 2204 Build 16. 0. 15128. 20240, 64-bit) for cleansing and outlier detection and elimination.

Descriptive statistics were also produced with this software. The Cox regression proportional hazards models were produced with the statistical package IBM Corp.® SPSS® Statistics, Armonk, NY, USA. Version: 28.0.1.1 (15).

Results

The descriptive statistics of the variables entered in the first model are shown in Table 2. The ANOVA test is highly significant ($F = 211.84$, 2df, $p < 0.001$) and all the means are significantly different ($p = 0.032$) between 'Pure Breed' and 'Cross Bred', and $p < 0.001$ for the other comparisons. The ANOVA table is presented as Table 3.

The first model differentiating 'Pure Breed', 'Cross Bred', and 'Mongrel' dogs was found to be highly significant (-2 Log Likelihood 567623, $\chi^2 = 595$, 2 df, $p < 0.001$) and the parameter is also significant (Wald 585, with 2 degrees of freedom, $p < 0.01$). The full model parameterization is shown in Table 4.

In a Cox proportional hazards model, a negative parameter is indicative of a decrease in the hazard ratio (HR) and an increase in survival in relation to the baseline. The exponential of the parameter (e^β) indicates the HR effect size and is interpreted as multiplicative effects on the hazard. Therefore, a 'Cross Bred' dog has a negative β , indicative of a decrease in the HR in relation to a 'Pure Breed' of $e^\beta = 0.838$ or 17.2%. 'Mongrel' dogs also have lower HR in relation to pure breed and the effect size is larger when compared to 'Cross Bred'. The decrease in the HR in mongrel dogs is 39%. These effects are evident in the survival functions plotted in Figure 1.

The second model differentiating dog breeds and using the 'Rates' calculated from the GISID was also found to be significant (-2 Log Likelihood 335993, $\chi^2 = 5013$ with 43 df, $p < 0.001$). The variable 'Breed' is significant (Wald 4314, with 42 df, $p < 0.01$), as it is the covariate 'Total Ratio' (Wald 64.58, with 1 df, $p < 0.001$). The full model parameterization is shown in Table 5.

In the second model, the lower the parameter associated with the breed, the higher the HR. Therefore, the breed Dogue de Bordeaux is the breed with the lower HR and higher survivability, while the Bulldog has the higher HR and the lower survivability. The breed Yorkshire Terrier is the baseline, therefore negative parameters are associated with lower HR, and positive parameters with higher HR, in relation to this breed. As the model includes over forty breeds any plot of the survival function becomes impossible to read, therefore it is not presented for this model.

The covariate 'Total Ratio' indicates that for an additional unit in the ratio, the HR is added by a factor of 17.5% while holding age constant. In other words, the higher the GISID, the higher the death hazard, therefore the lower the survivability. A negative correlation between GISID and survivability becomes, therefore, established within this model.

Discussion

An increase in dog's inbreeding coefficients is associated with a lower lifespan. In decreasing order, dogs with longer lives are 'Mongrel' (mean and 95%CI 12.761 [12.838, 12.684]), followed by 'Cross Bred' (11.225 [11.158, 11.293]), and 'Pure Bred' (11.086 [11.062, 11.110]). These results were reiterated by the Cox-regression, with a decrease in the HR in relation to pure breed dogs of 17.2% and 39% respectively for cross-bred and mongrel dogs.

Some previous studies have reported this relationship. In a study based on questionnaires filled by members of the Danish Kennel Club and representing a sample of 2928 dogs, Proschowsky et al. (2003) reported differences between mixed breed (median and IQR 11 [8, 13]) and several breeds with median varying between 7 and 11. These results were subject to a Kruskal-Wallis test and significant differences were found, however, no post hoc tests are reported and, therefore, the statement that differences between pure-breed and cross-bred dogs exist is not robust. Also, a study in Britain (Michell, 1999) using questionnaires, reports that mongrel dogs are between those with larger lifespans (median 14.0). This study, however, does not provide any inferential statistical test and anchors the statement in descriptive statistics only. Patronek et al. (1997) report statistical evidence of significant differences between mixed breeds and pure breeds (medians 8.5 and 6.7 years, respectively), using a large sample (23,535 dogs) of data collected in veterinary hospitals in the USA and Canada. Despite the high credibility of the study, there is no definition of what a mixed breed is; it could eventually be a mongrel, a cross between two pure breeds, or a cross between a mongrel and a pure breed.

Inbreeding is known by agglomerating homozygotic recessive genes in the genome of individuals (Mooney et al., 2021), and particularly in pure-breed dogs, it has been shown that higher inbreeding coefficients are associated with higher morbidity (Bannasch et al., 2021; Yordy et al., 2020), due to a larger accumulation of deleterious genes associated with the most common disorders (O'Neill et al., 2014).

The novelty in the present study is the clear differentiation between groups that are not pure breeds. This differentiation allows the definition of three groups with expected different inbreeding coefficients and therefore allows the clear establishment of a relationship between expected inbreeding coefficients and lifespan. Being pure-breed dogs bred in a closed population, they have higher inbreeding coefficients than mongrels; and cross-bred dogs with no more than one pure breed ancestor have, obviously, intermediate inbreeding coefficients. The

present study reports results based on a continuous variable allowing the comparison and the report of significantly different means in the three groups of dogs. Previous studies report median values only.

The life span differences between breeds have been the object of study and are well documented, e.g. Bannasch et al. (2021), O'Neill et al. (2013), Teng et al. (2022), and Yordy et al. (2020). The common report in lifespan differences between dog breeds is the size or weight of the breed. Larger dogs are reported to have higher morbidity, e.g. Bannasch et al. (2021), Greer et al. (2007), Michell (1999), and O'Neill et al. (2013). This trend is not evident in the present study. The limitations of this study are associated with the sample. The sample is based on dogs that attended veterinary practices in the UK, and for which both born, and death dates are known. The generalization of the results for other geographies must be done carefully. Also, not all dogs in the UK have records in veterinary practices and many that have records eventually don't show born and/or death dates. The sample may, therefore, be somehow biased.

Nevertheless, ethical dog breeding must be implemented with respect for dog welfare. Broeckx (2020) revised and elected two points of action: reduce the frequency of disorders and increase genetic diversity. For example in a study (Douglas et al., 2015) of policy comparison between the British Kennel Clube (BKC) and the German Kennel Club, Verband für das Deutsche Hundewesen (VDH) it was demonstrated that it is possible to tackle some of the most common problems faced by pure breeding by reducing the frequency of the disorder. Canine hip scoring can effectively be used in selection programs to reduce the incidence of hip dysplasia. By adopting VDH's system of mandatory hip scoring of breeding parents and only allowing those with low scores to breed, it was demonstrated that it is possible to reduce faster, hip scores in populations of pedigree dogs when compared with the BKC's voluntary system.

Conclusions

It was hypothesized that morbidity in dogs may be associated with inbreeding coefficients. The results of this study confirm the hypothesis, showing that survivability is higher in mongrel dogs followed by cross-bred with one of the ancestors only as a pure breed, and for the last pure breed dogs have the highest morbidity levels. Higher morbidity was associated with higher GISID scores, and therefore, higher levels of homozygotic recessive genes in the genomes of the individuals, and inbreeding coefficients. Future research may be directed to dog breeding, to decrease inbreeding coefficients and control deleterious genes.

References

- Adams, V. J., Evans, K. M., Sampson, J., & Wood, J. L. N. (2010). Methods and mortality results of a health survey of purebred dogs in the UK. *Journal of Small Animal Practice*, 51(10), 512-524. <https://doi.org/10.1111/j.1748-5827.2010.00974.x>
- Asher, L., Diesel, G., Summers, J. F., McGreevy, P. D., & Collins, L. M. (2009). Inherited defects in pedigree dogs. Part 1: Disorders related to breed standards. *The Veterinary Journal*, 182(3), 402-411. <https://doi.org/10.1016/j.tvjl.2009.08.033>
- Axelsson, E., Ljungvall, I., Bhoumik, P., Conn, L. B., Muren, E., Ohlsson, Å., . . . Lindblad-Toh, K. (2021). The genetic consequences of dog breed formation—Accumulation of deleterious genetic variation and fixation of mutations associated with myxomatous mitral valve disease in cavalier King Charles spaniels. *PLOS Genetics*, 17(9), e1009726. <https://doi.org/10.1371/journal.pgen.1009726>
- Bannasch, D., Famula, T., Donner, J., Anderson, H., Honkanen, L., Batcher, K., . . . Rebhun, R. (2021). The effect of inbreeding, body size and morphology on health in dog breeds. *Canine Medicine and Genetics*, 8(1), 12. <https://doi.org/10.1186/s40575-021-00111-4>
- Bosse, M., Megens, H.-J., Derks, M. F. L., de Cara, Á. M. R., & Groenen, M. A. M. (2019). Deleterious alleles in the context of domestication, inbreeding, and selection. *Evolutionary Applications*, 12(1), 6-17. <https://doi.org/https://doi.org/10.1111/eva.12691>
- Broeckx, B. J. G. (2020). The dog 2.0: Lessons learned from the past. *Theriogenology*, 150, 20-26.
- Douglas, C., Mata, F., & Menem, G. (2015/04/14-15). Hip scoring for canine hip dysplasia: A comparison of British and German breeding strategies. *Advances in Animal Biosciences, Science with Impact*. Proceedings of the British Society of Animal Science, Chester, UK.
- Fleming, J. M., Creevy, K. E., & Promislow, D. E. L. (2011). Mortality in North American Dogs from 1984 to 2004: An Investigation into Age-, Size-, and Breed-Related Causes of Death. *Journal of Veterinary Internal Medicine*, 25(2), 187-198. <https://doi.org/10.1111/j.1939-1676.2011.0695.x>
- Greer, K. A., Canterbury Sc Fau - Murphy, K. E., & Murphy, K. E. (2007). Statistical analysis regarding the effects of height and weight on life span of the domestic dog. *Research in Veterinary Science*, 82(2), 208-214. <https://doi.org/10.1016/j.rvsc.2006.06.005>

- Hodgman, S. F. J. (1963). Abnormalities and Defects in Pedigree Dogs—I. An Investigation into the Existence of Abnormalities in Pedigree Dogs in the British Isles. *Journal of Small Animal Practice*, 4(6), 447-456. <https://doi.org/10.1111/j.1748-5827.1963.tb01301.x>
- Horard-Herbin, M.-P., Tresset, A., & Vigne, J.-D. (2014). Domestication and uses of the dog in western Europe from the Paleolithic to the Iron Age. *Animal Frontiers*, 4(3), 23-31. <https://doi.org/10.2527/af.2014-0018>
- Janssens, L., Giemisch, L., Schmitz, R., Street, M., Van Dongen, S., & Crombé, P. (2018). A new look at an old dog: Bonn-Oberkassel reconsidered. *Journal of Archaeological Science*, 92, 126-138. <https://doi.org/10.1016/j.jas.2018.01.004>
- Mabunda, R. S., Makgahlela, M. L., Nephawe, K. A., & Mtileni, B. (2022). Evaluation of Genetic Diversity in Dog Breeds Using Pedigree and Molecular Analysis: A Review. *Diversity*, 14(12).
- Michell, A. R. (1999). Longevity of British breeds of dog and its relationships with-sex, size, cardiovascular variables and disease. *Veterinary Record*, 145(22), 625-629. <https://doi.org/10.1136/vr.145.22.625>
- Mooney, J. A., Yohannes, A., & Lohmueller, K. E. (2021). The impact of identity by descent on fitness and disease in dogs. *Proceedings of the National Academy of Sciences*, 118(16), e2019116118. <https://doi.org/10.1073/pnas.2019116118>
- Moore, G. E., Burkman, K. D., Carter, M. N., & Peterson, M. R. (2001). Causes of death or reasons for euthanasia in military working dogs: 927 cases (1993–1996). *Journal of the American Veterinary Medical Association*, 219(2), 209-214. <https://doi.org/10.2460/javma.2001.219.209>
- Nicholas, F., & Tammen, I. (2023). OMIA - Online Mendelian Inheritance in Animals
- O'Neill, D. (2022). Life tables of annual life expectancy and mortality for companion dogs in the United Kingdom [Mortality].
- O'Neill, D. G., Church, D. B., McGreevy, P. D., Thomson, P. C., & Brodbelt, D. C. (2013). Longevity and mortality of owned dogs in England. *The Veterinary Journal*, 198(3), 638-643. <https://doi.org/10.1016/j.tvjl.2013.09.020>
- O'Neill, D. G., Church, D. B., McGreevy, P. D., Thomson, P. C., & Brodbelt, D. C. (2014). Prevalence of Disorders Recorded in Dogs Attending Primary-Care Veterinary Practices in England. *PLOS ONE*, 9(3), e90501. <https://doi.org/10.1371/journal.pone.0090501>

- Patronek, G. J., Waters, D. J., & Glickman, L. T. (1997). Comparative Longevity of Pet Dogs and Humans: Implications for Gerontology Research. *The Journals of Gerontology: Series A*, 52A(3), B171-B178. <https://doi.org/10.1093/gerona/52A.3.B171>
- Proschowsky, H. F., Rugbjerg, H., & Ersbøll, A. K. (2003). Mortality of purebred and mixed-breed dogs in Denmark. *Preventive Veterinary Medicine*, 58(1), 63-74. [https://doi.org/10.1016/S0167-5877\(03\)00010-2](https://doi.org/10.1016/S0167-5877(03)00010-2)
- Salvin, H. E., McGreevy, P. D., Sachdev, P. S., & Valenzuela, M. J. (2012). The effect of breed on age-related changes in behavior and disease prevalence in cognitively normal older community dogs, *Canis lupus familiaris*. *Journal of Veterinary Behavior*, 7(2), 61-69. <https://doi.org/10.1016/j.jveb.2011.06.002>
- Summers, J. F., Diesel, G., Asher, L., McGreevy, P. D., & Collins, L. M. (2010). Inherited defects in pedigree dogs. Part 2: Disorders that are not related to breed standards. *The Veterinary Journal*, 183(1), 39-45. <https://doi.org/10.1016/j.tvjl.2009.11.002>
- Teng, K. T.-y., Brodbelt, D. C., Pegram, C., Church, D. B., & O'Neill, D. G. (2022). Life tables of annual life expectancy and mortality for companion dogs in the United Kingdom. *Scientific Reports*, 12(1), 6415. <https://doi.org/10.1038/s41598-022-10341-6>
- Worboys, M., Strange, J.-M., & Pemberton, N. (2018). *The invention of the modern dog: breed and blood in Victorian Britain*. Johns Hopkins University Press.
- Yordy, J., Kraus, C., Hayward, J. J., White, M. E., Shannon, L. M., Creevy, K. E., . . . Boyko, A. R. (2020). Body size, inbreeding, and lifespan in domestic dogs. *Conservation Genetics*, 21(1), 137-148. <https://doi.org/10.1007/s10592-019-01240-x>

Table 1(on next page)

Distribution by breed of the total number of dogs (N = 19172) used in the Cox-regression proportional hazards model for survival analysis of the different breeds using the Genetic Illness Severity Index for Dogs as covariate.

1

Breed	n	Breed	n
Alaskan Malamute	50	Hungarian Vizsla	44
Basset Hound	76	Irish Setter	39
Beagle	171	Labrador	2501
Bichon Frise	338	Lhasa Apso	282
Border Collie	1018	Miniature Schnauzer	211
Border Terrier	294	Poodle Miniature	102
Boxer	831	Poodle Standard	84
Bull Mastif	143	Poodle Toy	56
Bull Terrier	435	Pug	196
Bulldog	501	Rhodesian Ridgeback	68
Cairn Terrier	107	Rottweiler	505
Cavalier King Charles Spaniel	1004	Scottish Terrier	201
Chihuahua	405	Shar Pei	149
Cocker Spaniel	1063	Shetland Sheep Dog	61
Collie (Rough)	84	Shih Tzu	635
Dalmatian	149	Springer Spaniel	800
Doberman	148	Staffordshire Bull Terrier	2344
Dogue de Bordeaux	152	Tibetan Terrier	71
German Shepherd	1096	Weimaraner	129
German Short-haired Pointer	41	West Highland Terrier	1103
Golden Retriever	511	Whippet	125
Great Dane	83	Yorkshire Terrier	1060

2

Table 2 (on next page)

Descriptive statistics of the lifespan of the three types of dogs analyzed with model 1 (mongrel, cross-bred, and pure breed).

1

Statistic	Mongrel	Cross Bred	Pure Breed
Observations	2406	3962	24102
Minimum	0.008	0.003	0.003
1 st Quartile	11.006	8.909	9.126
Median	13.569	12.049	11.723
3 rd Quartile	15.198	14.350	13.722
Maximum	20.561	21.884	21.539
Mean	12.761	11.225	11.086
Standard deviation	3.762	4.276	3.729
Mean CI (95% upper)	12.838	11.293	11.110
Mean CI (95% lower)	12.684	11.158	11.062

2

3

Table 3 (on next page)

ANOVA table used to test significant differences of lifespan between types of dogs (mongrel, cross-bred and pure breed).

1

Type III Sum of					
Source	Squares	df	Mean Square	F	Sig.
Corrected Model	6140,12	2	3070,06	211,84	<.001
Intercept	1733638,69	1	1733638,69	119626,32	<.001
Dog group	6140,12	2	3070,06	211,84	<.001
Error	441531,33	30467	14,49		
Total	4294503,46	30470			
Corrected Total	447671,45	30469			

2

3

Table 4(on next page)

Parameters of the Cox proportional hazards model, modeling the survivability of three groups of dogs being studied (mongrel, cross-bred, and pure breed). The is highly significant (-2 Log Likelihood 567623, $\chi^2 = 595$, 2 df, $p < 0.001$). The parameter “group”

Pure breed is the baseline group.

1

	β	Wald	df	p-value	e^{β}	e^{β} 95% CI	
						Lower	Upper
Cross	-0.177	105.474	1	<.001	0.838	0.810	0.867
Mongrel	-0.494	527.621	1	<.001	0.610	0.585	0.637

Table 5 (on next page)

Parameters of the Cox proportional hazards model, modeling the survivability of different breeds of dogs with 'Rate Total' as a covariate. 'Rate Total' is a calculated rate based on the Genetic Illness Severity Index for Dogs. The model is highly significant

Yorkshire Terrier is the baseline breed.

1

							Exp (β) 95% CI	
		β	Wald	df	p-value	Exp(β)	Lower	Upper
Covariate	Rate Total	0.162	64.58	1	<0.001	1.175	1.130	1.223
Breed	Dogue de Bordeaux	-12.105	44.271	1	<0.001	6 ⁻⁶	1.565 ⁻⁷	1.960 ⁻⁴
	Whippet	-2.419	39.009	1	<0.001	0.089	0.042	0.190
	Rhodesian Ridgeback	-1.941	24.433	1	<0.001	0.144	0.067	0.310
	Staffordshire Bull Terrier	-1.806	40.143	1	<0.001	0.164	0.094	0.287
	Tibetan Terrier	0.483	13.633	1	<0.001	1.621	1.255	2.096
	Bull Mastiff	0.795	20.258	1	<0.001	2.214	1.566	3.129
	Shetland Sheepdog	1.546	61.335	1	<0.001	4.691	3.186	6.907
	Pug	1.599	411.244	1	<0.001	4.950	4.241	5.777
	Bull Terrier	1.629	102.357	1	<0.001	5.101	3.720	6.995
	Poodle Miniature	1.666	38.626	1	<0.001	5.292	3.129	8.950
	Poodle Toy	1.798	41.540	1	<0.001	6.037	3.494	10.429
	Labrador	1.874	57.486	1	<0.001	6.514	4.013	10.575
	West Highland Terrier	1.983	68.114	1	<0.001	7.265	4.536	11.634
	Irish Setter	2.103	78.054	1	<0.001	8.187	5.135	13.053
	Basset Hound	2.107	127.733	1	<0.001	8.225	5.707	11.853
	Cavalier King Chrls Spaniel	2.205	204.593	1	<0.001	9.071	6.705	12.270
	German Shorted-Hair Pointer	2.210	65.239	1	<0.001	9.119	5.333	15.592
	Chihuahua	2.286	182.833	1	<0.001	9.832	7.059	13.693
	Dalmatian	2.329	99.287	1	<0.001	10.270	6.495	16.239
	Springer Spaniel	2.423	85.607	1	<0.001	11.278	6.750	18.841
	Collie (Rough)	2.431	82.005	1	<0.001	11.376	6.721	19.255
	Hungarian Vizsla	2.444	82.177	1	<0.001	11.519	6.791	19.539
	Cairn Terrier	2.474	63.367	1	<0.001	11.865	6.453	21.815
	Border Collie	2.512	73.124	1	<0.001	12.334	6.935	21.938
	Shih Tzu	2.548	83.657	1	<0.001	12.779	7.402	22.059
	Poodle Standard	2.642	68.117	1	<0.001	14.036	7.496	26.285
	Bichon Frise	2.706	92.274	1	<0.001	14.963	8.615	25.987
	German Shepard	2.746	156.512	1	<0.001	15.582	10.134	23.959
	Shar Pei	2.813	349.604	1	<0.001	16.662	12.407	22.377
	Alaskan Malamute	3.012	132.368	1	<0.001	20.334	12.172	33.969
	Cocker Spaniel	3.030	96.623	1	<0.001	20.698	11.312	37.871
	Scottish Terrier	3.037	77.295	1	<0.001	20.845	10.591	41.024
	Lhasa Apso	3.346	89.761	1	<0.001	28.381	14.205	56.705
	Beagle	3.356	112.241	1	<0.001	28.661	15.406	53.320
	Golden Retriever	3.488	83.652	1	<0.001	32.730	15.499	69.118
	Miniature Schnauzer	3.747	87.736	1	<0.001	42.381	19.351	92.823
	Great Dane	3.912	235.326	1	<0.001	50.010	30.337	82.440
	Doberman	3.935	162.363	1	<0.001	51.146	27.923	93.682
	Boxer	3.983	130.334	1	<0.001	53.691	27.096	106.387
	Weimaraner	4.088	99.726	1	<0.001	59.640	26.734	133.048
	Rottweiler	4.247	151.288	1	<0.001	69.930	35.540	137.596
	Bulldog	4.446	201.831	1	<0.001	85.273	46.178	157.466

2

Figure 1

Representation of the survival functions of the Cox proportional hazards model, modeling the survivability of three groups of dogs being studied (mongrel, cross-bred, and pure breed).

