

# Identification of a pathogen causing fruiting body rot of *Sanghuangporus vaninii*

Weidong Yuan<sup>1,\*</sup>, Lin Ma<sup>2</sup>, Xingkun Chen<sup>3</sup>, Jiling Song<sup>1</sup> and Qing Chen<sup>4,\*</sup>

<sup>1</sup> Hangzhou Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

<sup>2</sup> Jiangsu Key Laboratory for Horticultural Crop Genetic Improvement, Institute of Vegetable Crops, Jiangsu Academy of Agricultural Sciences, Nanjing, Jiangsu, China

<sup>3</sup> College of Agronomy, Shandong Agricultural University, Taian, Shandong, China

<sup>4</sup> Zhejiang Agricultural Technology Extension Center, Hangzhou, Zhejiang, China

\* These authors contributed equally to this work.

## ABSTRACT

*Sanghuangporus vaninii* is a medicinal macrofungus that is increasingly cultivated in China. During cultivation, it was found that the fruiting body of *S. vaninii* was susceptible to pathogenic fungi, resulting in significant economic losses to the industry. The symptoms of the disease occur in the initial stage of fruiting body development. The isolate YZB-1 was obtained from the junction of the diseased and healthy areas of the fruiting body. In order to verify the pathogenicity of YZB-1, its purified spore suspension was inoculated into the exposed area nearby the developing fruiting body of *S. vaninii*. After 10 days, the same disease symptoms appeared in the inoculated area. Morphological identification and molecular analysis of rDNA ITS region confirmed that the isolate YZB-1 was identified as *Trichoderma virens*. The temperature stability assay revealed that the mycelia of YZB-1 grew the fastest at 25 °C, with growth slowing down gradually as the temperature increased or decreased. Dual-culture tests of *T. virens* and *S. vaninii* showed that the inhibition rate of *T. virens* on *S. vaninii* mycelium was the highest (79.01 ± 2.79%) at 25 °C, and more green spores were produced at the intersection of *T. virens* and *S. vaninii*.

Submitted 5 April 2023

Accepted 8 August 2023

Published 4 September 2023

Corresponding author

Lin Ma, 20090029@jaas.ac.cn

Academic editor

Aristóteles Góes-Neto

Additional Information and  
Declarations can be found on  
page 9

DOI 10.7717/peerj.15983

© Copyright  
2023 Yuan et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Agricultural Science, Microbiology, Mycology

**Keywords** *Sanghuangporus vaninii*, Green mould disease, Classification, *Trichoderma virens*

## INTRODUCTION

*Sanghuangporus vaninii* (Ljub.) Zhu et al. (2019) and Wu & Dai (2020) is a species of Basidiomycota, Hymenochaetales, Hymenochaetales, *Sanghuangporus*, of which fruiting body is commonly known as “Sanghuang” in China. Sanghuang has been recorded in historical studies such as “On Medicinal Properties” and “Compendium of Materia Medica” (Kim et al., 2004; Sun et al., 2006; Song et al., 2019). *Sanghuangporus vaninii* is considered as one of the medicinal macrofungi due to its excellent efficiency in treating dysentery and blood insidiousness, anti-tumor, hypoglycemic, anti-oxidative, and immune-enhancing effects (Song et al., 2020). It has been a hot topic in the research and development of pharmaceutical preparations and health products industries in China and some other countries (Che et al., 2005; Gao, Zhang & Yu, 2014). The development of

Sanghuang industry promotes the revitalization of rural economy in China. In 2021, the production of Sanghuang increased to 300 t, and the industry was attached great importance by the government (Yang et al., 2023a).

In China, Sanghuang and other mushrooms are grown using facilities cultivation techniques. Once the facilities are built, the same variety of mushroom is cultivated every year. Some even achieve annual cultivation in facilities by controlling temperature or rotating mushrooms suitable for different seasons, to improve facility utilization and obtain higher economic benefits (Yang et al., 2023b). However, as the cultivation years increase, the occurrence of diseases has a great impact on mushroom cultivation, reducing the quality and yield. A large number of diseases have been reported in mushroom cultivation, such as wet bubble disease caused by *Mycogone pernicioso* in white button mushrooms (*Agaricus bisporus*) (McGee, 2018; Yang et al., 2021), dry bubble disease caused by *Verticillium fungicola* in white button mushrooms and oyster mushroom (Murmu, Maurya & John, 2020), cobweb disease caused by *Cladosporium* spp. in oyster mushrooms (Oyebamiji et al., 2018; Gea, Navarro & Suz, 2019), and white mold disease caused by *Paecilomyces penicillatus* in morels (Yu et al., 2022). In addition to fungal pathogens, *Pseudomonas tolaasii* is consistently associated with mushroom brown blotch disease (Ghasemi et al., 2021), while *Ewingella americana* has been reported as a pathogenic bacterium of brown rot disease on shiitake mushroom (Na, Luo & Yu, 2021). However, despite the history of more than 2000 years of Sanghuang in China, diseases occurring during the process of *S. vaninii* cultivation have not been reported so far due to its short time of artificial cultivation.

In recent years, artificial cultivation of *S. vaninii* has made great progress and the cultivation scale is expanding (Yang et al., 2023a). However, the disease problem is becoming more prominent. From 2018 to 2021, we investigated cultivation companies where the disease occurred and found that the incidence of fungal disease in the cultivation bags of *S. vaninii* was as high as 30–70% in Hangzhou city, Zhejiang province of China. The symptoms of these diseases are basically the same, occurring in the initial or developing stage of *S. vaninii* fruiting bodies, preventing fruiting body formation, or causing brown to dark brown lesions on the fruiting body. The occurrence of this disease influences the quality and yield of Sanghuang, causing great economic losses to producers and becoming an important restriction factor of the Sanghuang industry.

In this study, we observed and described the symptoms of diseases in *S. vaninii* cultivation bags, isolated and identified pathogens using morphological characteristics and phylogenetic analysis with a combination of rDNA ITS genetic regions. The temperature stability of the pathogen was analyzed by *in vitro* test.

## MATERIALS AND METHODS

### Isolation and purification of pathogens

Disease symptoms of *S. vaninii* were observed in a greenhouse at Hangzhou Academy of Agricultural Sciences, located in Zhejiang province, China (120°0'88"E, 30°1'63"N) between late June and late July 2020. Ten diseased cultivation bags were collected, and samples were taken from the junction of the diseased and healthy areas of each bag and

plated onto potato dextrose agar (PDA) containing 0.25 g chloramphenicol. The plates were then incubated at 25 °C. After 7 days of incubation, agar blocks (5 mm in diameter) were cut from the growing edge of colonies and inoculated onto fresh PDA, and this process was repeated several times to obtain putative pure pathogens.

### Pathogenicity assay

To conduct the pathogenicity assay, we prepared a conidial suspension ( $1 \times 10^6$  spores/mL) using five representative isolates. At the end of the vegetative growth stage of *S. vaninii*, a semicircle was cut in the middle of the plastic bags to somatic part of the mycelia in the air. Then, 500  $\mu$ L of the pathogen's conidial suspension was inoculated into the areas surrounding the initial fruiting bodies of *S. vaninii*. The bags were incubated for 10 days at 25 °C and a relative humidity of 98%, and each isolate was tested in triplicate. Uninoculated bags were used as controls. Disease symptoms were observed and recorded, and the pathogens were isolated again from the diseased sites to confirm their morphological characteristics.

### Morphological identification

To identify the fungal pathogens, ten representative isolates were cultured on potato dextrose agar (PDA), CMD (cornmeal agar 20 g, dextrose 20 g, agar 20 g with 1 L distilled water) and SNA ( $\text{KH}_2\text{PO}_4$  1 g,  $\text{KNO}_3$  1 g,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  0.5 g, KCl 0.5 g, glucose 0.2 g, sucrose 0.2 g, agar 15 g with 1 L distilled water) (Jaklitsch, 2009), and incubated at 23 °C under a 12-h light/dark cycle. The structure of conidiophores, phialides, and conidia were observed and measured using a Zeiss Axiophot 2 microscope equipped with an AxioCam CCD camera and Axiovision digital imaging software (Axio-Vision Software Release 3.1., v.3–2002; Carl Zeiss Vision Imaging Systems, Jena, Germany), as previously described (Tomah et al., 2020).

### Molecular analysis

To analyze the ITS region and the genes involved in taxonomy, ten isolates of pathogens were grown in 100 mL potato dextrose broth (PDB) on a shaker at 180 rpm,  $25 \pm 1$  °C for 3 days. Genomic DNA was extracted using the Ezup Column Bacteria Genomic DNA Purification Kit (Sangon Biotech Co., Shanghai, China) according to the manufacturer's instructions. The ITS rDNA regions were amplified using the primer pairs ITS5 (5' GGAAG TAAAAGTCGTAACAAGG3') and ITS4 (5'TCCTCCGCTTATTGATATGC3') (Jiang et al., 2016). The purified PCR product was sequenced in both directions and edited by BioEdit 7.1.3.0. and compared with homologous sequences available in the NCBI databases using BLAST.

Multiple alignment of the ITS rDNA sequences of this study and sequences from NCBI database (type strains of *Trichoderma* species, containing some species reported to be harmful to edible mushroom and some species closely related to the isolated strains) was carried out using Clustal W and a phylogenetic tree was constructed using MEGA 6. The evolutionary history was inferred by using the maximum likelihood (ML) method based on the Jukes-Cantor model (da Silva et al., 2017). The ML method was used to

construct the phylogenetic tree with 1,000 bootstrap frequency. The type strain *Sphaerostilbella lutea* CBS 405.59 was used as the outgroup (Perera *et al.*, 2023).

### Temperature stability assay

Temperature stability was assessed by investigating *in vitro* mycelial growth at different temperatures. Isolate disks (5 mm diameter) were cultured on PDA plates and incubated in the dark at 5 °C, 15 °C, 25 °C, 30 °C, and 35 °C, each temperature treatment three replicates respectively. After 48 h, the diameters of the mycelial colonies were measured. Through diameter comparison, the temperature range suitable for the growth of the isolate was selected to continue the next experiment.

The inhibition of pathogenic isolate on mycelial growth of *S. vaninii* at different temperatures was observed by dual-culture test (Zang *et al.*, 2023). Disks (5 mm diameter) of *S. vaninii* were placed on one side of PDA plates and incubated in the dark at 15 °C, 25 °C, and 30 °C, each temperature treatment three replicates respectively. Seven days later (to compensate for the slower growth of *S. vaninii*), disks of pathogenic isolate were placed on the other side and continued to incubate at the same temperature. The plates with only one disk of *S. vaninii* without pathogenic isolate were used as controls. After another 9 days, the radius of the mycelial colonies of *S. vaninii* was measured.

Analysis of variance (ANOVA) was done using SPSS 20.0 software program (SPSS Inc., Chicago, IL, USA). Mean value and standard deviation of each experiment were grouped according to S-N-K multiple range test with significance level of 5%. Dunnett's test ( $P < 0.05$ ) was also used to compare treatment plots with positive and negative control plots in the experiments.


## RESULTS

### Disease symptoms and pathogen isolation

During the process of artificial cultivation, disease symptoms typically occurred around the timing of fruiting body production of *S. vaninii*. After the somatic growth of *S. vaninii* in a cultivation bag ended, a semi-circular area in the middle of the bag was cut to expose a part of mycelia for the development of fruiting bodies. Pathogen contamination manifested as white hyphae covering the surface of the exposed area or by infecting the initial small fruiting body. Subsequently, green spores appeared on the white mycelium (Fig. 1A). The entire exposed substrate or the fruiting body could be covered by the pathogen mycelium (Fig. 1B), thus preventing development or further development of the fruiting body. The disease symptoms were similar to those caused by *Tricoderma* spp. in green mold disease on other mushrooms. After purification, five representative single-spore isolates (YZB-1 to YZB-5) were collected for pathogenicity testing and identification.

### Pathogenicity tests

A spore suspension of the five isolates was inoculated into the exposed area nearby the developing fruiting body of *S. vaninii*, and white hyphae developed rapidly. Ten days after inoculation, a lot of hyphae with a green mold layer covered the exposed substrate and surrounded the developing fruiting body (Fig. 1C). All of the inoculated bags showed the



**Figure 1** Disease symptoms during the cultivation of *S. vaninii* and after artificial inoculation. (A and B) During the cultivation of *S. vaninii*. Pathogen hyphae covering the surface of the initial fruiting body and exposed substrate. (C) After inoculation. Hyphae inoculated with isolate YZB-1 covering the surface of the substrate and surrounding fruiting body. (D) Normally growing *S. vaninii* fruiting body. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac\_img.jpg\) DOI: 10.7717/peerj.15983/fig-1](https://doi.org/10.7717/peerj.15983/fig-1)


same symptoms as the natural incidence, whereas the control treatment remained symptomless. The five isolates were separated from the inoculated bag again (YZB-1-P to YZB-5-P).

### Morphological identification of pathogens

The colony characteristics of all ten isolates were similar. On PDA, the colonies were floccose with massive conidiation covering the whole surface of the plate (Fig. 2A). On CMD, isolates had a flat colony with aerial mycelium (Fig. 2B). Conidiophores and conidia were produced concentrically or near the margin of the plate. On SNA, they were relatively sparse (Fig. 2C). Conidiophores were gliocladium-like, arising from aerial hyphae, straight, 42–75  $\mu\text{m}$  long ( $n = 30$ ), generally unbranched (Fig. 2D), and sterile near the base, branching irregularly near the tip, with each branch terminating in a whorl of 3–6 phialides; metulae and phialides arose at narrow angles. Phialides were lageniform or ampulliform, 8.5–9.0  $\times$  3.9–4.2  $\mu\text{m}$  at the widest point. Conidia were green, smooth, subglobose, 4.2–4.5  $\times$  3.9–4.0  $\mu\text{m}$  (Fig. 2E). The isolates were similar to *T. virens* Gli 21, as described by *Chaverri, Samuels & Stewart (2001)*. They are markedly different from the reported *Trichoderma* species in terms of spore size, color and location of colonization, phialides morphology and number of branches, and so on (*Tomah et al., 2020; An et al., 2022*).


### Molecular analysis

The DNA from ten isolates was amplified using the primer pairs ITS5/ITS4. Sequence alignment results showed that the ITS nucleotide identity of all isolates was 100%. One isolate, YZB-1, was selected for subsequent analysis, and the ITS fragments were approximately 630 bp in length. The accession number in GenBank is [MZ220425.1](https://www.ncbi.nlm.nih.gov/nuclot/MZ220425.1). Phylogenetic analysis was performed using ITS sequences from 31 type strains of *Trichoderma* species and one outgroup type strain *Sphaerostilbella lutea*. The resulting phylogenetic tree showed that all strains were separated into different clades (Fig. 3), and most reference strains could be distinguished on the species level. Strain YZB-1 was clustered together with *T. virens*. These data confirmed that YZB-1 is a member of *T. virens*.



**Figure 2** Colonies and microscopic photographs of pathogenic fungi. YZB-1 grown on PDA, CMD or SNA in 9-cm-diam Petri dishes under 12 h darkness/12 h light for 7 d. (A) On PDA. (B) On CMD. (C) On SNA. (D and E) Conidiophores and phialides conidia. D = 100  $\mu\text{m}$ ; E = 10  $\mu\text{m}$ .

Full-size [DOI: 10.7717/peerj.15983/fig-2](https://doi.org/10.7717/peerj.15983/fig-2)




**Figure 3** The phylogenetic tree generated from the ITS sequences of *Trichoderma* spp. Branch values lower than 50% were omitted.

Full-size [DOI: 10.7717/peerj.15983/fig-3](https://doi.org/10.7717/peerj.15983/fig-3)


### Temperature stability assay

The mycelial growth of *T. virens* strain YZB-1 was significantly affected by different incubation temperatures (Fig. 4). The mycelia grew fastest at 25 °C, with an average colony diameter of  $57.67 \pm 2.52$  mm. At temperatures above or below 25 °C, mycelium growth gradually slowed down. At 5 °C, the mycelia stopped growing. Dual-cultures of *T. virens* and *S. vaninii* were performed at temperatures suitable for pathogen growth (15 °C, 25 °C, and 30 °C). The inhibition rate of *T. virens* on *S. vaninii* mycelium was highest when



**Figure 4** The diameters of *T. virens* strain YZB-1 at different temperatures. The error bars indicate the standard deviation, and different letters indicate significantly different values ( $P < 0.05$ ).

Full-size DOI: 10.7717/peerj.15983/fig-4



**Figure 5** The dual-culture of *T. virens* strain YZB-1 and *S. vaninii*. (A and B) *S. vaninii* (A) and *T. virens* × *S. vaninii* (B) were incubated at 25 °C. (C) The inhibition ratios of *S. vaninii* by *T. virens* at different temperatures. The error bars indicate the standard deviation, and different letters indicate significantly different values ( $P < 0.05$ ).

Full-size DOI: 10.7717/peerj.15983/fig-5

incubated at 25 °C ( $79.01 \pm 2.79\%$ ), with significant differences in inhibition rates at the three temperatures (Fig. 5C). *Trichoderma virens* not only occupied the medium surface more quickly with mycelial growth but also produced more green spores at the intersection of *T. virens* and *S. vaninii* (Fig. 5B).

## DISCUSSION

*Sanghuangporus vaninii* is a renowned oriental medicinal mushroom, known in China as “Sanghuang,” in Japan as “Meshimakobu,” and in Korea as “Sangwhang” (Chen *et al.*, 2019). Its fruiting body, also called yellow medicinal polyporus or basidiocarp, grows on the trunk of *Populus* sp. Linn., and is prized for its anti-tumor activity due to the bioactive protein-polysaccharide complex it contains (Oh & Han, 1993). However, Sanghuang occurs naturally in rare instances, making it highly valued. As a result, there has been

extensive research on the artificial cultivation of *S. vaninii* (Wang et al., 2018; Hur, 2008). To achieve the formation of fruiting bodies, indoor temperature ranging from 31–35 °C and over 96% relative humidity are ideal, conditions that are also suitable for the occurrence of diseases (Hong, Sung & Nam, 2004).

*Trichoderma* green mold in edible basidiomycetes has been well known for some time (Hatvani et al., 2012). Among the most significant diseases affecting the most commonly cultivated mushrooms worldwide, such as *P. ostreatus* and *A. bisporus*, are those caused by some *Trichoderma* species, including *T. guizhouense*, *T. harzianum*, *T. pleuroticola*, and *T. aggressivum* (Bisset et al., 2015; Chaverri et al., 2015; Kosanovic, Grogan & Kavanagh, 2020; Turgay et al., 2023). However, *T. virens* has been rarely reported to infect edible basidiomycetes. In this study, we found that *T. virens* colonized the mycelium of *S. vaninii*, with the infection being limited to the fruiting body stage. To our knowledge, this is the first report of green mold disease caused by *T. virens* in *S. vaninii* cultivation.

The antifungal mechanism of *Trichoderma* spp. against fungi has been reported because of their biocontrol functions. *Trichoderma* spp. control microorganisms through competition, parasitism, antibiotic action, synergistic antagonism, and other mechanisms (Contreras-Cornejo et al., 2016). Compared to pathogenic microorganisms, *Trichoderma* spp. have faster growth and reproduction rates, stronger decay ability, and wider adaptability. The optimal growth temperature for *Trichoderma* spp. for biocontrol is 25–30 °C (Daryaei et al., 2016). They achieve a fungistatic effect by competing for the living space and nutrient resources of pathogens (Alwathnani et al., 2012). When *T. harzianum* and *Fusarium solani* were co-cultured, *T. harzianum* parasitized *F. solani* from multiple contact points and led to its death (Amira et al., 2017). Additionally, the *Trichoderma* group can degrade the cell wall of pathogens and absorb their nutrients by secreting a series of hydrolases, such as cellulase, glucanase, chitinase, and protease (Mukherjee et al., 2013). *Trichoderma* is beneficial in plant cultivation, but harmful in edible mushroom cultivation (Kredics et al., 2021).

As macroscopic fungi, the growth of edible mushrooms is also inhibited by *Trichoderma* species as aforementioned antifungal mechanism (Velázquez-Cedeño et al., 2007; Abubaker, Sjaarda & Castle, 2013). The optimal growth environment for *Trichoderma* is consistent with the mycelia growth and fruiting body formation environment of most edible fungi, which leads to its infection and harm to edible fungi during the mycelium and fruiting body stages (Kosanovic et al., 2020; Ponnusamy et al., 2022). This was confirmed by the results of both fruiting body inoculation and hyphal dual-culture experiments in the present study. There are few reports on the pathogenic mechanism of *T. virens* infecting the fruiting body of edible mushrooms, which may be related to parasitism and antibiotic action. The control of *Trichoderma* mainly relies on environmental control methods for prevention. Some safe agents (Innocenti et al., 2019) or biocontrol microorganisms (Ma et al., 2019) can be used to control *Trichoderma* during the hypha growth stage. However, the agent may have the potential to cause phytotoxicity (Kwon et al., 2021) or residues (Li et al., 2022) during the fruiting body growth stage.



## CONCLUSIONS

This study has confirmed that the pathogen responsible for fruiting body rot in *S. vaninii* is the isolate YZB-1 through pathogenicity assays. Based on morphological identification and molecular analysis of the rDNA ITS region, the isolate YZB-1 was identified as *T. virens*. *Trichoderma virens* not only infects the fruiting body and causes abnormal growth but also inhibits hyphal growth. Further confirmation is required to determine whether its infection process and pathogenesis are consistent with the above mechanism. Finding safe and effective control methods for *Trichoderma* disease in *S. vaninii* is crucial for future studies.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was funded by a grant from the National Natural Science Foundation of China (No. 31901933) and the Sannongliufang Provincial Project of Zhejiang (CTZB-F190625LWZ-SN). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

National Natural Science Foundation of China: 31901933.

Sannongliufang Provincial Project of Zhejiang: CTZB-F190625LWZ-SN.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Weidong Yuan conceived and designed the experiments, performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Lin Ma conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Xingkun Chen performed the experiments, analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Jiling Song performed the experiments, prepared figures and/or tables, and approved the final draft.
- Qing Chen performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The isolate YZB-1 sequence is available at GenBank: [MZ220425.1](https://www.ncbi.nlm.nih.gov/nuclseq/MZ220425.1).

## Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplemental Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.15983#supplemental-information>.

## REFERENCES

- Abubaker SK, Sjaarda C, Castle JA. 2013.** Regulation of three genes encoding cell-wall-degrading enzymes of *Trichoderma aggressivum* during interaction with *Agaricus bisporus*. *Canadian Journal of Microbiology* **59**(6):417–424 DOI [10.1139/cjm-2013-0173](https://doi.org/10.1139/cjm-2013-0173).
- Alwathnani HA, Perveen K, Tahmaz R, Alhaqbani S. 2012.** Evaluation of biological control potential of locally isolated antagonist fungi against *Fusarium oxysporum* under in vitro and pot conditions. *African Journal of Microbiology Research* **6**(2):312–319 DOI [10.5897/AJMR11.1367](https://doi.org/10.5897/AJMR11.1367).
- Amira BM, Lopez D, Mohamed TA, Khouaja A, Chaar H, Fumanal B, Gousset-Dupont A, Bonhomme L, Label P, Goupil P, Ribeiro S, Pujade-Renaud V, Julien JL, Auguin D, Venisse JS. 2017.** Beneficial effect of *Trichoderma harzianum* strain Ths97 in biocontrolling *Fusarium solani* causal agent of root rot disease in olive trees. *Biological Control* **110**:70–78 DOI [10.1016/j.biocontrol.2017.04.008](https://doi.org/10.1016/j.biocontrol.2017.04.008).
- An XY, Cheng GH, Gao HX, Li XF, Yang Y, Li D, Li Y. 2022.** Phylogenetic analysis of *Trichoderma* species associated with green mold disease on mushrooms and two new pathogens on *Ganoderma sichuanense*. *Journal of Fungi* **8**(7):704 DOI [10.3390/jof8070704](https://doi.org/10.3390/jof8070704).
- Bisset J, Gams W, Jaklitsch W, Samuels GJ. 2015.** Accepted *Trichoderma* names in the year 2015. *IMA Fungus* **6**(2):263–295 DOI [10.5598/imafungus.2015.06.02.02](https://doi.org/10.5598/imafungus.2015.06.02.02).
- Chaverri P, Branco-rocha F, Jaklitsch W, Gazis R, Degenkolb T, Samuels GJ. 2015.** Systematics of the *Trichoderma harzianum* species complex and the reidentification of commercial biocontrol strains. *Mycologia* **107**:558–590 DOI [10.3852/14-147](https://doi.org/10.3852/14-147).
- Chaverri P, Samuels JG, Stewart LE. 2001.** *Hypocrea virens* sp. nov., the teleomorph of *Trichoderma virens*. *Mycologia* **93**:1113–1124 DOI [10.1080/00275514.2001.12063245](https://doi.org/10.1080/00275514.2001.12063245).
- Che HL, Meng FY, Du J, Li R. 2005.** Effect of extract from *Phellinus igniarius* on tumor growth and cell immunity function. *Chinese Journal of Public Health* **21**(1):79–81 DOI [10.3321/j.issn:1001-0580.2005.01.040](https://doi.org/10.3321/j.issn:1001-0580.2005.01.040).
- Chen WH, Tan HY, Liu Q, Zheng XH, Zhang H, Liu YH, Xu LC. 2019.** A review: the bioactivities and pharmacological applications of *Phellinus linteus*. *Molecules* **24**:1888 DOI [10.3390/molecules24101888](https://doi.org/10.3390/molecules24101888).
- Contreras-Cornejo HA, Macías-Rodríguez L, del-Val E, Larsen J. 2016.** Ecological functions of *Trichoderma* spp. and their secondary metabolites in the rhizosphere: interactions with plants. *FEMS Microbiology Ecology* **92**:fiw036 DOI [10.1093/femsec/fiw036](https://doi.org/10.1093/femsec/fiw036).
- da Silva AC, da Rachid CTCC, de Jesus HE, Rosado AS, Peixoto RS. 2017.** Predicting the biotechnological potential of bacteria isolated from Antarctic soils, including the rhizosphere of vascular plants. *Polar Biology* **40**(7):1393–1407 DOI [10.1007/s00300-016-2065-0](https://doi.org/10.1007/s00300-016-2065-0).
- Daryaei A, Jones EE, Ghazalibiglar H, Glare TR, Falloon RE. 2016.** Effects of temperature, light and incubation period on production, germination and bioactivity of *Trichoderma atroviride*. *Journal of Applied Microbiology* **120**(4):999–1009 DOI [10.1111/jam.13076](https://doi.org/10.1111/jam.13076).

- Gao WW, Zhang N, Yu SW. 2014. Research progress on antitumor effects and mechanisms of *Phellinus*. *China Journal of Chinese Materia Medica* 39:4165–4168 DOI 10.4268/cjcm20142116.
- Gea FJ, Navarro MJ, Suz LM. 2019. Cobweb disease on oyster culinary-medicinal mushroom (*Pleurotus ostreatus*) caused by the mycoparasite *Cladobotryum mycophilum*. *Journal of Plant Pathology* 101(2):349–354 DOI 10.1007/s42161-018-0174-z.
- Ghasemi S, Harighi B, Mojarrab M, Azizi A. 2021. Response of *Pseudomonas tolaasii*, the causal agent of mushroom brown blotch disease to the volatile compounds produced by endofungal bacteria. *BioControl* 66(3):421–432 DOI 10.1007/s10526-020-10071-6.
- Hatvani L, Sabolić P, Kocsubé S, Kredics L, Czifra D, Vágvölgyi C, Kaliterna J, Ivić D, Dermić E, Kosalec I. 2012. The first report on mushroom green mould disease in Croatia. *Archives of Industrial Hygiene and Toxicology* 63(4):481–487 DOI 10.2478/10004-1254-63-2012-2220.
- Hong IP, Sung GB, Nam HW. 2004. Artificial cultivation of medicinal mushroom, *Phellinus linteus* using mulberry log. *Journal of Huazhong Agricultural University* 23:72–77.
- Hur H. 2008. Cultural characteristics and log-mediated cultivation of the medicinal mushroom, *Phellinus linteus*. *Mycobiology* 36(2):81–87 DOI 10.4489/MYCO.2008.36.2.081.
- Innocenti G, Montanari M, Righini H, Roberti R. 2019. *Trichoderma* species associated with green mould disease of *Pleurotus ostreatus* and their sensitivity to prochloraz. *Plant Pathology* 68(2):392–398 DOI 10.1111/ppa.12953.
- Jaklitsch WM. 2009. European species of *Hypocrea* Part I. The green-spored species. *Studies in Mycology* 63:1–91 DOI 10.3114/sim.2009.63.01.
- Jiang Y, Wang JL, Chen J, Mao LJ, Feng XX, Zhang CL, Lin FC. 2016. *Trichoderma* biodiversity of agricultural fields in East China reveals a gradient distribution of species. *PLOS ONE* 11(8):e0160613 DOI 10.1371/journal.pone.0160613.
- Kim GY, Choi GS, Lee SH, Park YM. 2004. Acidic polysaccharide isolated from *Phellinus linteus* enhances through the up-regulation of nitric oxide and tumor necrosis factor-alpha from peritoneal macrophages. *Journal of Ethnopharmacology* 95(1):69–76 DOI 10.1016/j.jep.2004.06.024.
- Kosanovic D, Grogan H, Kavanagh K. 2020. Exposure of *Agaricus bisporus* to *Trichoderma aggressivum* f. *europaeum* leads to growth inhibition and induction of an oxidative stress response. *Fungal Biology* 124(9):814–820 DOI 10.1016/j.funbio.2020.07.003.
- Kosanovic D, Sheehan G, Grogan H, Kavanagh K. 2020. Characterisation of the interaction of *Pseudomonas putida* and *Pseudomonas tolaasii* with *Trichoderma aggressivum*. *European Journal of Plant Pathology* 156(1):111–121 DOI 10.1007/s10658-019-01867-z.
- Kredics L, Naeimi S, Hatvani L, Vágvölgyi C, Cai F, Druzhinina IS, Manczinger L. 2021. The Good, the Bad and the Ugly' in the shades of green: the genus *Trichoderma* in the spotlight. *Indian Phytopathology* 74(2):403–411 DOI 10.1007/s42360-021-00352-0.
- Kwon TH, Kim DB, Kim KW, Park MG, Roh GH, Lee BH. 2021. Scaled-up ethyl formate fumigation to replace methyl bromide on traded mushroom to disinfest mushroom fly (*Lycoriella mali*). *Applied Biological Chemistry* 64(1):41 DOI 10.1186/s13765-021-00635-0.
- Li Y, Qin G, He F, Zou K, Zuo B, Liu R, Zhang W, Yang B, Zhao G, Jia G. 2022. Investigation and analysis of pesticide residues in edible fungi produced in the mid-western region of China. *Food Control* 136(4):108857 DOI 10.1016/j.foodcont.2022.108857.
- Ma L, Qu S, Lin J, Jia J, Baird SM, Jiang N, Li H, Hou L, Lu S. 2019. The complete genome of the antifungal bacterium *Pseudomonas* sp. strain MS82. *Journal of Plant Diseases and Protection* 126(2):153–160 DOI 10.1007/s41348-019-00205-z.

- McGee CF. 2018. Microbial ecology of the *Agaricus bisporus* mushroom cropping process. *Applied Microbiology and Biotechnology* 102(3):1075–1083 DOI 10.1007/s00253-017-8683-9.
- Mukherjee PK, Horwitz BA, Herrera-Estrella A, Schmoll M, Kenerley CM. 2013. *Trichoderma* research in the genome era. *Annual Review of Phytopathology* 51(1):105–129 DOI 10.1146/annurev-phyto-082712-102353.
- Murmu R, Maurya AK, John V. 2020. Mycoflora of certain casing materials used in the production of white button mushroom (*Agaricus bisporus* (Lange) Imbach). *International Journal of Chemical Studies* 8(2):2863–2868 DOI 10.22271/chemi.2020.v8.i2ar.9185.
- Na W, Luo H, Yu JM. 2021. First report of bacterial brown rot disease on shiitake mushroom (*Lentinula edodes*) caused by *Ewingella americana* in Korea. *Journal of Plant Pathology* 103(4):1325–1326 DOI 10.1007/s42161-021-00897-z.
- Oh GT, Han SB. 1993. Immuno-stimulating activity of *Phellinus linteus* extracts to B-lymphocyte. *Archives of Pharmacal Research* 15(4):379–381 DOI 10.1007/bf02974118.
- Oyebamiji GH, Jonathan GS, Akinyemi DS, Popoola KK. 2018. Fungal and insect pests of the edible mushroom *Pleurotus ostreatus*. *Notulae Scientia Biologicae* 10(3):379–386 DOI 10.15835/nsb10310284.
- Perera RH, Hyde KD, Jones EBG, Maharachchikumbura SSN, Bundhun D, Camporesi E, Akulov A, Liu JK, Liu ZY. 2023. Profile of *Bionectriaceae*, *Calcarisporiaceae*, *Hypocreaceae*, *Nectriaceae*, *Tilachlidiaceae*, *Ijuhyaceae* fam. nov., *Stromatonectriaceae* fam. nov. and *Xanthonectriaceae* fam. nov. *Fungal Diversity* 118(1):95–271 DOI 10.1007/s13225-022-00512-1.
- Ponnusamy A, Ajis AH, Tan YS, Chai LC. 2022. Dynamics of fungal and bacterial microbiome associated with green-mould contaminated sawdust substrate of *Pleurotus pulmonarius* (grey oyster mushroom). *Journal of Applied Microbiology* 132(3):2131–2143 DOI 10.1111/jam.15327.
- Song JL, Lu N, Wang WK, Yuan WD, Li HB, Cheng JW, Kang XP, Yan J. 2019. Molecular identification of *Sanghuangporus* species *S.sanghuang* and *S.vaninii*. *Journal of Sichuan University* 56(4):765–770 DOI 10.1046/j.1365-2672.1999.00709.x.
- Song JL, Yuan WD, Zhou ZF, Wang WK, Lu N, Cheng JW, Yan J. 2020. Activities of extracellular enzymes and polysaccharides in liquid culture of *Sanghuangporus*. *Mycosystema* 39:352–361 DOI 10.13346/j.mycosystema.190211.
- Sun PL, Xu SY, Yang K, Zhang J, Song L. 2006. Advance of researches on *Phellinus* spp, a rare and precious medicinal fungus. *Microbiology China* 33:119–123 DOI 10.3969/j.issn.0253-2654.2006.02.025.
- Tomah AA, Alamer ISA, Li B, Zhang JZ. 2020. A new species of *Trichoderma* and gliotoxin role: a new observation in enhancing biocontrol potential of *T. virens* against *Phytophthora capsici* on chili pepper. *Biological Control* 145:104261 DOI 10.1016/j.biocontrol.2020.104261.
- Turgay G, Eren E, Eltem R, Çakır D, Aydemir BÇ. 2023. Molecular characterization of *Trichoderma* spp. isolates in mushroom-growing farms in Turkey and their effects on *Agaricus bisporus* production. *Plant Pathology* 16:1–11 DOI 10.1111/ppa.13741.
- Velázquez-Cedeño M, Farnet AM, Billette C, Mata G, Savoie JM. 2007. Interspecific interactions with *Trichoderma longibrachiatum* induce *Pleurotus ostreatus* defence reactions based on the production of laccase isozymes. *Biotechnology Letter* 29(10):1583–1590 DOI 10.1007/s10529-007-9445-z.
- Wang WH, Yang Y, Zhu LN, Jia W, Zhang JS, Liu YF, Yan MQ, Zhao XL, Zhang K, Zhang HN, Fan H. 2018. Inhibitory effects of ethanol extract of *Sanghuangporus sanghuang* fruiting bodies on SW620 colon cancer cells. *Mycosystema* 37:1751–1760 DOI 10.13346/j.mycosystema.180185.
- Wu SH, Dai YC. 2020. Species clarification of the medicinal fungus *Sanghuang*. *Mycosystema* 39:781–794 DOI 10.13346/j.mycosystema.190354.

- Yang Y, Chen X, Dai Y, Zhou L, Cai W, Guo L, Cui B, Li N, Lei P, Li C, Li Z, Shi W, Lu B, Qin S, Wang R, Li X, Song J, Yao S, Zhang A, Jiang N, Wan H, Ding F, Zhang H, Zheng Z, Zhang S, Zhou H, Fan Y, Chen W, Yu X, Tang S, Zhang Y, Zhang Q, Wang K, Yan J, Liu D, Wang C, Yang B, Chen W, Li T, Mo Y, Fu J, Wu D, Song B, Liu J, Wang J, Zhou W, Geng J, Zhao J, Li Q. 2023a. Sanghuang industry in China: current status, challenges and perspectives: the Qiandao Lake declaration for sanghuang industry development. *Mycosystema* 42:855–873 DOI 10.13346/j.mycosystema.220065.
- Yang Y, Sossah FL, Li Z, Hyde KD, Li D, Xiao S, Fu Y, Yuan X, Li Y. 2021. Genome-wide identification and analysis of chitinase GH18 gene family in *Mycogone pernicioso*. *Frontiers in Microbiology* 11:2029 DOI 10.3389/fmicb.2020.596719.
- Yang Y, Zhang Z, Liu J, Li C, Liu P, Li Y. 2023b. Comparison of cultivation characteristics between a wild *Sanghuangporus vaninii* strain and domestic three *Sanghuangporus* species. *Mycosystema* 42:896–906 DOI 10.13346/j.mycosystema.220456.
- Yu Y, Liu T, Liu L, Chen Y, Tang J, Peng W, Tan H. 2022. Application of the mushroom volatile 1-octen-3-ol to suppress a morel disease caused by *Paecilomyces penicillatus*. *Applied Microbiology and Biotechnology* 106(12):4787–4799 DOI 10.1007/s00253-022-12038-2.
- Zang C, Kong T, Liang B, Liu X, Xie J, Lin Y, Pei X, Yu S, Liang C. 2023. Evaluation of imide substance from *Streptomyces atratus* PY-1 for the biocontrol of Phytophthora blight. *European Journal of Plant Pathology* 165(4):725–734 DOI 10.1007/s10658-023-02648-5.
- Zhu L, Song J, Zhou JL, Si J, Cui BK. 2019. Species diversity, phylogeny, divergence time, and biogeography of the genus *Sanghuangporus* (Basidiomycota). *Frontiers in Microbiology* 10:812 DOI 10.3389/fmicb.2019.00812.