

Identification of a pathogen causing fruiting body rot of *Sanghuangporus vaninii* (#84125)

1

First submission

Guidance from your Editor

Please submit by **20 Apr 2023** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

5 Figure file(s)

1 Raw data file(s)



Custom checks

DNA data checks



Have you checked the authors [data deposition statement](#)?



Can you access the deposited data?



Has the data been deposited correctly?



Is the deposition information noted in the manuscript?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Identification of a pathogen causing fruiting body rot of *Sanghuangporus vaninii*

Weidong Yuan¹, Qing Chen², Xingkun Chen³, Jiling Song¹, Lin Ma^{Corresp. 4}

¹ Hangzhou Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

² Zhejiang Agricultural Technology Extension Center, Hangzhou, Zhejiang, China

³ College of Agronomy, Shandong Agricultural University, Taian, Shandong, China

⁴ Jiangsu Key Laboratory for Horticultural Crop Genetic Improvement, Institute of Vegetable Crops, Jiangsu Academy of Agricultural Sciences, Nanjing, Jiangsu, China

Corresponding Author: Lin Ma

Email address: 20090029@jaas.ac.cn

Sanghuangporus vaninii is a medicinal macrofungus that is increasingly cultivated in China. During cultivation, it was found that the fruiting body of *S. vaninii* was susceptible to pathogenic fungi, resulting in significant economic losses to the industry. The symptoms of the disease occur in the initial stage of fruiting body development. The isolate YZB-1 was obtained from a diseased fruiting body, and its spore suspension was inoculated into the exposed area nearby the developing fruiting body of *S. vaninii*. After 10 days, the same disease symptoms appeared in the inoculated area. Morphological identification and molecular analysis of rDNA ITS region confirmed that the isolate YZB-1 was identified as *Trichoderma virens*. The temperature stability assay revealed that the mycelia of YZB-1 grew the fastest at 25°C, with growth slowing down gradually as the temperature increased or decreased. Dual-culture tests of *T. virens* and *S. vaninii* showed that the inhibition rate of *T. virens* on *S. vaninii* mycelium was the highest ($78.65 \pm 2.90\%$) at 25°C, and more green spores were produced at the intersection of *T. virens* and *S. vaninii*.

Identification of a pathogen causing fruiting body rot of *Sanghuangporus vaninii*

Weidong Yuan¹, Qing Chen³, Xingkun Chen⁴, Jiling Song¹, Lin Ma^{2,*}

¹ Hangzhou Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

² Jiangsu Key Laboratory for Horticultural Crop Genetic Improvement, Institute of Vegetable Crops, Jiangsu Academy of Agricultural Sciences, Nanjing, Jiangsu, China

³ Zhejiang Agricultural Technology Extension Center, Hangzhou, Zhejiang, China

⁴ College of Agronomy, Shandong Agricultural University, Taian, Shandong, China

Corresponding Author:

Lin Ma²

Zhongling Str No.50, Nanjing, Jiangsu, 210014, China

Email address: 20090029@jaas.ac.cn

Abstract

Sanghuangporus vaninii is a medicinal macrofungus that is increasingly cultivated in China. During cultivation, it was found that the fruiting body of *S. vaninii* was susceptible to pathogenic fungi, resulting in significant economic losses to the industry. The symptoms of the disease occur in the initial stage of fruiting body development. The isolate YZB-1 was obtained from a diseased fruiting body, and its spore suspension was inoculated into the exposed area nearby the developing fruiting body of *S. vaninii*. After 10 days, the same disease symptoms appeared in the inoculated area. Morphological identification and molecular analysis of rDNA ITS region confirmed that the isolate YZB-1 was identified as *Trichoderma virens*. The temperature stability assay revealed that the mycelia of YZB-1 grew the fastest at 25°C, with growth slowing down gradually as the temperature increased or decreased. Dual-culture tests of *T. virens* and *S. vaninii* showed that the inhibition rate of *T. virens* on *S. vaninii* mycelium was the highest (79.01 ± 2.79%) at 25 °C, and more green spores were produced at the intersection of *T. virens* and *S. vaninii*.

Keywords: *Sanghuangporus vaninii*, green mould disease, classification, *Trichoderma virens*

Introduction

Sanghuangporus vaninii (Ljub.) L.W. Zhou & Y.C. Dai is a species of Basidiomycetes, Aphyllophorales, Hymenochaetaceae, *Sanghuangporus*, of which fruiting body is commonly known as "Sanghuang" in China (Zhu et al., 2019; Wu & Dai, 2020). Sanghuang has been recorded in historical works such as "On Medicinal Properties" and "Compendium of Materia Medica" (Kim et al., 2004; Sun et al., 2006; Song et al., 2019). *S. vaninii* is considered as one of the medicinal macrofungi due to its excellent efficiency in treating dysentery and blood insidiousness, anti-tumor, hypoglycemic, anti-oxidative, and immune-enhancing effects (Song et al., 2020). It has been a hot topic in the research and development of pharmaceutical preparations and health products industries in China and some other countries (Che et al., 2005; Gao et al., 2014).

In China, Sanghuang and other mushrooms are grown using facilities cultivation techniques. Once the facilities are built, the same variety of mushroom is cultivated every year. Some even achieve annual cultivation in facilities by controlling temperature or rotating mushrooms suitable for different seasons, to improve facility utilization and obtain higher economic benefits. However, as the cultivation years increase, the occurrence of diseases has a great impact on mushroom cultivation, reducing the quality and yield. A large number of diseases have been reported in mushroom cultivation, such as wet bubble disease caused by *Mycogone perniciosa* in white button mushrooms (*Agaricus bisporus*) (Regnier & Combrinck, 2010), dry bubble disease caused by *Verticillium fungicola* in white button mushrooms and oyster mushroom (Marlowe & Romaine, 1982; Largeteau & Savoie, 2008), cobweb disease caused by *Cladosporium* spp. in *Pleurotus* mushrooms (Back et al., 2012), and green mold disease caused by *Cladosporium* spp. (Komoń-Zelazowska et al., 2007). In addition to fungal pathogens, *Pseudomonas tolaasii* is consistently associated with brown-reddish blotches on *Pleurotus ostreatus* sporocarps (Cantore & Iacobellis, 2014), while *Pantoea* spp. has been reported as a pathogenic bacterium of soft rot disease with symptoms of watery lesions on the stipes and pileus of *P. eryngii* (Kim et al., 2007; Liu et al., 2013). However, despite the history of more than 2000 years of Sanghuang in China, diseases occurring during the process of *S. vaninii* cultivation have not been reported so far due to its short time of artificial cultivation.

In recent years, artificial cultivation of *S. vaninii* has made great progress and the cultivation scale is expanding. However, the disease problem is becoming more prominent. From 2018 to 2021, we investigated cultivation companies where the disease occurred and found that the incidence of fungal disease in the cultivation bags of *S. vaninii* was as high as 30% - 70% in Hangzhou city, Zhejiang province of China. The symptoms of these diseases are basically the same, occurring in the initial or developing stage of *S. vaninii* fruiting bodies, preventing fruiting body formation, or causing brown to dark brown lesions on the fruiting body. The occurrence of this disease influences the quality and yield of Sanghuang, causing great economic losses to producers and becoming an important restriction factor of the Sanghuang industry.

In this study, we observed and described the symptoms of diseases in *S. vaninii* cultivation bags, isolated and identified pathogens using morphological characteristics and phylogenetic analysis with a combination of rDNA ITS genetic regions. The temperature stability of the pathogen was analyzed by *in vitro* test.

Materials & Methods

Isolation and purification of pathogens

Disease symptoms of *S. vaninii* were observed in a greenhouse at Hangzhou Academy of Agricultural Sciences, located in Zhejiang province, China (120°0'88" E, 30°1'63" N) between late June and late July 2020. Ten diseased cultivation bags were collected, and samples were taken from the junction of the diseased and healthy areas of each bag and plated onto potato dextrose agar (PDA) containing 0.25 g chloramphenicol. The plates were then incubated at 25°C. After 7 days of incubation, agar blocks (5 mm in diameter) were cut from the growing edge of colonies and inoculated onto fresh PDA, and this process was repeated several times to obtain putative pure pathogens.

Pathogenicity assay

To conduct the pathogenicity assay, we prepared a conidial suspension (1×10^6 spores/mL) using five representative isolates. At the end of the vegetative growth stage of *S. vaninii*, a semicircle was cut in the middle of the plastic bags to expose part of the mycelia in the air. Then, 500 μ L of the pathogen's conidial suspension was inoculated into the areas surrounding the initial fruiting bodies of *S. vaninii*. The bags were incubated for 10 days at 25°C and a relative humidity of 98%, and each isolate was tested in triplicate. Uninoculated bags were used as controls. Disease symptoms were observed and recorded, and the pathogens were isolated again from the diseased sites to confirm their morphological characteristics.

Morphological identification

To identify the fungal pathogens, ten representative isolates were cultured on potato dextrose agar (PDA), CMD (cornmeal agar 20 g, dextrose 20 g, agar 20 g with 1 L distilled water) and SNA (KH_2PO_4 1 g, KNO_3 1 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g, KCl 0.5 g, glucose 0.2 g, sucrose 0.2 g, agar 15 g with 1 L distilled water) (Jaklitsch, 2009), and incubated at 23°C under a 12-hour light/dark cycle. The structure of conidiophores, phialides, and conidia were observed and measured using a Zeiss Axiophot 2 microscope equipped with an Axiocam CCD camera and Axiovision digital imaging software (Axio-Vision Software Release 3.1., v.3–2002; Carl Zeiss Vision Imaging Systems), as previously described (Tomah et al., 2020).

Molecular analysis

To analyze the ITS region and the genes involved in taxonomy, ten isolates of pathogens were grown in 100 mL potato dextrose broth (PDB) on a shaker at 180 rpm,

25 ± 1°C for 3 days. Genomic DNA was extracted using the Ezup Column Bacteria Genomic DNA Purification Kit (Sangon Biotech Co., Shanghai, China) according to the manufacturer's instructions. The ITS rDNA regions were amplified using the primer pairs ITS5 (5'GGAAG TAAAAGTCGTAACAAGG3') and ITS4 (5'TCCTCCGCTTATTGATATGC3') (Jiang *et al.*, 2016). The purified PCR product was sequenced in both directions and edited by BioEdit 7.1.3.0. Ambiguously aligned positions of sequences were eliminated using Gblocks 0.91b (Castresana, 2000). Maximum likelihood phylogenies were generated by raxmlGUI v. 1.5 (Silvestro & Michalak, 2012). ML bootstrap (ML-BS) analysis of each ML tree was completed with a fast 1000 bootstrap frequency using the Jukes-Cantor model of nucleotide substitution.

Temperature stability assay

Temperature stability was assessed by investigating *in vitro* mycelial growth at different temperatures. Isolate disks (5 mm diameter) were cultured on PDA plates and incubated in the dark at 5°C, 15°C, 25°C, 30°C, and 35°C, respectively. After 48 hours, the diameter of the mycelial colonies was measured. The inhibition of pathogenic isolate on mycelial growth of *S. vaninii* at different temperatures was observed by dual-culture test (Zang *et al.*, 2023). Disks (5 mm diameter) of *S. vaninii* were placed on one side of PDA plates and incubated in the dark at 15°C, 25°C, and 30°C, respectively. Seven days later (to compensate for the slower growth of *S. vaninii*), disks of pathogenic isolate were placed on the other side and continued to incubate at the same temperature. The plates with only one disk of *S. vaninii* without pathogenic isolate were used as controls. After another 9 days, the radius of the mycelial colonies of *S. vaninii* was measured. Three replicates of the plate assays were performed independently.

Results

Disease symptoms and pathogen isolation

During the process of artificial cultivation, disease symptoms typically occurred around the timing of fruiting body production of *S. vaninii*. After the vegetative growth of *S. vaninii* in a cultivation bag ended, a semi-circular area in the middle of the bag was cut to expose a part of mycelia for the development of fruiting bodies. Pathogen contamination manifested as white hyphae covering the surface of the exposed area or by infecting the initial small fruiting body. Subsequently, green spores appeared on the white mycelium (Figure 1A). The entire exposed substrate or the fruiting body could be covered by the pathogen mycelium (Figure 1B), thus preventing development or further development of the fruiting body. The disease symptoms were similar to those caused by *Tricoderma* spp. in green mold disease on other mushrooms. After purification, five representative single-spore isolates (YZB-1 to YZB-5) were collected for pathogenicity testing and identification.

Pathogenicity tests

A spore suspension of the five isolates was inoculated into the exposed area nearby the developing fruiting body of *S. vaninii*, and white hyphae developed rapidly. Ten days after inoculation, a lot of hyphae with a green mold layer covered the exposed substrate and surrounded the developing fruiting body (Figure 1C). All of the inoculated bags showed the same symptoms as the natural incidence, whereas the control treatment remained symptomless. The five isolates were separated from the inoculated bag again (YZB-1-P to YZB-5-P).

Morphological identification of pathogens

The colony characteristics of all ten isolates were similar. On PDA, the colonies were floccose with massive conidiation covering the whole surface of the plate (Figure 2A). On CMD, isolates had a flat colony with aerial mycelium (Figure 2B). Conidiophores and conidia were produced concentrically or near the margin of the plate. On SNA, they were relatively sparse (Figure 2C). Conidiophores were gliocladium-like, arising from aerial hyphae, straight, 42–75 µm long (n = 30), generally unbranched (Figure 2D), and sterile near the base, branching irregularly near the tip, with each branch terminating in a whorl of 3–6 phialides; metulae and phialides arose at narrow angles. Phialides were lageniform or ampulliform, 8.5–9.0 × 3.9–4.2 µm at the widest point. Conidia were green, smooth, subglobose, 4.2–4.5 × 3.9–4.0 µm (Figure 2E). The isolates were similar to *T. virens*, as described by Chaverri (Chaverri et al., 2001).

Molecular analysis

The DNA from ten isolates was amplified using the primer pairs ITS5/ITS4. Sequence alignment results showed that the ITS nucleotide identity of all isolates was 100%. One isolate, YZB-1, was selected for subsequent analysis, and the ITS fragments were approximately 630 bp in length. The accession number in GenBank is MZ220425.1. Phylogenetic analysis was performed using ITS sequences from 24 strains of *Trichoderma* species. The resulting phylogenetic tree showed that all strains were separated into different clades (Figure 3), and most reference strains could be distinguished on the species level. Strain YZB-1 was clustered with *T. virens* in a distinct clade with high bootstrap value (99.8%) support, confirming that YZB-1 is a member of *T. virens*.

Temperature stability assay

The mycelial growth of *T. virens* strain YZB-1 was significantly affected by different incubation temperatures (Figure 4). The mycelia grew fastest at 25°C, with an average colony diameter of 57.67 ± 2.52 mm. At temperatures above or below 25°C, mycelium growth gradually slowed down. At 5°C, the mycelia stopped growing. Dual-cultures of *T. virens* and *S. vaninii* were performed at temperatures suitable for pathogen growth (15°C, 25°C, and 30°C). The inhibition rate of *T. virens* on *S. vaninii* mycelium was highest when incubated at 25°C (79.01 ± 2.79%), with significant differences in inhibition rates at the three temperatures (Figure 5C). *T. virens* not only occupied the

medium surface more quickly with mycelial growth but also produced more green spores at the intersection of *T. virens* and *S. vaninii* (Figure 5B).

Discussion

S. vaninii is a renowned oriental medicinal mushroom, known in China as "Sanghuang," in Japan as "Meshimakobu," and in Korea as "Sangwhang" (Chen et al., 2019). Its fruiting body, also called yellow medicinal polyporus or basidiocarp, grows on the trunk of *Populus Linn*, and is prized for its anti-tumor activity due to the bioactive protein-polysaccharide complex it contains (Oh & Han, 1993). However, Sanghuang occurs naturally in rare instances, making it highly valued. As a result, there has been extensive research on the artificial cultivation of *S. vaninii* (Wang et al., 1993; Hur, 2008). To achieve the formation of fruiting bodies, indoor temperature ranging from 31-35°C and over 96% relative humidity are ideal, conditions that are also suitable for the occurrence of diseases (Hong et al., 2004).

Trichoderma green mold in edible basidiomycetes has been well known for some time (Hatvani et al., 2012). Among the most significant diseases affecting the most commonly cultivated mushrooms worldwide, such as *P. ostreatus* and *T. pleuroti*, are those caused by some *Trichoderma* species, including *T. guizhouense*, *T. harzianum*, *T. pleuroticola*, and *T. pleuroti* (Hatvani et al., 2007; Bisset et al., 2015; Chaverri et al., 2015; Woo et al., 2009). However, *T. virens* has been rarely reported to infect edible basidiomycetes. In this study, we found that *T. virens* colonized the mycelium of *S. vaninii*, with the infection being limited to the developing fruiting body. To our knowledge, this is the first report of green mold disease caused by *T. virens* in *S. vaninii* cultivation.

Trichoderma spp. are biocontrol fungi that can be used to control plant pathogenic microorganisms through competition, parasitism, antibiotic action, synergistic antagonism, and other mechanisms (Contreras-Cornejo et al., 2016). Compared to pathogenic microorganisms, *Trichoderma* spp. have faster growth and reproduction rates, stronger decay ability, and wider adaptability. The optimal growth temperature for *Trichoderma* spp. for biocontrol is 25-30 °C (Mukherjee & Liu, 1997). They achieve a fungistatic effect by competing for the living space and nutrient resources of pathogens (Alwathnani et al., 2012). Maroua et al. (Amira et al., 2017) found that when *T. harzianum* and *Fusarium solani* were co-cultured, *T. harzianum* parasitized *F. solani* from multiple contact points and led to its death. Additionally, the *Trichoderma* group can degrade the cell wall of pathogens and absorb their nutrients by secreting a series of hydrolases, such as cellulase, glucanase, chitinase, and protease (Mukherjee et al., 2013). The secondary metabolite harzianic acid (HA) produced by *T. harzianum* showed inhibition against the mycelium radial growth of soil-borne pathogens *Sclerotinia sclerotiorum* and *Rhizoctonia solani* (Vinale et al., 2014).

As macroscopic fungi, the growth of edible mushrooms is also inhibited by the aforementioned *Trichoderma* species (Velázquez-Cedeño et al., 2007; Abubaker et al., 2013). The optimal growth environment for *Trichoderma* is consistent with the mycelia growth

and fruiting body formation environment of most edible fungi, which leads to its infection and harm to edible fungi during the mycelium and fruiting body stages (Kosanovic *et al.*, 2020; Ponnusamy *et al.*, 2022). This was confirmed by the results of both fruiting body inoculation and hyphal dual-culture experiments in the present study. There are few reports on the pathogenic mechanism of *T. virens* infecting the fruiting body of edible mushrooms, which may be related to parasitism and antibiotic action. The control of *Trichoderma* mainly relies on environmental control methods for prevention. Some safe agents (Innocenti *et al.*, 2019) or biocontrol microorganisms (Ma *et al.*, 2019) can be used to control *Trichoderma* during the hypha growth stage. However, the agent may have the potential to cause phytotoxicity (Kwon *et al.*, 2021) or residues (Li *et al.*, 2022) during the fruiting body growth stage.

Conclusions

This study has confirmed that the pathogen responsible for fruiting body rot in *S. vaninii* is the isolate YZB-1 through pathogenicity assays. Based on morphological identification and molecular analysis of the rDNA ITS region, the isolate YZB-1 was identified as *T. virens*. *T. virens* not only infects the fruiting body and causes abnormal growth but also inhibits hyphal growth. Further confirmation is required to determine whether its infection process and pathogenesis are consistent with the above mechanism. Finding safe and effective control methods for *Trichoderma* disease in *S. vaninii* is crucial for future studies.

References


- Abubaker SK, Sjaarda C, Castle JA. 2013. Regulation of three genes encoding cell-wall-degrading enzymes of *Trichoderma aggressivum* during interaction with *Agaricus bisporus*. *Canadian Journal of Microbiology* **59**:417-424 DOI 10.1139/cjm-2013-0173.
- Alwathnani HA, Perveen K, Tahmaz R, Alhaqbani S. 2012. Evaluation of biological control potential of locally isolated antagonist fungi against *Fusarium oxysporum* under in vitro and pot conditions. *African Journal of Microbiology Research* **6**:312-319 DOI 10.5897/AJMR11.1367.
- Amira BM, Lopez D, Mohamed TA, Khouaja A, Chaar H, Fumanal B, Gousset-Dupont A, Bonhomme L, Label P, Goupil P, Ribeiro S, Pujade-Renaud V, Julien JL, Auguin D, Venisse JS. 2017. Beneficial effect of *Trichoderma harzianum* strain Ths97 in biocontrolling *Fusarium solani* causal agent of root rot disease in olive trees. *Biological Control* **110**:70-78 DOI j.biocontrol.2017.04.008.
- Back CG, Lee CY, Seo GS, Jung HY. 2012. Characterization of species of *Cladobotryum* which cause cobweb disease in edible mushrooms grown in Korea. *Mycobiology* **40**:189-194 DOI 10.5941/MYCO.2012.40.3.189.
- Bisset J, Gams W, Jaklitsch W, Samuels GJ. 2015. Accepted *Trichoderma* names in the year 2015. *IMA Fungus* **6**:263-295 DOI 10.5598/ima fungus.2015.06.02.02.
- Cantore PL, Iacobellis NS. 2014. Characterization of fluorescent pseudomonads responsible for the yellowing of oyster mushroom (*Pleurotus ostreatus*). *Phytopathologia Mediterranea* **53**:54-65 DOI 10.14601/Phytopathol_Mediterr-12292.

- 274 **Castresana J. 2000.** Selection of conserved blocks from multiple alignments for their use in
275 phylogenetic analysis. *Molecular Biology and Evolution* **17**:540-552 DOI [10.1093/oxfordjournals.molbev.a026334](https://doi.org/10.1093/oxfordjournals.molbev.a026334).
276
- 277 **Chaverri P, Branco-rocha F, Jaklitsch W, Gazis R, Degenkolb T, Samuels GJ. 2015.**
278 Systematics of the *Trichoderma harzianum* species complex and the reidentification of
279 commercial biocontrol strains. *Mycologia* **107**:558-590 DOI [10.3852/14-147](https://doi.org/10.3852/14-147).
280
- 281 **Chaverri P, Samuels JG, Stewart LE. 2001.** *Hypocrea virens* sp. nov., the teleomorph of
282 *Trichoderma virens*. *Mycologia* **93**:1113-1124 DOI [10.1080/00275514.2001.12063245](https://doi.org/10.1080/00275514.2001.12063245).
283
- 284 **Che HL, Meng FY, Du J, Li R. 2005.** Effect of extract from *Phellinus igniarius* on tumor growth
285 and cell immunity function. *Chinese Journal of Public Health* **21**:79-81 DOI
286 [10.3321/j.issn:1001-0580.2005.01.040](https://doi.org/10.3321/j.issn:1001-0580.2005.01.040).
287
- 288 **Chen WH, Tan HY, Liu Q, Zheng XH, Zhang H, Liu YH, Xu LC. 2019.** A review: the
289 bioactivities and pharmacological applications of *Phellinus linteus*. *Molecules* **24**:1888 DOI
290 [10.3390/molecules24101888](https://doi.org/10.3390/molecules24101888).
291
- 292 **Contreras-Cornejo HA, Macías-Rodríguez L, del-Val E, Larsen J. 2016.** Ecological functions
293 of *Trichoderma* spp. and their secondary metabolites in the rhizosphere: interactions with
294 plants, *FEMS Microbiology Ecology* **92**:fiw036 DOI [10.1093/femsec/fiw036](https://doi.org/10.1093/femsec/fiw036).
295
- 296 **Gao WW, Zhang N, Yu SW. 2014.** Research progress on antitumor effects and mechanisms of
297 *Phellinus*. *China Journal of Chinese Materia Medica* **39**:4165-4168 DOI
298 [10.4268/cjcmm20142116](https://doi.org/10.4268/cjcmm20142116).
299
- 300 **Hatvani L, Antal Z, Manczinger L, Szekeres A, Druzhinina IS, Kubicek CP, Nagy A, Nagy E,**
301 **Vágvölgyi C, Kredics L. 2007.** Green mould diseases of *Agaricus* and *Pleurotus* spp. are
302 caused by related but phylogenetically different *Trichoderma* species. *Phytopathology* **97**:532-
303 537 DOI [10.1094/PHYTO-97-4-0532](https://doi.org/10.1094/PHYTO-97-4-0532).
304
- 305 **Hatvani L, Sabolić P, Kocsubé S, Kredics L, Czifra D, Vágvölgyi C, Kaliterna J, Ivić D,**
306 **Đermić E, Kosalec I. 2012.** The first report on mushroom green mould disease in Croatia.
307 *Archives of Industrial Hygiene and Toxicology* **63**:481-487 DOI [10.2478/10004-1254-63-2012-
308 2220](https://doi.org/10.2478/10004-1254-63-2012-2220).
309
- 310 **Hong IP, Sung GB, Nam HW. 2004.** Artificial cultivation of medicinal mushroom, *Phellinus*
311 *linteus* using mulberry log. *Journal of Huazhong Agricultural University* **23**:72-77 DOI
312 [10.1093/j.cnki.hnlkxb.2004.01.019](https://doi.org/10.1093/j.cnki.hnlkxb.2004.01.019).
313
- 314 **Hur H. 2008.** Cultural characteristics and log-mediated cultivation of the medicinal mushroom,
315 *Phellinus linteus*. *Mycobiology* **36**:81-87 DOI [10.4489/MYCO.2008.36.2.081](https://doi.org/10.4489/MYCO.2008.36.2.081).
316
- 317 **Innocenti G, Montanari M, Righini H, Roberti R. 2019.** *Trichoderma* species associated with
318 green mould disease of *Pleurotus ostreatus* and their sensitivity to prochloraz. *Plant
Pathology* **68**:392-398 DOI [10.1111/ppa.12953](https://doi.org/10.1111/ppa.12953).
319
- 320 **Jaklitsch WM. 2009.** European species of *Hypocrea* Part I. The green-spored species. *Studies in
321 Mycology* **63**:1-91 DOI [10.3114/sim.2009.63.01](https://doi.org/10.3114/sim.2009.63.01).
322
- 323 **Jiang Y, Wang JL, Chen J, Mao LJ, Feng XX, Zhang CL, Lin FC. 2016.** *Trichoderma* biodiversity
324 of agricultural fields in East China reveals a gradient distribution of species. *PLoS ONE*
325 **11**:e0160613 DOI [10.1371/journal.pone.0160613](https://doi.org/10.1371/journal.pone.0160613).
326
- 327 **Kim GY, Choi GS, Lee SH, Park YM. 2004.** Acidic polysaccharide isolated from *Phellinus linteus*
328 enhances through the up-regulation of nitric oxide and tumor necrosis factor-alpha from
329 peritoneal macrophages. *Journal of Ethnopharmacology* **95**:69-76 DOI [10.1016/j.jep.2004.06.024](https://doi.org/10.1016/j.jep.2004.06.024).
330

- 319 **Kim MK, Ryu JS, Lee YH, Yun HD. 2007.** First report of *Pantoea* sp. induced soft rot disease of
320 *Pleurotus eryngii* in Korea. *Plant Diseases* 91:109-109 DOI [10.1094/PD-91-0109A](https://doi.org/10.1094/PD-91-0109A).
- 321 **Komoń-Zelazowska M, Bissett J, Zafari D, Hatvani L, Manczinger L, Woo S, Lorito M,**
322 **Kredics L, Kubicek CP, Druzhinina IS. 2007.** Genetically closely related but
323 phenotypically divergent *Trichoderma* species cause green mould disease in oyster
324 mushroom farms worldwide. *Applied and Environmental Microbiology* 73:7415-7426 DOI
325 [10.1128/AEM.01059-07](https://doi.org/10.1128/AEM.01059-07).
- 326 **Kosanovic D, Sheehan G, Grogan H, Kavanagh K. 2020.** Characterisation of the interaction
327 of *Pseudomonas putida* and *Pseudomonas tolaasii* with *Trichoderma aggressivum*. *European*
328 *Journal of Plant Pathology* 156:111-121 DOI [10.1007/s10658-019-01867-z](https://doi.org/10.1007/s10658-019-01867-z).
- 329 **Kwon Th, Kim Db, Kim Kw, Park Mg, Roh Gh, Lee Bh. 2021.** Scaled-up ethyl formate
330 fumigation to replace methyl bromide on traded mushroom to disinfest mushroom fly
331 (*Lycoriella mali*). *Applied Biological Chemistry* 64:64 DOI [10.1186/s13765-021-00635-0](https://doi.org/10.1186/s13765-021-00635-0).
- 332 **Largeteau ML, Savoie JM. 2008.** Effect of the fungal pathogen *Verticillium fungicola* on fruiting
333 initiation of its host, *Agaricus bisporus*. *Mycological Research* 112:825-828 DOI
334 [10.1016/j.mycres.2008.01.018](https://doi.org/10.1016/j.mycres.2008.01.018).
- 335 **Li Y, Qin G, He F, Zou K, Zuo B, Liu R, Zhang W, Yang B, Zhao G, Jia G. 2022.** Investigation
336 and analysis of pesticide residues in edible fungi produced in the mid-western region of
337 China. *Food Control* 136:108857 DOI [10.1016/j.foodcont.2022.108857](https://doi.org/10.1016/j.foodcont.2022.108857).
- 338 **Liu Y, Wang SX, Zhang DP, Wei SJ, Zhao S, Chen SF, Xu, F. 2013.** *Pantoea beijingensis* sp. nov.,
339 isolated from the fruiting body of *Pleurotus eryngii*. *Antonie van Leeuwenhoek* 104:1039-1047
340 DOI [10.1007/s10482-013-0024-0](https://doi.org/10.1007/s10482-013-0024-0).
- 341 **Ma L, Qu S, Lin J, Jia J, Baird SM, Jiang N, Li H, Hou L, Lu S. 2019.** The complete genome of
342 the antifungal bacterium *Pseudomonas* sp. strain MS82. *Journal of Plant Diseases and Protection*
343 126:153-160 DOI [10.1007/s41348-019-00205-z](https://doi.org/10.1007/s41348-019-00205-z).
- 344 **Marlowe A, Romaine CP. 1982.** Dry bubble of oyster mushroom caused by *Verticillium*
345 *fungicola*. *Plant Diseases* 66:859.
- 346 **Mukherjee PK, Raghu K. 1997.** Effect of temperature on antagonistic and biocontrol potential
347 of shape *Trichoderma* sp. on *Sclerotium rolfsii*. *Mycopathologia* 139:151-155 DOI
348 [10.1023/A:1006868009184](https://doi.org/10.1023/A:1006868009184).
- 349 **Mukherjee PK, Horwitz BA, Herrera-Estrella A, Schmoll M, Kenerley CM. 2013.** *Trichoderma*
350 research in the genome era. *Annual Review of Phytopathology* 51:105-129 DOI
351 [10.1146/annurev-phyto-082712-102353](https://doi.org/10.1146/annurev-phyto-082712-102353).
- 352 **Oh GT, Han SB. 1993.** Immuno-stimulating activity of *Phellinus linteus* extracts to B-
353 lymphocyte. *Archives of Pharmacal Research* 15:379-381 DOI [10.1007/bf02974118](https://doi.org/10.1007/bf02974118).
- 354 **Ponnusamy A, Ajis AH, Tan YS, Chai LC. 2022.** Dynamics of fungal and bacterial microbiome
355 associated with green-mould contaminated sawdust substrate of *Pleurotus*
356 *pulmonarius* (grey oyster mushroom). *Journal of Applied Microbiology* 132:2131-2143 DOI
357 [10.1111/jam.15327](https://doi.org/10.1111/jam.15327).
- 358 **Regnier T, Combrinck S. 2010.** *In vitro* and *in vivo* screening of essential oils for the control of
359 wet bubble disease of *Agaricus bisporus*. *South African Journal of Botany* 76:681-685 DOI
360 [10.1016/j.sajb.2010.07.018](https://doi.org/10.1016/j.sajb.2010.07.018).
- 361 **Silvestro D, Michalak I. 2012.** RaxmlGUI: a graphical frontend for RAxML. *Organisms Diversity*
362 *& Evolution* 12:335-337 DOI [10.1007/s13127-011-0056-0](https://doi.org/10.1007/s13127-011-0056-0).

- 363 Song JL, Lu N, Wang WK, Yuan WD, Li HB, Cheng JW, Kang XP, Yan J. 2019. Molecular
364 identification of *Sanghuangporus* species *S.sanghuang* and *S.vaninii*. *Journal of Sichuan*
365 *University (Natural Science Edition)* **56**:765-770 DOI [10.1046/j.1365-2672.1999.00709.x](https://doi.org/10.1046/j.1365-2672.1999.00709.x).
- 366 Song JL, Yuan WD, Zhou ZF, Wang WK, Lu N, Cheng JW, Yan J. 2020. Activities of
367 extracellular enzymes and polysaccharides in liquid culture of *Sanghuangporus*.
368 *Mycosystema* **39**:352-361 DOI [10.13346/j.mycosystema.190211](https://doi.org/10.13346/j.mycosystema.190211).
- 369 Sun PL, Xu SY, Yang K, Zhang J, Song, L. 2006. Advance of researches on *Phellinus* spp, a rare
370 and precious medicinal fungus. *Microbiology China* **33**:119-123 DOI [10.3969/j.issn.0253-2654.2006.02.025](https://doi.org/10.3969/j.issn.0253-2654.2006.02.025).
- 372 Tomah AA, Alamer ISA, Li B, Zhang JZ. 2020. A new species of *Trichoderma* and gliotoxin role:
373 A new observation in enhancing biocontrol potential of *T. virens* against *Phytophthora capsici*
374 on chili pepper, *Biological Control* **145**:104261 DOI [10.1016/j.biocontrol.2020.104261](https://doi.org/10.1016/j.biocontrol.2020.104261).
- 375 Vinale F, Manganiello G, Nigro M, Mazzei P, Piccolo A, Pascale A, Ruocco M, Marra R,
376 Lombardi N, Lanzuise S, Varlese R, Cavallo P, Lorito M, Woo SL. 2014. A novel fungal
377 metabolite with beneficial properties for agricultural applications. *Molecules* **19**:9760-9772
378 DOI [10.3390/molecules19079760](https://doi.org/10.3390/molecules19079760).
- 379 Velázquez-Cedeño M, Farnet AM, Billette C, Mata G, Savoie JM. 2007. Interspecific
380 interactions with *Trichoderma longibrachiatum* induce *Pleurotus ostreatus* defence reactions
381 based on the production of laccase isozymes. *Biotechnology Letter* **29**:1583-1590 DOI
382 [10.1007/s10529-007-9445-z](https://doi.org/10.1007/s10529-007-9445-z).
- 383 Wang WH, Yang Y, Zhu LN, Jia W, Zhang JS, Liu YF, Yan MQ, Zhao XL, Zhang K, Zhang
384 HN, Fan H. 2018. Inhibitory effects of ethanol extract of *Sanghuangporus sanghuang* fruiting
385 bodies on SW620 colon cancer cells. *Mycosystema* **37**:1751-1760 DOI
386 [10.13346/j.mycosystema.180185](https://doi.org/10.13346/j.mycosystema.180185).
- 387 Woo SL, Kubicek CP, Druzhinina IS, Vinale F, Cavallo P, Lorito M. 2009. Characterization of
388 *Trichoderma* species associated with the production of *Pleurotus ostreatus* in Italy. *Journal of*
389 *Plant Pathology* **91**:S4.94.
- 390 Wu SH, Dai YC. 2020. Species clarification of the medicinal fungus *Sanghuang*. *Mycosystema*
391 **39**:781-794 DOI [10.13346/j.mycosystema.190354](https://doi.org/10.13346/j.mycosystema.190354).
- 392 Zang C, Kong T, Liang B, Liu X, Xie J, Lin Y, Pei X, Yu S, Liang C. 2023. Evaluation of imide
393 substance from *Streptomyces atratus* PY-1 for the biocontrol of *Phytophthora* blight.
394 *European Journal of Plant Pathology* **165**:725-734 DOI [10.1007/s10658-023-02648-5](https://doi.org/10.1007/s10658-023-02648-5).
- 395 Zhu L, Song J, Zhou JL, Si J, Cui BK. 2019. Species diversity, phylogeny, divergence time, and
396 biogeography of the genus *Sanghuangporus* (Basidiomycota). *Frontiers in Microbiology* **10**:812
397 DOI [10.3389/fmicb.2019.00812](https://doi.org/10.3389/fmicb.2019.00812).

Figure 1

Figure 1. Disease symptoms during the cultivation of *S. vaninii* and after artificial inoculation. 

A-B: During the cultivation of *S. vaninii*. Pathogen hyphae covering the surface of the initial fruiting body and exposed substrate. C: After inoculation. Hyphae inoculated with isolate YZB-1 covering the surface of the substrate and surrounding fruiting body. D: Normally growing *S. vaninii* fruiting body.



Figure 2

Figure 2. Colonies and microscopic photographs of pathogenic fungi.

YZB-1 grown on PDA, CMD or SNA in 9-cm-diam Petri dishes under 12 h darkness /12 h light for 7d. A: On PDA. B: On CMD. C: On SNA. D-E: Conidiophores and phialides conidia. D =100 μm ; E =10 μm .




Figure 3

Figure 3. The phylogenetic tree generated from the ITS sequences of *Trichoderma* spp.

Branch length values shorter than 0.01 were omitted.




Figure 4

Figure 4. The diameters of *T. virens* strain YZB-1 at different temperatures.

The error bars indicate the standard deviation, and different letters indicate significantly different values ($P < 0.05$).




Figure 5

Figure 5. The dual-culture of *T. virens* strain YZB-1 and *S. vaninii*.

A-B: *S. vaninii* (A) and *T. virens* × *S. vaninii* (B) were incubated at 25 °C. C: the inhibition ratios of *S. vaninii* by *T. virens* at different temperatures. The error bars indicate the standard deviation, and different letters indicate significantly different values ($P < 0.05$).

