

Farmers' perception of the ecosystem services provided by diurnal raptors in arid Rajasthan (#75431)

1

First submission

Guidance from your Editor

Please submit by **18 Nov 2022** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

7 Figure file(s)
2 Table file(s)
1 Raw data file(s)
2 Other file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Farmers' perception of the ecosystem services provided by diurnal raptors in arid Rajasthan

Govind Tiwari¹, Puneet Pandey², Randeep Singh^{Corresp. 1}

¹ Amity Institute of Forestry and Wildlife, Amity University, Noida, Uttar Pradesh, India

² Enprotec India Foundation, Dehardun, Uttarakhand, India

Corresponding Author: Randeep Singh
Email address: randeep04@rediffmail.com

Understanding the farmer's perceptions, attitude, behaviour, and knowledge toward conservation is critical in developing an effective conservation programme in human-dominated landscapes. Farmers are the most important stakeholders in wildlife conservation in the agricultural landscape. We conducted semi-structured face-to-face interviews with 373 farmers to understand the farmer's perception of ecosystem services provided by diurnal raptors in the arid region of Rajasthan from July 2020 to February 2021 and from August 2021 to January 2022. We grouped ecosystem services and disservices into larger categories and estimated the correlation between them, finding that disservices are negatively correlated with benefits. Raptors were perceived as beneficial for their role in controlling rodents and pests, but negatively for poultry predation. In addition, we built a binomial generalised linear model with a logit function to better understand the factors that influence farmers' perceptions of raptors (positive or negative). We observed that males and females have different attitudes toward the ecosystem services provided by raptors. It is critical to understand social perceptions in order to conserve species that are rare on a global scale but may face negative perceptions on a local scale. Our study connects ecological information with socio-demographic factors, which can be useful in developing policy measures for raptor conservation.

To
The Editor
Peer J

Subject: Submission of the manuscript **Farmers' Perception of the Ecosystem Services Provided by Diurnal Raptors in Arid Rajasthan** for publication in Peer J Journal - Regd.

Dear Sir,
Submission of the manuscript **Farmers' Perception of the Ecosystem Services Provided by Diurnal Raptors in Arid Rajasthan**. We are submitting our manuscript as per Peer J celebrating 10-year completion and publishing paper related to conservation biology without any cost. Our manuscript belongs to conservation biology category and we are not unable to pay the APC charges.

Genesis of the manuscript:

Understanding the farmer's perceptions, attitude, behaviour, and knowledge toward conservation is critical in developing an effective conservation programme in human-dominated landscapes. Farmers are the most important stakeholders in wildlife conservation in the agricultural landscape. We conducted semi-structured face-to-face interviews with 373 farmers to understand the farmer's perception of ecosystem services provided by diurnal raptors in the arid region of Rajasthan from July 2020 to February 2021 and from August 2021 to January 2022. Our study connects ecological information with socio-demographic factors, which can be useful in developing policy measures for raptor conservation. We have uploaded the minimal data of collected during survey as a supporting information_S1. We have provided survey GPS location in raw data.

I am being the corresponding author would like to undertake that:

- The contents of this manuscript will not be copyrighted, submitted, or published elsewhere, while acceptance by the Journal is under consideration.
- All authors are agreed to the contents or views expressed in this manuscript and approved for submission of the present manuscript.
- There are no directly related manuscripts or abstracts, published or unpublished, by any authors of this paper;
- My Institute's authorities are fully aware of this submission.

We do not oppose any reviewers.

We look forward to future correspondence with you and the reviewers of this manuscript. Please feel free to contact me with any questions or requests regarding the submission of this manuscript. I hope the enclosed manuscript to be suitable for publication in your esteemed journal and look forward to hear you.

Sincerely yours,

Randeep Singh

Farmers' Perception of the Ecosystem Services Provided by Diurnal Raptors in Arid Rajasthan

Govind Tiwari^a, Puneet Pandey^b, Randeep Singh^{a*}

^aAmity Institute of Forestry and Wildlife, Amity University Uttar Pradesh, Gautam Buddha Nagar,
Uttar Pradesh, India

^bEnprotec India Foundation, Dehradun, Uttarakhand, India

*** Corresponding Author**

Randeep Singh: rsingh18@amity.edu

Contact:91-9643210059

Running Title: Perception of Farmers towards Raptors

Abstract

Understanding the farmer's perceptions, attitude, behaviour, and knowledge toward conservation is critical in developing an effective conservation programme in human-dominated landscapes.

Farmers are the most important stakeholders in wildlife conservation in the agricultural landscape.

We conducted semi-structured face-to-face interviews with 373 farmers to understand the farmer's perception of ecosystem services provided by diurnal raptors in the arid region of Rajasthan from July 2020 to February 2021 and from August 2021 to January 2022. We grouped ecosystem services and disservices into larger categories and estimated the correlation between them, finding that disservices are negatively correlated with benefits. Raptors were perceived as beneficial for their role in controlling rodents and pests, but negatively for poultry predation. In addition, we built a binomial generalised linear model with a logit function to better understand the factors that influence farmers' perceptions of raptors (positive or negative). We observed that males and females have different attitudes toward the ecosystem services provided by raptors. It is critical to understand social perceptions in order to conserve species that are rare on a global scale but may face negative perceptions on a local scale. Our study connects ecological information with socio-demographic factors, which can be useful in developing policy measures for raptor conservation.

Keywords: Arid region, community perception, ecosystem services, farmers, generalised linear model, raptors, India

Introduction

The importance of the social-ecological perspective or social dimensions (i.e., attitudes, beliefs, perceptions, or values) in human-dominated agricultural landscapes around the world is now recognised for biodiversity conservation and informing policymakers and land-use managers (Bennet et al. 2016, Pooley et al. 2017, Morales-Reyes et al. 2018). Agriculture is a dominant land use in many countries (Cai and Pettenella 2013), and agricultural landscapes provide refuge and habitat for a variety of wildlife species (Perrings et al. 2006). As a result, policymakers are constantly encouraging farmers to protect the habitat of many threatened species on their farms (Kross et al. 2018). The loss of biodiversity in agricultural landscapes is linked to the loss of benefits obtained from ecosystems (Perrings et al. 2006, Morandin et al. 2016). As a result, it is critical to comprehend the relationship between benefits obtained or ecosystem services and biodiversity conservation in agricultural areas (Gorosábel et al. 2022).

Many species (i.e., insects, birds, and rodents) are known crop pests in agricultural landscapes; they directly cause harm to farmers by damaging crops, which can result in reduced productivity or increased production costs (Zhang et al. 2007, Sekercioglu et al. 2016, Garcia et al. 2020). Raptors or birds of prey, on the other hand, are highly valued in agroecosystems because they significantly control pest abundance or activity and act as intraguild predators. They offer biological crop pest control (Belaire et al. 2015, Kross et al. 2016, Shave et al. 2018, Garcia et al. 2020), which benefits farmers indirectly (Kross et al. 2016). Raptors are a sustainable pesticide alternative that not only provides economic benefits but also reduces pest outbreaks (Naranjo et al. 2015). It also has a positive socio-ecological impact by lowering human health risks and preserving biodiversity (Gibbs et al. 2009, Sarwar 2015).

However, raptors face significant threats in agricultural landscapes due to a variety of anthropogenic activities such as intensive agriculture practices, the use of pesticides to maintain food production, land use change, widespread deforestation, habitat alterations, hunting, and trade (Gibbs et al. 2009). Indeed, anthropogenic threats are cited as one of the major causes of decline in the ecological or ecosystem services provided by raptors around the world (Emmerson et al. 2016, Rusch et al. 2016). Raptor conservation in agricultural landscapes is dependent on farmers' knowledge, behaviour, farm practices, and attitudes. Because they are the first stockholders, their direct and indirect involvement in raptor conservation decision-making is critical. Farmers' ecological knowledge can provide important information about raptor distributions, breeding, threats, and ecosystem services in agricultural landscapes (Gaston et al. 2018; Kross et al. 2018). As a result, raptor protection is heavily reliant on farmers' knowledge, attitudes, and perception. Few studies have documented farmers' knowledge and attitudes toward the ecological importance and value of raptors' ecosystem services. The relationships between functional traits of organisms and provisioning and regulating ecosystem services are well established, but the traits that underpin the benefits derived from cultural ecosystem services are not (Zoeller et al. 2020), and the contribution of raptors to cultural ecosystem services such as sense of place or education is unknown (Echeverri et al. 2018).

The current study aims to understand farmers' perceptions of raptors in Rajasthan's arid region, as well as the socioecological factors that influence whether raptors are viewed as a source of benefits or a source of damage by farmers (Fig. 1). Organic agriculture has gained popularity in the arid region of Rajasthan in recent years (Dangour et al. 2010), and its potential as a development strategy for rural communities is recognised (Panwar et al. 2010). Farmers who grow

organic crops have few pests control options, and raptors serve as a natural biological pest control agent in cropland (Costa et al. 2019, Van Bruggen et al. 2016).

As a result, we hypothesised that raptors are more beneficial to farmers growing organic crops in the region. Furthermore, rural communities in the study area rely primarily on agriculture for a living, with small poultry operations supplementing household income (Ithika et al. 2013). Male farmers are primarily responsible for livelihood (tourism, agriculture, crop protection, and animal husbandry), whereas female farmers are responsible for the household, livestock grazing, fodder/wood collection, and poultry (Kumar et al. 2021). Female farmers interact with livestock and poultry more directly than male farmers (Mohapatra and George 2021).

As a result, we hypothesised that male farmers interact with raptors more than female farmers, and thus have different perceptions of raptors. In addition, we assess farmers' attitudes toward other species (bats and perching birds) using the same criteria. The current study aimed to collect baseline data for the forest department and policymakers to use in developing conservation and management plans for raptors in the agriculture ecosystem. Through community outreach programmes, the forest department and conservation organisations can initiate education awareness programmes to improve farmers' knowledge of ecology and ecosystem services provided by raptors in agriculture ecosystems for future conservation initiatives in the region.

Females perceive raptors differently than males.

Figure 1. Flowchart explaining connection between ecosystem services and human perception (Zoeller et al. 2020).

Material and methods

Study area

In the hot arid region of Rajasthan, India, we studied a community of diurnal raptors (Fig. 2). The study area covered 0.198 million square kilometres and was located between 24°31' and 30°12' north latitudes and 69°15' to 76°42' east longitudes. The region is characterised by low and erratic rainfall, with an average annual rainfall of 500 mm, 90 percent of which falls during the monsoon season (Mohranna et al. 2012). Temperatures can range from 0°C in the winter to 50°C in the summer. The terrain is slightly undulating within the venue of sand deposited by inland drainage and streams, with salt lakes and limited water resources and arable lands (Sharma and Sharma 2004). Man's reliance on animal rearing, combined with a sparse and nomadic population (Singh and Kumar 2015). Northern tropical thorn forests (Champion and Seth Classification 6B), which include *Calligonum polygonoidis*, *Prosopis cineraria*, *Prosopis juliflora*, *Acacia capparis*, *Acacia Senegal*, *Acacia catechu*, *Anogeissus pendula*, *Butea monosperma*, and *Azadirachta indica*, cover the rolling arid landscape. Anthropogenic activities have an impact on the landscape because 22.5 million people live there, making it the world's most populous desert at a density of about 84 people per square kilometre (Singh and Kumar 2015). The majority of residents' occupations (70%) are farming, raising livestock, and mining. This area is home to numerous residents and migratory raptors despite its harsh climate and man-made limitations. The existence of so many raptor species in the arid region can be attributed to both socioeconomic and climatic factors (Chhangani 2007).

Figure 2: Location map of (a) study area and (b) sampling location in arid region Rajasthan during July 2020 to February 2022

Methods

Data Collection

From July 2020 to January 2022, we conducted face-to-face interviews with 373 respondents (Supporting information S2) using semi-structured questionnaires (Supplementary file 1). There were three main sections to the questionnaire: (A) sociodemographic profile of the respondents, (B) details on how farmers feel about raptors and the ecosystem services they provide, and (C) details on how they feel about other species in the area. On a five-point Likert scale (1-Strongly disagree to agree 5-Strongly) (Likert 1932), respondents were asked to rank raptors according to their subjective agreement with nine different statements. These items discussed the ecosystem benefits and harms that raptors provide (Martinez et al., 2020) and disservices that raptors provide (Echeverri et al. 2018, Zoeller et al. 2020).

To measure the attitudes of male and female respondents for ecosystem services offered by other avian species the data was gathered on a Likert scale (Likert 1932, Kross et al. 2018) and then compared with the ecosystem services offered by the raptors. The following questions and details were used to elicit responses regarding respondents' opinions on the trend of the raptor population: of the species that you see in your area, did you see them more, less, or about the same as you did in previous years: 0 (no change), 1 (increasing), and 1 (decreasing) (Morales Reyes et al. 2018). The survey was carried out utilising convenience sampling (Som 2020). People were informed of the purpose of the study before they participated in interviews, and only then did they give their informed consent.

Data analyses

We used regression analysis to examine the relationship between farmers' perceptions of species population trends and their perceptions of the services (Supporting information S2) offered by vultures and raptors (Morales reyes et al. 2018). The flow chart of analysis is described in Fig. 3. The responses to various ecosystem services were then divided into groups according to gender

and broad categories (Table 1). To verify the data's internal consistency, Cronbach's alpha for the variables was estimated (Cronbach 1951). To reduce the dimensionality of the variables, explanatory factor analysis was performed on Likert scale, which produced three different items that represented nine different services. Scree plots were used to estimate the number of factors (Supporting information S3, S4). Factanal function was used to divide the likert scale items into three major categories (Table 1) of ecosystem services provided by raptors (Echeverri et al. 2019, Zoeller et al. 2020). For both male and female respondents, we also calculated pairwise correlation across ecosystem services. We created a logit-based binomial generalised linear model (GLM) (Luoto and Hjort 2004, MacKenzie 2018). The sociodemographic data of respondent's was kept as a predictive variable, and their perception of raptors, whether they were helpful or harmful, was kept as an explanatory variable. ANOVA with a post-hoc-Tukey Honest Significant Difference (HSD) test was employed to determine whether there was a statistically significant difference between the perceptions of the male and female respondents about the ecosystem services that raptors provide. Tukey's HSD tests are conservative because they lessen the chance of a Type I error in addition to allowing comparisons between groups with multiple categories (Abdi and Williams 2010, Nanda et al. 2021). To compare respondents' perceptions with those of the other species present in the area (bats and perching birds), perceptions of the respondents were also collected for those species (Supporting information S5 - S10). The "CAR" (Fox and Weisberg 2019), "ggplot2" (Wickham 2016), "Psych" (Revelle 2022), and "Corrplot" (Wei and Simko 2021) packages were used to analyse all the data in R (R Core Team 2020). The open-source, free QGIS software was used to prepare the location map (QGIS 2021).

Results

Perception of Indirect Benefits: There was a significant difference between male and female respondents' perceptions of how raptors affect crop quality and production ($p = .0007$ and $p = .0001$, respectively). Regarding the impact of raptors on overall yield, there was no discernible difference between the opinions of the two categories of respondents ($p = .852$).

Perception of Services: There was no discernible difference between male and female respondents' perceptions of raptors' detrimental effects on pollinators ($p = .021$), poultry ($p = .013$), and livestock ($p = .002$) (Fig. 4).

Perception of Direct Benefits: There was no significant difference in male and female respondents' perceptions of the role of raptors in controlling rodents ($p = .013$) or insects ($p = .002$), but there was a significant difference in their perceptions of whether raptors can serve as an alternative to pesticides ($p = .002$) (Supporting information S11). Compared to conventional farmers, organic farmers were more tolerant of raptors (Fig. 5).

For both vultures and raptors, the regression plot (Fig. 6) shows that farmers frequently view species as advantageous if they believe their population is declining. The results of the factor analysis show that there are three factors among the items on the Likert scale (Table 1). These variables were interpreted as various categories that stood in for various ecosystem benefits and drawbacks. Disservices were found to be negatively correlated with ecosystem services and other categories for both male and female respondents, according to pairwise correlations across both categories. All the ecosystem services' advantages were adversely correlated with their disadvantages. The "Dr. Jekyll and Mr. Hyde paradox" is demonstrated by the respondents'

perceptions of the same species as both harmful and advantageous. For male respondents, the strength of the correlations was greater. According to GLM analysis (Table 2), growing fruit crops and seed crops were the main factors influencing people's favourable attitudes toward raptors ($p=0.02$ and $p0.001$, respectively).

Discussion

The concept of ecosystem services has gained widespread acceptance as a way for people to express the values they attach to different ecosystem functions (Ferreira et al 2018). Studies of locals' perceptions can offer crucial information for observing, comprehending, and interpreting the social impacts and ecological results of conservation. Our findings, which demonstrate how farmers view raptors, highlight the need for ongoing research, focused outreach efforts, and legislative measures that give farmers the information they need to choose wildlife-friendly agricultural practices (Kross et al. 2016). According to the impact they are having, raptors are seen by the respondents as both beneficial and harmful. According to our findings, they can improve pest control (Raimilla and Rau 2017), but they can also cause negative perceptions due to poultry predation. Additionally, raptors were viewed as being extremely beneficial for fruit growers' produce because they keep rodents and other pests off the farm. Putting up nest boxes to draw raptors can help reduce rodent populations on farms (Coles et al. 2019, Paz Luna et al. 2020). None of the respondents confirmed that any of the raptor species in the study area engaged in frugivory (Fitzsimons and Leighton, 2021).

Both male and female organic farmers perceived positive attitude towards raptors and were willing to spend for their conservation. Among respondents, most of the organic farmers believed that their cropping method can also be helpful in conservation of raptors, this point is also reflected in

the study completed by Kirk et al. (2020). Integrated pest management (IPM) is a decision-based process involving coordinated use of multiple tactics for optimizing the control of all classes of pests (insects, pathogens, weeds, vertebrates) in an ecologically and economically sound manner. It involves regular monitoring of pests, and their natural enemies (Ehler 2006). Raptors play a effective role in controlling damage to crops by feeding on pests (Peisley et al. 2017, Gorosabel 2022) and are important part of IPM (Zagorski 2019). The rodent population can be controlled by providing raptors with adequate conservation as suggested by Antkowiak (2004), which will also lower the cost of farming inputs (Machar et al. 2017). They can serve as an alternative to pesticides and reduce the impact of these harmful chemicals have on food chain (Maria et al. 1996, Hughes et al. 2013). As conventional farming is more common than organic farming, further research is needed to understand role of the raptors for controlling insect pests. Views of female respondents on the effect of raptors on livestock varied quite significantly from that of males as most of the female respondents spend more time with their livestock and were out with them in grazing areas for hours in search of fodder. Also, it was observed that perception of raptors and vultures as beneficial depends on level of rareness of species in terms of perceived population. Positive relationship between rareness of species and perception was firstly reported by (Courchamp et al. 2015, Hall et al. 2015). There is an opposite relationship between species rareness and perception of species as providers of ecosystem services (Morales Reyes et al. 2018). General public gives more value to rare species relative to common ones (Angulo and Courchamp 2009). It is a common belief that attitudes and perceptions towards a species are influenced by the degree of its rarity. Although, it was reported that rareness in terms of distribution cannot be a criterion in the decision for investing on conservation of the species (Martin-Lopez et al. 2007). Elusive species which are globally considered as endangered and are least known are rarely perceived as emblematic (Cortés-

Avizanda et al. 2022). Our results on rarity and perception towards a species are in accordance with study done by (Otsuka et al. 2016), which indicates that farmers have species specific view that incorporate cultural and aesthetic value of rare species and they prefer usefulness of these species over other.

Negative correlations between disservices and other cultural ecosystem services suggest that the categories are dependent on each other. People are influenced by general positive or negative effects when judging disservices and benefits. It suggests that likeability of respondents towards raptors was positively correlated with direct and indirect benefits while negatively correlated with the disservices. This “Dr. Jekyll and Mr. Hyde” paradox (Morales reyes et al. 2018) can be understood by socio economic characteristics of the respondents who are involved in poultry management cites raptor predation of chickens as a loss to their livelihood and livestock owners view raptors as a threat to the newborn cattle and a carrier of disease while fruit growing and seed growing farmers and those practicing organic agriculture perceive raptors as beneficial in their effect of controlling rodents and pest. Strength of correlation was slightly more for the male respondents. It may be explained by the fact that in this region male respondents are more involved in farming, poultry management, nature guides and transhumance and their interaction with raptors is more as compared to female respondents.

Implementing long term conservation plans needs taking socio perspective in consideration, wrong perception of a species can be detrimental for its survival (Ceríaco 2012).To change farmers’ behaviours toward more sustainable conservation of farmland biodiversity, instruments should aim to influence individual farmer’s motivation and behaviour. However, a lack of knowledge of farmers’ opinions toward wildlife can lead to poor integration of conservation measures (Katuwal et al. 2021, Kross et al. 2018). We should aim to place farmland biodiversity “in the hands and

minds” of farmers (Ahnström 2009). Without an appreciation of the human dimension to problems of conflict, sustaining species outside protected areas may be difficult (Lee and Priston 2005).

Conclusion

For the conservation of raptors, it required landscape-based approach beyond the protected areas. Very few resources and funding are allocated for the conservation of the raptors residing outside protected areas. Arid region of Rajasthan is home of many species of raptors but the overall conservation planning for raptors needs to include a socio-ecological perspective. Designing education and awareness programs along with community participation can reduce conflict with raptor in rural regions and will be beneficial for implementation of long-term conservation programs.

References

- Abdi, H., & Williams, L. J. (n.d.). *Tukey's Honestly Significant Difference (HSD) Test*.5.
- Ahnström, J. (2009). *Farmland Biodiversity—In the Hands and Minds of Farmers*.
- Angulo, E., & Courchamp, F. (2009). Rare species are valued big time. *PloS One*, 4(4), e5215.
<https://doi.org/10.1371/journal.pone.0005215>
- Antkowiak, K., & Hayes, T. (2004). *Rodent Pest Control Through the Reintroduction of an Extirpated Raptor Species*. 21(4), 4.
- Belaire, J. A., Westphal, L., Whelan, C., & Minor, E. (2015). Urban residents' perceptions of birds in the neighborhood: Biodiversity, cultural ecosystem services, and disservices. *The Condor*, 117, 192–202.
- Bennett, E., Cramer, W., Begossi, A., Cundill, G., Diaz, S., Egoh, B., Geijzendorffer, I. R., Krug, C., Lavorel, S., Lazos, E., Lebel, L., Martín-López, B., Meyfroidt, P., Mooney, H., Nel, J., Pascual, U., Payet, K., Pérez-Harguindeguy, N., Peterson, G., & Woodward, G. (2015). Linking biodiversity, ecosystem services, and human well-being:three challenges for designing research for sustainability. *Current Opinion in Environmental Sustainability*.
- Cai, M., & Pettenella, D. (2013). Protecting biodiversity outside protected areas: Can agricultural landscapes contribute to bird conservation on Natura 2000 in Italy? *Journal of Environmental Engineering and Landscape Management*, 21(1), 1–11.
<https://doi.org/10.3846/16486897.2012.663089>
- Ceríaco, L. M. (2012). Human attitudes towards herpetofauna: The influence of folklore and negative values on the conservation of amphibians and reptiles in Portugal. *Journal of Ethnobiology and Ethnomedicine*, 8, 8. <https://doi.org/10.1186/1746-4269-8-8>
- Chhangani, A. K. (2007). *Sightings and nesting sites of Red-headed Vulture Sarcogyps calvus in Rajasthan, India*. 3(6), 4.
- Coles, G., Wallis, R., & Brennan, D. (2019). *Using raptors to disperse pest birds in Victoria*. Vol. 36, no. (2019), pp. 132–136.

- 345 Cortés-Avizanda, A., Martín-López, B., Ceballos, O., & Pereira, H. M. (2018). Stakeholders
346 perceptions of the endangered Egyptian vulture: Insights for conservation. *Biological*
347 *Conservation*, 218, 173–180. <https://doi.org/10.1016/j.biocon.2017.09.028>
- 348 Costa, C. A., Guiné, R. P. F., Costa, D. V. T. A., Correia, H. E., & Nave, A. (2019). Chapter 3—
349 Pest Control in Organic Farming. In S. Chandran, M. R. Unni, & S. Thomas (Eds.),
350 *Organic Farming* (pp. 41–90). Woodhead Publishing. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-813272-2.00003-3)
351 [813272-2.00003-3](https://doi.org/10.1016/B978-0-12-813272-2.00003-3)
- 352 Courchamp, F., Angulo, E., Rivalan, P., Hall, R. J., Signoret, L., Bull, L., & Meinard, Y. (2006).
353 Rarity Value and Species Extinction: The Anthropogenic Allee Effect. *PLOS Biology*,
354 4(12), e415. <https://doi.org/10.1371/journal.pbio.0040415>
- 355 Cronbach, Lee J. (1951). "Coefficient alpha and the internal structure of tests".
356 *Psychometrika*. Springer Science and Business Media LLC. 16 (3): 297–334.
- 357 Dangour, A. D., Lock, K., Hayter, A., Aikenhead, A., Allen, E., & Uauy, R. (2010). Nutrition-
358 related health effects of organic foods: A systematic review. *The American Journal of*
359 *Clinical Nutrition*, 92(1), 203–210. <https://doi.org/10.3945/ajcn.2010.29269>
- 360 Echeverri, A., Karp, D. S., Naidoo, R., Tobias, J. A., Zhao, J., & Chan, K. M. A. (2020). Can avian
361 functional traits predict cultural ecosystem services? *People and Nature*, 2(1), 138–151.
362 <https://doi.org/10.1002/pan3.10058>
- 363 Ehler, L. E. (2006). Integrated pest management (IPM): Definition, historical development and
364 implementation, and the other IPM. *Pest Management Science*, 62(9), 787–789.
365 <https://doi.org/10.1002/ps.1247>
- 366 Emmerson, M., Morales, M., Oñate, J., Batary, P., Berendse, F., Liira, J., Aavik, T., Guerrero, I.,
367 Bommarco, R., Eggers, S., Pärt, T., Tschardtke, T., Weisser, W., Clement, L., &
368 Bengtsson, J. (2016). How Agricultural Intensification Affects Biodiversity and Ecosystem
369 Services. *Advances in Ecological Research*, 55, 43–97.
370 <https://doi.org/10.1016/bs.aecr.2016.08.005>

- 371 Ferreira, A., Zimmermann, H., Santos, R., & von Wehrden, H. (2018). A Social–Ecological
372 Systems Framework as a Tool for Understanding the Effectiveness of Biosphere Reserve
373 Management. *Sustainability*, 10(10), 3608.
- 374 Fitzsimons, J. A., & Leighton, J. (2021). Frugivory in Raptors: New Observations from Australia
375 and a Global Review. *Birds*, 2(4), 338–350.
- 376 Fox J, Weisberg S (2019). *An R Companion to Applied Regression*, Third edition. Sage, Thousand
377 Oaks CA. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- 378 Garcia, C. A., Bhagwat, S. A., Ghazoul, J., Nath, C. D., Nanaya, K. M., Kushalappa, C. G.,
379 Raghuramulu, Y., Nasi, R., & Vaast, P. (2010). Biodiversity Conservation in Agricultural
380 Landscapes: Challenges and Opportunities of Coffee Agroforests in the Western Ghats,
381 India. *Conservation Biology*, 24(2), 479–488.
- 382 Gaston, K. J., Cox, D. T. C., Canavelli, S. B., García, D., Hughes, B., Maas, B., Martínez, D.,
383 Ogada, D., & Inger, R. (2018). Population Abundance and Ecosystem Service Provision:
384 The Case of Birds. *BioScience*, 68(4), 264–272.
- 385 Gibbs, K. E., Mackey, R. L., & Currie, D. J. (2009). Human land use, agriculture, pesticides and
386 losses of imperiled species. *Diversity and Distributions*, 15(2), 242–253.
387 <https://doi.org/10.1111/j.1472-4642.2008.00543.x>
- 388 Gorosábel, A., Bernad, L., & Pedrana, J. (2022). Using the ecosystem services approach to link
389 raptors with human well-being in the Southeast Pampas of Argentina. *Biodiversity and
390 Conservation*, 1–22. <https://doi.org/10.1007/s10531-022-02372-2>
- 391 Hall, R. J., Milner-Gulland, E. J., & Courchamp, F. (2008). Endangering the endangered: The
392 effects of perceived rarity on species exploitation. *Conservation Letters*, 1(2), 75–81.
- 393 Hughes, J., Sharp, E., Taylor, M. J., Melton, L., & Hartley, G. (2013). Monitoring agricultural
394 rodenticide use and secondary exposure of raptors in Scotland. *Ecotoxicology*, 22(6), 974–
395 984. <https://doi.org/10.1007/s10646-013-1074-9>
- 396 Ithika, C.S., Singh, S.P. and Gautam, G. 2013. Adoption of Scientific Poultry Farming Practices
397 by the Broiler Farmers in Haryana. *Indian Journal of Animal Sciences*. 3(2): 417-422.

- Katuwal, H. B., Zhang, M., Baral, H. S., Sharma, H. P., & Quan, R.-C. (2021). Assessment of farmers' knowledge and perceptions towards farmland birds show the need of conservation interventions. *Global Ecology and Conservation*, 27, e01563. <https://doi.org/10.1016/j.gecco.2021.e01563>
- Kirk, D. A., Martin, A. E., & Freemark Lindsay, K. E. (2020). Organic farming benefits birds most in regions with more intensive agriculture. *Journal of Applied Ecology*, 57(6), 1043–1055. <https://doi.org/10.1111/1365-2664.13589>.
- Kross, S. M., Ingram, K. P., Long, R. F., & Niles, M. T. (2018). Farmer Perceptions and Behaviors Related to Wildlife and On-Farm Conservation Actions: Farmer perceptions of wildlife. *Conservation Letters*, 11(1), e12364. <https://doi.org/10.1111/conl.12364>
- Kross, S. M., Kelsey, T. R., McColl, C. J., & Townsend, J. M. (2016). Field-scale habitat complexity enhances avian conservation and avian-mediated pest-control services in an intensive agricultural crop. *Agriculture, Ecosystems & Environment*, 225, 140–149. <https://doi.org/10.1016/j.agee.2016.03.043>
- Kumar, R., Devi, I., Meena, R. L., & Kumar, A. (2021). Role of farm women in animal husbandry of semi-arid Rajasthan. *Indian Journal of Small Ruminants (The)*, 27(1), 118–122. <https://doi.org/10.5958/0973-9718.2021.00015.5>
- Lee, P., & Priston, N. (2005). Human attitudes to primates: Perceptions of pests, conflict and consequences for primate conservation. *Commensalism and Conflict: The Human-Primate Interface*. <https://doi.org/10.1002/047001539X.ch1>.
- Likert R. A technique for the measurement of attitudes. *Arch Psychology*. 1932;22(140):55
- Luoto, M., & Hjort, J. (2004). Generalized linear modelling in periglacial studies: Terrain parameters and patterned ground. *Permafrost and Periglacial Processes*, 15(4), 327–338. <https://doi.org/10.1002/ppp.482>
- Machar, I., Harmacek, J., Vrublova, K., Filippovova, J., & Brus, J. (2017). Biocontrol of Common Vole Populations by Avian Predators Versus Rodenticide Application. *Polish Journal of Ecology*, 65(3), 434–444.

- MacKenzie, D. I. (2018). *Occupancy estimation and modeling inferring patterns and dynamics of species occurrence*. Academic Press.
- Mohapatra, G., & George, M. (2021). Perception and adaptation of agricultural households to climate change in the semi-arid regions of Rajasthan – a gender perspective study. *Ecofeminism and Climate Change*, 2(3), 146–155. <https://doi.org/10.1108/EFCC-04-2021-0007>
- Morales-Reyes, Z., Martín-López, B., Moleón, M., Mateo-Tomás, P., Botella, F., Margalida, A., Donázar, J., Blanco, G., Pérez Ibarra, I., & Sánchez-Zapata, J. (2018). Farmer Perceptions of the Ecosystem Services Provided by Scavengers: What, Who, and to Whom. *Conservation Letters*, 11, e12392. <https://doi.org/10.1111/conl.12392>
- Morandin, L., & Winston, M. (2006). Pollinators provide economic incentive to preserve natural land in agroecosystems. *Agriculture, Ecosystems & Environment*, 116, 289–292.
- María, P., García-Fernández, A. J., Barba, A., Riboni, I., Romero, D., & Sánchez-García, J. A. (1996). Organochlorine pesticide accumulation in several species of raptors from Southeastern Spain. *Toxicology Letters*, 88, 80.
- Martínez-Sastre, R., García, D., Miñarro, M., & Martín-López, B. (2020). Farmers’ perceptions and knowledge of natural enemies as providers of biological control in cider apple orchards. *Journal of Environmental Management*, 266, 110589.
- Martín-López, B., Montes, C., & Benayas, J. (2007). The non-economic motives behind the willingness to pay for biodiversity conservation. *Biological Conservation*, 139, 67–82. <https://doi.org/10.1016/j.biocon.2007.06.005>
- Moharana PC, Sharma BM, Biswas DR, Dwivedi BS, Singh RV. Long-term effect of nutrient management on soil fertility and soil organic carbon pools under a 6-year-old pearl millet–wheat cropping system in an Inceptisol of subtropical India. *Field Crops Research*. 2012 Sep 20;136:32–41.
- Nanda, A., Mohapatra, Dr. B. B., Mahapatra, A. P. K., Mahapatra, A. P. K., & Mahapatra, A. P. K. (2021). Multiple comparison test by Tukey’s honestly significant difference (HSD): Do

- the confident level control type I error. *International Journal of Statistics and Applied Mathematics*, 6(1), 59–65.
- Naranjo, S. E., Ellsworth, P. C., & Frisvold, G. B. (2015). Economic Value of Biological Control in Integrated Pest Management of Managed Plant Systems. *Annual Review of Entomology*, 60(1), 621–645. <https://doi.org/10.1146/annurev-ento-010814-021005>
- Otsuka, K., Nakano, Y., & Takahashi, K. (2016). Contract Farming in Developed and Developing Countries. *Annual Review of Resource Economics*, 8.
- Panwar, N., Ramesh, P., Bahadur, A., & Ramana, S. (2010). Influence of Organic, Chemical, and Integrated Management Practices on Soil Organic Carbon and Soil Nutrient Status under Semi-arid Tropical Conditions in Central India. *Communications in Soil Science and Plant Analysis - COMMUN SOIL SCI PLANT ANAL*, 41, 1073–1083. <https://doi.org/10.1080/00103621003687166>
- Paz Luna, A., Bintanel, H., Viñuela, J., & Villanúa, D. (2020). Nest-boxes for raptors as a biological control system of vole pests: High local success with moderate negative consequences for non-target species. *Biological Control*, 146, 104267. <https://doi.org/10.1016/j.biocontrol.2020.104267>
- Peisley, R. K., Saunders, M. E., & Luck, G. W. (2017). Providing perches for predatory and aggressive birds appears to reduce the negative impact of frugivorous birds in vineyards. *Wildlife Research*, 44(4), 334–342. <https://doi.org/10.1071/WR17028>
- Perrings, C., Baumgärtner, S., Brock, W., Chopra, K., Conte, M., Costello, C., Duraiappah, A., Kinzig, A., Pascual, U., & Polasky, S. (2009). The economics of biodiversity and ecosystem services. *Biodiversity, Ecosystem Functioning, and Human Wellbeing: An Ecological and Economic Perspective*. Oxford University Press, Oxford, 230–247. <https://doi.org/10.1093/acprof:oso/9780199547951.003.0017>
- Pooley, S., Barua, M., Beinart, W., Dickman, A., Holmes, G., Lorimer, J., Loveridge, A., Macdonald, D., Marvin, G., Redpath, S., Sillero, C., Zimmermann, A., & Milner-Gulland, E. (2017). An interdisciplinary review of current and future approaches to improving human-predator relations. *Conservation Biology*, Early view.

- 480 QGIS.org, 2021. QGIS Geographic Information System. QGIS Association.
- 481 Revelle W (2022). psych: Procedures for Psychological, Psychometric, and Personality Research.
- 482 Northwestern University, Evanston, Illinois.
- 483 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for
- 484 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 485 Raimilla, V., & Rau, J. (2017). Human perceptions towards birds of prey: A synoptic review
- 486 centered in the customs and myths in the south-austral zone of Chile. *El Hornero*, 032(01),
- 487 139–149.
- 488 Rusch, A., Chaplin-Kramer, R., Gardiner, M. M., Hawro, V., Holland, J., Landis, D., Thies, C.,
- 489 Tschardtke, T., Weisser, W. W., Winqvist, C., Woltz, M., & Bommarco, R. (2016).
- 490 Agricultural landscape simplification reduces natural pest control: A quantitative synthesis.
- 491 *Agriculture, Ecosystems & Environment*, 221, 198–204.
- 492 <https://doi.org/10.1016/j.agee.2016.01.039>
- 493 Sarwar, A. K. M. G. (2019). Plant Biodiversity and Conservation of Higher Plants in Bangladesh:
- 494 Present Status and Future Prospects. In Sangeetha, J., Thangadurai, D., Goh, H.C. and
- 495 Islam, S. (eds.). Biodiversity and Conservation: Characterization and Utilization of Plants,
- 496 Microbes and Natural Resources for Sustainable Development and Ecosystem
- 497 Management.
- 498 Sekercioglu, C., Wenny, D., & Whelan, C. (2016). *Why Birds Matter: Avian Ecological Function*
- 499 *and Ecosystem Services*.
- 500 Sharma P, Sharma R. Groundwater Markets Across Climatic Zones: A Comparative Study of Arid
- 501 and Semi-Arid Zones of Rajasthan. Indian Journal of Agricultural Economics. 2004 Jan
- 502 1;59:138–50.
- 503 Shave, M. E., S. A. Shwiff, J. L. Elser, and C. A. Lindell. 2018. Falcons using orchard nest boxes
- 504 reduce fruit-eating bird abundances and provide economic benefits for a fruit-growing
- 505 region. *Journal of Applied Ecology* 55:2451–2460.
- 506 Singh RB, Kumar A. Climate variability and water resource scarcity in drylands of Rajasthan,
- 507 India. *Geoenvironmental Disasters*. 2015 Mar 13;2(1):7.

- SOM, R. K. (2020). *PRACTICAL SAMPLING TECHNIQUES*. CRC PRESS.
- van Bruggen, A. H., Gamliel, A., & Finckh, M. R. (2016). Plant disease management in organic farming systems. *Pest Management Science*, 72(1), 30–44.
- Wei T, Simko V (2021). R package 'corrplot': Visualization of a Correlation Matrix.
- Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
- Zagorski, M. E. (2019). *Winter ecology of raptors in cover-cropped agroecosystems in western Indiana* [Thesis, Purdue University Graduate School].
<https://doi.org/10.25394/PGS.11302805.v1>
- Zhang, W., Ricketts, T. H., Kremen, C., Carney, K., & Swinton, S. M. (2007). Ecosystem services and dis-services to agriculture. *Ecological Economics*, 64, 253–260.
<https://doi.org/10.1016/j.ecolecon.2007.02.024>
- Zoeller, K. C., Gurney, G. G., & Cumming, G. S. (2022). The influence of landscape context on the production of cultural ecosystem services. *Landscape Ecology*, 37(3), 883–894.
<https://doi.org/10.1007/s10980-022-01412-0>

Figure legends

Figure 1. Flowchart explaining connection between ecosystem services and human perception (Zoeller et al. 2020)

Figure 2: Location map of (a) study area and (b) sampling location in arid region Rajasthan during July 2020 to February 2022

Figure 3. Flowchart explaining steps followed for regression analysis (Morales reyes et al. 2018).

Figure 4. Perception of male and female respondents towards raptors in arid region Rajasthan
(**A Indirect Benefits:** (A1 Increases crop quality, A2 Increases yield, A3 Essential for crop production),
B Negative/Disservice: (B1 Causes damage to pollinators, B2 Causes damage to poultry, B3. Causes damage to livestock), **C Direct Benefits:** (C1 Controls Insects, C2 Controls Rodents, C3 Alternative to Pesticides)

Figure 5. Perception of respondents practicing conventional and organic agriculture towards raptors in arid region Rajasthan.

Figure 6. Regression plot showing perception of population of raptors/vultures vs. perception of ecosystem services by raptors/vultures in arid region Rajasthan.

Figure 7. Pairwise correlation between cultural ecosystem services as perceived by respondents in arid region Rajasthan.

Table legends

Table 1: Categorization of perception of farmers on ecosystem services in major categories (Echeverri et al. 2019, Zoeller et al. 2020).

Table 2. GLM analysis of Socio-Demographic variables and their effect on raptor perception in arid region Rajasthan.

Table 1(on next page)

Categorization of perception of farmers on ecosystem services in major categories (Echeverri et al. 2019, Zoeller et al. 2020).

- 1 Table 1: Categorization of perception of farmers on ecosystem services in major categories
- 2 (Echeverri et al. 2019, Zoeller et al. 2020)

Serial Code	Construct	Benefit and loss of raptors	Factor Loading		
A	Indirect Benefits (Cronbach's alpha=0.913)	A1. Increases crop quality	0.787	0.150	0.252
		A2. Increases yield	0.810	0.215	0.212
		A3. Essential for Crop Production	0.904	0.209	0.192
B	Negative/Disservice (Cronbach's alpha=0.769)	B1. Causes damage to pollinators	0.307		0.587
		B2. Causes damage to poultry	0.217	0.332	0.735
		B3. Causes damage to livestock	0.127	0.353	0.662
C	Direct Benefits (Cronbach's alpha=0.813)	C1. Controls Insects	0.137	0.769	0.132
		C2. Controls Rodents	0.197	0.734	0.237
		C3. Alternative to Pesticides	0.175	0.669	0.264

3
4
5

Table 2(on next page)

GLM analysis of Socio-Demographic variables and their effect on raptor perception in arid region Rajasthan.

- 1 Table 1: Categorization of perception of farmers on ecosystem services in major categories
- 2 (Echeverri et al. 2019, Zoeller et al. 2020)

Serial Code	Construct	Benefit and loss of raptors	Factor Loading		
A	Indirect Benefits (Cronbach's alpha=0.913)	A1. Increases crop quality	0.787	0.150	0.252
		A2. Increases yield	0.810	0.215	0.212
		A3. Essential for Crop Production	0.904	0.209	0.192
B	Negative/Disservice (Cronbach's alpha=0.769)	B1. Causes damage to pollinators	0.307		0.587
		B2. Causes damage to poultry	0.217	0.332	0.735
		B3. Causes damage to livestock	0.127	0.353	0.662
C	Direct Benefits (Cronbach's alpha=0.813)	C1. Controls Insects	0.137	0.769	0.132
		C2. Controls Rodents	0.197	0.734	0.237
		C3. Alternative to Pesticides	0.175	0.669	0.264

3
4
5

6 Table 2. GLM analysis of Socio-Demographic variables and their effect on Raptor perception

Coefficient	Estimate	Std. Error	Z value	Pr(>z)
(Intercept)	-16.99807	624.19538	-0.027	0.978275
Gender	0.08132	0.36984	0.220	0.825974
Fruit Crops	-0.76781	0.33034	-2.324	0.020111 *
Farming.method	1.08905	1.12765	0.966	0.334162
Education	14.83957	624.19389	0.024	0.981033
Seed Crops	-1.67898	0.43777	-3.835	0.000125 ***
Forage Crops	1.55572	0.40745	3.818	0.0546
Vegetable	0.55477	0.50588	1.097	0.272792
Livestock	0.83013	0.76202	1.089	0.275983
Null deviance: 501.08 on 372 degrees of freedom				
Residual deviance: 422.16 on 360 degrees of freedom				
AIC: 448.16, Number of Fisher Scoring iterations: 13				

7

Figure 1

Flowchart explaining connection between ecosystem services and human perception (Zoeller et al. 2020)

Flowchart explaining connection between ecosystem services and human perception (Zoeller et al. 2020)




Figure 1. Flowchart explaining connection between ecosystem services and human perception (Zoeller et al. 2020)

Figure 2

Location map of (a) study area and (b) sampling location in arid region Rajasthan during July 2020 to February 2022

Location map of (a) study area and (b) sampling location in arid region Rajasthan during July 2020 to February 2022




Figure 2: Location map of (a) study area and (b) sampling location in arid region Rajasthan during July 2020 to February 2022

Figure 3

Flowchart explaining steps followed for regression analysis (Morales reyes et al. 2018).

Flowchart explaining steps followed for regression analysis (Morales reyes et al. 2018).




Fig 3. Flowchart explaining steps followed for regression analysis (Morales reyes et al.2018).

Figure 4

Perception of male and female respondents towards raptors in arid region Rajasthan

A Indirect Benefits: (A1 Increases crop quality, A2 Increases yield, A3 Essential for crop production), **B Negative/Disservice:** (B1 Causes damage to pollinators, B2 Causes damage to poultry, B3. Causes damage to livestock), **C Direct Benefits:** (C1 Controls Insects, C2 Controls Rodents, C3 Alternative to Pesticides)




Figure 4. Perception of male and female respondents towards raptors in arid region Rajasthan

A Indirect Benefits: (A1 Increases crop quality, A2 Increases yield, A3 Essential for crop production), **B Negative/Disservice:** (B1 Causes damage to pollinators, B2 Causes damage to poultry, B3. Causes damage to livestock), **C Direct Benefits:** (C1 Controls Insects, C2 Controls Rodents, C3 Alternative to Pesticides)

Figure 5

Perception of respondents practicing conventional and organic agriculture towards raptors in arid region Rajasthan.

Perception of respondents practicing conventional and organic agriculture towards raptors in arid region Rajasthan.




Figure 5. Perception of respondents practicing conventional and organic agriculture towards raptors in arid region Rajasthan

Figure 6

Regression plot showing perception of population of raptors/vultures vs. perception of ecosystem services by raptors/vultures in arid region Rajasthan.

Regression plot showing perception of population of raptors/vultures vs. perception of ecosystem services by raptors/vultures in arid region Rajasthan.




Figure 6. Regression plot showing Perception of population of Raptors/Vultures vs. Perception of Ecosystem services by Raptors/Vultures in arid region of Rajasthan

Figure 7

Pairwise correlation between cultural ecosystem services as perceived by respondents in arid region Rajasthan.

Pairwise correlation between cultural ecosystem services as perceived by respondents in arid region Rajasthan.




Fig 7: Pairwise correlation between cultural ecosystem services as perceived by respondents in arid region of Rajasthan

A Indirect Benefits: (A1 Increases crop quality, A2 Increases yield, A3 Essential for crop production), **B Negative/Disservice:** (B1 Causes damage to pollinators, B2 Causes damage to poultry, B3. Causes damage to livestock), **C Direct Benefits:** (C1 Controls Insects, C2 Controls Rodents, C3 Alternative to Pesticides)