

A new gnathosaurine (Pterosauria, Archaeopteroedactyloidea) from the Late Jurassic of Portugal (#86679)

1

First submission

Guidance from your Editor

Please submit by **25 Jun 2023** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

6 Figure file(s)

2 Raw data file(s)



Custom checks

New species checks



Have you checked our [new species policies](#)?



Do you agree that it is a new species?



Is it correctly described e.g. meets ICZN standard?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

A new gnathosaurine (Pterosauria, Archaeopterygoidea) from the Late Jurassic of Portugal

Alexandra E. Fernandes^{1, 2, 3}, Victor Beccari^{2, 3}, Alexander WA Kellner⁴, Octávio Mateus^{Corresp. 3, 5}

¹ Department of Earth and Environmental Sciences, Ludwig-Maximilians-Universität München, Munich, Bayern, Germany

² SNSB, Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany

³ Museu da Lourinhã, Lourinhã, Portugal

⁴ Departamento de Geologia e Paleontologia, Laboratório de Sistemática e Tafonomia de Vertebrados Fósseis (LAPUG), Museu Nacional, Rio de Janeiro, Brazil

⁵ GEOBIOTEC, Department of Earth Sciences, Universidade Nova de Lisboa, Caparica, Portugal

Corresponding Author: Octávio Mateus

Email address: omateus@fct.unl.pt

An incomplete, yet remarkably-sized dentated rostrum and associated partial cervical vertebrae of a pterosaur (ML 2554) were recently discovered from the Late Jurassic (Late Kimmeridgian-Early Tithonian) Lourinhã Formation of Praia do Caniçal, of central west Portugal. This specimen exhibits features such as a spatulated anterior expansion of the rostrum, robust comb-like dentition, and pronounced rims of the tooth alveoli, indicating gnathosaurine affinities. Based on its further unique tooth and dentary morphology, a new genus and species, *Lusognathus almadrava* gen. et spec. nov., is proposed, making this the first **definitively-named** pterosaur species found within Portugal. The presence of this taxon adds yet **another ecological niche element** to the fluvio-deltaic lagoonal environment **that has been suggested to be representative** of Lourinhã Formation in the Late Jurassic, and **contributes to the diversity and distribution of gnathosaurines worldwide.**

A new gnathosaurine (Pterosauria, Archaeopterygiformes) from the Late Jurassic of Portugal

FERNANDES, Alexandra E.^{1,2,3}, BECCARI, Victor^{1,2}, KELLNER, A.W.A.⁴ & MATEUS, Octávio^{1,5}

1 - Museu da Lourinhã, Rua João Luis de Moura, 95, 2530-158 Lourinhã, Portugal

2 - SNSB, Bayerische Staatssammlung für Paläontologie und Geologie, Richard-Wagner-Str. 10, 80333 Munich, Germany

3 - Department of Earth and Environmental Sciences, Ludwig-Maximilians-Universität, Richard-Wagner-Str. 10, 80333 Munich, Germany

4 - Laboratório de Sistemática e Tafonomia de Vertebrados Fósseis (LAPUG), Departamento de Geologia e Paleontologia, Museu Nacional, Universidade Federal do Rio de Janeiro, 20940-040 Rio de Janeiro, Brazil

5 - GEOBIOTEC, Department of Earth Sciences, NOVA School of Science and Technology, Universidade NOVA de Lisboa, Campus de Caparica, Caparica, Portugal

Corresponding Author: Octávio Mateus omateus@fct.unl.pt

Submission Administrator: Alexandra Fernandes fernandes@snsb.de

Abstract:

An incomplete, yet remarkably-sized dentated rostrum and associated partial cervical vertebrae of a pterosaur (ML 2554) were recently discovered from the Late Jurassic (Late Kimmeridgian-Early Tithonian) Lourinhã Formation of Praia do Caniçal, of central west Portugal. This specimen exhibits features such as a spatulated anterior expansion of the rostrum, robust comb-like dentition, and pronounced rims of the tooth alveoli, indicating gnathosaurine affinities. Based on its further unique tooth and dentary morphology, a new genus and species, *Lusognathus almadrava* gen. et spec. nov., is proposed, making this the first definitively-named pterosaur species found within Portugal. The presence of this taxon adds yet another ecological niche element to the fluvio-deltaic lagoonal environment that has been suggested to be representative of Lourinhã Formation in the Late Jurassic, and contributes to the diversity and distribution of gnathosaurines worldwide.

Keywords: *Pterosauria*, *Pterodactyloidea*, *Gnathosaurinae*, *Jurassic*, *Portugal*

Introduction:

The known global distribution and diversity of pterosaurs **highly reinforces** their success as a group, as they are found in all continents including Antarctica (Barrett et al. 2008, Kellner

et al. 2019a), and yet their relatively sparse fossil record and often incomplete preservation (particularly when outside of Lagerstätten environments) can pose a challenge for further understanding their paleobiology, when compared with other vertebrates. Accordingly, the Jurassic of Portugal is a very productive and taxonomically diverse period concerning vertebrate fossils, especially for fishes, amphibians, plesiosaurs, ichthyosaurs, mosasaurs, dinosaurs, turtles, crocodylomorphs, and mammals (e.g. Antunes & Mateus, 2003; Guillaume et al. 2020; Malafaia et al., 2010; Mateus, 2006; Mocho et al., 2016; Pérez-García & Ortega, 2011; Puértolas-Pascual et al., 2021; Rauhut, 2001; Rotatori et al., 2020). However, despite this abundance, up to now, pterosaur material recovered from this deposit has been restricted to scant and often fragmentary isolated bones and teeth, hindering any confident taxonomic assignments. This is likely due to the physical bone fragility of pterosaurs, making their remains particularly susceptible to deterrent or destructive fossilization factors such as carcass scavenging or later taphonomic duress.

On a worldwide level, while ctenochasmatids occur with some regularity throughout the fossil record landscape for pterosaurs, gnathosaurines are significantly more rare (Barrett et al., 2008). The currently-known gnathosaurine temporal range spans from the Late Jurassic to the Early Cretaceous, and although their distribution has so far extended throughout Europe, Asia, and South America, most of the known occurrences are attributed to isolated teeth, which have been reported from England, Portugal, Morocco, Chile, Uruguay, China, and Japan (Barrett et al., 2008; Howse & Millner, 1995; Knoll, 2000; Martill et al., 2006; Perea et al., 2018; Soto et al. 2021; Sweetman & Martill, 2010; Zhou et al., 2016; Dong, 1982; Unwin et al., 2000).

The fossil material herein described now introduces a new taxon, which adds yet another ecological niche element to the fluvio-deltaic lagoonal environment that has been suggested to be representative of Lourinhã Formation in the Late Jurassic, and contributes to the diversity and distribution of gnathosaurines worldwide. The material is housed at the Museu da Lourinhã, in Lourinhã, Portugal, under the collection number ML 2554.

History and Record of Pterosaur Discoveries in Portugal:

The presence of pterosaurs in Portugal was first reported by Lapparent & Zbyszewski (1957: p. 58), in respect to four small, elongated vertebrae from the Campanian/Maastrichtian of Viso, which were initially identified as being similar to *Dimorphodon* or *Rhamphorhynchus*, and later ascribed to a maniraptoran theropod dinosaur by Galton (1994). Galton also claimed in the same work that a pterodactyloid cervical vertebra from the middle portion of the neck, likely the fifth (formerly MSGP N X 213, and now under the institutional acronym MG, belonging to the Museu Geológico in Lisbon, Portugal) that was discovered in the Lower Cretaceous Barremian of Serra Tiago dos Velhos was actually the first record of a pterosaur from Portugal, which he ascribed to cf. *Ornithocheirus*.

In 1968, from the Kimmeridgian beds of the Guimarota mine, Kühne initially attributed remains (teeth and a terminal manual phalanx) to two different groups of pterosaurs: aff. *Rhamphorhynchus* sp. and *Pterodactylus* sp. (Kühne, 1968; Thulborn, 1973; Wiechmann & Gloy, 2000). Later, this material, in conjunction with other isolated fragmentary remains composed of

a left scapula, wing? elements, and other appendicular elements could only be referred to the Pterodactyloidea, family incertae **sedis**. Guimarota has also yielded over 300 isolated pterosaur teeth, attributed to *Rhamphorhynchus* sp. and to the Pterodactyloidea (Wiechmann & Gloy, 2000). The fossils from Guimarota are also deposited in the Museu Geológico, in Lisbon, Portugal.

Other isolated teeth comprise the vast majority of pterosaur fossils from Portugal, and were recovered from the Areia do Mastro locality of the lower Barremian Papo Seco Formation of Cabo Espichel. They were assigned to the Ornithocheiridae and to the Ctenochasmatoidea (Figueiredo et al., 2020, 2022). From the Andrés fossil site of the Kimmeridgian Alcobaça Formation, isolated teeth were also attributed to Pterosauria indet. (Malafaia et al., 2010), based on their needle-like morphology (and similar in morphology to the *Rhamphorhynchus* sp. Guimarota teeth). Isolated teeth were also recovered in the area of Valmitão in Lourinhã (Guillaume et al., 2020), and teeth assigned to *Gnathosaurus* sp. were reported from the Sobral Formation (late Kimmeridgian-early Tithonian) locality of Praia Azul in Torres Vedras (Bertozzo et al., 2021).

Some fragmentary appendicular bones have been tentatively attributed to pterosaurs, including a large femur attributed to Dsungaripteroidea, from the locality Praia da Almagreira, Peniche (Bertozzo et al., 2021) of the upper Kimmeridgian/lower Tithonian. Some indeterminate pterosaur bone fragments were also reported from the Upper Cretaceous sediments of Taveiro, without any particular taxonomic assignment (Antunes & Pais, 1978).

About 400 *Pteraichnus* tracks have been collected from the Kimmeridgian/Tithonian Amoreira-Porto Novo Member locality of Peralta, which may correspond to the new taxon described in this work, although a more decisive attribution of these ichnofossils is beyond the scope of the current contribution. Trackways of *Pteraichnus* sp. (Mateus & Milán, 2010) are also reported from the Upper Jurassic (Kimmeridgian) Zambujal de Baixo locality of the Azóia Formation in Sesimbra and Porto das Barcas in Lourinhã (late Kimmeridgian-early Tithonian).

Overwhelmingly, the vast majority of previous pterosaur specimens from Portugal have been assigned to *Rhamphorhynchus* sp. However, this was likely a historically-generalized temporal attribution (rather than strictly based on morphology), and the known Portuguese material needs to be revised, especially considering that the range of paleobiogeographical variability has substantially grown since the time of these original attributions. In fact, no definitive non-tooth skeletal material of *Rhamphorhynchus* sp. has ever even been reported in Portugal to date, which further warrants the potential for misnomers.

Here we present and describe a new specimen found by Filipe Vieira in November 2018 at Praia do Caniçal, in the municipality of Lourinhã in central west Portugal. Further excavation efforts realized in March 2019 by members of the Museu da Lourinhã collected more material from this same specimen, which is the first pterosaur taxon named from this **country**.

Geographical and Geological Setting:

At Praia do Caniçal the outcropping rocks are of the Upper Jurassic of the Lourinhã Formation, namely the middle member of this unit: the Praia Azul Member. The Lourinhã Fm.

displays a succession of alternated terrestrial mudstones and sandstones, some paleosol levels, and three transgressive bioclastic brackish layers. The formation is mostly continental, as evidenced by the presence of terrestrial fauna such as lissamphibians, non-aquatic mammals, and dinosaurs. The paleoclimate was arid (Myers et al., 2012) and the paleolatitude, derived from paleolatitude.org (van Hinsbergen et al., 2015), is estimated as 28-29° North.

The Praia Azul Member is regarded to have been deposited near-shore, due to sea transgression. It is characterized by marls, mudstones, and sandstones, and is composed of three conspicuous carbonate levels (with the lower and upper levels used as the lithostratigraphic boundaries) and can be traced over distances of 20 km. The unit was mainly deposited by meandering fluvial systems flowing in a low-lying coastal plain, connected with transitional systems like deltas, sandy bay shorelines, and brackish lagoons. The brackish faunas of the shelly layers indicate short-time marine incursions (see Manuppella et al 1999; Hill, 1989; Martinius and Gowland, 2011; Myers et al., 2012; Taylor et al., 2014; Mateus et al., 2017). So far, all studies agree to an Upper Kimmeridgian-Lower Tithonian age of those layers, which is also supported by an only non-biostratigraphic inferred date, provided by strontium isotopes by Schneider et al. (2009). More precisely, the second and middle transgressive layer is at the Kimmeridgian-Tithonian boundary, which is currently at 149.2 Ma.

The Praia Azul Mb. is composed of three transgressive carbonate layers with different faunal assemblages: the lower level is categorized by *Isognomon lusitanica*, *Eomiodon securiformes*, *Arcomytilus morrisii*, dinosaur tracks, and ostreids (outcropping north of Caniçal, below the Paimogo fort). The middle level is comprised by *Jurassicorbula edwardsi*, *Isognomon lusitanica*, *Nerinea*, coprolites, fish, *Eomiodon securiformes*, echinoids (outcropping at the base of Praia do Caniçal), while the upper level fauna consists of *Jurassicorbula edwardsi*, abundant ostreids, pleurosternidae turtles, and *Isognomon lusitanica* (outcropping at the top of Praia do Caniçal). The member is known to bear abundant dinosaur remains, including bones, eggs, and tracks (Ribeiro et al., 2014; Mateus et al., 2011), and also crocodylomorphs, turtles, pterosaurs, mammals, lissamphibians, and **lepidosaurs**.

The specimen studied here was excavated from a stratigraphic layer at modern intertidal sea level, below the sand level of the beach itself, which is only exposed at low tide during the winter months (when sand is absent), which complicated its extraction. The fossil was preserved in a reddish micaceous fine sandstone, about 4 meters above the second (and middle) transgressive layer, and below the upper transgressive layer. The age of ML 2554 is thus estimated to be about 149 Ma.

Materials & Methods:

The preparation of the specimen was done mechanically, using PaleoTools Micro Jacks and a variety of manual tools, all performed at the Museu da Lourinhã in Lourinhã, Portugal. All bone surfaces were consolidated with 5% Paraloid B-72 diluted in acetone, and any breaks or deep fissures were glued and reinforced with 20% or 50% Paraloid B-72, as required.

CT scans were performed in Leiria, Portugal, with a microfocus CT system GE VtomeX M 240. Scan images were segmented and assembled utilizing Avizo and Meshlab softwares. (GE Sensing & Inspection Technologies GmbH., Wunstorf, Germany). This resulted in four stacks of

DICOM (.dicom) images (detailed information available at Morphobank Project 3968). The segmentation of the complete specimen was done using manual selection slice-by-slice in the software Avizo v9.1 (Thermo Fisher, Waltham, MA, USA). All meshes were exported as Wavefrontfiles (.obj) and treated in the open-source software Blender v3.4. All meshes were smoothed for rendering using the Smooth Laplacian modifier (Lambda factor = 1 and 10 repeats). Measurements were taken both directly from the physical specimen and digitally in Blender.

Phylogenetic analysis was conducted using TNT version 1.5 (Goloboff et al. 2008; Goloboff & Catalano, 2016) using the matrix by Andres et al., 2021, augmented by the additional Portuguese specimen described here (represented in the analysis by “Lusognathus_almadrava”). A basic traditional tree-search analysis was conducted with 1,000 random addition sequence replicates.

All specimen CT data, digital material, and phylogenetic matrix files are available at Morphobank Project 3968.

Nomenclatural Acts:

The electronic edition of this article conforms to the requirements of the amended International Code of Zoological Nomenclature, and hence the new names contained herein are available under that Code from the electronic edition of this article. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix “http://zoobank.org/”. The LSID for this publication is: urn:lsid:zoobank.org:pub:54B6D020-612A-4E23-ADEC-2CDE383B5C3B. The electronic edition of this work was published in a journal with an ISSN, and has been archived and is available from the following digital repositories: PubMed Central and LOCKSS.

RESULTS

Systematic Paleontology:

PTEROSAURIA Owen, 1842 [Andres & Padian, 2020a]

PTERODACTYLOIDEA Plieninger, 1901 [Andres & Padian, 2020b]

ARCHAEOPTERODACTYLOIDEA Kellner, 1996, 2001

CTENOCHASMATIDAE Nopsca, 1928 sensu Unwin, 2003

GNATHOSAURINAE Nopsca, 1928 sensu Unwin, 2002

Lusognathus, *gen. nov.*

L. almadrava, *sp. nov.*

Etymology: *Lusognathus* is derived from the Latin “Luso”, after Lusitania (the former name for the area of Portugal in Roman times) and “gnathus” meaning “jaw”; an “almadrava” is the name of a traditional Portuguese fishing trap for catching seafood.

Holotype: ML 2554, comprised of a fragment of the anterior part of a premaxillary rostrum, a fragment of a maxillary tooththrow, two isolated fragmentary teeth, and three (or four) partial fragments of cervical vertebrae.

Locality and Horizon: Praia de Caniçal, county of Lourinhã, Portugal. Lourinhã Formation, late Kimmeridgian-early Tithonian, about 149.2 Ma (Schneider et al., 2009).

Diagnosis: A gnathosaurine pterosaur with the following combination of characters: a rounded-triangular anterior expansion of the premaxilla, constriction of the maxilla directly posterior to the spatulate anterior expansion, robust laterally-projected teeth (spaced at 1.3 teeth per cm) with a subcircular to oval cross-section, and posterior teeth projected anterolaterally.

Specimen Description:

Specimen ML 2554, here named *Lusognathus almadrava*, is composed of a well-preserved three-dimensional maxillopremaxillary rostrum fragment with three(?) associated fragmentary mid-cervical vertebrae (two of which, although preserved in articulation, are too fragmentary to provide detailed anatomical information). The rostral fragments are divided between 3 blocks with only one side exposed. Two of the blocks connect and comprise the anteriormost premaxillary portion of the rostrum (Fig. 1), visible in ventral view, and the third block houses a more posterior maxillary section (of unknown exact position along the rostrum). No sutures are visible anywhere along the rostrum, although two longitudinal grooves are visible, running parallel to the tooththrow (difficult to distinguish due to slight dorsoventral taphonomic compression, likely from lithostatic pressure). For precise measurements of each individual element, see supplementary materials.

The anterior-most portion of the premaxilla is somewhat eroded. The increasingly anteromedial direction of the preserved roots of the anterior-most teeth indicate that the termination of the rostrum was not far beyond what is actually preserved (Fig. 1 & 2). In dorsoventral view, the rostrum begins to expand anterolaterally at approximately 28.0mm from the anteriormost preserved edge, into a spatulate A-line or rounded-triangle shape, with the widest point terminating at the alveolar collar of the largest anterior tooth. Posterior to this point of expansion, and with no sharp demarcation (but still a distinct constriction) of the rostrum, a gradual posterolateral expansion also begins towards its posterior edge, but with a more gradual degree of increase than the anterior spatulated expansion.

The third rostrum fragment, which contains the posterior-most portion of the preserved maxilla, still preserves tooth alveoli, indicating that it was likely positioned anterior to the nasoantorbital fenestra, however it should be noted that the actual percentage of the overall rostrum that is represented by this specimen cannot be determined.

There are twenty-nine teeth preserved in the rostrum, although in varying conditions, and none complete. Two additional isolated fragmentary teeth were also recovered from the surrounding sediment during preparation. Under these constraints, it is not possible to ascertain an exact overall tooth count for the taxon, although its tooth density can be calculated at approximately 1.3 teeth per cm. The teeth along the premaxilla and maxilla exhibit a clear thecodont dentition, with each tooth located in a single alveolus which forms a raised collar of bone at the base of each tooth. The rostrum exhibits at least sixteen well-preserved alveoli on each side. The depressions between the alveoli gives the rostrum a crenellated appearance in dorsal and ventral views. The teeth are robust, reaching up to 5 mm in width, show a slight anterolateral tilt, and become increasingly inclined laterally and anteriorly. From the anterior to posterior end of the toothrow, the teeth also diminish in dimension and begin to slightly recurve more posteriorly. The distance between individual teeth is greater than the tooth width, with the exception of the largest preserved tooth at the anterior end. In the transverse view, the gradation of the tooth row is not perfectly horizontal, and shifts slightly dorsally at the posterior region.

The preserved teeth have straight crowns, tapering towards the apex. They are subcircular to oval in cross-section, with a slight compression on their anteroventral and posterodorsal sides, resulting in keel-like attenuations running longitudinally. The tooth enamel is completely smooth and lacks any ornamentation (Fig. 2A). Where teeth have been broken, pulp cavities are readily visible.

The best-preserved vertebral fragment consists of the incomplete anterior portion of a cervical centrum (Fig. 3), exhibiting the prezygapophyseal and cotylar region. It exhibits slight taphonomic distortion but preserves its three-dimensionality. The neural spine is low. The neural canal, flanked by two depressions on either side, is visible as a deep divot centered on the anterior end. The bifid prezygapophyses project anterolaterally, and their articular faces turn anteromedially.

There are three (possibly four) remaining vertebral fragments. One block contains one or two fragmentary vertebrae preserved in the sediment, and are potentially the posterior end of the atlas-axis complex articulated with the first cervical vertebra. Of the two other fragmentary articulated fragments, one appears to be an elongated posterior part of a centrum, and the other an anterior prezygapophyseal region of another vertebra (with a pronounced neural spine). This material, however, is too damaged to provide much information. Judging by the height of the neural spine (when compared with the most complete vertebra), this spine appears slightly taller and more pronounced.

No sign of a premaxillary crest could be distinguished along the rostrum. However, since premaxillary crest in gnathosaurines usually begin to take shape in a more posterior position (e.g., *Gnathosaurus*), and the part of the rostrum preserved in *Lusognathus almadrava* is most likely well anterior to the nasoantorbital fenestra, we cannot be certain if this taxon would in fact bear one.

Phylogenetic Analysis:

Phylogenetic analysis was conducted analyzing the evolutionary relationships of *Lusognathus almadrava* and other pterosaurs using the data matrix of Andres et al. (2021). The analysis resulted a single most parsimonious tree (Fig. 4). The Gnathosaurinae was phylogenetically defined as the least inclusive group including *Gnathosaurus subulatus* von Meyer 1834 and *Huanhepterus quingyangensis* Dong 1982 (Andres, 2021), and accordingly, *Lusognathus almadrava* falls within the Gnathosaurinae.

Discussion/Taxonomic Assignment:

Typically, the defining feature of the Ctenochasmatidae (Nopsca, 1928) has been the tip of the rostrum being dorsoventrally depressed and rounded, whereas the designation for the gnathosaurines therein is the significant lateral expansion of the anterior end of the rostra (Unwin, 2002). However, this expanded feature is not exclusive to gnathosaurines alone, as is also presents to a degree in anhanguerids (e.g., Kellner, 2003), istiodactylids (e.g., Lu et al. 2008), and ornithocheirids (e.g., Unwin, 2002, but see Rodrigues & Kellner, 2013 for the status of Ornithocheiridae). What does make this feature unique for gnathosaurs (beyond the aforementioned dorsoventral rostral composition of the larger Ctenochasmatidae) is the pronounced extent and variation of the spatula, e.g., the spoon-shaped spatulas of *Tucudactylus luciae* (Soto et al., 2021), *Gnathosaurus macrurus* (Howse & Milner, 1995), and *Gnathosaurus subulatus* (Meyer, 1833) (Fig. 5A), and the more bulbously circular end of *Plataleorhynchus streptophorodon* (Howse & Milner, 1995) (Fig. 5C). The extent of the diversity of these shapes has only been revealed as more specimens have been discovered (e.g., Soto et al., 2021). Specimen ML 2554 sustains damage at its anterior edge, and therefore is more difficult to precisely distinguish in rostrum shape, but it seems to exhibit a rounded-triangular anterior rostrum shape (Fig. 5B). ML 2554 also shares in common a constriction of the rostrum just posterior to the spatulated end with *G. macrurus* (Howse & Milner, 1995), a demarcation of the spatula which is more pronounced than in the other afore-mentioned gnathosaurines. This overall variability in rostrum shape and dental apparatus is likely linked to contrasting functional feeding movements, ecological niches, and perhaps even prey differences (Ósi et al., 2011; Kellner et al. 2019b), just as modern spoonbills today also vary in bill shape (e.g., the Yellowbilled Spoonbill *Platalea flavipes* has a longer bill and narrower spoon than the Royal Spoonbill *Platalea regia*, with both occupying the same habitat but consuming different prey [Hancock et al., 1992]).

Other defining features of the Ctenochasmatidae pertain to their dentition; namely, 25 or more teeth per side (with seven or more teeth in the premaxilla), the most rostrally-situated teeth being elongate with cylindrical crowns that project laterally from the dental border (Unwin, 2003). ML 2554 exhibits at least sixteen preserved teeth per side, with at least 8 preserved in the spatula (damage impeding exact further determinations), making for an assignation therein, and making it further comparable with the tooth distribution of *G. macrurus* and *T. luciae*. Furthermore, if calculations of maximum tooth density are made based on the preserved portion, we can infer approximately 1.3 teeth per centimeter, making ML 2554 the gnathosaurine with the least dense dentition known so far. Tooth density is also

informative in the Gnathosaurinae, and even often autapomorphic within the Ctenochasmatidae, given that it can also potentially differentiate feeding specializations.

As in ctenochasmatids as a whole, gnathosaurines have **conspicuous teeth** set in the alveoli, but gnathosaurines have a raised collar of bone around their tooth bases (Perea, et al. 2018), which is much more pronounced than in other ctenochasmatids, a feature well exhibited by ML 2554. This feature also contributes to the appearance of an inter-dental crenelation in ML 2554, what has been previously reported in *P. streptophorodon*, *G. subulatus*, and *T. luciae* (Howse & Milner, 1995; Perea, et al. 2018; Soto et al. 2021). This collar also differs in being much more subtle than the distinctive alveolar parapet typically used to describe lonchodectids, which exhibit a more raised margin of the alveoli, usually elevated above the medial portion of the occlusal surface (Unwin, 2001; Avarianov, 2020), and which also have vertically-inclined teeth.

Similar to all ctenochasmatids, the teeth of ML 2554 show a sub-circular cross-section, with smooth enamel. However, one significant difference of gnathosaurines from other ctenochasmatids is in their tooth robustness. Ctenochasmatids are **traditionally** filter-feeding, foraging in shallow waters with “needle-like” thin teeth, indicative of smaller prey items (Wang et al. 2007; Bestwick et al., 2018; Paul 2022). The marked robustness of individual gnathosaur teeth suggests different feeding habits for their teeth being substantially larger and too widely-spaced for a true “filtration” functionality (Bestwick et al. 2018; Knoll, 2000), instead likely more associated to larger prey (e.g., piscivory). **This is not dissimilar to modern gavials, whose teeth are conical and slender, with pointed apices, the shape of which is widely associated with fish-piercing, since the pointed apex would facilitate penetration of prey items, and the slenderness of the teeth would enable teeth to slide between the bones of prey, minimizing breakage (Massare, 1987). Accordingly, the finer teeth of ctenochasmatids (other than gnathosaurines) would therefore be more utile in squashing or gripping prey items rather than for directly piercing them.**

Another differentiation in the dentition of the Gnathosaurinae can be made in that their teeth taper overall, from base to tip (whereas ctenochasmatid teeth are only most apically tapered) (Knoll, 2000). Additionally, tooth projection itself can also be a relevant differentiator of feeding function, as the torsional stresses applied to teeth throughout prey capture and feeding also vary, depending on the nature and agility of that prey. Whereas procumbent dentition is found in *Rhamphorhynchus* sp. and also in most ctenochasmatids (e.g. *Gegepterus changi*, *Ctenochasma elegans*, *G. subulatus*, and *T. luciae*), lateral projection is also found in certain rhamphorhynchids (e.g. *Angustinariperus longicephalus*, *Qinglongopterus guoi*, *Sericipterus wucaiwansensis*) and in ctenochasmatids (*P. streptophorodon*, *Liaodactylus primus*, *C. elegans* and *T. luciae*). ML 2554 has laterally-projected teeth in the spatular region, and also anterolaterally projected teeth posterior to the spatula, which is a feature that is found in other taxa such as *T. luciae* and *G. macrurus*, but different from taxa with posterolateral projections as inferred in *P. streptophorodon* (Howse & Milner, 1995). **The recurvature of dentition could aid in maneuvering prey back towards the gullet, as in modern gavials consuming fish (Massare, 1987), however,** ontogeny could also potentially change tooth orientation (Bennett, 2007), potentially allowing for feeding niches to change somewhat over the course of an animal’s lifespan, and so this feature should be considered in a fluctuating capacity. Making any

deductions on teeth alone should also be made prudently, since many taxonomic attributions for members of the clade are based on isolated teeth, and so may not be a wholly reliable metric.

Regarding vertebrae, the condition exhibited in one the partial vertebra of ML 2554 appears to follow the anteroposterior elongation typically found in both ctenochasmatids and azhdarchids (e.g., Howse, 1986; Martill et al., 1998; Unwin, 2003; Andres and Ji, 2008), although its fragmentary state of preservation does not allow for any kind of insightful overall measurement. However, it is still informative in that there are no obvious sutures between the vertebral centrum and the neural arch, suggesting that it belonged to an osteologically-mature individual (e.g., Bennett, 1993).

Portrayals of pterosaurs throughout the Triassic and Jurassic have traditionally been as relatively small animals, with wingspans constrained to around 1.6–1.8 m or less, whereas Cretaceous pterosaurs reached well into and above 3 m (O’Sullivan et al, 2013; Jagielska et al. 2022). Of late, it has been increasingly postulated that Jurassic pterosaurs have been previously underestimated for their size range (and particularly when considering the larger end of the spectrum). Some previously-known larger-sized examples include the 1.73 minimum wingspan of *Sericipterus* (Andres et al., 2010), the 1.8 m wingspans of *Camplognathoides* (Padian, 2008) and *Rhamphorhynchus* (Wellnhofer, 1975), the 1.9 m wingspan of the rhamphorhynchine pterosaur from the Whitby Mudstone Formation (O’Sullivan et al, 2013), and the 2.5 m wingspan of the Middle Jurassic sub-adult *Dearc sgiathanach* Jagielska et al. 2022. There are also the proposed (but not wholly reliable) over 2.5 m wingspans of *Harpactognathus gentryi* Carpenter et al., 2003 (the fossil has been removed from museum collections since its publication) and the gnathosaurine *Huanhepterus quingyangensis* Dong, 1982 (of unreliable age), the 3.3 m wingspan of a very fragmentary potential *Rhamphorhynchus* (Spindler & Irfam, 2021), and the 3.5 m–5 m wingspan of a specimen described by Meyer and Hunt, (1999), but which may be scaled using an uncertain attribution, according to O’Sullivan et al. (2013). Pterosaurs over the 2.5 m wingspan have therefore not typically been found to be older than the Early Cretaceous (Meyer & Hunt 1999).

Notwithstanding these larger size estimates, the pterosaurs of the Jurassic of Portugal are especially remarkable for the time period (i.e., the 4 m estimated wingspan of the fossil femur described by Bertozzo et al., 2021), with ML 2554 being no exception. The minimum overall length of the preserved rostrum of *Lusognathus almadrava* begins at 20.2 cm (when all three rostral blocks are lined up) but could potentially be even larger. This means that if we scale the specimen with the overall skull dimensions of *Gnathosaurus subulatus* (following the skull/skeleton scaling by Carpenter et al. 2003), where the calculated total minimum skull length of ML 2554 would be 60.8 cm, then *Lusognathus almadrava* would have achieved a minimum wingspan starting at 3.6 m. Additionally, the coeval pterosaur track site of Peralta, less than 10 km to the south of Praia do Caniçal, records about 400 *Pteraichnus*-like tracks of different sizes (half manus/half pes). The pes track length varies from 5.5 to 15 cm, which indicates the occurrence of very large pterosaurs with wingspans of four meters or more, preliminarily making *Lusognathus almadrava* a likely candidate for the trackmaker. Based on ML 2554, *Lusognathus almadrava* could potentially reach these larger sizes, making it one of

the largest Jurassic pterosaurs known, and the largest known Jurassic gnathosaurine, having a size matched only by some Cretaceous records of the group.

Conclusion:

A new pterosaur taxon has been found from the Late Jurassic of Portugal, introduced here as *Lusognathus almadrava* gen et sp. nov. Several characters ascribe to this specimen to the Gnathosaurinae, namely its spatulate rostrum, robust comb-like dentition, and the pronounced rims of its tooth alveoli. The presence of this taxon in the fluvio-deltaic lagoonal environment that has been suggested to be representative of Lourinhã Formation in the Late Jurassic (Fig. 6) is in keeping with the modern analog of modern-day spoonbills, whose habitats generally include more shallow-water estuarine, tidal flat, coastal marshland, or even inland lake areas (areas where there are mud, clay, or fine-sand bottoms) with inflows of fresh, brackish, or salt water (Hancock et al., 1992).

The paleobiological implications of such an exceptionally large-sized pterosaur in the Jurassic of Portugal denote and reinforce a thriving ecosystem, abundant with prey, in this case perhaps fish (as indicated by the robust teeth of ML 2554). Although it has been generally agreed upon that pterosaur body size steadily increased throughout and up to the end of the Cretaceous, ML 2554 adds more evidence for body sizes already having increased substantially by the end of the Late Jurassic (when compared with earlier forms); this growth having been a response to filling a different ecological niche than their competitors, the birds (Benson et al. 2014; Tennant et al., 2017). Although transitioning to this larger size may have inadvertently contributed to their downfall at the end of the Cretaceous (since fewer niches are available for larger animals, making big species more likely predisposed to decline [Cardillo et al., 2005]), and since larger terrestrial animals were disproportionately affected by the Cretaceous–Paleogene extinction (Benson et al. 2014), at least in the Late Jurassic paleoenvironments this extraordinary growth seemed advantageous for their flourishing success.

Acknowledgements: Immense gratitude is extended to Filipe Viera for his invaluable scientific discovery and donation of the fossil to the Museu da Lourinhã. Thank you to the Lourinhã Paleoteam for their good-humored willingness to excavate a partially-underwater specimen (in winter, and at a moment's notice), with a special thank you to Alexandre Audigane and Miguel Moreno-Azanza for assisting in mobilizing the excavation permits (also in winter, and at a moment's notice). Heartfelt thanks are also extended to Carla Alexandra Tomás, Micael Martinho, Laura de Jorge, and Carla Hernandez for their impeccable preparation of the specimen, and to the rest of the staff of the Museu da Lourinhã. Thank you to the DinoParque da Lourinhã and the Museu da Lourinhã, for their generosity in providing the Horacio Mateus (PIIHM) Grant, and to Bruno Pereira and Maria Rios for being coordinators of the Departamento de Investigação do GEAL. Many thanks to Bruno Camillo Silva of the Sociedade de História Natural de Torres Vedras, Pedro Aquino, Flávio Domingues, and Tiago Ferreira at

Micronsense Metrologia Industrial, Lda. for their invaluable CT scan implementation. Oliver Rauhut is thanked for constructive suggestions. AWAK acknowledges funding from the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Rio de Janeiro (FAPERJ #E-26/201.095/2022) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq #313461/2018-0, #406779/2021-0, #406902/2022-4). OM was funded by the Fundação para Ciência e a Tecnologia, I. P. with National Funds from the Ministério da Ciência, Tecnologia e Ensino Superior. (Grant numbers: UIDB/04035/2020, GeoBioTec).

References Cited

- Andres B. 2021. Phylogenetic systematics of *Quetzalcoatlus* Lawson 1975 (Pterodactyloidea: Azhdarchoidea). *Journal of Vertebrate Paleontology* 41(1):203-217.
- Andres B, Ji Q. 2008. A new pterosaur from the Liaoning Province of China, the phylogeny of the pterodactyloidea, and convergence in their cervical vertebrae. *Palaeontology* 51(2):453-469.
- Andres B, Clark JM, Xing X. 2010. A new rhamphorhynchid pterosaur from the Upper Jurassic of Xinjiang, China, and the phylogenetic relationships of basal pterosaurs. *Journal of Vertebrate Paleontology* 30(1):163–187.
- Andres B, Clark J, Xu X. 2014. The earliest pterodactyloid and the origin of the group. *Current Biology* 24(9):1011-1016.
- Andres B, Padian K. 2020a. Pterosauria R. Owen 1842. In: de Queiroz K, Cantino PD, Gauthier JA, eds. *Phylonyms*. Boca Raton: CRC Press, 1201-1204.
- Andres B, Padian K. 2020b. Pterodactyloidea F. Plieninger 1901. In: de Queiroz K, Cantino PD, Gauthier JA, eds. *Phylonyms*. Boca Raton: CRC Press, 1205-1207.
- Antunes MT, Pais J. 1978. Notas sobre depósitos de Taveiro: estratigrafia, paleontologia, idade, paleoecologia [Notes on the Taveiro deposits: stratigraphy, paleontology, age, paleoecology]. *Ciências da Terra* 4:108-128.
- Averianov AO. 2020. Taxonomy of the Lonchodectidae (Pterosauria, Pterodactyloidea). *Proceedings of the Zoological Institute RAS*, 324(1):41–55.
- Barrett PM, Butler RJ, Edwards NP, Milner AR. 2008. Pterosaur distribution in time and space: an atlas. *Zitteliana* B(28):61-107.
- Bell CM, Padian K. 1995. Pterosaur fossils from the Cretaceous of Chile: evidence for a pterosaur colony on an inland desert plain. *Geological Magazine* 132:31-38.

- Bertozzo F, Camilo da Silva B, Martill D, Vorderwuelbecke EM, Aureliano T, Schouten R, Aquino P. 2021. A large pterosaur femur from the Kimmeridgian, Upper Jurassic of Lusitanian Basin, Portugal. *Acta Palaeontologica Polonica* 66(4):815–825.
- Bennett SC. 2007. A review of the pterosaur *Ctenochasma*: taxonomy and ontogeny. *N. Jb. Geol. Paelont. Abh.* 245(1):23–31.
- Benson R, Frigot R, Goswami A, Andres B, Butler RJ. 2014. Competition and constraint drove Cope's rule in the evolution of giant flying reptiles. *Nature Communications* 5(1):3567.
- Bestwick J, Unwin DM, Butler RJ, Henderson DM, Purnell MA. 2018. Pterosaur dietary hypotheses: a review of ideas and approaches. *Biol. Rev.* 93:2021–2048.
- Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds OR, Sechrest W, Orme CD, Purvis A. 2005. Multiple causes of high extinction risk in large mammal species. *Science* 309(5738):1239–1241.
- Carpenter K, Unwin D, Cloward K, Miles C, Miles C. 2003. A new scaphognathine pterosaur from the Upper Jurassic Morrison Formation of Wyoming, USA. In: Buffetaut E, Mazin J-M, eds. *Evolution and Palaeobiology of Pterosaurs*. London: Geological Society, Special Publications 217(1):45–54.
- Dong Z. 1982. A new pterosaur (*Huanhepterus quingyangensis* gen. et sp. nov.) from Ordos, China. *Vertebrata Palasiatica* 20(2):115–121.
- Figueiredo SD, Rosina P, Strantzali IB, Antunes V, Figueiredo S. 2020. Paleoenvironmental approach on the Lower Cretaceous sequences of Areia do Mastro (Cabo Espichel, Southern Portugal). *Journal of Environmental Science and Engineering A*(9):66–71.
- Figueiredo SD, Cunha PP, Pereda-Suberbiola X, de Carvalho CN, de Souza Carvalho I, Buffetaut E, Tong H, Sousa MF, Antunes V, Anastácio R. 2022. The dinosaur tracksite from the lower Barremian of Areia do Mastro Formation (Cabo Espichel, Portugal): Implications for dinosaur behavior. *Cretaceous Research* 137:105219.
- Galton, PM. 1994. Notes on Dinosauria and pterodactylia from the Cretaceous of Portugal. *Neues Jahrbuch für Geologie und Paläontologie*. December 194(2/3):253–267.
- Goloboff P, Catalano S. T.N.T., version 1.5, with a full implementation of phylogenetic morphometrics [computer software]. Cladistics; 2016. doi:10.1111/cla.12160.
- Goloboff PA, Farris JS, Nixon KC. 2008. T.N.T., a free program for phylogenetic analysis. *Cladistics* 24:774–786.

- Guillaume ARD, Moreno-Azanza M, Puértolas-Pascual E, Mateus O. 2020. Palaeobiodiversity of crocodylomorphs from the Lourinhã Formation based on the tooth record: insights into the palaeoecology of the Late Jurassic of Portugal. *Zoological Journal of the Linnean Society* 189(2):549–583.
- Hancock, James, et al. *Storks, Ibises and Spoonbills of the World*, Bloomsbury Publishing Plc, 1992.
- Hill G. 1989. *The sedimentology and lithostratigraphy of the Upper Jurassic Lourinhã Formation, Lusitanian Basin, Portugal*. [Doctoral dissertation, The Open University]. United Kingdom. <http://oro.open.ac.uk/54566/1/328171.pdf>
- Howse SCB. 1986. On the cervical vertebrae of the Pterodactyloidea (Reptilia: Archosauria). *Zoological Journal of the Linnean Society* 88:307-328.
- Howse SCB, Milner AR. 1995. The pterodactyloids from the Purbeck Limestone Formation of Dorset. *Bulletin of the Natural History Museum of London, Geology Series* 51(1):73-88.
- Jagielska N, O’Sullivan M, Funston GF, Butler IB, Challands TJ, Clark NDL, Fraser NC, Penny A, Ross DA, Wilkinson M, Brusatte SL. 2022. A skeleton from the Middle Jurassic of Scotland illuminates an earlier origin of large pterosaurs. *Current Biology* 32:1446–1453.
- Kellner AWA. 1996. Description of new material of Tapejaridae and Anhangueridae (Pterosauria, Pterodactyloidea) and discussion of pterosaur phylogeny. Ph.D. diss., Columbia University (available through UMI - University Microfilms International; registered 1997 - 091855442). 347p.
- Kellner AWA. 2001. New hypothesis of Pterosaur Phylogeny. In: Barros, L. M., Nuvens, P. C. & Filgueira, J. B. M. (Eds.) I e II Simpósios sobre a Bacia do Araripe e bacias interiores do Nordeste, Comunicações. Crato: 249-258.
- Kellner AWA. 2003. Pterosaur phylogeny and comments on the evolutionary history of the group. In: Buffetaut E, Mazin J-M, eds. *Evolution and Palaeobiology of Pterosaurs*. London: Geological Society, Special Publications, 217(1): 105-137.
- Kellner AWA, Rodrigues T, Costa FR, Weinschütz LC, Flgueiredo RG, Souza GAD, Brum AS, Eleutério LHS, Mueller C, Sayão JM. 2019. Pterodactyloid pterosaur bones from Cretaceous deposits of the Antarctic Peninsula. *Anais da Academia Brasileira de Ciências* 91(Suppl. 2) e20191300.
- Kellner AWA, Caldwell MW, Holgado B, Dalla Vecchia FM, Nohra R, Sayão JM, Currie PJ. 2019. First complete pterosaur from the Afro-Arabian continent: insight into pterodactyloid diversity. *Scientific Reports* 9:17875.

- Kühne WG. 1968. History of discovery, report on the work performed, procedure, technique and generalities. *Contribuição para a fauna do Kimeridgiano da Mina de lignito Guimarães (Leiria, Portugal), I parte*. Lisboa: Mem Serv. Geol. Portugal, (N.S.) 14:7-20.
- Knoll F. 2000. Pterosaurs from the Lower Cretaceous (?Berriasian) of Anoual, Morocco. *Annales de Paléontologie* 86:157–164.
- Longrich NR, Martill DM, Andres B. 2018. Late Maastrichtian pterosaurs from north Africa and mass extinction of Pterosauria at the Cretaceous-Paleogene boundary. *PLoS Biology* 16(3):1-38
- Lü J, Xu L, Ji Q. 2008. Restudy of *Liaoxipterus* (Istiodactylidae: Pterosauria), with comments on the Chinese istiodactylid pterosaurs. *Zitteliana* B28:229-241.
- Malafaia E, Ortega F, Escaso F, Dantas P, Pimentel N, Gasulla JM, Ribeiro B, Barriga F, Sanz JL. 2010. Vertebrate fauna at the *Allosaurus* fossil-site of Andrés (Upper Jurassic), Pombal, Portugal. *Journal of Iberian Geology* 36(2):193-204.
- Manuppella G, Antunes M, Pais J, Ramalho M, Rey J. 1999. Notícia explicativa da folha 30-A (Lourinhã) da Carta Geológica de Portugal. Esc. 1/50 000. Instituto Geológico e Mineiro.
- Martill DM, Sadaqah RM, Khoury HN. 1998. Discovery of the holotype of the giant pterosaur *Titanopteryx philadelphiae* Arambourg 1959, and the status of Arambourgiania and Quetzalcoatlus. *Neues Jahrbuch für Geologie und Paläontologie* 207(1):57-76.
- Martill DM, Frey E, Bell CM, Diaz GC. 2006. Ctenochasmatid pterosaurs from early Cretaceous deposits in Chile. *Cretaceous Research* 27(5):603-610.
- Martinius AW, Gowland S. 2011. Tide-influenced fluvial bedforms and tidal bore deposits (late Jurassic Lourinhã Formation, Lusitanian Basin, Western Portugal). *Sedimentology* 58(1):285-324.
- Massare JA. 1987. Tooth Morphology and Prey Preference of Mesozoic Marine Reptiles. *Journal of Vertebrate Paleontology* 7(2):121-137.
- Mateus O, Milàn J. 2010. First records of crocodile and pterosaur tracks in the Upper Jurassic of Portugal. *New Mexico Museum of Natural History and Science Bulletin* 51:83–87.
- Mateus O, Milàn J, Romano M, Whyte MA. 2011. New finds of stegosaur tracks from the Upper Jurassic Lourinhã Formation, Portugal. *Acta Palaeontologica Polonica* 56(3):651-658.

- Mateus O, Dinis J, Cunha PP. 2017. The Lourinhã Formation: the Upper Jurassic to lower most Cretaceous of the Lusitanian Basin, Portugal—landscapes where dinosaurs walked. *Ciências da Terra/Earth Sciences Journal* 19(1):75-97.
- Meyer Hv. 1834. *Gnathosaurus subulatus*, ein Saurus aus den lithographischen Schiefer von Solnhofen. *Museum Senckenbergianum* 1:1-26.
- Meyer CA, Hunt AP. 1999. The first pterosaur from the Late Jurassic of Switzerland: Evidence for the largest Jurassic flying animal. *ORYCTOS* 2:111-116.
- Mocho P, Royo-Torres R, Malafaia E, Escaso F, Ortega F. 2016. Systematic review of late Jurassic sauropods from the Museu Geológico Collections (Lisboa, Portugal). *Journal of Iberian Geology* 42(2):227-250.
- Myers TS, Tabor NJ, Jacobs LL, Mateus O. 2012. Estimating soil p CO2 using paleosol carbonates: implications for the relationship between primary productivity and faunal richness in ancient terrestrial ecosystems. *Paleobiology* 38(4):585-604.
- Nopsca FJ. 1928. The genera of reptiles. *Palaeobiologica* 1:163-188.
- O’Leary MA, Kaufman SG. 2012. MorphoBank 3.0: Web Application for Morphological Phylogenetics and Taxonomy [computer software]. Available online: <http://morphobank.org/permalink/?3968>
- O’Sullivan M, Martill DM, Grocock D. 2013. A pterosaur humerus and scapulocoracoid from the Jurassic Whitby Mudstone Formation, and the evolution of large body size in early pterosaurs. *Proc. Geol. Assoc.* 124(6):973-981.
- Ósi, A. 2011: Feeding-related characters in basal pterosaurs: implications for jaw mechanism, dental function and diet. *Lethaia*, Vol. 44, pp. 136–152.
- Owen R. Report on British Fossil Reptiles, Part II. In: Proceedings of the 11th Meeting of the British Association for the Advancement of Science, Plymouth, UK, 24 July 1842; pp. 60–204.
- Perea D, Soto M, Toriño P, Mesa V, Maisey J. G. 2018. A Late Jurassic–earliest Cretaceous ctenochasmatid (Pterosauria, Pterodactyloidea): The first report of pterosaurs from Uruguay. *Journal of South American Earth Sciences* 85:298-306.
- Pérez-García A, Ortega F. 2011. *Selenemys lusitanica*, gen. et sp. nov., a new pleurosternid turtle (Testudines: Paracryptodira) from the upper Jurassic of Portugal. *Journal of Vertebrate Paleontology* 31(1):60-69.
- Puértolas-Pascual E, Marx M, Mateus O, Saleiro A, Fernandes AE, Marinheiro J,

- Tomás C, Mateus S. 2021. A new plesiosaur from the lower Jurassic of Portugal and the early radiation of Plesiosauroidea. *Acta Palaeontologica Polonica* 66(2):369–388.
- Rauhut OWM. 2001. Herbivorous dinosaurs from the Late Jurassic (Kimmeridgian) of Guimarota, Portugal. *Proceedings of the Geologists' Association* 112:275-283.
- Ribeiro V, Mateus O, Holwerda F, Araújo R, Castaninha R. 2014. Two new theropod egg sites from the Late Jurassic Lourinhã Formation, Portugal. *Historical Biology* 26(2):206-217.
- Rodrigues T, Kellner AWA. 2013. Taxonomic review of the Ornithocheirus complex (Pterosauria) from the Cretaceous of England. *ZooKeys (Monographs)* 308: 1-112 (doi: 10.3897/zookeys.308.5559 (ISSN 1313-2970 / 1313-2989).
- Rotatori FM, Moreno-Azanza M, Mateus O. 2020. New information on ornithopod dinosaurs from the Late Jurassic of Portugal. *Acta Palaeontologica Polonica* 65(1):35–57.
- Schneider S, Fürsich FT, Werner W. 2009. Sr-isotope stratigraphy of the Upper Jurassic of central Portugal (Lusitanian Basin) based on oyster shells. *International Journal of Earth Sciences* 98:1949-1970.
- Soto M, Montenegro F, Toriño P, Mesa V, Perea D. 2021. A new ctenochasmatid (Pterosauria, Pterodactyloidea) from the late Jurassic of Uruguay. *Journal of South American Earth Sciences* 111:1-11.
- Spindler F, Iffrim C. 2021. Die Spur einer Spur – ein möglicher erster Flugsaurier aus Ettling. *Archaeopteryx* 37:75–83.
- Sweetman SC, Martill DM. 2010. Pterosaurs of the Wessex Formation (Early Cretaceous, Barremian) of the Isle of Wight, southern England: a review with new data. *Journal of Iberian Geology* 36(2):225-242.
- Taylor AM, Gowland S, Leary S, Keogh KJ, Martinius AW. 2014. Stratigraphical correlation of the Late Jurassic Lourinhã Formation in the Consolação Sub-basin (Lusitanian Basin), Portugal. *Geological Journal* 49(2):143-162.
- Tennant JP, Mannion PD, Upchurch P, Sutton MD, Price GD. 2017. Biotic and environmental dynamics through the Late Jurassic–Early Cretaceous transition: evidence for protracted faunal and ecological turnover. *Biol Rev* 92:776-814.
- Thulborn, RA. 1973. Teeth of ornithischian dinosaurs from the Upper Jurassic of Portugal with description of a Hypsilophodontid (*Phyllodon henkeli* gen. et sp. nov.) from the Ouimarota lignite. *Contribuição para o conhecimento da fauna do Kimeridgiano da mina de lignito Ouimarota (Leiria, Portugal), III parte*. Lisboa: Memórias dos Serviços Geológicos de Portugal (N.S.) 22:89–134.

- Unwin DM. 2001. An overview of the pterosaur assemblage from the Cambridge Greensand (Cretaceous) of Eastern England. *Fossil Record* 4:189–221.
- Unwin DM. 2002. On the systematic relationships of *Cearadactylus atrox*, an enigmatic Early Cretaceous pterosaur from the Santana Formation of Brazil. *Fossil Record* 5: 239-263.
- Unwin DM. 2003. On the phylogeny and evolutionary history of pterosaurs. In: Buffetaut E, Mazin J-M, eds. *Evolution and Palaeobiology of Pterosaurs*. London: Geological Society, Special Publications 217(1):139-190.
- Unwin DM, Lü J, Bakhurina NN. 2000. On the systematic and stratigraphic significance of pterosaurs from the Lower Cretaceous Yixian Formation (Jehol Group) of Liaoning, China. *Mitteilungen aus dem Museum für Naturkunde in Berlin, Geowissenschaftlichen Reihe* 3:181–206.
- van Hinsbergen DJJ, de Groot LV, van Schaik SJ, Spakman W, Bijl PK, Sluijs A, Langereis CG, Brinkhuis H. 2015. A Paleolatitude Calculator for Paleoclimate Studies. *PLOS ONE* 10(6): e0126946.
- Vidovic SU, Martill DM. 2017. The taxonomy and phylogeny of *Diopecephalus kochi* (Wagner, 1837) and *Germanodactylus rhamphastinus* (Wagner, 1851). *Geological Society, London, Special Publications* 455:125-147.
- Wang X, Kellner AWA, Zhou Z, Campos DA. 2007. A new pterosaur (Ctenochasmatidae, Archaeopterygidae) from the Lower Cretaceous Yixian Formation of China. *Cretaceous Research*, 28: 245-260.
- Wellnhofer P. 1975. Die Rhamphorhynchoidea (Pterosauria) der Oberjura-Plattenkalke Süddeutschlands. Teil II Systematische Beschreibung.– *Palaeontographica A* 148(4–6):132–186.
- Wiechmann MF, Gloy U. 2000. Pterosaurs and Urvogels from the Guimarota mine. In: Verlag,FP, ed. *Guimarota: A Jurassic Ecosystem*. München: Verlag Dr. Friedrich Pfeil, 83-86.
- Zhou CF, Gao KQ, Yi, H, Xue, J, Li Q, Fox RC. 2016. Earliest filter-feeding pterosaur from the Jurassic of China and ecological evolution of Pterodactyloidea. *Royal Society Open Science* 4(2):160672.

Supplementary Material:

1. Measurements Table
2. Phylogenetic Data Matrix

770
771
772
773
774
775

Figure 1

Figure 1. The upper jaw of the holotype (ML 2554) of *Lusognathus almadrava* gen. et sp. nov. (ML 2554).

A, photography of both jaw fragments; B, reconstruction of the jaw in dorsal view, showing minimum possible preserved length, in dorsal view; C, ventral view; D, left lateral view; E, right lateral view; F, anterior view.



Figure 2

Figure 2. Tooth morphology of *Lusognathus almadrava* gen. et sp. nov. holotype (ML 2554).

A, photography of the best-preserved tooth in labial view; B, transparent 3D representation of the upper jaw in ventral view, showing the insertion of teeth, with cross-section cuts (C1 and C2) of the best- preserved tooth.

A



B



C1

Labial

C2



Lingual



10 mm

Figure 3

Figure 3. Photographs and virtual three-dimensional renderings of the mid-cervical vertebra of *Lusognathus almadrava* gen. et sp. nov. holotype (ML 2554).

A-B, photograph and 3D rendering in dorsal view; C and F, photograph and 3D rendering in anterior view; D and G, photograph and 3D rendering in ventral view; E and H, photograph and 3D rendering in left lateral view. Abbreviations: cot, cotyle; nc, neural canal; ns, neural spine; pf, pneumatic foramen; prz, prezygapophysis.




Figure 4

Figure 4. The relationship of *Lusognathus almadrava* gen. et sp. nov. holotype, and the Pterodactyloidea, after the data matrix by Andres et al., 2021.



Figure 5

Figure 5. Comparison of the ventral rostra of:

A, *Gnathosaurus subulatus* Meyer, 1833 (JME-SOS 4580); B, *Lusognathus almadrava* n. gen. et sp. (ML 2554), and C, *Plataleorhynchus streptophorodon* Howse & Milner, 1995 (BMNH R 11957).

A



B



C



20mm

Figure 6

Figure 6. Reconstruction of *Lusognathus almadrava* and its paleoenvironment by © Jason Brougham

