

# Distribution of multidrug-resistant bacterial infections in diabetic foot ulcers and risk factors for drug resistance: a retrospective analysis

Huihui Guo<sup>Equal first author, 1</sup>, Qiwei Song<sup>Equal first author, 1</sup>, Siwei Mei<sup>1</sup>, Zhengqiang Xue<sup>1</sup>, Junjie Li<sup>Corresp., 1</sup>, Tao Ning<sup>Corresp. 1</sup>

<sup>1</sup> Department of Orthopaedic Surgery, Fuyang People's Hospital, Fuyang, China

Corresponding Authors: Junjie Li, Tao Ning

Email address: ljj3610@126.com, doctorning87@163.com

## Abstract

**Objective** To investigate the distribution, drug resistance and risk factors of multi-drug resistant bacterias MDROs in patients with Type 2 diabetic foot ulcers DFU.

**Method** The clinical data, foot secretions, pathogenic microorganisms and drug sensitivity tests of 147 patients with type 2 diabetes admitted to our department from January 2018 to December 2021 were analyzed. Patients were divided into two groups according to whether they had been infected with MDROs or not. 71 cases were infected with MDROs as the case group, and the remaining 76 cases were the control group. Chi-square test and t-test were used to analyze the results of MDROs infection and DFU, and logistic multivariate regression was used to evaluate the risk factors of MDROs infection.

**Results** A total of 71 strains were isolated from the MDROs-positive group, with the top three being Staphylococcus aureus (46.48%), Escherichia coli (22.53%), and Pseudomonas aeruginosa (18.31%), respectively. Logistic multifactorial regression analysis showed that history of previous antimicrobial exposure, neuroischemic wound, Wagner grade 3-5, and combined osteomyelitis were associated with Type 2 diabetic foot infection MDROs ( $P < 0.05$ ).

**Conclusion** Previous history of antimicrobial exposure, neuroischemic wounds, Wagner grade 3-5, and combined osteomyelitis are independent risk factors for MDROs, which can identify the risk factors for MDROs at an early stage and help to identify people at high risk of MDROs infection and take relevant comprehensive treatment in time to slow down the development of the disease.

# Abstract

**Objective:** To investigate the distribution, drug resistance and risk factors of multi-drug resistant bacterias (MDROs) in patients with Type 2 diabetic foot ulcers (DFU) .

**Method:** The clinical data, foot secretions, pathogenic microorganisms and drug sensitivity tests of 147 patients with type 2 diabetes admitted to our department from January 2018 to December 2021 were analyzed. Patients were divided into two groups according to whether they had been infected with MDROs or not. 71 cases were infected with MDROs as the case group, and the remaining 76 cases were the control group. Chi-square test and t-test were used to analyze the results of MDROs infection and DFU, and logistic multivariate regression was used to evaluate the risk factors of MDROs infection.

**Results:** A total of 71 strains were isolated from the MDROs-positive group, with the top three being *Staphylococcus aureus* (46.48%), *Escherichia coli* (22.53%), and *Pseudomonas aeruginosa* (18.31%), respectively. Logistic multifactorial regression analysis showed that history of previous antimicrobial exposure, neuroischemic wound, Wagner grade 3-5 , and combined osteomyelitis were associated with Type 2 diabetic foot infection MDROs ( $P < 0.05$ ).

**Conclusion:** Previous history of antimicrobial exposure, neuroischemic wounds, Wagner grade 3-5, and combined osteomyelitis are independent risk factors for MDROs, which can identify the risk factors for MDROs at an early stage and help to identify people at high risk of MDROs infection and take relevant comprehensive treatment in time to slow down the development of the disease.

**Keywords:**Diabetic foot,multidrug-resistant bacteria,infection, risk-factors

27

## 28 **Background**

29 Diabetic foot ulcer (DFU) is a common complication of Type 2 diabetes, and approximately  
 30 10%-25% of people with diabetes will develop DFU over the course of their lifetime (Zubair &  
 31 Ahmad, 2019). DFU can cause serious health problems, diminish patients' quality of life, and  
 32 increase the incidence of infections among diabetics, imposing a significant social, psychological,  
 33 and economic burden on patients and the health care system (Raspovic & Wukich, 2014; Armstrong  
 34 et al., 2017). The greatest barrier to DFU is increased vulnerability to several possible  
 35 pathogens, which can result in significant effects such as infection, gangrene, osteomyelitis,  
 36 amputation, and even death. (Hitam et al., 2019).

37 DFU infections are primarily microbial infections, and many of the complications of DFU  
 38 are caused by bacterial infections (Hawkins et al., 2022); thus, DFU can be reduced through early  
 39 and appropriate intervention, as well as glycemic control. Early detection of specific pathogens  
 40 and their bacterial susceptibility drug sensitivity results allow for early antibiotic treatment.  
 41 Furthermore, because the susceptibility of different bacteria to the same drug varies and bacterial  
 42 resistance shows different trends depending on the time of drug administration and geographic  
 43 area, the clinic must still adjust the drug regimen and prevent antibiotic misuse through drug  
 44 sensitivity identification.

45 Multidrug-resistant organisms (MDROs) are common pathogens in patients with DFU  
 46 infection (Kandemir et al., 2007; Adeyemo et al., 2021). The situation of MDROs has deteriorated  
 47 in recent years as a result of excessive use of clinical antibiotics, and the problem of multidrug-  
 48 resistant bacteria in DFU patients infected with pathogens is particularly serious. As a result, the  
 49 correct and reasonable application of antibiotics is the key to improve the cure rate of diabetic  
 50 foot infection, reduce the incidence of adverse reactions and reduce the occurrence of bacterial  
 51 resistance. Due to the rising occurrence of MDROs and the importance of early and effective  
 52 antimicrobial therapy, current national guidelines recommend the use of broad-spectrum  
 53 antibiotics as an empirical treatment in patients with moderate to severe DFU (Lipsky et al.,

2012). Given that the risk of MDROs infection is increasing year after year, combined with the complex and variable strain resistance, the infection is difficult to control, treatment costs are high, the amputation rate and mortality rate are increasing, reducing treatment effectiveness and quality of life in DFU patients. In order to reduce the occurrence of infections caused by drug-resistant bacteria and improve patient prognosis, numerous studies have reported risk factors for infections in MDROs in patients with DFU (Noor et al., 2017; Saltoglu et al., 2018; Datta et al., 2019; Pessoa et al., 2020; Matta-Gutierrez et al., 2021; Liu et al., 2022). In recent years, with the increase of studies related to MDROs infection in DFU patients, more and more attention has been paid to the related risk factors, but due to the interference of relevant factors such as sample size, study population, and different geographical areas, the conclusions reached are inconsistent and sometimes even contradictory, leading to a decrease in the strength and credibility of the study (Dai et al., 2020). The aim of this study was to pool and analyze the traumatic bacterial cultures of DFU patients in our hospital in recent years and the antimicrobial susceptibility drugs of these bacterial isolates in order to understand the relationship between MDROs infections and DFU patients in our region. An attempt was made to identify risk factors associated with MDROs infections in DFU patients in our region for early treatment with appropriate antibiotics.

## **Data and methods**

### **Data**

A total of 147 patients with Type 2 diabetic foot were selected from the Diabetic Foot Treatment Center of Fuyang People's Hospital from January 2018 to December 2021 with complete clinical cases and follow-up data, including 93 males and 54 females, aged  $62.62 \pm 11.71$  years. Diabetes course  $12.13 \pm 6.90$  years; All cases met the criteria of the International Diabetic Foot Task Force and the relevant diagnosis of diabetic foot of the American Society of Infectious Diseases. The diagnostic criteria for diabetic foot infections were the 2012 Special Guidelines of the International Diabetic Foot Working Group on the treatment of diabetic foot infections (Lipsky et al., 2012).

The Wagner grading scale was used to grade the diabetic foot in this study: 0: risk factors for the development of foot ulcers but no current ulcers Grade 1:superficial foot ulcers without signs of infection, penetrating the superficial or total skin layer; grade 2:ulcers deep to the ligaments, tendons, joint capsule, or deep fascia, without abscess or osteomyelitis; grade 3:ulcers deep to the ligaments, tendons, joint capsule, or deep fascia, with abscess, osteomyelitis, or sepsis; Grade 4: ischemic gangrene (toe, heel, or forefoot dorsum) usually combined with neuropathy; Grade 5: total foot gangrene (Monteiro-Soares et al., 2020).

Inclusion criteria: (1) Clinically confirmed Type 2 diabetic foot ulcer; (2) Complete clinical cases and follow-up data;(3)Patients with positive bacterial culture. Exclusion criteria: (1) patients with type 1 diabetes mellitus; (2) Foot ulcers caused by other diseases were excluded from the study, such as vasculitis, varicose ulcers, or those wounds caused by specific infections or tumors. (3) Patients taking glucocorticoids for a long time; (4) Incomplete clinical data.

### **Grouping mode**

Patients with MDROs infection were divided to the MDROs infection group based on bacterial culture and drug sensitivity results, whereas the rest of the patients were assigned to the NMDROs infection group. MDROs was defined as acquired non-susceptibility to at least one agent in three or more antimicrobial categories (Magiorakos et al., 2012). If drug-resistant bacteria coexisted with sensitive bacteria in the same patient, the patient was classified as having MDROs.

### **Index of observations**

Clinical data of the patient were collected, including gender, age, blood pressure, diabetes course, glycosylated hemoglobin at admission, cholesterol, triglycerides, inflammatory indicators, ankle-brachial index(ABI), previous antibiotic exposure history, amputation, osteomyelitis, diabetes-related complications, Wagner's grade, and hospitalization of the same ulcer surface > 2 times/year. Enter and build database. The two groups were respectively compared to see whether there was statistical significance in the differences of various indicators.

### **Isolation and identification of specimens**

Specimen collection was done after the patient was admitted to the hospital and prior to the administration of antimicrobial drugs. The surrounding skin was disinfected with iodophor before collecting specimens, and the ulcer foci were rinsed with sterile saline. Tissues from infected wounds were collected in sterile test tubes with sterilised cotton swabs at the ulcer's base and sent for analysis within 1 hour. The VITEK 2 Compact fully automated microbiological analysis system from bioMérieux, France, was used for bacterial identification, and the reagents were Gram-negative and Gram-positive bacteria from bioMérieux's bacterial identification card. The drug sensitivity test results were evaluated using the American Clinical Laboratory Standardisation Institute's (CLSI) standards.

# **Ethical approval**

This clinical study was reviewed and approved by the Ethics Committee of Fuyang People's Hospital (project number :([2022.182])).

# **Statistical methods**

SPSS 25.0 software was used for statistical analysis of the data. Count data were expressed as cases or percentages, and two independent data t-test was used for comparison of two samples; chi-square test was used for comparison of sample rates; univariate analysis with statistical significance was included in multifactor analysis, and unconditional logistic regression model was used for multifactor analysis, and risk factors with  $P < 0.05$  in multivariate analysis were independent risk factors.  $p < 0.05$  indicated that the comparison of two groups of data had Statistical significance.

# **Results**

## **Comparison of the general conditions of the two groups of patients**

A total of 147 DFU patients meeting the inclusion criteria were collected between the two groups. There were no statistically significant differences between the two groups in gender, age, duration of diabetes, glycosylated haemoglobin (HbA1c (%)) ,Erythrocyte Sedimentation Rate (ESR) ,C-reactive protein (CRP), Procalcitonin(PCT), ABI were not statistically significant ( $P > 0.05$ ). See Table 1.

# **Pathogenic bacteria distribution in DFU patients**

A total of 154 strains of pathogenic bacteria were isolated from 147 DFU patients, and 7 patients were cultured with 2 strains of bacteria in the same culture. Among the 154 pathogenic bacteria, the number of gram-positive bacteria (G+) was 89, accounting for 57.79%, and the number of gram-negative bacteria (G-) was 65, accounting for 42.21%. According to the bacterial culture results of DFU patients, the number of G- cases was significantly lower than that of G+, which was different from the results of previous studies, which may be related to regional differences.(Liu et al., 2022; Wu et al., 2018), and the G- bacteria were dominated by *Pseudomonas aeruginosa* (13.00%) and *Escherichia coli* (14.29%), while the G+ bacteria were dominated by *Staphylococcus aureus* (27.92%).See Table 2 and figure 1.

# **The detection rate and location of positive MDROs in DFU patients' secretions.**

A total of 71 MDROs (48.30%)were isolated in this study, with 33 (46.48%) of Gram-positive bacteria dominated by *S. aureus*, followed by 2 (2.82%) of *S. epidermidis*, and 16 (22.53%) of Gram-negative bacteria dominated by *Escherichia coli*, followed by 13 (18.31%) of *Pseudomonas aeruginosa* , *Klebsiella pneumoniae pneumoniae* subspecies 5(7.04%), *Citrobacter burgdorferi* 1(1.41%), and *Acinetobacter baumannii* 1 (1.41%).See Table 3 and figure 2.

# **Resistance of main MDROs to antibiotics**

In MDROs, *S. aureus* was resistant to clindamycin (96.97%), erythromycin (96.97%) and penicillin (96.97%), but *S. aureus* was sensitive to furantoin, quinuputin-dafoputin, vancomycin and cefoxitin. *Pseudomonas aeruginosa* were mainly resistant to ciprofloxacin (100%), levofloxacin (76.92%) and ceftazidime (76.92%), but *Pseudomonas aeruginosa* were sensitive to cefoxitin, linezolid, moxifloxacin,  $\beta$ -lactamase, quinuputin-dafodine, rifampicin, tegacycline and vancomycin. *Escherichia coli* was mainly sensitive to cotrimoxazole (93.75%), cefuroxime (75%), cefuroxime axetil (75%) and levofloxacin (75%). *Escherichia coli* was sensitive to tetracycline, tegacycline and vancomycin. The sensitivity rate to vancomycin in MDROs was 100%.See Table 4.

# **Results of a univariate analysis of wound infection in patients with diabetic foot.**

In the univariate analysis of risk factors for the occurrence of MDROs infection, nine possible risk factors were analyzed univariately, of which five factors, including history of previous antimicrobial exposure, hospitalization for the same infected wound >2 times/year, neuroischemic wound, Wagner classification, and combined osteomyelitis, constituted multiple drug resistance risk factors. See Table 5

### **Multifactorial analysis of risk factors for the occurrence of MDROs infection.**

Multifactorial logistic regression analysis revealed that the occurrence of MDROs in DFU patients was associated with previous antimicrobial exposure, duration of exposure, neuroischemic wounds, and osteomyelitis; it was not associated with more than 2 hospitalizations/year for the same infected wound. See Table 6.

## **Discussion**

Today, Type 2 diabetes is a worldwide epidemic affecting about 400 million people, or approximately 10 of the world's population (Li et al., 2018), and DFU is a common and serious complication in people with Type 2 diabetes, with DFU occurring in about 6.3% (5.4% -7.3%) of people with Type 2 diabetes (Zhang et al., 2017), with a prevalence of more than 15% in some developed countries (USA) (Menke et al., 2015; Nelson et al., 2018). DFU is one of the most significant and expensive complications in Type 2 diabetics. It is predicted that the incidence of DFU in diabetic foot patients will increase year by year and even reach 50% (Hurlow et al., 2018). MDROs is a common pathogen in DFU, and MDROs can lead to DFU aggravation, while improper treatment of DUF can easily cause MDROs, thus forming a vicious cycle. In recent years, with the abuse and misuse of antibiotics, bacterial resistance has become more and more serious, especially the problem of Type 2 diabetic foot infection caused by MDROs. The correct and reasonable application of antibiotics is the key to improve the efficacy of diabetic foot infection, reduce the incidence of adverse reactions and reduce the occurrence of bacterial resistance.

In the present study, G+ were predominantly *Staphylococcus aureus* (46.48%) and G- bacteria were predominantly *Escherichia coli* (22.53%) and *Pseudomonas aeruginosa* (18.31%)



in the MDROs of DFU inpatients, which is the same as the recent data reported nationally and internationally (Du F et al., 2022; Atlaw et al., 2022).

In this study, the number of G+ was slightly higher than that of G-(G+89 strains, G-65 strains), but some studies showed that the proportion of G- in the bacterial culture of DFU patients gradually increased (Ma et al., 2021), which may be related to the widespread use of broad-spectrum antibacterial drugs, particularly third-generation cephalosporins, in recent years, because the antibacterial spectrum of third-generation cephalosporins primarily targets G-. It has been shown that G- infections are positively associated with amputation and negatively associated with DFU healing (de Vries et al., 2014). Therefore, special attention should be paid to infections caused by G- in patients with DFU. Furthermore, Logistics analysis revealed that previous antimicrobial exposure, neuroischemic wounds, Wagner grade 3-5, and combined osteomyelitis were the most important independent risk factors for MDROs infections. This study shows that the history of antibacterial drug exposure is a risk factor for multiple drug resistance. Repeated use of multiple antimicrobials for a long time can easily induce mutation of drug resistance genes, and drug resistance genes can form a complex of multi-drug resistance genes through the transfer of drug resistance gene elements, resulting in the emergence of multi-drug resistance. The function of the defense mechanism of the autoimmune immune system of diabetic patients is weakened compared with that of normal people, so diabetic patients are prone to concurrent infection, which is not easy to control. In clinical practice, broad-spectrum antibiotics are often used to control infection, and long-term application of broad-spectrum antibiotics is likely to cause bacterial imbalance and increase drug resistance (Xia et al., 2021), which will not only prolong the treatment time, but also prolong the treatment time. It will also limit the use of multiple antibiotics, greatly increasing the difficulty of treatment, and have to use higher levels of antibiotics to treat infections, and over time, the formation of a vicious cycle. ERTUGRUL et al. (Ertugrul et al., 2012) found that a history of antibiotic exposure within 30 days was associated with a four-fold increase in MDROs infection. Therefore, in clinical work, antibiotics should be used rationally and the importance of standardized diagnosis and treatment

should be emphasized to patients to reduce the increase of the risk of drug resistance caused by improper drug use. Studies have shown that the incidence of infection in hospitalized patients with Type 2 diabetic foot is between 9.68% and 11.25% (Zhou et al., 2021).

This study also shows that the frequency of hospitalization for the same infected wound is also a risk factor for MDROs, suggesting that MDROs are more likely to develop infection in hospital. For patients with the same ulcer surface repeatedly hospitalized, due to repeated infection, wound debridement and other operations, the possibility of cross-infection of ulcers will be increased. After antibacterial treatment, some drug-resistant bacteria may occur on the same ulcer surface, and cross-infection will promote the growth of drug-resistant bacteria. Studies have confirmed that the incidence of hospital-associated MDROs infection can reach 67%, and compared with community-associated MDROs infection, hospital-associated MDROs infection is more adverse to patient prognosis and treatment outcomes (Wang et al., 2010). A history of antimicrobial exposure increases the risk of exposure to MDROs, which can cause iatrogenic infections when nursing or touching patients. This suggests that in addition to rational use of antibiotics, aseptic operation, hand hygiene and other links should be strictly controlled to reduce cross infection between different patients.

Some studies have found that nerve defect wound is a risk factor for MDROs infection, which is consistent with the results of this survey (Laakso et al., 2017; Datta et al., 2019; Lazaro-Martinez et al., 2022). Amin et al. found that DFU patients with nerve ischemic wounds had a 7-fold increased risk of MDROs infection (Amin & Doupis, 2016). Neuroischemic wounds, in contrast to ordinary wounds, are frequently characterised by vascular (including microvascular and macrovascular) circulation disorders and neuropathy (Khanolkar et al., 2008), and blood perfusion at the ulcer site is obstructed, making it difficult for antimicrobial drugs to reach the lesion site, resulting in a decrease in the concentration of local antimicrobial drugs and a consequent weakening of the antimicrobial effect, easily inducing the generation of drug-resistant bacteria. Furthermore, ischemic nerve wounds frequently have reduced leukocyte phagocytosis function and abnormal expression of

cytokines and inflammatory factors, increasing the difficulty of wound healing and increasing the chance of wound repeated infection (Zhang et al., 2014), so that patients for the same wound repeated visits, followed by longer and higher intensity of antibacterial drugs, so that in the past, It also comes at the cost of increasing the number of drug-resistant bacteria.

The Wagner grade was also discovered to be a risk factor for MDROs in this investigation. Studies have shown that the higher the Wagner grade, the more severe the degree of tissue destruction, infection, and ischemia of patients is, and the probability of pathogenic bacteria spreading deep is increased (Xie et al., 2017), making bacteria removal difficult and prolonging treatment time. Exposing the wound to the multidrug-resistant bacteria-prone milieu of the hospital increases the likelihood of MDROs infection in patients (McComb, 2023). Simultaneously, the bacterial composition of the wounds of patients with mixed infections was complicated, which increased the synergistic action and lethality of the bacteria and boosted the level of medication resistance.

Concurrent osteomyelitis has recently been identified as an independent risk factor for MDROs infection. (Feng et al., 2013; Garcia et al., 2020; Yan et al., 2022). The results of the present study also showed that osteomyelitis can increase the risk of MDROs infection. Patients with osteomyelitis are more difficult to treat clinically because bacteria are usually able to invade the reticular structure of the bone cell space and evade debridement and antibiotic action (Ji et al., 2014; de Mesy et al., 2018; Masters et al., 2019). In addition, these bacterial colonies can form abscesses in the skin, which act as a physical barrier and prevent immune cells from entering and killing the bacteria, allowing them to survive for a long time, thus inducing MDROs infection. At present, long-term conservative antimicrobial therapy is still the main treatment for osteomyelitis, which induces MDROs. Therefore, when DFU patients are accompanied by osteomyelitis, the use of antibiotics should be appropriately reduced, and the use of bacteria-sensitive antibiotics can avoid the occurrence of MDROs (Lipsky & Uckay, 2021).

## Conclusion

To summarise, Previous antibiotic exposure, nerve ischemia wound, Wagner grade, and

osteomyelitis were the most relevant independent risk variables for MDROs infection in this investigation. As a result, medical staff should pay attention to key risk factors when patients are admitted, which can assist us to identify high-risk groups of MDROs early and implement appropriate treatment measures as soon as possible, reducing the occurrence of MDROs infection. Furthermore, DFU can infect one or more diseases and generate a significant number of MDROs. Because drug resistance is increasing year after year, it is critical to perform bacterial biological analysis and drug sensitivity testing before utilising antibacterial medications.

## Limitations

Although our study contributed to the identification of medicines that should be taken frequently following bacterial infection in DFU patients in our location, it has several drawbacks. Firstly, the study's sample was drawn from a tertiary care institution, and the sample size was tiny. is not a complete representation of all Chinese DFU sufferers; Secondly, because the collection time node of patients' wound secretions and relevant clinical data is the day of admission, it is impossible to determine whether patients were infected with MDROs prior to admission, so it is unclear whether some factors are the cause of MDROs infection or the outcome caused by MDROs infection. Furthermore, cross-sectional studies lack the power to demonstrate causality between factors, hence prospective cohort studies are required for additional evidence.

## Author contributions

Huihui Guo and Qiwei Song have the same contribution.

Junjie Li and Tao Ning are co-corresponding authors.

Huihui Guo is responsible for writing the thesis; Huihui Guo and Qiwei Song are responsible for the main operation; Tao Ning puts forward the research idea of this chapter; Siwei Mei and Zhengqiang Xue is responsible for collecting and processing data; Junjie Li put forward a lot of suggestions for the later article modification and the places to be improved.

## **Ethics statement**

This study was approved by the Institutional Review Board of Fuyang People's Hospital([2022.182]).The study was a non-invasive retrospective study that collected patients using the hospital information system's electronic medical records, did not interfere with patients' usual care, did not damage any medical rights of patients, and did not increase patients' risk of disease. Because the research effort did not concern personal privacy or economic interests, informed patient permission was not obtained. To properly protect patients' personal privacy, the identities of included patients were coded, and the medical records were saved in a special computer at Fuyang People's Hospital's Diabetic Foot Clinic for only the investigators' access.

This study was supported financially by Fuyang People's Hospital. We thank all participants of this research.

## **Funding**

This study was supported financially by Fuyang People's Hospital. We thank all participants of this research. Masquelet technique combined with transverse bone transfer technique for the treatment of refractory Wagner III and IV diabetic foot (FY2021-027), and tibial periosteum lateral extension for Wagner III and IV diabetic foot in a multidisciplinary collaborative mode (2022xkj225) all provided funding for this study. The funders had no involvement in the study's design, data collection and analysis, publication decision, or manuscript writing.

## **Conflict of interest**

There is no conflict of interest.

## **Consent for publication**

The author(s) read and approved the final manuscript.

## **Availability of data and materials**

All the data were from Fuyang People's Hospital, and the data were true and legally liable.

## **Data availability**

Data are available. All data used in the study have been uploaded.

# References:

- Adeyemo, A.T., Kolawole, B., Rotimi, V.O., and Aboderin, A.O. 2021. Multicentre study of the burden of multidrug-resistant bacteria in the aetiology of infected diabetic foot ulcers. *African Journal of Laboratory Medicine* 10:1261. 10.4102/ajlm.v10i1.1261
- Amin, N., and Doupis, J. 2016. Diabetic foot disease: From the evaluation of the "foot at risk" to the novel diabetic ulcer treatment modalities. *World Journal of Diabetes* 7:153-164. 10.4239/wjd.v7.i7.153
- Armstrong, D.G., Boulton, A., and Bus, S.A. 2017. Diabetic Foot Ulcers and Their Recurrence. *NEW ENGLAND JOURNAL OF MEDICINE* 376:2367-2375. 10.1056/NEJMra1615439
- Atlaw, A., Kebede, H.B., Abdela, A.A., and Woldeamanuel, Y. 2022. Bacterial isolates from diabetic foot ulcers and their antimicrobial resistance profile from selected hospitals in Addis Ababa, Ethiopia. *Frontiers in Endocrinology* 13:987487. 10.3389/fendo.2022.987487
- Dai, J., Jiang, C., Chen, H., and Chai, Y. 2020. Assessment of the Risk Factors of Multidrug-Resistant Organism Infection in Adults With Type 1 or Type 2 Diabetes and Diabetic Foot Ulcer. *Canadian Journal of Diabetes* 44:342-349. 10.1016/j.jcjd.2019.10.009
- Datta, P., Chander, J., Gupta, V., Mohi, G.K., and Attri, A.K. 2019. Evaluation of various risk factors associated with multidrug-resistant organisms isolated from diabetic foot ulcer patients. *Journal of Laboratory Physicians* 11:58-62. 10.4103/JLP.JLP\_106\_18
- de Mesy, B.K., MacDonald, A., Schwarz, E.M., and Oh, I. 2018. Chronic Osteomyelitis with Staphylococcus aureus Deformation in Submicron Canaliculi of Osteocytes: A Case Report. *JBJS Case Connect* 8:e8. 10.2106/JBJS.CC.17.00154
- de Vries, M.G., Ekkelenkamp, M.B., and Peters, E.J. 2014. Are clindamycin and ciprofloxacin appropriate for the empirical treatment of diabetic foot infections? *EUROPEAN JOURNAL OF CLINICAL MICROBIOLOGY & INFECTIOUS DISEASES* 33:453-456. 10.1007/s10096-013-1977-7
- Du F, Ma, J., Gong, H., Bista, R., Zha, P., Ren, Y., Gao, Y., Chen, D., Ran, X., and Wang, C. 2022. Microbial Infection and Antibiotic Susceptibility of Diabetic Foot Ulcer in China: Literature Review. *Frontiers in Endocrinology* 13:881659. 10.3389/fendo.2022.881659
- Ertugrul, B.M., Oncul, O., Tulek, N., Willke, A., Sacar, S., Tunccan, O.G., Yilmaz, E., Kaya, O., Ozturk, B., Turhan, O., Yapar, N., Ture, M., and Akin, F. 2012. A prospective, multi-center study: factors related to the management of diabetic foot infections. *EUROPEAN JOURNAL OF CLINICAL MICROBIOLOGY & INFECTIOUS DISEASES* 31:2345-2352. 10.1007/s10096-012-1574-1
- Feng, S.H., Chu, Y.J., Wang, P.H., Jun, X., Min, D., and Li, X.M. 2013. Risk factors and gene type for infections of MRSA in diabetic foot patients in Tianjin, China. *International Journal of Lower Extremity Wounds* 12:106-112. 10.1177/1534734613489991
- Garcia, Z.V., Hernandez, T.A., Garcia, V.E., Soria, C.T., Canteras, J.M., Ruiz, G.J., Gomez, G.J., Hernandez, M.A., and Barberan, J. 2020. Risk factors for methicillin-resistant Staphylococcus aureus and extended-spectrum ss-lactamase producing Enterobacterales in patients with diabetic foot infections requiring hospital admission.

361 *Revista Espanola de Quimioterapia* 33:430-435. 10.37201/req/101.2020

362 Hawkins, B.K., Barnard, M., Barber, K.E., Stover, K.R., Cretella, D.A., Wingler, M., and Wagner, J.L. 2022.

363 Diabetic foot infections: A microbiologic review. *Foot (Edinb)* 51:101877. 10.1016/j.foot.2021.101877

364 Hitam, S., Hassan, S.A., and Maning, N. 2019. The Significant Association between Polymicrobial Diabetic Foot

365 Infection and Its Severity and Outcomes. *Malaysian Journal of Medical Sciences* 26:107-114.

366 10.21315/mjms2019.26.1.10

367 Hurlow, J.J., Humphreys, G.J., Bowling, F.L., and McBain, A.J. 2018. Diabetic foot infection: A critical

368 complication. *International Wound Journal* 15:814-821. 10.1111/iwj.12932

369 Ji, X., Jin, P., Chu, Y., Feng, S., and Wang, P. 2014. Clinical characteristics and risk factors of diabetic foot ulcer

370 with multidrug-resistant organism infection. *International Journal of Lower Extremity Wounds* 13:64-71.

371 10.1177/1534734614521236

372 Kandemir, O., Akbay, E., Sahin, E., Milcan, A., and Gen, R. 2007. Risk factors for infection of the diabetic foot

373 with multi-antibiotic resistant microorganisms. *JOURNAL OF INFECTION* 54:439-445.

374 10.1016/j.jinf.2006.08.013

375 Khanolkar, M.P., Bain, S.C., and Stephens, J.W. 2008. The diabetic foot. *QJM-AN INTERNATIONAL JOURNAL*

376 *OF MEDICINE* 101:685-695. 10.1093/qjmed/hcn027

377 Laakso, M., Honkasalo, M., Kiiski, J., Ala-Houhala, M., Haapasalo, H., Laine, H.J., Luukkaala, T., Lahtela, J., and

378 Kaartinen, I. 2017. Re-organizing inpatient care saves legs in patients with diabetic foot infections. *DIABETES*

379 *RESEARCH AND CLINICAL PRACTICE* 125:39-46. 10.1016/j.diabres.2017.01.007

380 Lazaro-Martinez, J.L., Garcia-Madrid, M., Garcia-Alamino, J.M., Bohbot, S., Garcia-Klepzig, J.L., and Garcia-

381 Alvarez, Y. 2022. Increasing Transcutaneous Oxygen Pressure in Patients With Neuroischemic Diabetic Foot

382 Ulcers Treated With a Sucrose Octasulfate Dressing: A Pilot Study. *International Journal of Lower Extremity*

383 *Wounds* 21:450-456. 10.1177/1534734620952244

384 Li, J., Parrott, S., Sweeting, M., Farmer, A., Ross, J., Dack, C., Pal, K., Yardley, L., Barnard, M., Hudda, M.,

385 Alkhalidi, G., and Murray, E. 2018. Cost-Effectiveness of Facilitated Access to a Self-Management Website,

386 Compared to Usual Care, for Patients With Type II Diabetes (HeLP-Diabetes): Randomized Controlled Trial.

387 *JOURNAL OF MEDICAL INTERNET RESEARCH* 20:e201. 10.2196/jmir.9256

388 Lipsky, B.A., Berendt, A.R., Cornia, P.B., Pile, J.C., Peters, E.J., Armstrong, D.G., Deery, H.G., Embil, J.M.,

389 Joseph, W.S., Karchmer, A.W., Pinzur, M.S., and Senneville, E. 2012. 2012 Infectious Diseases Society of America

390 clinical practice guideline for the diagnosis and treatment of diabetic foot infections. *CLINICAL INFECTIOUS*

391 *DISEASES* 54:e132-e173. 10.1093/cid/cis346

392 Lipsky, B.A., and Uckay, I. 2021. Treating Diabetic Foot Osteomyelitis: A Practical State-of-the-Art Update.

393 *Medicina-Lithuania* 57. 10.3390/medicina57040339

394 Liu, X., Ren, Q., Zhai, Y., Kong, Y., Chen, D., and Chang, B. 2022. Risk Factors for Multidrug-Resistant

395 Organisms Infection in Diabetic Foot Ulcer. *Infection and Drug Resistance* 15:1627-1635. 10.2147/IDR.S359157

396 Ma, Y.N., Zhang, L.X., Hu, Y.Y., and Shi, T.L. 2021. Nomogram Model for Predicting the Risk of Multidrug-

397 Resistant Bacteria Infection in Diabetic Foot Patients. *Infection and Drug Resistance* 14:627-637.

398 10.2147/IDR.S287852

399 Magiorakos, A.P., Srinivasan, A., Carey, R.B., Carmeli, Y., Falagas, M.E., Giske, C.G., Harbarth, S., Hindler, J.F.,

400 Kahlmeter, G., Olsson-Liljequist, B., Paterson, D.L., Rice, L.B., Stelling, J., Struelens, M.J., Vatopoulos, A., Weber,

401 J.T., and Monnet, D.L. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an

international expert proposal for interim standard definitions for acquired resistance. *CLINICAL MICROBIOLOGY AND INFECTION* 18:268-281. 10.1111/j.1469-0691.2011.03570.x

Masters, E.A., Trombetta, R.P., de Mesy, B.K., Boyce, B.F., Gill, A.L., Gill, S.R., Nishitani, K., Ishikawa, M., Morita, Y., Ito, H., Bello-Irizarry, S.N., Ninomiya, M., Brodell, J.J., Lee, C.C., Hao, S.P., Oh, I., Xie, C., Awad, H.A., Daiss, J.L., Owen, J.R., Kates, S.L., Schwarz, E.M., and Muthukrishnan, G. 2019. Evolving concepts in bone infection: redefining "biofilm", "acute vs. chronic osteomyelitis", "the immune proteome" and "local antibiotic therapy". *Bone Research* 7:20. 10.1038/s41413-019-0061-z

Matta-Gutierrez, G., Garcia-Morales, E., Garcia-Alvarez, Y., Alvaro-Afonso, F.J., Molines-Barroso, R.J., and Lazaro-Martinez, J.L. 2021. The Influence of Multidrug-Resistant Bacteria on Clinical Outcomes of Diabetic Foot Ulcers: A Systematic Review. *Journal of Clinical Medicine* 10. 10.3390/jcm10091948

McComb, W.J. 2023. Wagner Grading of Diabetic Foot Ulcers and National Pressure Injury Advisory Panel Staging of Pressure Injuries: A Comparison for Clinical Use. *Advances in Skin & Wound Care* 36:278-279. 10.1097/01.ASW.0000922688.68275.b1

Menke, A., Casagrande, S., Geiss, L., and Cowie, C.C. 2015. Prevalence of and Trends in Diabetes Among Adults in the United States, 1988-2012. *JAMA-JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION* 314:1021-1029. 10.1001/jama.2015.10029

Monteiro-Soares, M., Boyko, E.J., Jeffcoate, W., Mills, J.L., Russell, D., Morbach, S., and Game, F. 2020. Diabetic foot ulcer classifications: A critical review. *DIABETES-METABOLISM RESEARCH AND REVIEWS* 36 Suppl 1:e3272. 10.1002/dmrr.3272

Nelson, A., Wright-Hughes, A., Backhouse, M.R., Lipsky, B.A., Nixon, J., Bhogal, M.S., Reynolds, C., and Brown, S. 2018. CODIFI (Concordance in Diabetic Foot Ulcer Infection): a cross-sectional study of wound swab versus tissue sampling in infected diabetic foot ulcers in England. *BMJ Open* 8:e19437. 10.1136/bmjopen-2017-019437

Noor, S., Borse, A.G., Ozair, M., Raghav, A., Parwez, I., and Ahmad, J. 2017. Inflammatory markers as risk factors for infection with multidrug-resistant microbes in diabetic foot subjects. *Foot (Edinb)* 32:44-48. 10.1016/j.foot.2017.05.001

Pessoa, E.C.T., Duarte, B., Joao, A.L., Coelho, M., Formiga, A., Pinto, M., and Neves, J. 2020. Multidrug-resistant bacteria in diabetic foot infections: Experience from a portuguese tertiary centre. *International Wound Journal* 17:1835-1839. 10.1111/iwj.13473

Raspovic, K.M., and Wukich, D.K. 2014. Self-reported quality of life and diabetic foot infections. *Journal of Foot & Ankle Surgery* 53:716-719. 10.1053/j.jfas.2014.06.011

Saltoglu, N., Ergonul, O., Tulek, N., Yemisen, M., Kadanali, A., Karagoz, G., Batirel, A., Ak, O., Sonmezer, C., Eraksoy, H., Cagatay, A., Surme, S., Nemli, S.A., Demirdal, T., Coskun, O., Ozturk, D., Ceran, N., Pehlivanoglu, F., Sengoz, G., Aslan, T., Akkoyunlu, Y., Oncul, O., Ay, H., Mulazimoglu, L., Erturk, B., Yilmaz, F., Yoruk, G., Uzun, N., Simsek, F., Yildirmak, T., Yasar, K.K., Sonmezoglu, M., Kucukardali, Y., Tuna, N., Karabay, O., Ozgunes, N., and Sargin, F. 2018. Influence of multidrug resistant organisms on the outcome of diabetic foot infection. *INTERNATIONAL JOURNAL OF INFECTIOUS DISEASES* 70:10-14. 10.1016/j.ijid.2018.02.013

Wang, S.H., Sun, Z.L., Guo, Y.J., Yang, B.Q., Yuan, Y., Wei, Q., and Ye, K.P. 2010. Meticillin-resistant *Staphylococcus aureus* isolated from foot ulcers in diabetic patients in a Chinese care hospital: risk factors for infection and prevalence. *JOURNAL OF MEDICAL MICROBIOLOGY* 59:1219-1224. 10.1099/jmm.0.020537-0

Wu, M., Pan, H., Leng, W., Lei, X., Chen, L., and Liang, Z. 2018. Distribution of Microbes and Drug Susceptibility in Patients with Diabetic Foot Infections in Southwest China. *Journal of Diabetes Research* 2018:9817308.



10.1155/2018/9817308

Xia, W., He, W., Luo, T., and Tang, N. 2021. Risk factors for multidrug-resistant bacterial infections in patients with diabetic foot ulcers: a systematic review and meta-analysis. *Annals of Palliative Medicine* 10:12618-12630. 10.21037/apm-21-3406

Xie, X., Bao, Y., Ni, L., Liu, D., Niu, S., Lin, H., Li, H., Duan, C., Yan, L., Huang, S., and Luo, Z. 2017. Bacterial Profile and Antibiotic Resistance in Patients with Diabetic Foot Ulcer in Guangzhou, Southern China: Focus on the Differences among Different Wagner's Grades, IDSA/IWGDF Grades, and Ulcer Types. *International Journal of Endocrinology* 2017:8694903. 10.1155/2017/8694903

Yan, X., Song, J.F., Zhang, L., and Li, X. 2022. Analysis of risk factors for multidrug-resistant organisms in diabetic foot infection. *BMC Endocrine Disorders* 22:46. 10.1186/s12902-022-00957-0

Zhang, J., Chu, Y., Wang, P., Ji, X., Li, X., Wang, C., and Peng, Y. 2014. Clinical outcomes of multidrug resistant pseudomonas aeruginosa infection and the relationship with Type III secretion system in patients with diabetic foot. *International Journal of Lower Extremity Wounds* 13:205-210. 10.1177/1534734614545878

Zhang, P., Lu, J., Jing, Y., Tang, S., Zhu, D., and Bi, Y. 2017. Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis (dagger). *ANNALS OF MEDICINE* 49:106-116. 10.1080/07853890.2016.1231932

Zhou, Y., Chi, J., Lv, W., and Wang, Y. 2021. Obesity and diabetes as high-risk factors for severe coronavirus disease 2019 (Covid-19). *DIABETES-METABOLISM RESEARCH AND REVIEWS* 37:e3377. 10.1002/dmrr.3377

Zubair, M., and Ahmad, J. 2019. Potential risk factors and outcomes of infection with multidrug resistance among diabetic patients having ulcers: 7 years study. *Diabetes & Metabolic Syndrome-Clinical Research & Reviews* 13:414-418. 10.1016/j.dsx.2018.10.014

**Table 1**(on next page)

Comparison of general clinical data between the two groups of patients.

Comparison of general clinical data between the two groups of patients.

**Table 1**

Comparison of general clinical data between the two groups of patients.

Index	MDROs Group (n=71)	NMDROs Group(n=76)	Statistics	P Value
Gender				
Male	42	51	$\chi^2=0.793$	0.373
Female	29	25		
Age	64.60±11.25	62.00±11.45	t=1.354	0.181
Hospital stays (days)	12.42±7.35	11.68±6.21	t=0.757	0.452
HbA1c (%)	8.83±1.96	8.52±1.64	t=1.038	0.303
Systolic pressure	144.87±19.54	143.55±18.81	t=0.378	0.706
Diastolic pressure	83.4±10.77	82.31±12.93	t=0.578	0.565
Cholesterol(mmol/L)	4.62±1.26	4.22±1.59	t=1.642	0.105
TG(mmol/L)	1.40±0.59	1.34±0.71	t=0.585	0.561
ESR(mm/h)	65.04±24.75	60.12±28.62	t=1.250	0.215
CRP(mg/L)	40.34±17.76	36.65±22.20	t=1.191	0.238
PCT(pg/ml)	260.20±48.53	245.62±44.63	t=1.862	0.067
ABI	0.75±0.28	0.82±0.29	t=1.834	0.071

Abbreviations:

HbA1c:Hemoglobin A1c;CHO:Cholestero;TG ;Triglyceride;TG:Triglyceride;ESR:Erythrocyte Sedimentation Rate;CRP:C-reactive protein, PCT:Procalcitonin;ABI:Ankle-Brachial Index.

\*P < 0.05 indicated statistically significant difference.

**Table 2**(on next page)

Distribution of pathogenic bacteria in patients with DFU.

Distribution of pathogenic bacteria in patients with DFU.

**Table 2**

Distribution of pathogenic bacteria in patients with DFU.

Bacteria	MDROs Group(n=71)	NMDROs Group(n=76)	Total	Percentage (%)
Gram-positive bacteria				
Staphylococcus aureus	33	10	43	27.92
Other Staphylococcus	2	6	8	5.19
Enterococcus	0	8	8	5.19
Streptococcus	0	11	11	7.14
Others	0	19	19	12.33
Gram-negative bacteria				
Pseudomonas aeruginosa	13	7	20	13.00
Klebsiella pneumoniae	5	5	10	6.50
E. coli	16	6	22	14.29
Enterobacteriaceae	0	3	3	1.95
Aspergillus singularis	0	6	6	3.89
Others	2	2	4	2.60
Total	71	83	154	100

**Table 3**(on next page)

Distribution characteristics of MDROs

Distribution characteristics of MDROs

**Table 3**

Distribution characteristics of MDROs

Bacteria	Frequency	Percentage (%)
Staphylococcus aureus	33	46.48
E. coli	16	22.53
Pseudomonas aeruginosa	13	18.31
Klebsiella pneumoniae subspecies	5	7.04
Staphylococcus epidermidis	2	2.82
Citrobacter brucei	1	1.41
Acinetobacter baumannii	1	1.41
Total	71	100

**Table 4**(on next page)

Drug resistance rate of main MDROs to antibiotics in patients with DFU and positive secretion.

Drug resistance rate of main MDROs to antibiotics in patients with DFU and positive secretion.



# Table 4

Drug resistance rate of main MDROs to antibiotics in patients with DFU and positive secretion.

Antibacterial drugs	Staphylococcus aureus (n=33)		Pseudomonas aeruginosa (n=13)		E. coli (n=16)	
	Frequency	Percentage	Frequency	Percentage	Frequency	Percentage
		(%)		(%)		(%)
Ciprofloxacin	8	24.24	13	100	1	6.25
Clindamycin	32	96.97	3	23.08	-	-
Erythromycin	32	96.97	3	23.08	-	-
Cefoxitin	2	6.06	0	0	6	37.50
Inducible clindamycin resistance	22	66.67	3	23.08	-	-
Linezolid	1	3.03	0	0	-	-
Levofloxacin	8	24.24	10	76.92	12	75
Moxifloxacin	1	3.03	0	0	-	-
Furantoin	0	0	6	46.15	1	6.25
Benzocillin	31	93.94	3	23.08	-	-
$\beta$ -lactamase	1	3.03	0	0	-	-
Penicillin	32	96.97	3	23.08	-	-
Quinupristin-Dafopristin	0	0	0	0	-	-
Rifampin	1	3.03	0	0	-	-
Cotrimoxazole	7	21.21	1	7.69	15	93.75
Tetracycline	13	39.39	3	23.08	0	0
Tigecycline	1	3.03	0	0	0	0
Vancomycin	0	0	0	0	-	-
Ceftazidime	-	-	10	76.92	11	68.75
Ceftriaxone	-	-	6	46.15	17	106.25
Cefuroxime	-	-	-	-	12	75
Cefepime	-	-	1	7.69	9	56.25
Cefoxitin	0	0	-	-	1	6.25
Imipenem	-	-	1	7.69	0	0
Cefoperazone / Sulbactam	-	-	0	0	2	12.50
Piperacillin-tazobactam	-	-	1	7.69	1	6.25
Amoxicillin-rod acid	-	-	-	-	6	37.50
Amikacin	-	-	-	-	2	12.50
Ampicillin	-	-	6	46.15	6	37.50
Aminotransol	-	-	-	-	5	31.25
Cefotetan	-	-	5	38.46	0	0
Cefazolin	-	-	6	46.15	6	37.50

Tobramycin	-	-	0	0	6	37.50
Ampicillin-Sulbactam	-	-	6	46.15	6	37.50
Polymyxin E	-	-	0	0	-	-
Meropenem	-	-	0	0	-	-
Ticarcillin-rod acid	-	-	4	30.77	-	-

Note: "-" means not measured.

**Table 5**(on next page)

Univariate analysis of risk factors for the infection of MDROs.

Univariate analysis of risk factors for the infection of MDROs.

**Table 5.**

Univariate analysis of risk factors for the infection of MDROs.

Exposure factor	MDROs Group (n=71)		NMDRO Group (n=83)		OR-value	95% CI	P-value
	+	-	+	-			
Previous antimicrobial exposure	63	8	35	41	8.944	(2.349,34.049)	0.001
Combined osteomyelitis	51	20	14	62	4.140	(1.219,14.064)	0.023
Concomitant neuropathy	60	11	55	21	0.297	(0.078,1.122)	0.073
Concomitant retinopathy	20	51	21	55	2.141	(0.612,7.484)	0.233
Complicated nephropathy	32	41	31	45	0.468	(0.135,1.623)	0.232
The same infected wound >2 times per year	19	52	18	58	0,143	(0.030,0.682)	0.015
Whether to amputate toe/limb	16	55	10	66	2.746	(0.642,11.744)	0.173
Wanger grade 3-5	50	21	27	49	0.157	(0.030,0.808)	0.027
Ischemic nerve wound	48	23	25	51	36.790	(8.386,161.395)	0.000

**Table 6**(on next page)

Multivariate Logistic regression analysis for the infection of MDROs.

Multivariate Logistic regression analysis for the infection of MDROs.

**Table 6**


Multivariate Logistic regression analysis for the infection of MDROs.

Selected variable	$\beta$	SE	Wald	P-value	OR( 95% CI)
Previous antimicrobial exposure	1.925	0.606	10.085	0.001	6.853(2.090,22.477)
Combined osteomyelitis	1.294	0.577	5.030	0.025	3.649(1.177,11.311)
Wanger grade 3-5	1.708	0.777	4.833	0.028	0.181(0.040,0.831)
Ischemic nerve wound	3.074	0.659	21.762	0.000	21.629(5.945,78.697)
The same infected wound >2 times per year	1.288	0.681	3.579	0.059	0.276(0.073,1.047)

# Figure 1

Figuer1 Distribution of pathogenic bacteria in DFU patients

A:Bacterial distribution;B:G+ bacterial distribution;C:G- bacterial distribution.



# Figure 2

Figuer2 Bacteria distribution of MDROS.

Bacteria distribution of MDROS.

