

Comprehensive genomic characterisation of the NAC transcription factor family and its response to drought stress in *Eucommia ulmoides*

Qi Wang¹, FengCheng Hu², ZhaoQun Yao³, XinFeng Zhao², GuangMing Chu^{Corresp., 1}, Jing Ye^{Corresp. 1}

¹ Laboratory of Forestry Department, Agricultural College, Shihezi University, Shihezi, China

² Lveyang County Forest Tree Seedling Workstation, Forestry Bureau of Lveyang County, Lveyang, China

³ Laboratory of Plant Protection Department, Agricultural College, Shihezi University, Shihezi, China

Corresponding Authors: GuangMing Chu, Jing Ye

Email address: chgmjx@163.com, yejing_shz@foxmail.com

The NAC transcription factor family enhances plant adaptation to environmental challenges by participating in signalling pathways triggered by abiotic stressors and hormonal cues. We identified 69 NAC genes in the *Eucommia ulmoides* genome and renamed them according to their chromosomal distribution. These EuNAC proteins were clustered into 13 sub-families and distributed on 16 chromosomes and 2 scaffolds. The gene structures suggested that the number of exons varied from 2 to 8 among these *EuNACs*, with a multitude of them containing three exons. Duplicated events resulted in a large gene family; 12 and 4 pairs of *EuNACs* were the result of segmental and tandem duplicates, respectively. The drought-stress response pattern of 12 putative *EuNACs* was observed under drought treatment, revealing that these *EuNACs* could play crucial roles in mitigating the effects of drought stress responses and serve as promising candidate genes for genetic engineering aimed at enhancing the drought stress tolerance of *E. ulmoides*. This study provides insight into the evolution, diversity, and characterisation of NAC genes in *E. ulmoides* and will be helpful for future characterisation of putative *EuNACs* associated with water deficit.

Comprehensive genomic characterisation of the NAC transcription factor family and its response to drought stress in *Eucommia ulmoides*

Qi Wang¹, FengCheng Hu², ZhaoQun Yao³, XinFeng Zhao², GuangMing Chu^{1*}, Jing Ye^{1*}

¹ Laboratory of Forestry Department, Agricultural College, Shihezi University, Shihezi, Xinjiang, China

² Lveyang County Forest Tree Seedling Workstation, Forestry Bureau of Lveyang County, Lveyang, Shanxi, China

³ Laboratory of Plant Protection Department, Agricultural College, Shihezi University, Shihezi, Xinjiang, China

Corresponding Author:

GuangMing Chu¹, Jing Ye¹

Beiwu Road Shihezi University Beiyuan New Area, Shihezi, Xinjiang, 832003, China

Email address: chgmxi@163.com, yejing_shz@foxmail.com

Abstract

The NAC transcription factor family enhances plant adaptation to environmental challenges by participating in signalling pathways triggered by abiotic stressors and hormonal cues. We identified 69 *NAC* genes in the *Eucommia ulmoides* genome and renamed them according to their chromosomal distribution. These *EuNAC* proteins were clustered into 13 sub-families and distributed on 16 chromosomes and 2 scaffolds. The gene structures suggested that the number of exons varied from 2 to 8 among these *EuNACs*, with a multitude of them containing three exons. Duplicated events resulted in a large gene family; 12 and 4 pairs of *EuNACs* were the result of segmental and tandem duplicates, respectively. The drought-stress response pattern of 12 putative *EuNACs* was observed under drought treatment, revealing that these *EuNACs* could play crucial roles in mitigating the effects of drought stress responses and serve as promising candidate genes for genetic engineering aimed at enhancing the drought stress tolerance of *E. ulmoides*. This study provides insight into the evolution, diversity, and characterisation of *NAC* genes in *E. ulmoides* and will be helpful for future characterisation of putative *EuNACs* associated with water deficit.

Keywords: *Eucommia ulmoides*, NAC family, drought-responsive, gene expression, phylogenetic analysis

Introduction

Eucommia ulmoides Oliver is a highly valued tertiary relict perennial tree species native to China (Deng et al. 2022). It is widely used in industry because it not only produces wood but is also a valuable raw biomaterial for extracting the active ingredients of Chinese medicine and *trans*-rubber (Wei et al. 2021; Zhu & Sun 2018). Compared with *cis*-rubber, *trans*-rubber has unique

characteristics, such as high hardness, resistance to acid and alkali corrosion, good insulation, and a low thermal expansion/contraction coefficient (Enoki et al. 2003; Kent & Swinney 1966; Rose & Steinbuchel 2005). To improve the viscosity, resilience, elasticity, weather resistance, and tensile strength of *trans*-rubber, it can be made into *cis*-rubber by vulcanisation (Yan 1995). The vulnerability of *Hevea brasiliensis* to pests and diseases, as well as its narrow habitat, has led to significant challenges for the rubber industry (Tang et al. 2016). *Eucommia ulmoides* has wide adaptability, few pests and diseases, and its leaves, bark, and pericarp are rich in *trans*-rubber. Therefore, *E. ulmoides* is considered an ideal alternative or complementary tree species to *H. brasiliensis* (Wuyun et al. 2018). During the growth of *E. ulmoides*, some environmental factors, such as drought and low-temperature stress, can prevent its full genetic potential, resulting in reduced yield and even plant death (Zuo et al. 2022). The identification and utilisation of resistance genes is the basis for breeding resistant varieties. Transcription factors (TFs), which activate or inhibit their expression by specifically binding to *cis*-acting elements on the promoters of target genes, play an important role in many biological processes (Yuan et al. 2020). As a plant-specific supergene family, *NAC* has been demonstrated to play a key role in plant growth and development and in the response to abiotic stress (Du et al. 2022b; Hussain et al. 2017). Notably, *NAC* is very important in plant adaptation to land (Xu et al. 2014). Therefore, *NAC* family genes have been widely studied in many species. However, the identification and analysis of *NAC* genes in *E. ulmoides* have not been emphasised.

The *NAC* [no apical meristem (NAM), *Arabidopsis* transcription activator (ATAF1/ATAF2), and cup-shaped cotyledon (CUC2)] family is one of the largest gene families in plants. The N-terminus region has a highly conserved NAM domain, and the C-terminus consists of variable transcriptional regulatory regions, the latter of which have been implicated in specific biological functions (Shao et al. 2015). Increasing evidence suggests that *NAC* TFs have multiple functions in plant-responses to biotic and abiotic stress. *SINAC1* is involved in the process of fruit softening and fruit pigmentation based on the phytohormone pathway (Ma et al. 2014). *NAC13* has important significance in popular responses to salt stress (Zhang et al. 2019). In wheat, overexpression of *TaNAC1-D1* and *TaNAC071-A* improves resistance to *Fusarium* head blight (Perochon et al. 2019) and drought (Mao et al. 2022), respectively; *TaNAC30* negatively regulates stripe rust (Wang et al. 2018). Furthermore, *TaNAC29* improves salt tolerance by strengthening the antioxidant system (Xu et al. 2015). *GhirNAC2* regulates ABA biosynthesis and stomatal closure by regulating *GhNCED3a/3c* expression, thus playing an active role in cotton drought resistance (Shang et al. 2020). Although *NAC* TFs are related to various developmental processes and stress responses in plants, the specific functions of most *NAC* genes remain obscure, especially in *E. ulmoides*.

The chromosome-level genome of *E. ulmoides* was recently sequenced (Li et al. 2020); this provides the opportunity to systematically study the *NAC* gene family and to explore the potential functional involved in *E. ulmoides* biotic and abiotic responses. In the present study, we performed genome-wide identification and characterisation of *NAC* proteins based on the genome of *E. ulmoides*. In addition, we surveyed their expression under drought stress by transcriptome

sequencing. This study will lay the foundation for further studies of the molecular mechanisms of NAC TFs in *E. ulmoides* response to drought stress.

Materials & Methods

Identification of EuNAC proteins from the *E. ulmoides* genome

The complete genome data of *E. ulmoides* were obtained from the Gene Warehouse (<https://ngdc.cncb.ac.cn/gwh/Assembly/25206/show>). The NAC protein sequences of *Arabidopsis* were obtained from *Arabidopsis* Information Resources (TAIR, <https://www.arabidopsis.org/index.jsp>), and the protein sequences of *poplar* and rice were both derived from the Ensembl Plants website (<http://plants.ensembl.org/index.html>). The Hidden Markov model (HMM) files of the NAC domain (PF01849) and the NAM domain (PF02365) were obtained from the Pfam database (<https://pfam.sanger.ac.uk>), which were used for identification analysis. HMMER 3.3.2 (<http://hmmmer.org/>) was then employed to scan the NAC proteins from the *E. ulmoides* genome with the default parameters. The candidate *EuNACs* were further validated by the NCBI Conserved Domain Search Service (CD Search) (<https://www.ncbi.nlm.nih.gov>), SMART (<http://smart.embl-heidelberg.de>), and Pfam database. Proteins without NAC and NAM domains and duplicates were manually deleted. The molecular weight (MW) and isoelectric point (pI) of each protein were analysed using the Expasy pI/Mw tool (<https://www.expasy.org>).

Phylogenetic analysis of EuNAC proteins

The *Arabidopsis* NAC protein sequences were downloaded from the TAIR database (<http://www.Arabidopsis.org>). Full-length protein sequence multiple alignments were performed using the ClustalW programme (Larkin et al. 2007). MEGA 6.0 software (Hall. 2013) was employed to construct an unrooted phylogenetic tree of *E. ulmoides* and *Arabidopsis* NAC proteins using the neighbour-joining (NJ) method with 1000 bootstrap iterations. All *EuNAC* proteins were classified according to the NAC protein classification criteria in *Arabidopsis* (Ooka et al. 2003a).

Conserved motif and gene structure analysis of *EuNAC* genes

The Gene Structure Display Server (GSDS; <http://gsds.cbi.pku.edu.cn/>) programme was used to explore the exon/intron structure pattern of the *EuNAC* genes by comparing their predicted coding sequence with the corresponding full-length gDNA sequence. Multiple Expectation Maximization for Motif Elicitation (MEME) (<http://meme-suite.Org/>) programmes were employed to identify the conserved domains for candidate *EuNAC* proteins with default parameters. The conserved motifs and exon/intron structure were visualised using Tbtools (Chen et al. 2020).

Genome distribution, selective pressure, and syntenic analysis of *EuNAC* genes

The location of each *EuNAC* gene on the chromosome was determined based on the *E. ulmoides* genome annotation file and visualised using TBtools software (Chen et al. 2020). MCScan X software (Wang et al. 2012) with default parameters was employed to analyse duplication events, and the intra-species and inter-species collinearity relationships. Circos software (Krzywinski et al. 2009) and TBtools were used for visualisation. TBtools were also used to calculate the nonsynonymous (*Ka*) and synonymous (*Ks*) rates of *EuNAC* homologous genes. The selection pressure acting on the gene pairs was calculated based on the *Ka/Ks* ratio, and the

dates of each duplication event were further deduced with the formula $T = Ks/2\lambda$, the mean synonymous substitution rate (λ) was assumed to be 6.5×10^{-9} (Liu et al. 2021; Lynch & Conery 2000a).

Promoter region analysis of *EuNAC* genes

The 2-kb promoter sequences upstream of the *EuNAC* genes start codon (ATG) were extracted, and the cis-acting elements and their potential related functions were predicted with the PlantCARE online server (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>). The 20 cis-acting elements with the highest frequency were visualised using Tbttools (Chen et al. 2020).

Expression analysis of *EuNAC* genes under drought stress

Two-year-old ‘Qinzhong 1’ grafted potted plants with consistent growth were placed in the Agricultural College of Shihezi University and well managed in the natural environment. Three months later, they were subjected to drought stress treatment. For drought treatment, the soil was saturated with water, and watering was terminated. Leaves were collected at 0, 15, 30, and 45 d and labelled CK, D15, D30, and D45, respectively. Each sample was pooled from six individual plants, and three biological replicates were set for each treatment. Samples were quickly cleaned with distilled water, immediately frozen in liquid nitrogen, and stored at -80°C for future use.

The above samples were submitted to Beijing Novogen Bioinformatics Technology Co., Ltd. (Beijing, China) for cDNA library construction and transcriptome sequencing. The data were uploaded to the National Center for Biotechnology Information (NCBI) Sequence Read Archive, with accession number PRJNA961078. Gene expression levels were normalised with FPKM (fragment per kilobase per million mapped reads) and visualised with TBtools.

Results

Identification of *EuNAC* proteins from the *E. ulmoides* genome

We identified 69 *NAC* genes in the *E. ulmoides* genome, named *EuNAC1* TO *EuNAC69*, according to their order on the chromosomes (Table S1). The *EuNAC* proteins varied significantly in length and molecular weight. The length of proteins encoded by *EuNAC* genes ranged from 86 (*EuNAC50*) to 617 (*EuNAC64*) amino acids (aa), and the molecular weight varied from 9.81 (*EuNAC50*) to 70.46 kDa (*EuNAC64*); the isoelectric points (pIs) ranged from 4.51 (*EuNAC59*) to 10.01 (*EuNAC2*). This study also analysed other basic information about 69 *NAC* genes, including homologous genes in *Arabidopsis*, open reading frame (ORF) length, location coordinates, chromosomal positions, and exon numbers (Table S1).

Phylogenetic analysis and classification of *EuNAC* proteins

To investigate the evolutionary relationships among *EuNAC* family genes, MEGA6.0 software was employed to construct a neighbour-joining phylogenetic tree based on full-length protein sequences of *NAC* in *E. ulmoides* and *Arabidopsis* (Figure 1). According to the classification system for *Arabidopsis*, *NACs* from *E. ulmoides* were divided into 13 distinct subfamilies, namely *NAC2*, *ONAC022*, *AtNAC3*, *NAP*, *ANAC063*, *ANAC011*, *ONAC003*, *NAM*, *NAC1*, *OSNAC7*, *OSNAC8*, *TIP*, and *ANAC011*, using a phylogenetic tree; however, no *EuNAC* members were identified in the *ATAF* subfamily. Of these 13 subfamilies, *OSNAC7* had 11 *EuNAC*

members, which was the most abundant, followed by *NAM*, which had eight members. *ANAC001* was the least frequent with only two members. Phylogenetic analysis revealed that *EuNAC* proteins had evolved in some diversity, similar to a report in *Arabidopsis* (Ooka et al. 2003a).

Conserved motif and gene structure analysis of *EuNAC* genes

To gain insight into the functional regions of *EuNAC*s, the conserved motifs for each *EuNAC* protein were analysed using the MEME programme. A total of 10 conserved motifs were identified and named motifs 1–10 (Table S2). These conserved motifs had large variations in length, with a distribution range of 11–50 amino acid residues. As shown in Figure 2A, motif 3 had the highest frequency in the *EuNAC* family, and it existed in almost all members except *EuNAC2*, *19*, and *33*. In addition, motifs 1, 2, 4, 5, and 6 were very abundant in the *EuNAC* family, but none of the *ONAC003* subfamily members had these motifs. Most of the conserved motifs were distributed in the N-terminus of the *NAC* proteins, indicating that the N-terminal region plays an important role in *NAC* gene function. In addition, similar motif compositions existed among different members of the same subfamily, indicating that members of the same subfamily had similar functions.

To investigate the structural features of *EuNAC*s, we analysed the intron/exon distribution patterns of each *EuNAC* gene. The exon distribution within the *EuNAC* genes varied from 2 to 8 (Table S1, Figure 2B). Forty-five (65.2%) genes had three exons, 10 (14.5%) genes had six exons, and *EuNAC45* had the largest number of exons, with 8 exons (Figure 2B, Table S1). Forty-nine (71.0%) *EuNAC* genes possessed less than three exons, indicating a low structural diversity among *EuNAC* genes.

Genome distribution, selective pressure, and synteny analysis of *EuNAC* genes

To investigate the distribution of *EuNAC* genes on the chromosomes of *E. ulmoides*, TBtools software was employed to map the chromosome locations for all *EuNAC*s identified in this study (Figure 3). The 69 *EuNAC*s were unevenly scattered on 16 chromosomes and two scaffolds, and the length of each chromosome showed no correlation with the number of genes contained. Chromosomes 8 and 12 had the most *EuNAC*s, both with seven genes. Only one *EuNAC* was distributed on chromosomes 3 and 16, and no *EuNAC*s were distributed on chromosome 11. Notably, most *EuNAC*s were distributed near the ends of the chromosome.

Furthermore, the duplication events of *EuNAC* gene family members were examined using MCScanX software. A total of 12 pairs of segmental duplications were identified in the *EuNAC* family, which were distributed across 15 chromosomes, except for chromosomes 3 and 11, and four pairs of tandem replications (*EuNAC7/8*, *EuNAC30/31*, *EuNAC48/49*, and *EuNAC53/54*) were identified, which were distributed on chromosomes 2, 7, 12, and 13, respectively (Figure 4, Table S3). The results showed that segmental duplication events might be the crucial driving force in *EuNAC* gene family expansion. To evaluate the selection pressure of *EuNAC*s, the *Ka*, *Ks*, and *Ka/Ks* for duplicated gene pairs were calculated. In general, a *Ka/Ks* > 1 indicates positive selection, *Ka/Ks* = 1 indicates neutral selection, while *Ka/Ks* < 1 indicates purifying selection (Vahdati & Lotfi 2013). The *Ka/Ks* ratios of 16 replicated *EuNAC* gene pairs were all less than 1, indicating that the evolution of the *EuNAC* gene family was subjected to purification selection (Table S4).

To further understand the phylogenetic mechanisms of the *EuNAC* gene family, we constructed a comparative homologous map of *NAC* genes in *E. ulmoides*, *Arabidopsis*, and rice. In total, 31 *EuNACs* had a collinear relationship with 27 *AtNACs* and 10 *OsNACs*. Thirty-one and eleven pairs of *NAC* homologous gene pairs were formed between *E. ulmoides* and *Arabidopsis* and between *E. ulmoides* and rice, respectively (Figure 5, Table S5). The results indicate that the *NAC* genes underwent significant evolution and replication after differentiation in monocotyledonous and dicotyledonous plants.

Promoter region analysis of *EuNAC* genes

To investigate the potential functions of *EuNACs*, we employed PlantCARE to predict the cis-acting elements within the 2.0 kb sequence upstream of the initiation codon (ATG) of *EuNACs* (Table S6). As expected, both TATA and CAAT boxes with good characteristics were found in the results; we also found several other CIS regulators (Table S7 and Figure S1). They were mined in the promoter region of *EuNACs*. As shown in Figure 6, we divided the homeopathic elements into four categories according to their functions. The first category was phytohormone-responsive elements, such as CGTCA-motif, TGACG, TCA-element, and ABRE, wherein ABREs have been associated with ABA responses and TCA-element have been associated with salicylic acid responsiveness. The second category was elements of cis-regulation related to the response to external or environmental pressure. This category includes low-temperature response elements (LTR), which respond to external abiotic stress, abundant cis-regulatory elements (AREs), which are required for anaerobic induction, and MYB binding site (MBS) elements. Notably, 29 of the 69 *EuNAC* promoters contained MBS elements that were involved in drought induction as MYB binding sites and could be predicted based on their responses to drought stress treatments. The third category included light-responsive elements, such as G-box, Box-4, and GT1-motif, in which at least one photo-responsive element was detected in almost every promoter region of *EuNACs*. The last category was cis-regulatory elements related to growth and development. CAT-box and O2-site were mainly detected, which also indicated that most *EuNACs* may be involved in *E. ulmoides* meristem expression, zein metabolism regulation, and cell cycle regulation. Finally, based on the above results, *EuNACs* may be involved in stress response, light, hormones, and growth pathways.

Expression analysis of *EuNAC* genes under drought stress

To further investigate the potential function of *EuNACs* in response to drought stress, comparative transcriptomics of *E. ulmoides* under drought stress were used to analyse the expression patterns of *EuNACs*. Of the 69 *EuNAC* genes, 20 showed high expression levels with $FPKM \geq 20$, including *EuNAC1*, 2, 12, 15, 20, and 25. Forty-three *EuNACs* showed low expression levels, with $FPKM \leq 20$, including *EuNAC10*, 11, 13, 14, 16, 17, and 18. In addition, *EuNAC19*, 47, 52, 54, 55, and 58 were not expressed in *E. ulmoides* leaves (Table S8). Differential gene expression (DEG) analysis showed that 12 *EuNAC* genes were significantly differentially expressed, of which *EuNAC2*, 3, 11, and 14 were upregulated after drought stress treatment, and *EuNAC1*, 36, 37, 64, and 66 were downregulated. The expression levels of *EuNAC8* and 13 first decreased and then increased with prolonged treatment time; in contrast, the expression level of

EuNAC61 first increased and then decreased (Figure 7, Table S8). The variable expression patterns of *EuNACs* may indicate their differential roles in the drought stress response of *E. ulmoides*. In particular, differentially expressed *EuNACs* may play a crucial role in *E. ulmoides*' response to drought stress.

Discussion

The NAC transcription factor family is one of the largest gene families in plants. These factors are involved in regulating hormone signalling pathways, biotic and abiotic stress responses, and plant growth and development (Yuan et al. 2020; Zhang et al. 2019). Several plant genomes have defined the *NAC* gene family. The evolutionary connection and duplication patterns of the *NAC* gene family in *E. ulmoides* can be better understood thanks to the genome of this organism. Previous studies have shown that genes with close evolutionary relationships often share similar functions (Lynch & Conery 2000b). Therefore, by studying the evolutionary relationships across gene families, we can learn more about and perhaps even anticipate how genes operate (Balazadeh et al. 2011; Zhang et al. 2019).

According to our study, the genome of *E. ulmoides* included 69 *NAC* genes, which is fewer than that of *A. thaliana* (117 *NAC* genes) (Ooka et al. 2003b), but similar to *K. obovate* (79 *NAC* genes) (Du et al. 2022a). Our results indicate that the majority of *EuNACs* did not experience environmental selection-induced elimination, but rather demonstrated a high level of conservation throughout evolution, underlining the necessity for more research from an evolutionary standpoint. All 69 *NAC* proteins were divided into 13 subgroups based on their sequence homology and classification relative to *Arabidopsis* (Ooka et al. 2003b). *NACs* in *Arabidopsis* exhibit a high degree of similarity among members of the same class or *NAC* subgroup. Four *EuNACs* in the *ANAC011* subgroup are orthologous to *Arabidopsis* genes, including *AtNAC071* and *AtNAC096*, which are in charge of tissue reunification, dehydration, and other processes.

Our research indicates that the *NAM* subgroup has 8 *EuNACs* that are orthologous to *AtNAC054* and *AtNAC059* in *Arabidopsis*, which are known to be crucial for organ development, programmed cell death, secondary wall building, and biotic and abiotic stress responses (Kim et al. 2007). The 7 *EuNACs* in subgroup *NAP* are orthologous to *AtNAC018*, *AtNAC025*, and *AtNAC56* and are essential for leaf senescence (Guo & Gan 2006). Three genes of the *EuNAC* gene family's subgroup *TIP* are orthologs of the *Arabidopsis* genes *AtNAC060* and *AtNAC091*. These orthologous genes have been demonstrated to be crucial in the stress response and abscisic acid (ABA) signalling (Donze et al. 2014; Jeong et al. 2008; Li et al. 2014). Four *EuNACs* in the *ANAC011* subgroup are orthologous to *Arabidopsis* genes, including *AtNAC071* and *AtNAC096*, which are in charge of tissue reunification, dehydration, and osmotic stress (Asahina et al. 2011; Yang et al. 2020). Five orthologous *EuNACs* to *AtNAC016*, which are known to be involved in chlorophyll degradation, are found in the *NAC2* subgroup. This indicates that this subgroup of *EuNACs* may also control how chlorophyll degrades in plants (Sakuraba et al. 2015). Similar to *AtNAC003* and *AtNAC068*, the *ANAC001* subgroup also has two *EuNACs* that are orthologous to

them. These genes control salt and osmotic stress tolerance, in addition to DNA damage responses (Xu et al. 2013; Yoshiyama et al. 2014).

Except for individuals from the *ONAC003* and *ANAC063* subgroups in this investigation, the N-terminus of the EuNAC protein had motifs 1, 2, 3, 4, 5, and 6. *ONAC7* comprised motifs 7 and 10, indicating that NAC transcription factors had a highly conserved N-terminus and a very varied C-terminus. The range of *EuNAC* introns was 2 to 8, which is comparable to the number observed in many plants (Du et al. 2022a; Liu et al. 2019). In general, the deletion or insertion of introns can have diverse effects on gene function. Sometimes, intron deletions can lead to the loss of gene function if they result in a frameshift mutation or the removal of critical regulatory sequences. Our analysis supported the findings of Jeffares' research, which demonstrated that genes susceptible to abrupt changes in stress expression levels have much less intron density (Jeffares et al. 2008); *EuNACs* with fewer introns merit higher consideration if the study objective is to concentrate on genes that react instantly to environmental stress.

As a consequence of our findings, which included the identification of 12 segmental duplications and four tandem duplications in 69 *EuNACs*, we concluded that segmental duplication served as the primary catalyst for the growth of the *EuNAC* gene family, in agreement with research on *K. obovata* (Du et al. 2022a). In addition, larger segmental duplications account for most of the *A. thaliana* genome, and at least four large-scale replication events occurred during the formation of angiosperm diversity (100–200 million years ago) (Vision et al. 2000). This might explain why there are more *NAC* members in *Arabidopsis*, while having a smaller genome than *E. ulmoides*. Only 10 genes, according to our research, have collinear connections between *O. sativa* and *E. ulmoides*. However, we discovered 27 orthologous pairings in *Arabidopsis*, a dicotyledonous plant. These findings demonstrate a closer evolutionary link between dicotyledons and *EuNACs* than between monocotyledons.

Cis-acting elements are specific DNA sequences located in the promoter region of genes that serve as binding sites for transcription factors (Liu et al. 2016; Kaur et al. 2017). In this study, more than half of the 69 *EuNAC* promoters included ABRE homeopathic elements, suggesting that these genes may operate via the control of ABA. MBS elements were found in 29 *EuNACs*, indicating that these genes may be crucial in response to drought stress.

Research on gene function can benefit from understanding the patterns of gene expression. According to RNA-seq studies, drought stress drastically altered the expression levels of a few *EuNACs* in the leaves of *E. ulmoides*. Significant differential expression was seen in 12 *EuNACs*. *EuNAC1*, *EuNAC8*, and *EuNAC36* are identical to *ANAC019* (*AT1G52890*), *ANAC055* (*AT3G15500*), and *ANAC072* (*AT4G27410*), which are members of the *AtNAC3* subgroup. Their expression is variably expressed during drought treatment, caused by drought, and stimulated by ABA (Tran et al. 2004). Therefore, we hypothesise that genes *EuNAC1*, *EuNAC8*, and *EuNAC36* belong to the same subgroup and are drought-responsive genes that control *E. ulmoides*' survival ability in drought-stressed environments. Additionally, *ANAC054* (*At3g15170*) and *ANAC059* (*At3g29035*) were identical to *EuNAC3*, *EuNAC11*, and *EuNAC61* and were grouped into the *NAM* subgroup, suggesting that they may be crucial in *E. ulmoides*' response to drought stress.

Conclusions

In summary, 69 *NACs* were identified in *E. ulmoides* in this study. We studied the characteristics of *EuNAC* genes at the genomic level and analysed tissue expression patterns and responses to drought stress. These TFs can be divided into 13 subgroups according to the *NAC* classification method of *Arabidopsis*. Chromosomal localisation and homology analysis showed that segmental duplication was the main driving force for *EuNAC* gene amplification. Genome-wide expression analysis of *NAC* genes in response to drought provides an opportunity to further understand the strong tolerance mechanism of *E. ulmoides* to drought.

Acknowledgements

The authors would like to thank YongJie Yang, GuoTao Song and YiXue Sun for their help in our experiment.

References

- Asahina M, Azuma K, Pitaksaringkarn W, Yamazaki T, Mitsuda N, Ohme-Takagi M, Yamaguchi S, Kamiya Y, Okada K, Nishimura T, Koshiba T, Yokota T, Kamada H, and Satoh S. 2011. Spatially selective hormonal control of RAP2.6L and ANAC071 transcription factors involved in tissue reunion in *Arabidopsis*. *Proceedings of the National Academy of Sciences* 108:16128-16132.
- Balazadeh S, Kwasniewski M, Caldana C, Mehrnia M, Zanol MI, Xue GP, and Mueller-Roeber B. 2011. ORS1, an H₂O₂-responsive NAC transcription factor, controls senescence in *Arabidopsis thaliana*. *Mol Plant* 4:346-360.
- Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, and Xia R. 2020. TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Molecular plant* 13:1194-1202.
- Deng P, Wang Y, Hu F, Yu H, Liang Y, Zhang H, Wang T, Zhou Y, and Li Z. 2022. Phenotypic Trait Subdivision Provides New Sight Into the Directional Improvement of *Eucommia ulmoides* Oliver. *Frontiers in Plant Science* 13.
- Donze T, Qu F, Twigg P, and Morris TJ. 2014. Turnip crinkle virus coat protein inhibits the basal immune response to virus invasion in *Arabidopsis* by binding to the NAC transcription factor TIP. *Virology* 449:207-214.
- Du Z, You S, Yang D, Tao Y, Zhu Y, Sun W, Chen Z, and Li J. 2022a. Comprehensive analysis of the NAC transcription factor gene family in *Kandelia obovata* reveals potential members related to chilling tolerance. *Front Plant Sci* 13:1048822.
- Du Z, You S, Yang D, Tao Y, Zhu Y, Sun W, Chen Z, and Li J. 2022b. Comprehensive analysis of the NAC transcription factor gene family in *Kandelia obovata* reveals potential members related to chilling tolerance. *Frontiers in Plant Science* 13:4735.
- Enoki M, Doi Y, and Iwata T. 2003. Oxidative degradation of cis-and trans-1, 4-polyisoprenes and vulcanized natural rubber with enzyme-mediator systems. *Biomacromolecules* 4:314-320.
- Guo Y, and Gan S. 2006. AtNAP, a NAC family transcription factor, has an important role in leaf senescence. *Plant J* 46:601-612.

- Hall BG. 2013. Building phylogenetic trees from molecular data with MEGA. *Molecular biology and evolution* 30:1229-1235.
- Hussain RM, Ali M, Feng X, and Li X. 2017. The essence of NAC gene family to the cultivation of drought-resistant soybean (*Glycine max* L. Merr.) cultivars. *BMC plant biology* 17:1-11.
- Jeffares DC, Penkett CJ, and Bähler J. 2008. Rapidly regulated genes are intron poor. *Trends Genet* 24:375-378.
- Jeong R-D, Chandra-Shekara A, Kachroo A, Klessig D, and Kachroo P. 2008. HRT-Mediated Hypersensitive Response and Resistance to Turnip crinkle virus in *Arabidopsis* Does Not Require the Function of *TIP*, the Presumed Guardee Protein. *Molecular plant-microbe interactions : MPMI* 21:1316-1324.
- Kaur A, Pati PK, Pati AM, and Nagpal AK. 2017. In-silico analysis of cis-acting regulatory elements of pathogenesis-related proteins of *Arabidopsis thaliana* and *Oryza sativa*. *PLoS One* 12:e0184523.
- Kent E, and Swinney F. 1966. Properties and applications of trans-1, 4-polyisoprene. *Industrial & Engineering Chemistry Product Research and Development* 5:134-138.
- Kim SG, Kim SY, and Park CM. 2007. A membrane-associated NAC transcription factor regulates salt-responsive flowering via FLOWERING LOCUS T in *Arabidopsis*. *Planta* 226:647-654.
- Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, and Marra MA. 2009. Circos: an information aesthetic for comparative genomics. *Genome research* 19:1639-1645.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, and Lopez R. 2007. Clustal W and Clustal X version 2.0. *bioinformatics* 23:2947-2948.
- Li P, Zhou H, Shi X, Yu B, Zhou Y, Chen S, Wang Y, Peng Y, Meyer RC, Smeekens S, and Teng S. 2014. The ABI4-Induced *Arabidopsis ANAC060* Transcription Factor Attenuates ABA Signaling and Renders Seedlings Sugar Insensitive when Present in the Nucleus. *PLoS Genetics* 10.
- Li Y, Wei H, Yang J, Du K, Li J, Zhang Y, Qiu T, Liu Z, Ren Y, and Song L. 2020. High-quality de novo assembly of the *Eucommia ulmoides* haploid genome provides new insights into evolution and rubber biosynthesis. *Horticulture Research* 7.
- Liu J, Wang X, Chen Y, Liu Y, Wu Y, Ren S, and Li L. 2021. Identification, evolution and expression analysis of WRKY gene family in *Eucommia ulmoides*. *Genomics* 113:3294-3309.
- Liu M, Ma Z, Sun W, Huang L, Wu Q, Tang Z, Bu T, Li C, and Chen H. 2019. Genome-wide analysis of the NAC transcription factor family in Tartary buckwheat (*Fagopyrum tataricum*). *BMC Genomics* 20:113.
- Liu Y, Sun J, and Wu Y. 2016. *Arabidopsis ATAF1* enhances the tolerance to salt stress and ABA in transgenic rice. *J Plant Res* 129:955-962. 10.1007/s10265-016-0833-0
- Lynch M, and Conery JS. 2000a. The evolutionary fate and consequences of duplicate genes. *Science* 290:1151-1155.
- Lynch M, and Conery JS. 2000b. The evolutionary fate and consequences of duplicate genes. *Science* 290:1151-1155.

- Ma N, Feng H, Meng X, Li D, Yang D, Wu C, and Meng Q. 2014. Overexpression of tomato SINAC1 transcription factor alters fruit pigmentation and softening. *BMC plant biology* 14:1-14.
- Mao H, Li S, Chen B, Jian C, Mei F, Zhang Y, Li F, Chen N, Li T, and Du L. 2022. Variation in cis-regulation of a NAC transcription factor contributes to drought tolerance in wheat. *Molecular plant* 15:276-292.
- Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N, Kawai J, and Carninci P. 2003a. Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*. *DNA research* 10:239-247.
- Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N, Kawai J, Carninci P, Hayashizaki Y, Suzuki K, Kojima K, Takahara Y, Yamamoto K, and Kikuchi S. 2003b. Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*. *DNA Res* 10:239-247.
- Perochon A, Kahla A, Vranić M, Jia J, Malla KB, Craze M, Wallington E, and Doohan FM. 2019. A wheat NAC interacts with an orphan protein and enhances resistance to Fusarium head blight disease. *Plant biotechnology journal* 17:1892-1904.
- Rose K, and Steinbuchel A. 2005. Biodegradation of natural rubber and related compounds: recent insights into a hardly understood catabolic capability of microorganisms. *Applied and environmental microbiology* 71:2803-2812.
- Sakuraba Y, Piao W, Lim JH, Han SH, Kim YS, An G, and Paek NC. 2015. Rice *ONAC106* Inhibits Leaf Senescence and Increases Salt Tolerance and Tiller Angle. *Plant Cell Physiol* 56:2325-2339. 10.1093/pcp/pcv144
- Shang X, Yu Y, Zhu L, Liu H, Chai Q, and Guo W. 2020. A cotton NAC transcription factor GhirNAC2 plays positive roles in drought tolerance via regulating ABA biosynthesis. *Plant Science* 296:110498.
- Shao H, Wang H, and Tang X. 2015. NAC transcription factors in plant multiple abiotic stress responses: progress and prospects. *Frontiers in Plant Science* 6:902.
- Tang C, Yang M, Fang Y, Luo Y, Gao S, Xiao X, An Z, Zhou B, Zhang B, and Tan X. 2016. The rubber tree genome reveals new insights into rubber production and species adaptation. *Nature plants* 2:1-10.
- Tran LS, Nakashima K, Sakuma Y, Simpson SD, Fujita Y, Maruyama K, Fujita M, Seki M, Shinozaki K, and Yamaguchi-Shinozaki K. 2004. Isolation and functional analysis of *Arabidopsis* stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. *Plant Cell* 16:2481-2498.
- Vahdati K, and Lotfi N. 2013. Abiotic stress tolerance in plants with emphasizing on drought and salinity stresses in walnut. *Abiotic stress—Plant responses and applications in agriculture* 10:307-365.
- Vision TJ, Brown DG, and Tanksley SD. 2000. The origins of genomic duplications in *Arabidopsis*. *Science* 290:2114-2117.
- Wang B, Wei J, Song N, Wang N, Zhao J, and Kang Z. 2018. A novel wheat NAC transcription factor, TaNAC30, negatively regulates resistance of wheat to stripe rust. *Journal of integrative plant biology* 60:432-443.
- Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee T-h, Jin H, Marler B, and Guo H. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Research* 40:e49-e49.

- Wei X, Peng P, Peng F, and Dong J. 2021. Natural polymer *Eucommia ulmoides* rubber: A novel material. *Journal of agricultural and food chemistry* 69:3797-3821.
- Wuyun T-n, Wang L, Liu H, Wang X, Zhang L, Bennetzen JL, Li T, Yang L, Liu P, and Du L. 2018. The hardy rubber tree genome provides insights into the evolution of polyisoprene biosynthesis. *Molecular plant* 11:429-442.
- Xu B, Ohtani M, Yamaguchi M, Toyooka K, Wakazaki M, Sato M, Kubo M, Nakano Y, Sano R, and Hiwatashi Y. 2014. Contribution of NAC transcription factors to plant adaptation to land. *Science* 343:1505-1508.
- Xu Z, Wang C, Xue F, Zhang H, and Ji W. 2015. Wheat NAC transcription factor TaNAC29 is involved in response to salt stress. *Plant Physiology and Biochemistry* 96:356-363.
- Xu ZY, Kim SY, Hyeon do Y, Kim DH, Dong T, Park Y, Jin JB, Joo SH, Kim SK, Hong JC, Hwang D, and Hwang I. 2013. The Arabidopsis NAC transcription factor ANAC096 cooperates with bZIP-type transcription factors in dehydration and osmotic stress responses. *Plant Cell* 25:4708-4724.
- Yan R. 1995. Prospects and research progress on *Eucommia ulmoides* gum. *Progress in Chemistry* 7:65.
- Yang JH, Lee KH, Du Q, Yang S, Yuan B, Qi L, and Wang H. 2020. A membrane-associated NAC domain transcription factor XVP interacts with TDIF co-receptor and regulates vascular meristem activity. *New Phytol* 226:59-74.
- Yoshiyama KO, Kimura S, Maki H, Britt AB, and Umeda M. 2014. The role of SOG1, a plant-specific transcriptional regulator, in the DNA damage response. *Plant Signal Behav* 9:e28889.
- Yuan C, Li C, Lu X, Zhao X, Yan C, Wang J, Sun Q, and Shan S. 2020. Comprehensive genomic characterization of NAC transcription factor family and their response to salt and drought stress in peanut. *BMC plant biology* 20:1-21.
- Zhang X, Cheng Z, Zhao K, Yao W, Sun X, Jiang T, and Zhou B. 2019. Functional characterization of poplar NAC13 gene in salt tolerance. *Plant Science* 281:1-8.
- Zhu M-Q, and Sun R-C. 2018. *Eucommia ulmoides* Oliver: a potential feedstock for bioactive products. *Journal of agricultural and food chemistry* 66:5433-5438.
- Zuo Y, Li B, Guan S, Jia J, Xu X, Zhang Z, Lu Z, Li X, and Pang X. 2022. EuRBG10 involved in indole alkaloids biosynthesis in *Eucommia ulmoides* induced by drought and salt stresses. *Journal of Plant Physiology* 278:153813.

Figures captions

Figure 1. Phylogenetic relationships among *NACs* identified in *E. ulmoides* and *Arabidopsis thaliana*. The unrooted phylogenetic tree was constructed by MEGA 6.0 software using the Neighbor-Joining (NJ) method with 1,000 bootstrap iterations. Each subfamily was distinguished by different colors.

Figure 2. Motif compositions and DNA structures of *NAC* gene family in *E. ulmoides*. A. The conserved motif distribution of *EuNAC* proteins. Different motifs were distinguished by different colored boxes, and black lines represent non-conserved regions. B. Gene structure of the *EuNAC* gene. Green boxes represent non-coding regions, yellow boxes represent exons, and black lines represent introns.

493 Figure 3. Distribution of 69 *EuNAC*s on 16 chromosomes and two scaffolds. Vertical bars represent the
 494 chromosomes of *E. ulmoides*. The chromosome number is on the left of each chromosome. The
 495 scale on the left represents the length of the chromosome.

496 Figure 4. Schematic representations of the interchromosomal relationships of *EuNAC* genes. The deep red
 497 line represents the *EuNAC* gene pairs replicated in tandem, while the remaining colored lines
 498 represent the *EuNAC* gene pairs replicated in segments.

499 Figure 5. Synteny analysis of *NAC* genes between *E. ulmoides* and two representative plant species
 500 (*Arabidopsis thaliana* and *Oryza sativa*). Green and purple lines represent syntenic *NAC* gene
 501 pairs of *E. ulmoides* and *A. thaliana* and *O. sativa*, respectively.

502 Figure 6. The number of each type of cis-acting element in the promoter region of each *EuNAC* gene.

503 Figure 7. Expression levels of 69 *NAC* genes under drought stress in *E. ulmoides* of leaves. The expression
 504 level was presented based on the transformed data of \log_2 (FPKM+1) values.

Figure 1

Phylogenetic relationships among *NACs* identified in *E. ulmoides* and *Arabidopsis thaliana*.

The unrooted phylogenetic tree was constructed by MEGA 6.0 software using the Neighbor-Joining (NJ) method with 1,000 bootstrap iterations. Each subfamily was distinguished by different colors.




Figure 2

Motif compositions and DNA structures of *NAC* gene family in *E. ulmoides*.

A. The conserved motif distribution of EuNAC proteins. Different motifs were distinguished by different colored boxes, and black lines represent non-conserved regions. B. Gene structure of the *EuNAC* gene. Green boxes represent non-coding regions, yellow boxes represent exons, and black lines represent introns.



Figure 3

Distribution of 69 *EuNACs* on 16 chromosomes and two scaffolds.

Vertical bars represent the chromosomes of *E. ulmoides*. The chromosome number is on the left of each chromosome. The scale on the left represents the length of the chromosome.




Figure 4

Schematic representations of the interchromosomal relationships of *EuNAC* genes.

The deep red line represents the *EuNAC* gene pairs replicated in tandem, while the remaining colored lines represent the *EuNAC* gene pairs replicated in segments.




Figure 5

Synteny analysis of *NAC* genes between *E. ulmoides* and two representative plant species (*Arabidopsis thaliana* and *Oryza sativa*)

Green and purple lines represent syntenic *NAC* gene pairs of *E. ulmoides* and *A. thaliana* and *O. sativa*, respectively.




Figure 6

The number of each type of cis-acting element in the promoter region of each *EuNAC* gene.




Figure 7

Expression levels of 69 *NAC* genes under drought stress in *E. ulmoides* of leaves

The expression level was presented based on the transformed data of \log_2 (FPKM+1) values.

