

Competitive effects of the macroalga *Caulerpa taxifolia* on key physiological processes in the scleractinian coral *Turbinaria peltata* under thermal stress (#85054)

1

Second revision

Guidance from your Editor

Please submit by **17 Oct 2023** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)
1 Rebuttal letter(s)
5 Figure file(s)
2 Table file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Competitive effects of the macroalga *Caulerpa taxifolia* on key physiological processes in the scleractinian coral *Turbinaria peltata* under thermal stress

JianRong Fu¹, Jie Zhou¹, JiaLi Zhou¹, YanPing Zhang¹, Li Liu^{Corresp. 1}

¹ Fisheries College, Guangdong Ocean University, Zhanjiang, Guangdong, China

Corresponding Author: Li Liu

Email address: zjoulili@163.com

An increased abundance of macroalgae has been observed in coral reefs damaged by climate change and local environmental stressors. Macroalgae have a sublethal effect on corals that includes the inhibition of their growth, development, and reproduction. Thus, this study explored the effects of the macroalga, *Caulerpa taxifolia*, on the massive coral, *Turbinaria peltata*, under thermal stress. We compared the responses of the corals' water-mediated interaction with algae (the co-occurrence group) and those in direct contact with algae at two temperatures. The results show that after co-culturing with *C. taxifolia* for 28 days, the density and chlorophyll *a* content of the dinoflagellate endosymbionts was not influenced by the presence of *C. taxifolia* at ambient temperature (27 °C). The protein content of *T. peltata* decreased by 37.2% in the co-occurrence group and 49.0% in the direct contact group compared to the control group. Meanwhile, the growth rate of *T. peltata* decreased by 57.7% in the co-occurrence group and 65.5% in the direct contact group compared to the control group. The activity of the antioxidant enzymes significantly increased, and there was a stronger effect of direct coral contact with *C. taxifolia* than the co-occurrence group. At 30 °C, the endosymbiont density, chlorophyll *a* content, and growth rate of *T. peltata* significantly decreased compared to the control temperature; the same pattern was seen in the increase in antioxidant enzyme activity. Additionally, when the coral was co-cultured with macroalgae at 30 °C, there was no significant decrease in the density or chlorophyll *a* content of the endosymbiont compared to the control. However, the interaction of macroalgae and elevated temperature was evident in the feeding rate, protein content, superoxide dismutase (SOD), and catalase (CAT) activity compared to the control group. The direct contact of the coral with macroalga had a greater impact than water-mediated interactions. Hence, the competition between coral and macroalga may be more intense under thermal stress.

Competitive effects of the macroalga *Caulerpa taxifolia* on key physiological processes in the scleractinian coral *Turbinaria peltata* under thermal stress

Jian Rong Fu¹, Jie Zhou¹, Jia Li Zhou¹, Yan Ping Zhang¹, Li Liu¹

¹ Fisheries College, Guangdong Ocean University, Zhanjiang, China, 524088;

Corresponding author:

Li Liu¹

Haida road, Zhanjiang, Guangdong Province, China

Email address: corresponding_Li Liu_zjouliuli@163.com

Abstract

An increased abundance of macroalgae has been observed in coral reefs damaged by climate change and local environmental stressors. Macroalgae have a sublethal effect on corals that includes the inhibition of their growth, development, and reproduction. Thus, this study explored the effects of the macroalga, *Caulerpa taxifolia*, on the massive coral, *Turbinaria peltata*, under thermal stress. We compared the responses of the corals' water-mediated interaction with algae (the co-occurrence group) and those in direct contact with algae at two temperatures. The results show that after co-culturing with *C. taxifolia* for 28 days, the density and chlorophyll *a* content of the dinoflagellate endosymbionts was not influenced by the presence of *C. taxifolia* at ambient temperature (27 °C). The protein content of *T. peltata* decreased by 37.2% in the co-occurrence group and 49.0% in the direct contact group compared to the control group. Meanwhile, the growth rate of *T. peltata* decreased by 57.7% in the co-occurrence group and 65.5% in the direct contact group compared to the control group. The activity of the antioxidant enzymes significantly increased, and there was a stronger effect of direct coral contact with *C. taxifolia* than the co-occurrence group. At 30 °C, the endosymbiont density, chlorophyll *a* content, and growth rate of *T. peltata* significantly decreased compared to the control temperature; the same pattern was seen in the increase in antioxidant enzyme activity. Additionally, when the coral was co-cultured with macroalgae at 30 °C, there was no significant decrease in the density or chlorophyll *a* content of the endosymbiont compared to the control. However, the interaction of macroalgae and elevated

temperature was evident in the feeding rate, protein content, superoxide dismutase (SOD), and catalase (CAT) activity compared to the control group. The direct contact of the coral with macroalga had a greater impact than water-mediated interactions. Hence, the competition between coral and macroalga may be more intense under thermal stress.

Introduction

The recent effects of climate change and other anthropogenic impacts have caused the severe degradation of coral reefs worldwide (Leggat et al., 2022). The first mass coral bleaching event was observed in 1998 and it killed approximately 8% of the world's coral; an additional 14% of corals were lost between 2009 and 2018 (Souter 2021).

Many studies have asserted that ocean warming is a major factor in the reduction of coral cover (Hughes et al., 2017, 2019; Lough et al., 2018; Leggat et al., 2022). For example, the successive bleaching events in 2016-2017 devastated Australia's Great Barrier Reef and resulted in an 89% decline in larval recruitment in 2018 compared to historical levels (Hughes et al., 2017,2019; Lough et al., 2018). During this period, 31% of reefs experienced 8–16 degree heating weeks (DHWs, °C-weeks). A decline in coral cover may lead to an increase in the cover of other benthic organisms in the reefs, such as macroalgae (Fulton et al., 2019). The impact of herbivorous fishes is mostly ignored, although some studies assert that overfishing and nutrient pollution is the main cause of phase shifts towards macroalgae (Barott et al., 2012). Research had shown that prior to 2011, the estimated global average cover of algae ~~at the global scale~~ was low (~16%) and stable for 30 years. Since 2011, the amount of algae on the world's coral reefs has increased by about 20% (Souter, 2021). Thus, the coral reef ecosystem is undergoing an ecological phase transition to that of an ecosystem dominated by macroalgae.

Macroalgae are functional communities that are important for stabilizing reef structure (Fulton et al., 2019), generating primary productivity (Fulton et al., 2014; 2019), maintaining nutrient cycling in reef areas, and providing food sources for herbivores (Dubinsky and Stambler, 2011). However, there is competition between macroalgae and corals. Macroalgae harm corals through direct contact (Coyer et al., 1993; Manikandan et al., 2021) and allelopathy (Roberta et al., 2014; Fong et al., 2020), weakening the photosynthetic performance of symbiodiniaceae (Rasher et al., 2011), causing the retraction of polyps (Jompa and Mccook, 2003), increasing the number of pathogenic microorganisms (Clements et al., 2020; Rasher and Hay, 2010), triggering coral bleaching (Roberta et al., 2014), and resulting in the reduced calcification and coral growth,

fecundity, survival rate, and settlement rate (Fong et al., 2020; Jason and Tanner, 1995; Leong et al., 2018; Rasher and Hay, 2010). Specifically, macroalgae affect coral feeding, endosymbiont function, tissue recovery, and oxidative stress response. Morrow and Carpenter (2008) found that *Dictyopteris undulata* weakened the particle capture rates of *Corynactis californica* by redirecting particles around polyps and causing contraction of the feeding tentacles. The dissolved organic carbon (DOC) and terpenoids released by macroalgae decreased photosynthesis and the density of endosymbionts (Rasher et al., 2011; Smith et al., 2006; Diaz-Pulido and Barrón, 2020). Bender (2012) asserted that the green filamentous macroalga, *Chlorodesmis fastigiata*, significantly reduced tissue recovery in *Acropora pulchra* and led to the infection of *A. pulchra* with ciliates. High levels of reactive oxygen species (ROS, a toxic byproduct of biological aerobic metabolism) could cause damage to cells (Blanckaert et al., 2021). Shearer et al. (2012) found that the oxidative stress response of *Acropora millepora* was activated in response to ROS by altering the transcription factors after contact with the macroalga *Chlorodesmis fastigiata* and its hydrophobic extract over a short-term period (1 h and 24 h). The oxidative imbalance results in rapid protein degradation and eventually to apoptosis and/or necrosis when compensatory transcriptional action by the coral holobiont insufficiently mitigates damage.

In addition, the combined effects of ocean warming, acidification, and macroalgae contact could significantly alter the physiological response of corals (Chadwick et al., 2011; Kornder et al., 2018; Brown et al., 2019; Rölfer et al., 2021). Rölfer et al. (2021) have shown that light enhanced calcification (LEC) rates of *Porites lobata* were negatively affected after contact with *Chlorodesmis fastigiata* in an ocean warming and acidification scenario, compared to coral under ambient conditions. Typically, the coral-algal competition is related to seasonal and temporal cycles. These, in turn, may be related to the abundance, biomass, and composition of macroalgae, as well as the seasonal dynamics of temperature, $p\text{CO}_2$, and light intensity (Brown et al., 2019; 2020). The sensitivity of various macroalgae to environmental stressors is also different. For example, intermediate levels of ocean warming could enhance the growth and production of *Laurencia* sp. and *Lobophora* sp., which was not the case for *Sargassum* sp. (Fulton et al., 2014; Hernández et al., 2018). Additionally, overfishing and eutrophication have been shown to lead to an increased growth rate of some kinds of macroalgae (Lapointe and Bedford, 2010), which may indirectly enhance the competitive ability of macroalgae. Therefore, to better understand the resilience of coral reef ecosystems in the future, it is necessary to determine how coral-algal interactions which are impacted by global and local stressors.

According to the China Ocean Climate Monitoring bulletin (www.oceanguide.org.cn), the average seawater temperature in the Xuwen Sea area was 27-30 °C from May to September in 2020. The DHW is an accumulation of instantaneous heat stress (coral bleaching hotspots) over a 12-week running window (Skirving et al., 2020). When compared with the 2021-2022 DHW from December to April in the Great Barrier Reef, the average seawater temperature is 28-30 °C (Spady et al., 2022). It is plausible that the physiological responses of corals in the Xuwen Coral Reef National Nature Reserve of China may be affected by thermal stress. During thermal stress, coral feeding rates are drastically reduced and more energy is needed for depleting protein to maintain biological processes (DNA repair etc.) to resist heat stress (Ferrier-Pagès et al., 2010; Chakravarti et al., 2020). Triggered by thermal stress, ROS may be produced by the endosymbionts mainly due to PS II dysfunction caused by damage to the D1 protein (Warner, 1999) and host cells (Nii and Muscatine, 1997). The increase in the production of ROS is a stress signaling mechanism that can potentially trigger an oxidative stress and apoptotic cascade in coral cells (Hensley et al., 2000; Drury et al., 2022).

The relationship between macroalgae and corals under climate change conditions remains inconclusive. To investigate the effects of macroalgae on hermatypic coral under ocean warming, the massive coral, *Turbinaria peltata*, and macroalga, *Caulerpa taxifolia*, which are common species with frequent interactions in the Xuwen Sea, were selected as study species. *C. taxifolia* is a multinucleate siphonous green alga and is known to have great invasive potential worldwide (Zubia et al., 2020). Furthermore, it has been found that *C. taxifolia* can produce potential allelochemicals, such as monoterpenes and sesquiterpenes (Guerriero et al., 1992, 1993). Given that *C. taxifolia* usually grows on various hard substrates close to large numbers of live coral colonies, the physical and chemical impact has to be better understood. To evaluate the effect of its chemical and physical effects, indirect contact group was used to investigate chemical effects, and the direct-contact group was used to explore the combined effects of physical and chemical processes. In this study, we show that increased temperature, representing the ocean-warming range projected for this century (Spady et al., 2022), enhances the ability of seaweeds to impact the physiology of corals, potentially shifting the competitive interaction between corals and seaweeds in favour of seaweeds. It provides a reliable basis for the evolution of competition between corals and macroalgae under future global changes.

Materials & Methods

Part of this text were previously published as part of a preprint

(<https://www.researchsquare.com/article/rs-1878684/v1>).

Sample collection

T. peltata (with a skeleton size of length:width:height = 20:18:12 cm, one colonies) and *C. taxifolia* (5 kg) were collected from the Xuwen Coral Reef National Nature Reserve (109° 55' E, 20° 16' N) at a depth of approximately 4 m. The samples were transported to the laboratory and cultured in two 200-L tanks at a temperature of 26.5 °C, pH of 8.0, salinity of 33, and 200 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ with a 12: 12 h light/dark cycle for 3 months. Blue lighting was provided by a photosynthetically active radiation (PAR) lamp (Aqua Knight M029). An illuminometer (UNI-T UT383) was used for light measurement. After 3 months of acclimation, the corals from the same colony were cut into 54 pieces, approximately 4 cm in diameter, and were fixed on a ceramic base with aqua rubber (Aron Alpha GEL-10). The samples were then placed in another 200-L tank for one week until the anthopolyp had stretched naturally.

Experimental design

After acclimation, 54 coral nubbins were randomly allocated into 10 L experimental tanks. To mimic the coral-macroalgal interaction in coral reefs, the same amount of macroalgae was co-cultured with the coral samples in the following ways: (1) no algae were added to the tank, i.e., the control group (Figure 1A). (2) Algae (25 g) were cultured in external algae boxes with no direct contact between the algae and the coral samples. The allelopathic substances produced by algae could enter the tank with the corals by water flow to determine whether the algae had a water-mediated interaction in the co-occurrence group (Figure 1B). (3) The algae and corals were co-cultured in the same tank with direct contact, and a panel made by polymethyl methacrylate (PMMA) was used to fix the height of the algae parallel to the coral samples; this was referred to as the direct contact group (Figure 1C). The experimental tanks were subjected to ambient conditions (27 °C) and the shared socioeconomic paths (SSPs) scenario SSP2-4.5 (30 °C) (Zhongming et al., 2021), to mimic the typical temperature range that the corals experienced at Xuwen Coral Reef Nature Reserve in order to explore the coral-macroalgae contacts at different temperatures. Each treatment contained three replicate tanks, within which three coral nubbins were placed per tank.

The temperature in each experimental group was increased to the set temperature by 1 °C per day. The first three days were the temperature adjustment period. Some algae tips decomposed due to metabolism or blue light intolerance. Therefore, the algae were checked and replaced each day to ensure that the experimental group had 25 g (0.0025 g cm^{-3}) of fresh algae, which is the

amount of algae with the density of 0.0022 g per cubic meter of water surveyed from the inshore reef of Xuwen Coral Reef Nature Reserve. Fifty percent of the seawater was replaced every three days in each tank. Organisms were kept under treatment conditions for a period of 28 days and physiological measurements were subsequently performed.

Endosymbiont density and Chl *a* content

At the end of the experiment, coral tissue were removed from the nubbins using a waterpick (0.45 µm filtered seawater), and the slurry was homogenized. Six 15 ml samples were taken from the slurry, centrifuged (4,000 rpm min⁻¹, 4 °C, 10 min), and the supernatant was removed. Part of the pellet was suspended in 5 mL formaldehyde to count the endosymbiont density under an inverted microscope (DMI 6001B, magnification eyepiece × magnification objective: 10×20) with a blood counting plate (CKSLAB CB30). Another portion was resuspended in 8 mL methanol. The pigments were extracted at 4 °C for 24 h. The extract was centrifuged (4,000 rpm min⁻¹, 4 °C, 10 min), and Chl *a* was determined according to the method described by Ritchie (2006). Data were normalized to skeletal surface with the aluminium foil technique (Marsh 1970).

Feeding rate

A total of 1ml of *Artemia* solution was added to 1 L of water and a plankton counter was used to measure the mixed solution and to determine the individuals of the *Artemia* solution per milliliter. Nubbins were moved into the feeding tanks (1 L) with an *Artemia* concentration of ~ 2 ind mL⁻¹, while one tank served as a control (without coral). After an incubation period of 1 h, the coral nubbins were rinsed with seawater and returned to their respective positions in the experimental tanks. The feeding rate was calculated as the decline in *Artemia* concentration in the feeding tanks and normalized per polyp. The measurement was performed once a week between 11:00-12:00 a.m. The number of polyps in each nubbins were visually counted before the experiment.

Growth rate

The coral nubbins were weighed on a balance (UW 2200H, accuracy = 0.01 g) using the buoyant weight technique (Davies, 1989). Before each measurement, the surface of the coral ceramic base was lightly brushed with a toothbrush to remove algae. A glass beaker was filled with 1 L of filtered seawater (27 °C, salinity 32). Then the coral nubbins were placed on the bottom of the beaker and the weight (minus the weight of the beaker without samples) as measured. The growth rate (g d⁻¹) was calculated as $(M_{t_i} - M_{t_0}) / T_i$, where M_{t_0} represents the nubbin weight at the beginning of experiment, M_{t_i} the weight at the measuring time point and T_i represents the duration

in days. The measurement was repeated every 7 d. Data were normalized to skeletal surface area determined with the aluminium foil technique (Marsh, 1970).

SOD and CAT

The homogenized coral tissue slurry that was used to measure the endosymbiont density and Chl *a* content was centrifuged (4,000 rpm min⁻¹, 10 min, 4 °C), and the supernatant was collected to measure the SOD and CAT activities. These measurements were determined in the dilution using kits (A001-1-1, A007-1-1, Nanjing Jicheng, China). A BCA (Bicinchoninic Acid Assay) kit was used to determine the protein concentration (A045-3-1, Nanjing Jicheng, China). The enzyme activities were normalized to total protein content as U mg prot⁻¹.

Data analysis

The results are presented as the means ± standard deviations. Data were tested for homogeneity of variance (visual inspection of residuals vs. fitted values), and the normality of the residuals was tested using the Shapiro–Wilk normality test. All response data of corals were tested using a two-factor analysis of variance (ANOVA) with “temperature (27 °C, 30 °C)” and “algae (control, direct contact, indirect interaction)” as fixed factors, including the interaction term. When significant effects of factors occurred, ANOVAs were followed by a post-hoc Fisher's least significant difference (LSD) multiple comparisons test to identify significant groups. The data was analysed and graphed using GraphPad Prism 8.0 and $p < 0.05$ was considered a significant difference.

Results

Seawater chemistry monitoring

The temperature, pH, and salinity values were measured continuously during the experiment (Figure 2).

Endosymbiont density and Chl *a* content

Figures 3A and 3B show that the density and pigment content of the endosymbiont were significantly influenced by temperature. Without algae, elevated temperature resulted in an average 44.6% (from 1.3 ± 0.1 to $0.72 \pm 0.07 \times 10^6$ cells cm⁻²) decrease in endosymbiont density ($p < 0.05$) and a average 58% (from 17.1 ± 1.5 to 7.2 ± 2.6 µg cm⁻²) decrease in Chl *a* content ($p < 0.01$). Compared with the control group at ambient temperature, the co-occurrence groups and direct contact groups in both temperature treatments showed lower mean values of endosymbiont density. However, there were not significantly different between the control, co-occurrence and

direct contact groups. The only difference in endosymbiont density was a significant difference in Chl *a* concentration between the control and direct contact groups at ambient temperature ($p = 0.03$).

Feeding rate

The results of the feeding rate are displayed in Figure 4A. Primarily, elevated temperature had a significantly detrimental effect on the feeding rate among the treatments. There was no significant difference between the three algae treatments at 27 °C. At 30°C, direct contact with algae caused the feeding rate to fall to a minimum of 12.8 ± 1.1 ind polyp⁻¹ h⁻¹. The interaction between temperature and algae was significant ($F = 4.7$, $p = 0.04$, Table 1).

Protein content

The results of protein content was shown in Figure 4B. At 27 °C, algae caused a loss of 37.5% of protein content in coral tissue in the co-occurrence group ($p = 0.01$) and 49.0% protein content in the direct contact group ($p < 0.01$). Although contact with algae further decreased the mean of protein content, the difference was not significant with the co-occurrence group. At 30 °C, the direct contact with algae resulted in the lowest protein content of 1.2 ± 0.3 mg cm⁻² ($p < 0.01$, $p < 0.01$) compared with the control and co-occurrence groups. No significant difference was observed between the temperature treatments in the presence or absence of algae ($p = 0.11$, Table 1). There was no significant interaction between algae and the temperature treatment ($F = 2.86$, $p = 0.12$, Table 1).

Growth rate

As indicated by the change in buoyant weight, the growth rate was affected by algae at 27°C and the effect of elevated temperature was only seen in the control group (Figure 4C). At 27 °C, the growth rate of corals in the control group was highest, with a mean value of 4.1 ± 1.4 mg cm⁻² d⁻¹. Compared with control group, the coculture with macroalgae decreased the growth rate of coral by 57.7% in the co-occurrence group ($p > 0.05$) and 65.5% in the direct contact group ($p = 0.03$). But no significant difference between co-occurrence and direct contact group. Elevated temperature had an inhibitory effect on the growth rate in the coral culture system without algae. The growth rate in this group was 83.4% lower than the control group ($p < 0.01$). The elevated temperature combined with direct contact with algae resulted in the lowest coral growth rate, with a value of 0.57 ± 0.45 mg cm⁻² d⁻¹. However, the differences among algae treatments at an elevated temperature were not significant. The interaction between temperature and algae was significant ($F = 28$, $p < 0.01$, Table 1).

SOD and CAT

As shown in Figure 5A, macroalgae treatments increased the SOD antioxidant capacity of corals under both temperature conditions. At 27 °C, co-occurrence with algae increased the SOD activity of coral 1.85-fold compared with control group ($p = 0.03$). Moreover, the SOD activity was higher for the direct contact group (288.1 ± 16.6 U mgprot⁻¹) than control groups ($p < 0.01$) and co-occurrence group ($p = 0.03$). At 30 °C, the mean SOD activity of coral without the presence of algae increased by 1.77-fold compared with its counterpart at ambient temperature ($p = 0.03$). In the direct contact group at 30 °C, SOD activity increased to the highest level of 354.3 ± 59.56 U mg prot⁻¹ compared with control group ($p < 0.01$). However, in the co-occurrence and direct contact system, there was no significant difference caused by elevated temperature, indicating that both factors did not interact ($F = 2.37$, $p = 0.16$, Table 1).

The CAT activity also increased after algae interaction and elevated temperature, as shown in Figure 5B. At 27 °C, co-occurrence with algae caused the CAT activity in coral tissue to rise 5.3-fold ($p < 0.05$), which is comparable to the level of CAT activity in the direct contact group. At 30 °C, the CAT activity in the control also increased by 7.1-fold ($p < 0.01$). Moreover, when cultured in contact with the algae, the CAT activity further doubled compared with the control group ($p = 0.03$). The combined effect of temperature and macroalgae was significant ($F = 5.13$, $p = 0.04$, Table 1).

Discussion

This study explored the crucial issue of how the physiology and oxidative stress response of a hermatypic coral are affected by macroalgae at elevated temperatures. We set up three treatments of the macroalga *C. taxifolia* (direct contact, indirect, water-mediated presence, no alga) to act on the coral *T. peltata* at ambient temperatures (27 °C) and elevated temperature (30 °C). The results demonstrated that macroalgal presence increased the antioxidant activity at 27 °C. In addition, combined with elevated temperature, there was a remarkably synergistic effect that macroalgae impacted the feeding rate, protein contain and further increased the oxidative stress of the coral, in which contact with algae had a more severe effect than indirect interaction.

Effects of *C. taxifolia* on endosymbiont of *T. peltata*

Algae was found to influence the average endosymbiont density and chl *a* content of *T. peltata*, however, no bleaching occurred. A number of studies have reported that coral's photosynthetic efficiency decreased (Fv/Fm), or bleaching occurred, when there was direct or

indirect contact with macroalgae. However, not all coral species are equally susceptible to algae and not all algae will have deleterious effects on corals (Smith et al., 2006; Rasher et al., 2010; Fong et al., 2020). Rasher et al. (2010) suggested that *Padina perindusiata* and *Sargassum* sp. did not inhibit photosynthetic efficiency or induce bleaching of *Porites porites*, which might be explained by the fact that the 20-day interaction period was too short to impact the coral health. Additionally, *T. peltata* is a massive coral that could resist environmental pressure by increasing its basic metabolism (Loya et al., 2001). This may explain why there was no significant effects of macroalgae on endosymbiont density. While, there was a stronger effect for direct contact compared to co-occurrence in chl *a* concentration at 27 °C, suggesting that physical mechanisms are still the main way in which macroalgae affect corals.

Effects of *C. taxifolia* and thermal stress on the physiology of *T. peltata*

There were no significant effects of *C. taxifolia* on the feeding rate of *T. peltata* at ambient temperature. However, the feeding rate was affected by elevated temperature. Johannes and Tepley (1974) also found that the feeding rate of coral decreased in heat stress because of the polyp contraction or a loss of nematocyst function. Our results suggested that a decrease in chl *a* content and endosymbiont density was the reason why the feeding rate was impacted in elevated temperature. Endosymbionts provide photosynthate to host cells (van Oppen and Blackall, 2019). A decrease in the endosymbiont density at high temperatures may result in reduced energy expenditure to maintain normal physiological functions and reduce resistance to predation. This study showed that contact with *C. taxifolia* resulted in the greatest reduction in the feeding rate at 30 °C. In summary, thermal stress might a crucial factor affecting the feeding ratio of *T. peltata*, which became more severe when in contact with macroalgae.

Macroalgae can induce reduced protein content in corals. Damage to coral tissue by contact with macroalgae has been documented in many studies. Bender et al. (2012) asserted that *Acropora* sp. lost tissue and decreased its growth rate due to allelopathy mechanisms after coming into contact with *Chlorodesmis fastigiata*. In fact, macroalgae may transfer many allelopathic substances to corals, altering the structure of the microbial community and impacting the physiological processes of corals (Fong et al., 2020). This damage may ultimately reduce the protein content. Under stress, massive corals with thicker tissues may overcome the effects of endosymbiont loss through catabolism (DeCarlo and Harrison, 2019). Macroalgae may affect coral tissue by creating anoxic zones. Barott et al. (2009) demonstrated that after the interactions between corals (*Pocillopora verrucosa*, *Montipora* sp.) and some species of macroalgae (e.g.

Gracilaria sp., *Bryopsis* sp., and various turf algae), the characteristic patterning of coral pigments and polyps was altered and the tissue appeared damaged.

The growth rate of coral was altered by the macroalgae and temperature, the results were consistent with previous studies (Tanner 1995; Rölfer et al., 2021; Rebecca et al., 2012; Vermeij et al., 2009). At 27 °C, the growth rate of the direct contact group was lower than that of the co-occurrence group. Therefore, the effects of seaweed on coral growth may require direct contact at ambient temperatures (Clements et al., 2020). Brown et al. (2019) also demonstrated that coral growth was reduced or even negative at 30 °C when in contact with algae. Longo et al. (2015) determined that corals showed signs of stress at 30 °C, and contact with *Halimeda heteromorpha* further contributed to a decreased growth rate and increased mortality rate. These results may be due to the simultaneous decline in the autotrophic and heterotrophic activities of corals under the impacts of thermal stress or macroalgae, resulting in a drop of protein content and ultimately affecting the growth rate.

Effects of *C. taxifolia* and thermal stress on oxidative stress of *T. peltata*

Corals under thermal stress may produce ROS (Blanckaert et al., 2021). Downs (2002) documented that when exploring the varied oxidative stress response of coral under seasonal change, the SOD in summer was three times higher than that in winter. This study determined that SOD in corals was higher when macroalgae were present (the effect was stronger for direct contact compared to co-occurrence group), the increased temperature, or there was the synergistic effect of both. Thus, weakened corals were found to be more vulnerable to competition from algae, which was also supported by the results of Diaz-Pulido et al (2010). The level of both antioxidant enzyme activities was similar to thermal stress alone when *C. taxifolia* indirectly contacted *T. peltata*. These results indicate that the stress triggered by macroalgal allelochemicals on coral was equivalent to that induced by increased temperature. The temperature effect was more significant in CAT activity compared with SOD activity under direct contact, which may be related to the reduced protein content in coral tissues caused by elevated temperature under the direct contact treatment. Due to the evident decrease in the protein content of coral tissues in the direct contact group, the amount of antioxidant enzymes produced by coral is not enough to resist the damage of ROS.

Conclusions

The shift from coral dominance to algal dominance that has been observed in many reefs due to global climate change and overfishing. Coral dominance is sensitive to key algal groups and other benthic groups, and shifts in ecosystem phases have a noticeable impact (Tebbett et al., 2023). The results of this study showed that *C. taxifolia* negatively affected the growth rate and protein content of *T. peltata* and increased the antioxidant activity at 27 °C. The combination of elevated temperature and macroalgae interactions may further exacerbate the adverse effects on corals. Future studies are needed to explore the interactions of multiple coral-macroalgal species under climate change. Because of the vulnerability and sensitivity of coral reef ecosystem, relevant entities should take urgent steps to prevent CO₂ emissions that exceed the goals of the Paris Climate Agreement. Herbivorous fish populations should also be restored to improve macroalgae management in reefs.

References

- Barott K, Smith J, Dinsdale E, Hatay M, Sandin S, Rohwer F (2009) Hyperspectral and physiological analyses of coral-algal interactions. PLoS One, 4(11): e8043.
- Barott KL, Williams GJ, Vermeij MJA, Harris J, Smith JE, Rohwer FL, Sandin SA (2012) Natural history of coral–algae competition across a gradient of human activity in the Line Islands. Mar Ecol Prog Ser 460:1-12.
- Bender D, Diaz-Pulido G, Dove S (2012) Effects of macroalgae on corals recovering from disturbance. Journal of Experimental Marine Biology and Ecology, 429: 15-19.
- Blanckaert A, Marangoni L, Rottier C, Grover R, Ferrier-Pagès C (2021) Low levels of ultra-violet radiation mitigate the deleterious effects of nitrate and thermal stress on coral photosynthesis. Marine Pollution Bulletin, 167:112257.
- Brown KT, Bender-Champ D, Kenyon TM, R'émond C, Hoegh-Guldberg O, Dove S (2019) Temporal effects of ocean warming and acidification on coral–algal competition. Coral Reefs 38, 297–309.
- Brown KT, Bender-Champ D, Hoegh-Guldberg O, Dove S (2020) Seasonal shifts in the competitive ability of macroalgae influence the outcomes of coral – algal competition. Royal

378 Society open science,7(12):201797.

379 Chadwick NE, Morrow KM (2011) Competition among sessile organisms on coral reefs. Coral
380 reefs: an ecosystem in transition, 347-371.

381 Chakravarti LJ, Buerger P, Levin RA, van Oppen MJH (2020) Gene regulation underpinning
382 increased thermal tolerance in a laboratory-evolved coral photosymbiont. *Mol Ecol*, 29(9):1684-
383 1703.

384 Clements CS, Burns AS, Stewart FJ, Hay ME (2020) Seaweed-coral competition in the field:
385 effects on coral growth, photosynthesis and microbiomes require direct contact. *Proceedings of*
386 *the Royal Society B*, 287(1927):20200366.

387 Coyer JA, Ambrose RF, Engle JM, Carroll JC (1993) Interactions between corals and algae on a
388 temperate zone rocky reef: mediation by sea urchins. *Journal of Experimental Marine Biology and*
389 *Ecology*, 167(1):21-37.

390 Davies, P.S (1989) Short-term growth measurements of corals using an accurate buoyant weighing
391 technique. *Marine biology*, 101, 389–395.

392 DeCarlo TM, Harrison HB (2019) An enigmatic decoupling between heat stress and coral
393 bleaching on the Great Barrier Reef. *PeerJ*, 7: e7473.

394 Diaz-Pulido G, Gouezo M, Tilbrook B, Dove S, Anthony KR (2010) High CO₂ enhances the
395 competitive strength of seaweeds over corals. *Ecology Letters*, 14(2):156-162.

396 Diaz-Pulido G, Barrón C (2020) CO₂ enrichment stimulates dissolved organic carbon release in
397 coral reef macroalgae. *Journal of phycology*, 56(4): 1039-1052.

398 Downs CA, Fauth JE, Halas JC, Dustan P, Bemiss J, Woodley CM (2002) Oxidative stress and
399 seasonal coral bleaching. *Free Radical Biology and Medicine*, 33(4): 533-543.

400 Drury C, Dilworth J, Majerová E, Caruso C, Greer JB (2022) Expression plasticity regulates
401 intraspecific variation in the acclimatization potential of a reef-building coral. *Nature*
402 *communications*, 13(1):4790.

403 Dubinsky Z, Stambler N (2011) *Coral Reefs: An Ecosystem in Transition*. Springer Science &
404 Business Media.

405 Ferrier-Pagès C, Rottier C, Beraud E, Levy O (2010) Experimental assessment of the predation
406 effort of three scleractinian coral species during a thermal stress: Effect on the rates of
407 photosynthesis. *Journal of Experimental Marine Biology and Ecology*, 390(2): 118-124.

408 Fong J, Deignan LK, Bauman AG, Steinberg PD, McDougald D, Todd PA (2020) Contact-and
409 water-mediated effects of macroalgae on the physiology and microbiome of three indo-pacific

410 coral species. *Frontiers in Marine Science*, 6(831).

411 Fulton CJ, Depczynski M, Holmes TH, Noble MM, Radford B, Wernberg T, Wilson SK (2014)

412 Sea temperature shapes seasonal fluctuations in seaweed biomass within the Ningaloo coral reef

413 ecosystem. *Limnology and Oceanography*, 59(1), 156-166.

414 Fulton CJ, Abesamis RA, Berkström C, Depczynski M, Graham NAJ, Holmes TH, Kulbicki M,

415 Noble MM, Radford BT, Tano S, Tinkler P, Wernberg T, Wilson SK (2019) Form and function of

416 tropical macroalgal reefs in the Anthropocene. *Functional Ecology*, 33(6): 989-999.

417 Guerriero A, Meinesz A, D'Ambrosio M, Pietra F (1992) Isolation of Toxic and Potentially Toxic

418 Sesqui- and Monoterpenes from the Tropical Green Seaweed *Caulerpa taxifolia* Which Has

419 Invaded the Region of Cap Martin and Monaco. *Helvetica Chimica Acta*, 75: 689-695.

420 Guerriero A, Marchetti F, D'Ambrosio M, Senesi S, Dini F, Pietra F (1993) New

421 ecotoxicologically and biogenetically relevant terpenes of the tropical green seaweed *Caulerpa*

422 *taxifolia* which is invading the Mediterranean. *Helvetica Chimica Acta*, 76(2): 855-864.

423 Gouvêa LP, Schubert N, Martins CDL, Sissini M, Ramlov F, de Oliveira Rodrigues ER, Bastos

424 EO, Freire VC, Maraschin M, Simonassi JC, Varela DA, Franco D, Cassano V, Fonseca AL,

425 Barufi J B, Horta PA (2017) Interactive effects of marine heatwaves and eutrophication on the

426 ecophysiology of a widespread and ecologically important macroalga. *Limnology and*

427 *Oceanography*, 62(5): 2056-2075.

428 Hensley K, Robinson KA, Gabbita SP, Salsman S, Floyd RA (2000) Reactive oxygen species, cell

429 signaling, and cell injury. *Free Radical Biol Med* 28:1456–1462.

430 Hernández CA, Sangil C, Fanai A, Hernández JC (2018) Macroalgal response to a warmer ocean

431 with higher CO₂ concentration. *Marine Environmental Research*, 136, 99–105.

432 Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH,

433 Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE,

434 Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF,

435 Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs JA, Hoogenboom MO, Kennedy EV, Kuo

436 CY, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM,

437 Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld

438 DR, Willis BL, Wilson SK (2017) Global warming and recurrent mass bleaching of corals. *Nature*,

439 543(7645): 373-377.

440 Hughes TP, Kerry JT, Baird AH, Connolly SR, Chase TJ, Dietzel A, Hill T, Hoey AS,

441 Hoogenboom MO, Jacobson M, Kerswell A, Madin JS, Mieog A, Paley AS, Pratchett MS, Torda

442 G, Woods RM (2019) Global warming impairs stock–recruitment dynamics of corals. *Nature*,
443 568(7752):1-4.

444 Tanner JE (1995) Competition between scleractinian corals and macroalgae: An experimental
445 investigation of coral growth, survival and reproduction. *Journal of Experimental Marine Biology*
446 *and Ecology*, 190(2):151-168.

447 Johannes RE, Tepley L (1974) Examination of feeding of the reef coral *Porites lobata* in situ using
448 time lapse photography. *Proceeding of the 2nd International Coral Reef Symposium*, 1, 127–131.

449 Jompa J, Mccook LJ (2003) Contrasting effects of turf algae on corals: massive *Porites spp.* are
450 unaffected by mixed-species turfs, but killed by the red alga *Anotrichium tenue*. *Marine Ecology*
451 *Progress Series*, 258:79-86.

452 Kornder NA, Riegl BM, Figueiredo J (2018) Thresholds and drivers of coral calcification
453 responses to climate change. *Glob Chang Biol*, 24(11):5084-5095.

454 Lapointe BE, Bedford BJ (2010) Ecology and nutrition of invasive *Caulerpa brachypus f.*
455 *parvifolia* blooms on coral reefs off southeast Florida, U.S.A. *Harmful Algae*, 9(1):1-12.

456 Leong RC, Marzinelli EM, Low J, Bauman AG, Lim EWX, Lim CY, Steinberg PD, Guest JR
457 (2018) Effect of Coral-Algal Interactions on Early Life History Processes in *Pocillopora acuta* in
458 a Highly Disturbed Coral Reef System. *Frontiers in Marine Science*, 5(385):1-11.

459 Leggat W, Heron SF, Fordyce A, Suggett DJ, Ainsworth TD (2022) Experiment Degree Heating
460 Week (eDHW) as a novel metric to reconcile and validate past and future global coral bleaching
461 studies. *Journal of Environmental Management*, 301: 113919.

462 Longo GO, Hay ME (2015) Does seaweed–coral competition make seaweeds more palatable?
463 *Coral Reefs*,34(1): 87-96.

464 Lough JM, Anderson KD, Hughes TP (2018) Increasing thermal stress for tropical coral reefs:
465 1871 – 2017. *Scientific Reports*, 8(1):6079.

466 Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, van Woesik R (2001) Coral bleaching: the
467 winners and the losers. *Ecology Letters*, 4(2), 122–131.

468 Marsh JA (1970) Primary Productivity of Reef-Building Calcareous Red Algae. *Ecology*,51.

469 Manikandan B, Padelkar A A, Ravindran J, Joseph S (2021) Histopathological investigation of the
470 reef coral *Goniastrea* sp. affected by macroalgal abrasion. *Marine Biology*, 168: 1-7.

471 Morrow KM , Carpenter RC (2008) Macroalgal morphology mediates particle capture by the
472 corallimorpharian *Corynactis californica*. *Marine Biology*, 155(3):273-280.

473 Nii CM, Muscatine L (1997) Oxidative stress in the symbiotic sea anemone *Aiptasia pulchella*

474 (Carlgren, 1943): contribution of the animal to superoxide ion production at elevated temperature.
 475 The Biological Bulletin, 192(3):444-456.

476 Rasher DB, Hay ME (2010) Chemically rich seaweeds poison corals when not controlled by
 477 herbivores. Proceedings of the National Academy of Sciences, 107(21): 9683-9688.

478 Rasher DB, Stout EP, Engel S, Kubanek J, Hay ME (2011) Macroalgal terpenes function as
 479 allelopathic agents against reef corals. Proceedings of the National Academy of
 480 Sciences, 108(43):17726-17731.

481 Vega Thurber R, Burkepile DE, Correa AM, Thurber AR, Shantz AA, Welsh R, Pritchard C,
 482 Rosales S (2012) Macroalgae Decrease Growth and Alter Microbial Community Structure of the
 483 Reef-Building Coral, *Porites astreoides*. Plos One, 7(9):e44246.

484 Ritchie RJ (2006) Consistent Sets of Spectrophotometric Chlorophyll Equations for Acetone,
 485 Methanol and Ethanol Solvents. Photosynthesis Research, 89(1):27-41.

486 Bonaldo RM, Hay ME (2014) Seaweed-Coral Interactions: Variance in Seaweed Allelopathy,
 487 Coral Susceptibility, and Potential Effects on Coral Resilience. PLoS ONE, 2014,9(1):e85786.

488 Rölfer L, Reuter H, Ferse SCA, Kubicek A, Dove S, Hoegh-Guldberg O, Bender-Champ D (2021)
 489 Coral-macroalgal competition under ocean warming and acidification. Journal of Experimental
 490 Marine Biology and Ecology, 534: 151477.

491 Shearer T, Rasher D, Snell T, Hay M (2012) Gene expression patterns of the coral *Acropora*
 492 *millepora* in response to contact with macroalgae. Coral Reefs, 31(4):1177-1192.

493 Smith JE, Shaw M, Edwards RA, Obura D, Pantos O, Sala E, Sandin SA, Smriga S, Hatay M,
 494 Rohwer FL (2006) Indirect effects of algae on coral: algae-mediated, microbe-induced coral
 495 mortality. Ecology Letters, 9(7).

496 Skirving W, Marsh B, De La Cour J, Liu G, Harris A, Maturi E, Geiger E, Eakin CM (2020)
 497 CoralTemp and the Coral Reef Watch Coral Bleaching Heat Stress Product Suite Version 3.1.
 498 Remote Sensing, 12(23):3856.

499 Souter D, Planes S, Wicquart J, Logan M, Obura D, Staub F (2021) Status of coral reefs of the
 500 world: 2020 report. Global Coral Reef Monitoring Network (GCRMN)/International Coral Reef
 501 Initiative (ICRI). Accessed: <https://gcrmn.net/2020-report/>

502 Spady B L, Skirving W J, Liu G, De La Cour J L, McDonald C J, Manzello D P (2022).
 503 Unprecedented early-summer heat stress and forecast of coral bleaching on the Great Barrier Reef,
 504 2021-2022. F1000Research, 11, 127.

505 Tebbett S B, Crisp S K, Evans R D, Fulton C J, Pessarrodona A, Wernberg T, Wilson S K,

506 Bellwood D R (2023) On the Challenges of Identifying Benthic Dominance on Anthropocene
507 Coral Reefs. *Bioscience*, 73(3), 220–228.

508 Van Oppen MJH, Blackall LL (2019) Coral microbiome dynamics, functions and design in a
509 changing world. *Nature Reviews Microbiology*, 17(9): 557-567.

510 Vermeij MJ, Smith JE, Smith CM, Vega Thurber R, Sandin SA (2009) Survival and settlement
511 success of coral planulae: independent and synergistic effects of macroalgae and microbes.
512 *Oecologia*, 2009,159(2):325-336.

513 Warner ME (1999) Damage to photosystem II in symbiotic dinoflagellates: A determinant of coral
514 bleaching. *Proceedings of the National Academy of Sciences of the United States of America*,
515 96(14):8007-8012.

516 Zhongming Z, Linong L, Xiaona Y (2021) AR6 Climate Change 2021: The Physical Science
517 Basis. *Chemistry International* 43(4):22-23.

518 Zubia M, Draisma S, Morrissey KL, Varela-Álvarez E, De Clerck O (2020) Concise review of the
519 genus *Caulerpa* J.V. Lamouroux. *Journal of Applied Phycology*, 32(1).

Figure 1

Experimental operating system

System control with a feedback loop to adjust the conditions. Seawater at 27 °C (blue) and 30 °C (red) was heated in different collection sumps (36 L) and then fed into each tank (10 L). There is a drain at the bottom of each tank. The control group had a separate sump and the algae treatment group shared a sump. Macroalgae treatments were applied to (A) *T. peltata* (B) *T. peltata* indirect interaction with *C. taxifolia*. Water flowed first into the algae box (yellow) and then into the tanks to deliver the allelopathic substances secreted by the algae. (C) *T. peltata* in direct contact with *C. taxifolia*. Panel made of PMMA was used to fix the height of the algae parallel to the coral samples. Each treatment contained three replicate tanks, within which three coral nubbins were placed.




Figure 2

The variability in the temperature, pH, and salinity over the course of the 4 weeks experiment.

Seawater at 27 °C (A) temperature, (C) pH, (E) salinity. Seawater at 30 °C (B) temperature, (D) pH, (F) salinity. The dotted lines represent control, direct contact and co-occurrence groups. The solid line represents the average.




Figure 3

The effect of 27 °C (grey) and 30 °C (blue) temperatures on the (A) endosymbiont density and (B) Chl *a* concentration of corals in direct contact or co-occurrence with macroalgae.

Upper case letters represent differences between temperatures, lower case letters represent differences between algal treatments at the same temperature, and different letters represent significant differences ($p < 0.05$). Data are expressed in terms of the mean \pm standard deviation, $n=3$.




Figure 4

The effect of 27 °C (grey) and 30 °C (blue) temperatures on the (A) feeding rate, (B) protein content and (C) growth rate of corals in direct contact or co-occurrence with macroalgae.

Upper case letters represent differences between temperatures, lower case letters represent differences between algal treatments at the same temperature, and different letters represent significant differences ($p<0.05$). Data are expressed in terms of the mean \pm standard deviation, $n=3$.




Figure 5

The effect of 27 °C (grey) and 30 °C (blue) temperatures on the (A) superoxide dismutase activity, (B) catalase activity of corals in direct contact or co-occurrence with macroalgae.

Upper case letters represent differences between temperatures, lower case letters represent differences between algal treatments at the same temperature, and different letters represent significant differences ($p < 0.05$). Data are expressed in terms of the mean \pm standard deviation, $n=3$.




Table 1 (on next page)

Two-way ANOVA output of different variables for *T. peltate*.

The bold values indicate the significant effects on the variable. F=F value; p =p value (significant <0.05).

Variable	Source of variation	F	p
Growth Rate	Algae	$F(2,8) = 28.80$	<0.01
	Temperature	$F(1,4) = 7.78$	0.049
	Interaction	$F(2,8) = 28.00$	<0.01
Feeding Rate	Algae	$F(2,8) = 14.57$	<0.01
	Temperature	$F(1,4) = 119.80$	<0.01
	Interaction	$F(2,8) = 4.70$	0.04
Endosymbiont density	Algae	$F(2,8) = 0.75$	0.5
	Temperature	$F(1,4) = 21.94$	0.01
	Interaction	$F(2,8) = 9.05$	0.01
Chl <i>a</i>	Algae	$F(2,8) = 1.58$	0.27
	Temperature	$F(1,4) = 90.92$	<0.01
	Interaction	$F(2,8) = 7.29$	0.02
Protein	Algae	$F(2,8) = 31.87$	<0.01
	Temperature	$F(1,4) = 4.28$	0.11
	Interaction	$F(2,8) = 2.86$	0.12
SOD	Algae	$F(2,8) = 38.81$	<0.01
	Temperature	$F(1,4) = 16.01$	0.02
	Interaction	$F(2,8) = 2.37$	0.16
CAT	Algae	$F(2,8) = 10.01$	<0.01
	Temperature	$F(1,4) = 64.48$	<0.01
	Interaction	$F(2,8) = 5.13$	0.04