

# Seasonal availability of edible underground and aboveground carbohydrate resources to human foragers on the Cape south coast, South Africa

Jan C De Vynck, Richard M Cowling, Alastair J Potts, Curtis W Marean

The coastal environments of South Africa's Cape Floristic Region (CFR) provide some of the earliest and most abundant evidence for the emergence of cognitively modern humans. In particular, the south coast of the CFR provided a uniquely diverse resource base for hunter-gatherers, which included marine shellfish, game, and carbohydrate-bearing plants, especially those with underground storage organs (USOs). It has been hypothesized that these resources underpinned the continuity of human occupation in the region since the Middle Pleistocene. Very little research has been conducted on the foraging potential of carbohydrate resources in the CFR. This study focuses on the seasonal availability of plants with edible carbohydrate at six-weekly intervals over a two-year period in four vegetation types on South Africa's Cape south coast. Different plant species were considered available to foragers if the edible carbohydrate was directly (i.e. above-ground edible portions) or indirectly (above-ground indications to below-ground edible portions) visible to an expert botanist familiar with this landscape. A total of 52 edible plant species were recorded across all vegetation types. Of these, 33 species were geophytes with edible USOs and 21 species had aboveground edible carbohydrates. Limestone Fynbos had the richest flora, followed by Strandveld, Renosterveld and lastly, Sand Fynbos. The availability of plant species differed across vegetation types and between survey years. The number of available USO species was highest for a six-month period from winter to early summer (Jul-Dec) across all vegetation types. Months of lowest species' availability were in mid-summer to early autumn (Jan-Apr); the early winter (May-Jun) values were variable, being highest in Limestone Fynbos. However, even during the late summer carbohydrate "crunch", 25 carbohydrate bearing species were visible across the four vegetation types. To establish a robust resource landscape will require additional spatial mapping of plant species abundances. Nonetheless, our results demonstrate that plant-based carbohydrate resources available to Stone Age foragers of the Cape south coast, especially USOs belonging to the Iridaceae family, are likely to have comprised a reliable and nutritious source of calories over most of the year.

# Seasonal availability of edible underground and aboveground carbohydrate resources to human foragers on the Cape south coast, South Africa

Jan C. De Vynck<sup>1</sup>, Richard M. Cowling<sup>1</sup>, Alastair J. Potts<sup>1</sup> and Curtis W. Marean<sup>1,2</sup>

<sup>1</sup>Centre for Coastal Palaeosciences, Nelson Mandela Metropolitan University, PO Box 77 000, Port Elizabeth, 6031, Eastern Cape, South Africa

<sup>2</sup>Institute of Human Origins, School of Human Evolution and Social Change, PO Box 872402, Arizona State University, Tempe, Arizona 85287-2402, USA

Corresponding author:

Jan De Vynck

jandevynck@vodamail.co.za

# Abstract

The coastal environments of South Africa's Cape Floristic Region (CFR) provide some of the earliest and most abundant evidence for the emergence of cognitively modern humans. In particular, the south coast of the CFR provided a uniquely diverse resource base for hunter-gatherers, which included marine shellfish, game, and carbohydrate-bearing plants, especially those with underground storage organs (USOs). It has been hypothesized that these resources underpinned the continuity of human occupation in the region since the Middle Pleistocene. Very little research has been conducted on the foraging potential of carbohydrate resources in the CFR. This study focuses on the seasonal availability of plants with edible carbohydrate at six-weekly intervals over a two-year period in four vegetation types on South Africa's Cape south coast. Different plant species were considered available to foragers if the edible carbohydrate was directly (i.e. above-ground edible portions) or indirectly (above-ground indications to below-ground edible portions) visible to an expert botanist familiar with this landscape. A total of 52 edible plant species were recorded across all vegetation types. Of these, 33 species were geophytes with edible USOs and 21 species had aboveground edible carbohydrates. Limestone Fynbos had the richest flora, followed by Strandveld, Renosterveld and lastly, Sand Fynbos. The availability of plant species differed across vegetation types and between survey years. The number of available USO species was highest for a six-month period from winter to early summer (Jul-Dec) across all vegetation types. Months of lowest species' availability were in mid-summer to early autumn (Jan-Apr); the early winter (May-Jun) values were variable, being highest in Limestone Fynbos. However, even during the late summer carbohydrate "crunch", 25 carbohydrate bearing species were visible across the four vegetation types. To establish a robust resource landscape will require additional spatial mapping of plant species abundances. Nonetheless, our results demonstrate that plant-based carbohydrate resources available to Stone Age foragers of the Cape south coast, especially USOs belonging to the Iridaceae family, are likely to have comprised a reliable and nutritious source of calories over most of the year.

# Introduction

The Cape south coast has likely been occupied by hominins for at least the last 1 million years and the earliest archaeological remains attributable to *Homo sapiens* dates to approximately 160,000 years ago (Jerardino & Marean, 2010). The archaeological record on the Cape south coast for the period between 160,000 and 50,000 years is unusually data rich and well dated. This region and time period is crucial to our understanding of modern human origins as it provides some of the earliest evidence for the emergence of complex behaviours associated with cognitively modern humans, making it highly significant to human origins studies (Marean et al., 2014). The archaeological record documents human occupation during periods of glacial maxima, such as Marine Isotope Stages 6 and 4, when climatic conditions over much of the rest of Africa were too harsh for human occupation or could sustain only very small populations (Bar-Matthews et al., 2010; Marean et al., 2014). It has been hypothesised that the richness of the record and continuity of occupation along the Cape south coast, and the Cape Floristic Region (CFR) more generally, is a consequence of an unusually rich resource base unique to this area (Parkington 2001, 2003, 2006; Marean, 2010, 2011). The coastline in this region offers a highly productive inter-tidal zone for shellfish collection for human foragers. During glacial phases, an extensive plain was exposed off the current coast that supported a diverse plains game fauna, which would have offered excellent hunting opportunities (Klein, 1983; Marean, 2010). The CFR is also home to a globally exceptional flora with many species offering harvestable edible carbohydrates (Deacon, 1970, Parkington & Poggenpoel, 1971; Van Wyk & Gericke, 2000; Van Wyk, 2002; Schwegler, 2003; Dominy et al., 2008; De Vynck, Van Wyk & Cowling, 2016). These include geophytic underground storage organs (USOs) that are both highly diverse and locally abundant (Goldblatt, 1978; Procheş, Cowling & du Preez, 2005; Procheş et al., 2006; Singels et al., 2015), as well as many species with aboveground carbohydrates such as fruit, vegetables, seed pods and seeds (De Vynck, Van Wyk & Cowling, 2016). Together these resources may have provided a complementary set of protein and carbohydrate-rich foods to a human forager, thus explaining the continuity of human occupation through glacial maxima.

However, to date the actual availability and productivity of these potential food resources to a human forager has been largely based on conjecture. For example, Marean's (2010) argument that CFR geophyte diversity directly resulted in a wide range of collectable plant foods would not hold if most of those plants were poisonous, very low in caloric returns, very costly to procure, or unavailable for large parts of the year. To better understand the record for hunter-gatherer foraging in this region, we must develop robust understandings of the foraging potential of the plant foods. This includes analyses of their nutritional character (Kyriacou et al., 2014), availability in the landscape and importantly their seasonal availability to a forager.

This study solely focuses on the temporal availability of edible carbohydrate from a range of plant species in four vegetation types on South Africa's Cape south coast. Availability is estimated based on the visibility of plant parts that directly or indirectly lead to edible carbohydrate. We focus on visibility as a proxy for availability as human foragers primarily rely on sight and have a poor sense of smell. Thus, above-ground edible carbohydrate may be readily visible and available when plants are in fruit, but below-ground carbohydrate can only be found if there are above-ground indicators such as leaves, flowers, or dried stalks. To the best of our knowledge at this time, above ground visibility of plant foliage is the most reliable determinant that a human forager could positively identify and extract the underground resource. The ultimate goal is to combine these observations with studies of nutrition, abundance estimates and return rates, so as to contribute to the resourcescape for a paleoscape model for the Cape south coast (Marean et al., 2015). More generally, our paper adds to a growing literature on the importance of geophytes and aboveground carbohydrates to hunter-gatherer diet worldwide (Kaye & Moodie, 1978; Hatley & Kappelman, 1980; Vincent, 1985; Murray et al., 2001; Bird, Bliege Bird & Parker, 2005; Bliege Bird et al., 2008), and expands the range of variation of those data to a region that is megadiverse in plant species yet relatively unstudied in regards to plants, and geophytes in particular, as a food resource.

## Methods

# Study area

The study area is situated in the coastal plain between the Breede and Gouritz rivers on the Cape south coast (Fig. 1). The rainfall regime shows little seasonality and rain may fall at any time of the year although slight rainfall peaks are observed in March–April and with more pronounced peaks during August and October–November (Engelbrecht et al., 2014). The overall climate of the study area is semi-arid to sub-humid with annual rainfall ranging from 350 to 550 mm. The three summer months (Dec-Feb) are the most stressful for plant growth, owing to generally lower rainfall and persistently higher temperatures.

[Fig. 1 here]

Vegetation of the Cape coastal lowlands is under strong edaphic control (Thwaites & Cowling, 1988; Rebelo et al., 1991; Cowling and Potts, 2015) and the study area has a wide range of geologies which generate different soil types. These include Table Mountain Group sandstones (visible on the coast), Bokkeveld shales (exposed on the inland margin of the study area), Cretaceous Enon Formation conglomerates and mudstones (~25 km from the coast), and Bredasdorp Formation limestones (Rogers, 1984; Malan, 1987). In addition, near the coastal margin aeolian sands of marine origin mantle the geology and this varies in pH with age; younger sands are alkaline and older sands are leached and acidic (Rebelo et al., 1991; Abanda, Compton & Hannigan, 2011). Shale- and mudstone-derived soils are moderately fertile, while those associated with leached sands are infertile. The calcareous sands associated with limestone, calcrete and coastal dunes are also relatively infertile due to their high alkalinity and subsequent low levels of plant-available phosphorus (Thwaites & Cowling, 1988).

The Cape Floristic Region (CFR) is home to four biomes, namely Fynbos, Renosterveld, Forest and Subtropical Thicket (Bergh et al. 2014). The vegetation types chosen for this study are representative of dissected coastal margin of the southern CFR, namely Fynbos (two types), Renosterveld and Subtropical Thicket (Rebelo et al., 1991). Generally, Fynbos communities are richest in species, especially local endemics (Cowling, 1990); Renosterveld harbours a high

diversity of USO-bearing geophytes (Proches et al., 2006 ), while Subtropical Thicket has a high diversity of fruit-bearing trees and shrubs (Cowling et al., 1997).

This study monitored the phenological phase of edible plants - periodic life cycle events - growing in single plots located in Renosterveld, Sand Fynbos, Limestone Fynbos, and Strandveld (a form of Subtropical Thicket). Renosterveld occurs on the relatively fertile and fine-grained soils derived from shales and mudstones, and is a fire-prone grassy shrubland often dominated by *Elytropappus rhinocerotis* (renosterbos). Sand Fynbos occurs on infertile acid soils and is a fire-prone heath-like shrubland, characterised by the presence of Restionaceae and Proteaceae. Limestone-derived soils support Limestone Fynbos, a highly endemic-rich vegetation type (Willis et al., 1996). Marine sands are associated with Subtropical thicket, either in its solid form or as thicket clumps in a matrix of Fynbos. This vegetation is colloquially known as Strandveld. Plant compositional change, or beta diversity, between these edaphically differentiated vegetation types is extremely high; consequently few species are shared among these four vegetation types and regional-scale plant richness is very high (Cowling, 1990).

# **Data collection**

The monitoring period was from May 2010 until April 2012. In the year prior to monitoring (2009), the rainfall was far below average across all plots (~70% of the mean annual rainfall). The effects of the previously dry year were still evident when monitoring started. Above average rainfall was experienced over the two years of monitoring (see Fig. S1 in supplementary materials for climate diagrams and Fig. S2 for spatial relation of weather stations to survey plots).

Monitoring plots were located in representative areas of each of the major vegetation types (one plot per vegetation type) described above: Renosterveld, Sand Fynbos, Limestone Fynbos, and Strandveld (Fig. 1). These plots were located within protected areas and were considered to be in a pristine condition. Further biophysical data for the plots are provided in the supplementary materials (Table S1). The Sand Fynbos site had burnt four years before the start of the survey

period and this would likely have enhanced the visibility of USO species (Deacon, 1993), many of which flower more profusely in the early post-fire years (Le Maitre & Midgley, 1992).

Each plot was divided into six  $20 \times 300$  m transects (3.6 ha in total). Monitoring consisted of surveying each plot every six weeks over a two-year period by one of the authors (JC De Vynck), who is a trained field botanist familiar with the vegetation in this landscape. Along each transect, the following were counted: 1) individuals of species bearing underground storage organs (USOs) which would be apparent to a forager (i.e. in a phenological phase where one or more aboveground organs were visible) and, 2) individuals of species with edible aboveground carbohydrates; these included fruits, leaves, seed pods, seeds, and inflorescences. In our sampling approach we adopted a forager's perspective by only including species known to be edible (De Vynck, Van Wyk & Cowling, 2016) and excluding any plants considered too small to harvest. We included as edible any USOs that required cooking in order to render them edible (e.g. tubers of *Rhoicissus digitata* and corms of *Chasmanthe aethiopica* and *Watsonia* spp.) (Wells, 1965; Parkington & Poggenpoel, 1971; Deacon, 1976, 1979; Liengme, 1987; Opperman & Heydenrych, 1990; Skead, Manning & Anthony, 2009).

As stated above, availability of edible carbohydrates was based on direct or indirect visibility of the resource. For example, USOs in their dormant phase are not visible aboveground and can therefore not be procured by foragers. However, visible phases of USOs include any above indicator of their presence, such as green leaves, flowers or dry wilted leaves. The nutritional content may vary through these phases, but is not the focus of this study. The majority of species with aboveground carbohydrates (e.g. fruiting species) are perennially visible, but were only recorded as 'available' to foragers when edible carbohydrates were visible.

## Data analysis

For each plot, the number of visible, and hence available, species with edible carbohydrates over time was determined; this was calculated as the number of individuals observed in a given survey that would provide access to edible carbohydrate divided by the maximum number of individuals observed across all surveys (for a given species within a plot). In order to calculate



the number of species with edible carbohydrate for a given period within a plot, the continuous proportions of individuals with edible carbohydrate per species were converted to binary presence or absence categories using a 10% threshold. Thus, we considered edible carbohydrate offered by a given species readily visible and available in the landscape if at least 10% of the maximum observed individuals were visible with direct or indirect access to carbohydrates. The number of species with edible carbohydrate considered available in each plot through time was calculated. All analyses were conducted in R version 2.15 (R Development Core Team, 2014)

## Results

Within the four 3.6 ha plots spread across the four vegetation types, 52 edible plant species were recorded. Of these, 33 species were geophytes with edible underground storage organs (USOs) and 21 species had aboveground edible carbohydrates (Table 1; see Table S3 and S4 in supplementary materials for full species list per type). Note that some species had more than one edible part. Richness of edible species varied across the vegetation types (Table 1): Limestone Fynbos had the richest flora, followed by Strandveld, Renosterveld and lastly, Sand Fynbos.

[Table 1 here]

Species varied in the length of time they were available through the year [Fig. 2; see supplementary materials for full list of species phenology diagrams (Table S2.1 to S2.8) and phenphase synchronicity among the species (Fig. S3)]. Species with USOs were available for longer periods of the year relative to those with edible aboveground carbohydrates. The availability of USO species differed across vegetation types and between survey years (Fig. 3). Nonetheless, the number of available USO species was highest for a six-month period from winter to early summer (Jul-Dec) across all vegetation types. Months of lowest species' availability were in mid-summer to early autumn (Jan-Apr); the early winter (May-Jun) values were variable, being highest in Limestone Fynbos. In the wetter second year, the summer "crunch" period – where few USO species were available – was at least one month shorter than in the first year. The number of species with available edible aboveground carbohydrates also

varied across vegetation types and sample years. Species richness peaked in spring (Sep-Nov) for all vegetation types; relatively high availability extended into summer (Dec-Feb) but autumn and early winter were lean months for harvesting aboveground carbohydrates in all vegetation types, especially Renosterveld and Sand Fynbos. The presence of two *Carpobrotus* species, which bear ripe fruits during the drier months, was a key factor for the extension of aboveground availability period in Limestone Fynbos.

An impressive 25 species provided edible carbohydrate during the late summer (Feb – Mar) “crunch period (Table 2). Twelve of these were USOs and Limestone Fynbos supported the most species (16) with available carbohydrate present at this time.

[Fig. 2 here]

[Fig. 3 here]

[Table 2 here]

## Discussion

Substantial archaeological evidence exists for the use of underground storage organs (USOs), fruits and leaves by Late Stone Age peoples in southern Africa (Deacon & Deacon, 1963; Parkington & Poggenpoel, 1971; Deacon, 1970, 1976, 1984; Opperman & Heydenrych, 1990; Deacon & Deacon, 1999). This evidence is substantiated by direct observations of contemporary hunter-gatherer communities in Africa (Lee, 1969, 1973, 1984; Silberbauer, 1981; Youngblood, 2004; Berbesque & Marlowe, 2009; Marlowe & Berbesque, 2009). The diversity and abundance of edible plants, especially USOs, along the Cape coast, together with a rich source of both marine and terrestrial based protein, has been hypothesised to be key components facilitating the persistence of Middle Stone Age (MSA) people in the region during glacial phases when other African regions may have been resource poor (Marean, 2010). However, very little research has been conducted on the potential availability of food plants to hunter-gatherers on the Cape south coast to corroborate this hypothesis. In the same study area, Singels et al. (2015) found

surprisingly high edible biomass values for USOs (maximum values range from 600 kg/ha in Sand Fynbos to 5 000 kg/ha in Limestone Fynbos), although these were restricted to occasional biomass hotspots within a matrix of much lower biomass. Also, these USO hotspots were found within all vegetation types. Here we address the temporal availability of belowground (i.e. USOs) and aboveground sources of carbohydrates across the four principal vegetation types of the Cape south coast. We use this to speculate on the importance of carbohydrates as fallback foods for coastal hunter-gatherers, and what role this may have played in the emergence of cognitively modern people in the region (Marean, 2010).

The number of species with edible carbohydrate resources that are visible and available to foragers was highest between winter and early summer in the study area. This is consistent with the dominant cool-season phenology of plants in the Cape Floristic Region (Pierce, 1984). This six-month period provides a diversity of USOs associated with corms belonging to petaloid geophytes, mostly members of the Iridaceae (e.g. *Babiana*, *Freesia*, *Gladiolus*, *Watsonia*). These species provide relatively large (10-100 g) starch-rich and low-fibre food parcels that are inexpensive to harvest (Parkington, 1977; Deacon, 1993; Singels et al., 2015), and many do not require cooking for digestion (Youngblood, 2004; Dominy et al., 2008; J. De Vynck pers. obs., 2011). Also available during the cooler and mostly wetter months are fruits borne largely by Subtropical thicket species (e.g. *Carissa*, *Diospyros*, *Olea*, *Searsia*) as well as leaf crops (*Trachyandra* spp.). There are currently no data on the biomass, nutritional value and foraging returns for aboveground sources of carbohydrate in the Cape Floristic Region. Fruit loads of mature thicket shrubs and trees range from tens of thousands of fruits per plant for *Sideroxylon inerme* (fruit diameter 10 mm) and *Searsia* spp (3 mm) to fewer than 100 fruits for *Euclea racemosa* (7 mm), *Cassine tetragona* (8 mm) and *Osyris compressa* (20 mm) (Cowling et al., 1997). Mat-forming *Carpobrotus* species may bear several tens of large (35 mm diameter) fruits (J. De Vynck pers. obs., 2011).

Late summer to early autumn periods have considerably fewer available edible species than in the other times of the year. This is a period when all traces of leaves and inflorescences of the dominant deciduous geophyte component have disappeared (Deacon, 1993). However, even during this relatively warm and dry period, we recorded some 25 available species across the

four vegetation types (Table 2). These include USOs such as hysteroanthous, autumn-flowering *Gladiolus* (cormous) and *Pelargonium* (tuberous) spp, the corms of evergreen *Watsonia* spp., and the fibrous tubers of the evergreen liana, *Rhoicissus digitata*. Also apparent are the fruits *Carpobrotus* spp, the fruits of many thicket shrubs and trees, and the leaf crop, *Tetragonia decumbens*. Nonetheless, the late summer – early autumn months could represent a carbohydrate “crunch” for foragers: at this time the number of edible plant species is at its lowest and the high-biomass items available to foragers (e.g. *Pelargonium* spp., *R. digitata*) are fibrous and require cooking for digestion (Deacon, 1995; Wandsnider, 1997; Laden & Wrangham, 2005; Dominy et al., 2008; Schnorr et al., 2015).

Overall, the plant-based carbohydrate resources available to Stone Age foragers of the Cape south coast, especially USOs belonging to the Iridaceae (Deacon, 1976, 1993), are likely to have comprised a reliable and nutritious source of calories over most of the year. Moreover, availability of USOs showed little between-year variation, most likely due the existence of sufficient storage reserves to enable at least leaf growth every year (Ruiters and McKenzie, 1994) despite variation in rainfall. In an assessment of foraging potential of six USO species growing in our study area, Singels et al. (2015) showed that 50% of foraging events conducted yielded enough calories to meet the daily requirements of a hunter-gatherer of small stature within two hours.

The juxtaposition within a 10 kilometre foraging radius of four major vegetation types, belonging to three regional biomes (Fynbos, Renosterveld and Subtropical Thicket; Bergh et al., 2014), would have enabled humans to forage in very different resourcescapes on a daily basis. While the Limestone Fynbos and Strandveld – the two vegetation types closest to the coast – are likely to have offered the best foraging returns for much of the year, Renosterveld provides an abundance of Iridaceae corms in the spring and Sand Fynbos harbours evergreen *Watsonia* spp., which can be harvested during the late summer-autumn “crunch” (Singels et al., 2015). Ethnographic evidence suggest that the harvesting of Iridaceae corms (*uintjies*) in spring was an important event for the San of the Cape west coast (Van Vuuren, 2014).

328 Given the temporal and spatial availability of edible plant species in the Cape, we argue that is  
 329 highly likely that USOs, fruit, seedpods, seeds, inflorescences and leaf crops were harvested as  
 330 fallback foods by Stone Age people living in this region. The likely preferred food for south  
 331 Cape coastal hunter-gatherers comprised the region's diverse and abundant marine resources  
 332 (Marean et al., 2007; Jerardino & Marean, 2010; Parkington, 2010), and a diverse ungulate  
 333 plains fauna, including in the Pleistocene, several species of now extinct megafauna, associated  
 334 with the submerged Agulhas Bank (Klein, 1983; Parkington, 2001, 2003; Matthews, Marean &  
 335 Nilssen, 2009; Marean, 2010; Faith, 2011). However, these resources were not always available  
 336 to harvesters and hunters, and the contraction and expansion of the Agulhas Plain ecosystem and  
 337 its ungulate communities must have been a major driver of changing foraging patterns on the  
 338 south coast (Marean et al., 2014). It has been hypothesized that the mammal fauna formed a  
 339 migratory community that moved west during the winter rains and east to intercept the summer  
 340 rains. Thus, the local abundance of many of the larger ungulates may have plummeted during  
 341 the winter months when populations migrated west to graze winter-growing grasses of the west  
 342 coast. Marine invertebrates, harvested from the intertidal, comprised the most reliable and  
 343 accessible source of protein for hunter-gatherers living on the Cape south coast (Marean, 2011).  
 344 Evidence for their use has been found in MSA sites such as Pinnacle Point (PP) 13B dating back  
 345 to ~160 ka (Marean et al., 2007; Jerardino & Marean, 2010) and at early modern human sites that  
 346 date between 110-50 ka such as Blombos Cave (Henshilwood et al., 2001; Langejans et al.,  
 347 2012), and Klasies River Mouth (Voigt, 1973; Thackeray, 1988). Late Stone Age sites suggest an  
 348 increase in the intensity of intertidal foraging (Marean et al., 2014) and indications of resource  
 349 depletion (Klein & Steele, 2013). Using experienced foragers of Khoe-San descent, J. De Vynck  
 350 et al. (unpublished data) showed exceptionally high peak return rates (~4,500 kcal hr<sup>-1</sup>) from the  
 351 Cape south coast intertidal under ideal harvesting conditions. However, owing to tidal  
 352 constraints, and the fierce sea conditions experienced there, harvesting was only possible for 10  
 353 days a month, for 2-3 hours on each day; lowest returns were recorded in winter and spring – a  
 354 time of strong winds and high seas – and highest returns in summer and autumn, when sea  
 355 conditions were calmer (J. De Vynck et al., unpublished data). Consequently, there would have  
 356 been periods of various lengths – ranging from days to weeks – when hunter-gatherers depended  
 357 on, or fell back upon carbohydrates for sustenance. As pointed out above, the winter and early  
 358 spring months likely coincided with a scarcity of protein but an abundance of carbohydrates. At

these times, plant carbohydrates, especially USOs, may have comprised 100% of dietary intake, which would categorise them as a staple fallback food (Marshall & Wrangham, 2007).

It has been hypothesized that the persistence of a small group of hominins on the Cape south coast – as opposed to their widespread extinction elsewhere in Africa during Marine Isotope Stage 6 (MIS6, 193 000 -125 000 BP) (Foley, 1998; Lahr & Foley, 1998; Fagundes et al., 2007; Basell, 2008; Masson-Delmotte et al., 2010) – was a consequence of the Cape’s relatively moderate climate during the largely glacial MIS6 and its rich and diverse resource base. The persistently warm Agulhas Current reduced the regional impact of glacial cooling substantially (Negre et al., 2010; Zahn et al., 2010). Marean (2010) has hypothesised that during strong glacial environments, such as those experienced in MIS6, the Cape south coast provided a unique juxtaposition of resources important for hominin persistence, namely a diverse USO flora and a rich and productive marine ecosystem. At that time the exposed Agulhas Plain (Fisher, Barr-Matthews & Marean, 2010) was mantled in substrata that likely supported Renosterveld, Limestone Fynbos and Strandveld (Cawthra et al., 2015), offering a wide array of USOs, fruit and leaf crops which would comprise reliable fallback foods when it was not possible to forage in the intertidal and game was scarce. The cognitive challenges of exploiting marine resources (e.g. comprehending lunar cycles), and defending them against competition from adjacent groups, led to a coastal adaptation that may have contributed to the emergence of cognitively modern *Homo sapiens* (Marean, 2011). Similarly, the ability to recognise which and when vegetation types are most productive for carbohydrates, identifying hotspots of productivity and distinguishing between edible and toxic USOs, must have been challenging (Deacon, 1995). Here we have established the temporal availability of plant species with edible carbohydrates across four dominant vegetation types along the south coast. Much additional research must be done to evaluate more comprehensively the role of above- and belowground carbohydrates in the ecology and evolution of the human lineage in the Cape Floristic Region and elsewhere. Work is currently underway to establish the return rates of carbohydrate resources harvested by contemporary subjects of Khoe-San descent, in the different vegetation types and in different seasons; and on the rates of depletion of resources in successively harvested areas. This needs to be complemented with data on the nutritional value of the consumed parts of the species selected. Ultimately, we aim to use these data to populate the carbohydrate resourcescape in an

agent-based model aimed at predicting the effects of spatial and temporal variability – governed by changes in climate and the resource base over the seasonal cycle as well as the glacial-interglacial cycle of the Pleistocene – on the population size and structure, mobility, social organization, territoriality, and technology of Cape hunter-gatherers (Marean et al., 2015).

## Acknowledgements

We thank the Cape Nature team – Rhett Heismann, Jean Du Plessis and Leandi Wessels – for access, support, information and GIS assistance. We also thank the Hessequa Municipality, and in particular Hendrik Visser, for their help and support. The authors are also grateful for the climate data supplied by South African Weather Service. Peter Henzi and an anonymous reviewer provided valuable comments that improved the manuscript.

## References

- Abanda PA, Compton JS, Hannigan RE. 2011. Soil nutrient content, above-ground biomass and litter in a semi-arid shrubland, South Africa. *Geoderma* 164 (3-4): 128-137.
- Bar-Matthews M, Marean CW, Jacobs Z, Karkanas P, Fisher EC, Herries AIR, Brown K, Williams HM, Bernatchez J, Ayalon, Nilssen PJ. 2010. A high resolution and continuous isotopic speleothem record of paleoclimate and paleoenvironment from 90 to 53 ka from Pinnacle Point on the south coast of South Africa. *Quaternary Science Reviews*, 29 (17): 2131-2145.
- Basell LS. 2008. Middle Stone Age (MSA) site distributions in eastern Africa and their relationship to Quaternary environmental change, refugia and the evolution of *Homo sapiens*. *Quaternary Science Reviews* 27 (27): 2484-2498.

418 Bergh NG, Verboom GA, Rouget M, Cowling RM. 2014. Vegetation types of the Greater Cape  
419 Floristic Region. In: Allsopp N, Colville JF, Verboom T, eds. *Fynbos: Ecology, Evolution, and*  
420 *Conservation of a Megadiverse Region.* , Oxford: Oxford University Press, 1-25.  
421

422 Berbesque JC, Marlowe FW. 2009. Sex differences in food preferences of Hadza Hunter-  
423 Gatherers. *Evolutionary Psychology* 7 (4): 601-616.  
424

425 Bird DW, Bliege Bird RB, Parker CH. 2005. Aboriginal burning regimes and hunting strategies  
426 in Australia's Western Desert. *Human Ecology* 33 (4): 443-464.  
427

428 Bliege Bird RB, Bird DW, Coddig BF, Parker CH, Jones JH. 2008. The "fire stick farming"  
429 hypothesis: Australian Aboriginal foraging strategies, biodiversity, and anthropogenic fire  
430 mosaics. *Proceedings of the National Academy of Sciences* 105: 14796-14801.  
431

432 Cawthra HC, Compton JS, Fisher EC, MacHutchon MR, Marean CW. 2015. Submerged  
433 shorelines and landscape features offshore of Mossel Bay, South Africa. *Geological Society,*  
434 *London, Special Publications* 411: SP411-11.  
435

436 Cowling RM. 1990. Diversity components in a species-rich area of the Cape Floristic Region.  
437 *Journal of Vegetation. Science* 1: 699-710.  
438

439 Cowling RM, Kirkwood D, Midgley JJ, Pierce SM. 1997. Invasion and persistence of bird-  
440 dispersed, subtropical thicket and forest species in fire-prone fynbos. *Journal of Vegetation*  
441 *Science* 8: 475-488.  
442

443 Cowling RM, Potts AJ. 2015. Climatic, edaphic and fire regime determinants of biome  
444 boundaries in the eastern Cape Floristic Region. *South African Journal of Botany* 101: 73-81.  
445

446 Deacon HJ. 1970. Plant remains from Melkhoutboom Cave, South Africa. *Proceedings Transkei*  
447 *and Ciskei Research Society* 1: 13-15.  
448



449 Deacon HJ. 1976. *Where hunters gathered: a study of Holocene Stone Age people in the Eastern*  
 450 *Cape*. Claremont: South African Archaeological Society.  
 451

452 Deacon HJ. 1979. Excavations at Boomplaas cave-a sequence through the upper Pleistocene and  
 453 Holocene in South Africa. *World Archaeology* 10 (3): 241-257.  
 454

455 Deacon HJ. 1993. Planting an Idea: An Archaeology of Stone Age Gatherers in South Africa.  
 456 *The South African Archaeological Bulletin* 48: 86-93.  
 457

458 Deacon HJ. 1995. Two late Pleistocene-Holocene archaeological depositories from the southern  
 459 Cape, South Africa. *The South African Archaeological Bulletin* 121-131.  
 460

461 Deacon J. 1984. Later Stone Age people and their descendants in southern Africa. In: Klein RG  
 462 ed. *Southern African Prehistoric and Paleoenvironments*. Rotterdam/Boston: A. A. Balkema,  
 463 220-328.  
 464

465 Deacon HJ, Deacon J. 1963. Scott's Cave: a late Stone Age site in the Gamtoos Valley Annals of  
 466 the Cape Provincial Museums. *Natural History* 3: 96-112.  
 467

468 Deacon HJ, Deacon J. 1999. *Human beginnings in South Africa: Uncovering the secrets of the*  
 469 *Stone Age*. Cape Town: David Philip Publishers (Pty) Ltd.  
 470

471 De Vynck JC, Van Wyk BE, Cowling RM. 2016. Indigenous Edible Plant Use by Contemporary  
 472 Khoe-San Descendants of South Africa's Cape South Coast. *South African Journal of Botany*  
 473 102: 60-69.  
 474

475 Dominy NJ, Vogel ER, Yeakel JD, Constantino P, Lucas PW. 2008. Mechanical properties of  
 476 plant underground storage organs and implications for dietary models of early Hominins.  
 477 *Evolutionary Biology* 35: 159-175.  
 478

Engelbrecht CJ, Landman WA, Engelbrecht FA, Malherbe J. 2014. A synoptic decomposition of  
rainfall over the Cape south coast of South Africa. *Climate Dynamics* 44 (9-10): 2589-2607.

Fagundes NJ, Ray N, Beaumont M, Neuenschwander S, Salzano FM, Bonatto SL, Excoffier L.  
2007. Statistical evaluation of alternative models of human evolution. *Proceedings of the  
National Academy of Sciences* 104 (45): 17614-17619.

Faith JT. 2011. Ungulate community richness, grazer extinctions, and human  
subsistence behavior in southern Africa's Cape Floral Region. *Palaeogeography,  
Palaeoclimatology, Palaeoecology* 306 (3): 219-227.

Fisher EC, Bar-Matthews MJA, Marean CW. 2010. Middle and Late Pleistocene paleoscape  
modeling along the southern coast of South Africa. *Quaternary Science Reviews* 29 (11): 1382-  
1398.

Foley R. 1998. The context of human genetic evolution. *Genome Research* 8 (4): 339-347.

Goldblatt P. 1978. An analysis of the flora of Southern Africa: its characteristics, relationships,  
and origins. *Annals of the Missouri Botanical Gardens* 65: 369-436.

Hatley T, Kappelman J. 1980. Bears, pigs, and plio-Pleistocene Hominids: A case for the  
exploitation of belowground food resources. *Human Evolution* 8: 371-387.

Henshilwood CS, Sealy JC, Yates R, Cruz-Urbe K, Goldberg P, Grine FE, Klein RG,  
Poggenpoel C, van Niekerk K, Watts I. 2001. Blombos Cave, southern Cape, South Africa:  
preliminary report on the 1992–1999 excavations of the Middle Stone Age levels. *Journal of  
Archaeological Science* 28 (4): 421-448.

Jerardino A, Marean CW. 2010. Shellfish gathering, marine paleoecology and modern human  
behavior: perspectives from cave PP13B, Pinnacle Point, South Africa. *Journal of Human  
Evolution* 59 (3-4): 412-424.

510

511 Kaye B, Moodie DW. 1978. The Psoralea food resource of the Northern Plains. *Plains*  
512 *Anthropologist* 23: 329-36

513

514 Klein RG. 1983. Palaeoenvironmental implications of Quaternary large mammals in the fynbos  
515 region. In: Deacon HJ, Hendey QB, Lambrechts JJN, eds. *Fynbos Paleoecology: a preliminary*  
516 *synthesis*. Pretoria: South African National Scientific Programmes Report, 75: 116-138.

517

518 Klein RG, Steele TE. 2013. Archaeological shellfish size and later human evolution in Africa.  
519 *Proceedings of the National Academy of Science* 110 (27): 10910-10915.

520

521 Kyriacou K, Parkington JE, Marais AD, Braun DR. 2014. Nutrition, modernity and the  
522 archaeological record: Coastal resources and nutrition among Middle Stone Age hunter-gatherers  
523 on the western Cape coast of South Africa. *Journal of Human Evolution* 77: 64-73.

524

525 Laden G, Wrangham R. 2005. The rise of the hominids as an adaptive shift in fallback foods:  
526 plant underground storage organs (USOs) and australopith origins. *Journal of Human Evolution*  
527 49 (4): 482-498.

528

529 Lahr MM, Foley RA. 1998. Towards a theory of modern human origins: geography,  
530 demography, and diversity in recent human evolution. *Yearbook of Physical Anthropology*, 41:  
531 137-176.

532

533 Langejans GHJ, van Niekerk KL, Dusseldorp GL, Thackeray JF. 2012. Middle Stone Age  
534 shellfish exploitation: Potential indications for mass collecting and resource intensification at  
535 Blombos Cave and Klasies River, South Africa. *Quaternary International* 270: 80-94.

536

537 Lee RB. 1969. !Kung Bushman subsistence: An input-output analysis. In: Vayda AP, ed.  
538 *Environment and Cultural Behaviour*. Austin: University of Texas Press, 47-79.

539

- 540 Lee RB. 1973. Mongongo: The ethnography of a major wild food resource. *Ecology of Food and*  
541 *Nutrition* 2 (4): 213-307.
- 542
- 543 Lee RB. 1984. *The Dobe !Kung*. New York: Holt, Rinehart and Winston, Inc., 40-44.
- 544
- 545 Le Maitre DC, Midgley JJ. 1992. Plant reproductive ecology. In: Cowling RM, ed. *The ecology*  
546 *of Fynbos: nutrients, fire and diversity*. Cape Town: Oxford University Press, 135-174.
- 547
- 548 Liengme C. 1987. Botanical remains from archaeological sites in the Western Cape. *Paper in the*  
549 *prehistory of the Western Cape*. Oxford: BAR International Series, 237-261.
- 550
- 551 Malan JA. 1987. The Bredasdorp Group in the area between Gans Bay and Mossel Bay. *South*  
552 *African Journal of Science* 83 (8): 506-507.
- 553
- 554 Marean CW. 2010. Pinnacle Point Cave 13B (Western Cape Province, South Africa) in context:  
555 the Cape Floral kingdom, shellfish, and modern human origins. *Journal of Human Evolution* 59  
556 (3-4): 425-443.
- 557
- 558 Marean CW. 2011. Coastal South Africa and the coevolution of the modern human lineage and  
559 the coastal adaptation. In: Bicho NS, Haws JA, Davis LG, eds. *Trekking the shore.*  
560 *Interdisciplinary contributions to Archaeology*. New York: Springer, 421-440.
- 561
- 562 Marean CW, Bar-Matthews M, Bernatchez J, Fisher E, Goldberg P, Herries AIR, Jacobs Z,  
563 Jerardino A, Karkanas P, Minichillo T, Nilssen PJ, Thompson E, Watts I, Williams HM. 2007.  
564 Early human use of marine resources and pigment in South Africa during the Middle  
565 Pleistocene. *Nature* 449 (7164): 905-908.
- 566
- 567 Marean CW, Cawthra HC, Cowling, RM, Esler KJ, Fisher E, Milewski A, Potts AJ, De Vynck  
568 JC. 2014. Stone Age People in a Changing South African Greater Cape Floristic Region. In:  
569 Allsopp N, Colville JF, Verboom T, eds. *Fynbos: Ecology, Evolution, and Conservation of a*  
570 *Megadiverse Region*. Oxford: Oxford University Press, 164-199.

571

572 Marean CW, Anderson RJ, Bar-Matthews M, Braun K, Cawthra HC, Cowling RM, Engelbrecht  
573 F, Esler KJ, Fisher E, Franklin J, Hill K, Janssen M, Potts AJ, Zahn R. 2015. A new research  
574 strategy for integrating studies of paleoclimate, paleoenvironment, and paleoanthropology.  
575 *Evolutionary Anthropology: Issues, News, and Reviews* 24 (2): 62-72.

576

577 Marshall A, Wrangham R. 2007. Evolutionary consequences of fallback foods. *International*  
578 *Journal of Primatology* 28: 1219-1235.

579

580 Marlowe FW, Berbesque JC. 2009. Tubers as fallback foods and their impact on Hadza Hunter-  
581 Gatherers. *American Journal of Physical Anthropology* 140 (4): 751-758.

582

583 Masson-Delmotte V, Stenni B, Pol K, Braconnot P, Cattani O, Falourd S, Kageyama M, Jouzel  
584 J, Landais A, MinsterB, Barnola JM, Chappellaz J, Krinner G, Johnsen S, Rothlisberger R,  
585 Hansen J, Mikolajewicz U, Otto-Bliesner B. 2010. EPICA Dome C record of glacial and  
586 interglacial intensities. *Quaternary Science Reviews* 29(1): 113-128.

587

588 Matthews T, Marean CW, Nilssen, PJ. 2009. Micromammals from the Middle Stone Age (92  
589 000 – 167 000 ka) at Cave PP13B, Pinnacle Point, south coast, South Africa. *Paleontologia*  
590 *Africana* 4: 112-120.

591

592 Murray SS, Schoeninger MJ, Bunn HT, Pickering TR, Marlett, J. A. 2001. Nutritional  
593 Composition of Some Wild Plant Foods and Honey Used by Hadza Foragers of Tanzania.  
594 *Journal of Food Composition and Analysis* 14: 3-13.

595

596 Negre C, Zahn R, Thomas AL, Masqué P, Henderson GM, Martínez-Méndez G, Hall IR, Mas  
597 JL. 2010. Reversed flow of Atlantic deep water during the Last Glacial Maximum. *Nature* 468  
598 (7320): 84-88.

599

- Opperman H, Heydenrych B. 1990. A 22 000 Year-Old Middle Stone Age camp site with plant food remains from the North-Eastern Cape. *The South African Archaeological Bulletin* 45 (152): 93-99.
- Parkington JE. 1977. Soaqua: Hunter-fisher-gatherers of the Olifants River valley Western Cape. *South African Archaeological Society* 32: 150-157.
- Parkington JE. 2001. Milestones: the impact of the systematic exploitation of marine foods on human evolution. In: Tobias PV, Raath MA, Moggi-Cecchi J, Doyle GA, eds. *Humanity from African naissance to coming Millennia - colloquia in human biology and palaeoanthropology*. Firenze: Firenze University Press, 327-336.
- Parkington J. 2003. Middens and moderns: shellfishing and the Middle Stone Age of the Western Cape, South Africa. *South African Journal of Science* 99 (5/6): 243.
- Parkington J. 2006. Shorelines, strandloppers and shell middens: Archaeology of the Cape Coast. Cape Town: Krakadouw.
- Parkington J. 2010. Coastal diet, encephalization, and innovative behaviors in the late Middle Stone Age of southern Africa. *Human Brain Evolution: The Influence of Freshwater and Marine Food Resources*. New York: Wiley-Blackwell, 189-202.
- Parkington JE, Poggenpoel C. 1971. Excavations at de Hangen, 1968. *The South African Archaeological Bulletin* 26: 3-26.
- Pierce SM. 1984. *A synthesis of plant phenology in the Fynbos Biome*. South African National Scientific Programmes Report no. 88. Pretoria: Graphics art division of the CSIR, 29.
- Procheş S, Cowling RM, du Preez DR. 2005. Patterns of geophyte diversity and storage organ size in the winter-rainfall region of southern Africa. *Diversity and Distributions* 11 (1): 101-109.

631 Procheş S, Cowling RM, Goldblatt P, Manning JC, Snijman DA. 2006. An overview of the Cape  
632 geophytes. *Biological Journal of the Linnean Society* 87 (1): 27-43.  
633

634 R Development Core Team. 2014. R: A language and environment for statistical computing.  
635 Vienna, Austria: R Foundation for Statistical computing, ISBN 3-900051-07-0, URL  
636 <http://www.R-project.org>  
637

638 Rebelo AG, Cowling RM, Campbell BM, Meadows M. 1991. Plant communities of the  
639 Riversdale plain. *South African Journal of Botany* 57: 10-28.  
640

641 Rogers J. 1984. *Cenozoic geology of the southern Cape coastal plain between Cape Agulhas and*  
642 *Mossel Bay, focussing on the area between the Kafferkuils and Gouritz rivers*. Technical report  
643 No. 15: University of Cape Town: Marine Geoscience Unit.  
644

645 Ruiters C, McKenzie B. 1994. Seasonal allocation and efficiency patterns of biomass and  
646 resources in the perennial geophyte *Sparaxis grandiflora* subspecies *fimbriata* (Iridaceae) in  
647 lowland coastal fynbos, South Africa. *Annals of Botany* 74 (6): 633-646.  
648

649 Schnorr SL, Crittenden AN, Venema K, Marlowe FW, Henry AG. 2015. Assessing digestibility  
650 of Hadza tubers using a dynamic in-vitro model. *American Journal of Physical Anthropology*  
651 158 (3): 371-385.  
652

653 Schwegler M. 2003. *Medicinal and Other Uses of Southern Overberg Fynbos Plants*. Cape  
654 Town: M. Schwegler.  
655

656 Silberbauer GB. 1981. *Hunter and habitat in the central Kalahari Desert*. Cambridge:  
657 Cambridge University Press.  
658

659 Singels E, Potts AJ, Esler KJ, Cowling RM, Marean CW, De Vynck JC. 2015. Foraging potential  
660 of underground storage organ plants in the southern Cape, South Africa. *PeerJ PrePrints* 3:e1962  
661 <https://doi.org/10.7287/peerj.preprints.1573v1>

662

663 Skead CJ, Manning JC, Anthony NC. 2009. *Historical plant incidence in southern Africa: a*  
664 *collection of early travel records in southern Africa*. Pretoria: South African National  
665 Biodiversity Institute.

666

667 Thackeray JF. 1988. Molluscan Fauna from Klasies River, South Africa. *South African*  
668 *Archaeological Bulletin* 43: 27-32.

669

670 Thwaites RN, Cowling RM. 1988. Soil-vegetation relationships on the Agulhas plain, South  
671 Africa. *Catena* 15 (3): 333-45.

672

673 Van Vuuren H. 2014. ‘A song sung by the star !Gaunu, and especially by Bushman women’: the  
674 blossoming of the uintjieblom. In: Deacon J, Skotnes P. *The courage of //kaboo. Celebrating the*  
675 *100th anniversary of the publication of Specimens of Bushman Folklore*. Cape Town: UCT  
676 Press, 317-328.

677

678 Van Wyk BE. 2002. A review of ethnobotanical research in South Africa. *South African Journal*  
679 *of Botany* 68: 1–13.

680

681 Van Wyk BE, Gericke N. 2000. *People’s plants: a guide to useful plants of southern Africa*.  
682 Johannesburg: Briza Publications.

683

684 Vincent AS. 1985. *Underground Plant Foods and Subsistence in Human Evolution*. California:  
685 University of California at Berkeley.

686

687 Voigt E, 1973. Klasies River Mouth Cave: An exercise in shell analysis. *Bulletin of the*  
688 *Transvaal Museum* 14: 14-15.

689

690 Wandsnider L. 1997. The roasted and the boiled: food composition and heat treatment with  
691 special emphasis on pit-hearth cooking. *Journal of Anthropological Archaeology*, 16 (1): 1-48.

692



693 Wells MJ. 1965. An analysis of plant remains from Scott's Cave in the Gamtoos Valley. *The*  
694 *South African Archaeological Bulletin* 79-84.  
695  
696 Willis CK, Lombard AT, Cowling RM, Heydenrych BJ, Burgers CJ. 1996. Reserve systems for  
697 limestone endemic flora of the Cape lowland fynbos: iterative versus linear programming.  
698 *Biological Conservation* 77 (1): 53-62.  
699  
700 Youngblood D. 2004. Identification and quantification of edible plant foods in the Upper (Nama)  
701 Karoo, South Africa. *Economic Botany* 58 (1): 43-65.  
702  
703 Zahn R, Lutjeharms J, Biastoch A, Hall I, Knorr G, Park W, Reason C. 2010. Investigating the  
704 global impacts of the Agulhas Current. *Eos, Transactions American Geophysical Union* 91 (12):  
705 109-110.  
706  
707

Table 1: Summary of edible species in 3.6 ha plots situated in four dominant vegetation types along the Cape south coast.

	USOs	Fruit	Other <sup>1</sup>	All
Renosterveld	8	6	2	16
Sand Fynbos	5	4	1	10
Limestone Fynbos	21	11	7	39
Strandveld	15	8	5	28
Across all types <sup>2</sup>	33	14	8	52


<sup>1</sup> 'Other' includes species with edible: seed pods, seeds, leaves, and inflorescences.

<sup>2</sup> Note that this is the number of unique species (i.e. some species are shared between vegetation types or have more than one edible part).

716 Table 2: Species available during the ‘carbohydrate-crunch’ late summer period (February-  
717 March) in both survey years.

718

Vegetation type	Carbohydrate category	Species
Renosterveld	Underground storage organ	<i>Babiana patula</i> ; <i>Cyphia digitata</i> ; <i>Watsonia meriana</i>
	Aboveground	<i>Diospyros dichrophylla</i> (fruit); <i>Osyris compressa</i> (fruit); <i>Sideroxylon inerme</i> (fruit)
Limestone Fynbos	Underground storage organ	<i>Cyphia digitata</i> ; <i>Gladiolus exilis</i> ; <i>Pelargonium lobatum</i> ; <i>Pelargonium triste</i> ; <i>Rhoicissus digitata</i> ; <i>Watsonia fergusoniae</i>
	Aboveground	<i>Carissa bispinosa</i> (fruit); <i>Carpobrotus accinaciformis</i> (fruit); <i>Carpobrotus edulis</i> (fruit); <i>Cynanchum obtusifolium</i> (seedpods); <i>Euclea racemosa</i> (fruit); <i>Osyris compressa</i> (fruit); <i>Searsia glauca</i> (fruit); <i>Sideroxylon inerme</i> (fruit); <i>Tetragonia decumbens</i> (leaves); <i>Zygophyllum morskana</i> (seed)
Sand Fynbos	Underground storage organ	<i>Gladiolus guthriei</i> ; <i>Watsonia fourcadei</i>
	Aboveground	<i>Carpobrotus edulis</i>
Strandveld	Underground storage organ	<i>Chasmanthe aethiopica</i> ; <i>Ferraria crispa</i> ; <i>Rhoicissus digitata</i>
	Aboveground	<i>Carissa bispinosa</i> (fruit); <i>Carpobrotus accinaciformis</i> (fruit); <i>Osteospermum moniliferum</i> (fruit); <i>Schotia afra</i> (seed); <i>Tetragonia decumbens</i> (leaves)




719

720 Figure 1: a) The location and major vegetation types of the study region and the plot localities [1:  
 721 Renosterveld (purple); 2: Limestone Fynbos (green); 3: Sand Fynbos (yellow); and 4: Strandveld  
 722 (orange; restricted to the coastal margin; see Tabe S1 for further plot details)]. b) A Walter-Leith

723 climate diagram from the town of Still Bay (~5 km from plot 2). c) Photos taken within the four  
724 plots in the different vegetation types.


725



726

727

728 Figure 2: A breakdown of the number of months in which different plant species (with edible  
729 carbohydrates) are visible through the year separated into a) underground storage organs (USOs)  
730 and b) aboveground carbohydrate (e.g. fruit, seed pods, seeds, leaves or inflorescences).



732 Figure 3: The seasonal availability of edible species visible to a human forager in four vegetation  
 733 types dominant along the Cape south coast. Underground storage organs are geophytes that have  
 734 tubers, corms, bulbs or rhizomes, while above-ground carbohydrate includes specie with edible  
 735 fruit, seed pods, seeds, leaves or inflorescences. The number of new species observed since the  
 736 previous survey is shown in each circle; this provides an indication of species turnover.  
 737