Evaluation of sesame (Sesamum indicum L.) varieties for drought tolerance using agromorphological traits and drought tolerance indices (#85584)

First submission

Guidance from your Editor

Please submit by 26 May 2023 for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the <u>materials page</u>.

- 3 Figure file(s)
- 8 Table file(s)
- 3 Other file(s)

i

Structure and Criteria



Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

- 1. BASIC REPORTING
- 2. EXPERIMENTAL DESIGN
- 3. VALIDITY OF THE FINDINGS
- 4. General comments
- 5. Confidential notes to the editor
- You can also annotate this PDF and upload it as part of your review

When ready submit online.

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your guidance page.

BASIC REPORTING

- Clear, unambiguous, professional English language used throughout.
- Intro & background to show context.
 Literature well referenced & relevant.
- Structure conforms to <u>PeerJ standards</u>, discipline norm, or improved for clarity.
- Figures are relevant, high quality, well labelled & described.
- Raw data supplied (see <u>PeerJ policy</u>).

EXPERIMENTAL DESIGN

- Original primary research within Scope of the journal.
- Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
- Rigorous investigation performed to a high technical & ethical standard.
- Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

- Impact and novelty not assessed.

 Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
- All underlying data have been provided; they are robust, statistically sound, & controlled.



Conclusions are well stated, linked to original research question & limited to supporting results.



Standout reviewing tips



The best reviewers use these techniques

Τ	p

Support criticisms with evidence from the text or from other sources

Give specific suggestions on how to improve the manuscript

Comment on language and grammar issues

Organize by importance of the issues, and number your points

Please provide constructive criticism, and avoid personal opinions

Comment on strengths (as well as weaknesses) of the manuscript

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Your introduction needs more detail. I suggest that you improve the description at lines 57-86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

- 1. Your most important issue
- 2. The next most important item
- 3. ...
- 4. The least important points

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.



Evaluation of sesame (Sesamum indicum L.) varieties for drought tolerance using agromorphological traits and drought tolerance indices

Getahun Yemata Corresp., 1, Tewachew Bekele 2

Corresponding Author: Getahun Yemata Email address: gyemata12@gmail.com

Sesame (Sesamum indicum L.) is an important cash crop cultivated under rain-fed condition and contributing a significant proportion of Ethiopia's foreign exchange earnings. However, its productivity is constrained by drought stress. The present study aimed to identify drought tolerant varieties using agromorphological traits and drought tolerance indices. The sesame varieties were evaluated under well watered (WW) and water stressed (WS) conditions in a field with a factorial design laid down in randomized complete block design in three replications. The results revealed a significant variation in agromorphological and drought tolerance indices due to water levels, varieties and interactive effect. On average, a 21.8, 49.6, 48.4, 47.9 and 21.7% reduction was recorded in plant height, number of leaves, leaf length, leaf width and relative growth rate (RGR), respectively under WS condition. Similarly, a significant reduction in shoot biomass, root biomass, biological yield, numbers of pods per plant and seed yield was found under WS condition in the study varieties. These traits showed an average reduction of 52.2, 72.5, 54.0, 51.9 and 52.8%, respectively as compared to WW condition. The highest yield reduction was recorded in wollega, while the lowest was in abasena. Wollega variety produced the highest seed yield ha⁻¹ under WW condition, while gondar-1 and humera-1 had the highest yield ha-1 under WS condition. Under both water levels, abasena produced the lowest yield ha⁻¹. Moreover, gondar-1 and humera-1 varieties had a comparatively higher values of stress tolerance index (STI), yield stress score index (YSSI), yield potential score index (YPSI), geometric mean productivity (GMP) and mean productivity (MP) that are significantly and positively correlated with yield under WS, indicating higher yield performance under water stress. The biplot analysis clustered the varieties as low yielding (abasena) and relatively above average performing varieties (humera-1, gondar-1 and wollega). According to the rank sum of all indices, humera-1 was identified as drought

PeerJ reviewing PDF | (2023:05:85584:0:0:CHECK 5 May 2023)

¹ College of Science, Department of Biology, Bahir Dar University, Bahir Dar, Ethiopia

² Biology, North Achefer District, Liben Senior Secondary and Preparatory School, Liben, Amhara National Regional State, Ethiopia



tolerant, while abasena as the most susceptible and low yielding varieties. Thus, humera-1 followed by gondar-1 are found to be drought tolerant and high yielding varieties. However, further studies focusing on drought tolerance mechanisms of the varieties are recommended.



- 1 Evaluation of sesame (Sesamum indicum L.) varieties for drought tolerance using
- 2 agromorphological traits and drought tolerance indices

4 Getahun Yemata¹ and Tewachew Bekele²

5

- 6 ¹College of Science, Department of Biology, Bahir Dar University, P. O. Box 79, Bahir Dar,
- 7 Ethiopia; ²Biology, North Achefer District, Liben Senior Secondary and Preparatory School,
- 8 Biology Department, Liben, Amhara National Regional State, Ethiopia

9

- 10 Corresponding author: gyemata12@gmail.com; Getahun.Yemata@bdu.edu.et
- 11 Abstract
- 12 Sesame (Sesamum indicum L.) is an important cash crop cultivated under rain-fed condition and
- contributing a significant proportion of Ethiopia's foreign exchange earnings. However, its
- productivity is constrained by drought stress. The present study aimed to identify drought
- tolerant varieties using agromorphological traits and drought tolerance indices. The sesame
- varieties were evaluated under well watered (WW) and water stressed (WS) conditions in a field
- 17 with a factorial design laid down in randomized complete block design in three replications. The
- 18 results revealed a significant variation in agromorphological and drought tolerance indices due to
- water levels, varieties and interactive effect. On average, a 21.8, 49.6, 48.4, 47.9 and 21.7%
- 20 reduction was recorded in plant height, number of leaves, leaf length, leaf width and relative
- 21 growth rate (RGR), respectively under WS condition. Similarly, a significant reduction in shoot
- biomass, root biomass, biological yield, numbers of pods per plant and seed yield was found
- 23 under WS condition in the study varieties. These traits showed an average reduction of 52.2,
- 72.5, 54.0, 51.9 and 52.8%, respectively as compared to WW condition. The highest yield



reduction was recorded in wollega, while the lowest was in abasena. Wollega variety produced 25 the highest seed yield ha-1 under WW condition, while gondar-1 and humera-1 had the highest 26 yield ha⁻¹ under WS condition. Under both water levels, abasena produced the lowest yield ha⁻¹. 27 Moreover, gondar-1 and humera-1 varieties had a comparatively higher values of stress 28 tolerance index (STI), yield stress score index (YSSI), yield potential score index (YPSI), 29 30 geometric mean productivity (GMP) and mean productivity (MP) that are significantly and positively correlated with yield under WS, indicating higher yield performance under water 31 stress. The biplot analysis clustered the varieties as low yielding (abasena) and relatively above 32 average performing varieties (humera-1, gondar-1 and wollega). According to the rank sum of 33 all indices, humera-1 was identified as drought tolerant, while abasena as the most susceptible 34 and low yielding varieties. Thus, humera-1 followed by gondar-1 are found to be drought 35 tolerant and high yielding varieties. However, further studies focusing on drought tolerance 36 mechanisms of the varieties are recommended. 37 **Keywords:** Abasena, drought tolerance indices, humera-1, gondar-1, wollega, stress tolerance 38 index, seed yield 39 Introduction 40 Sesame (Sesamum indicum L.) is an annual oil seed crop widely grown in arid and semiarid 41 42 tropical and sub-tropical regions (Weiss, 1983). It belongs to Pedaliaceae family and Sesamum genus. The genus consists of 20 species native to Africa and Asia (Bedigian, 2015). However, S. 43 indicum has been recognized as a cultivated species. It is one of the oldest and most traditional 44 oilseed crops, valued for its high-quality seed composed of 44–57% oil, 18–25% protein, 13– 45 14% carbohydrates (*Borchani et al.*, 2010). Evidences about the origin of sesame are debatable. 46



India (Alegbejo et al., 2003). 48 In Ethiopia, the production of sesame is rain-fed, characterized by intensive labour and low 49 levels of inputs (Coates et al., 2011). It grows in a wide variety of soil types. However, the crop 50 thrives best on well drained and medium textured fertile soil with pH range of 5 to 8. It also 51 needs adequate moisture for germination and early growth. Precipitation of 300-800 mm per 52 season is necessary for reasonable yields (*Terefe et al., 2012*). Sesame is temperature sensitive 53 that requires hot conditions during growth to produce maximum yields. It shows optimum 54 development and yield at 25 to 37°C temperature throughout its growth period. Generally, the 55 56 crop grows to a height of 1.5 to 2.0 m depending on the variety and growing conditions (Terefe et al., 2012). 57 Sesame is produced in different parts of Ethiopia starting from an elevation of 1500 meter above 58 sea level. The major producers that contribute over 83% to the national production (CSA, 2011), 59 are located in the regions of Tigray (West Tigray), Amhara (North Gondar) and most recently, in 60 Benishangul-Gumuz Region (Metekel). In the years 2005-2012, on average, almost 37% of the 61 country's total seed production is contributed by the Amhara Regional state, 30% from Tigray 62 and 16% from Oromia (CSA, 2013). Sesame production shows an increasing trend through the 63 years. In 2014/15 cropping season, 464 000 metric tons sesame seed was produced. This 64 65 increased to 487 000 metric tons in 2015/16. It showed 5% increment. The increment has been brought mainly by expansion of production area (Francom, 2016) due to the fast growing nature 66 of the oilseed sector in the country. It has become the second largest source of foreign exchange 67 earnings after coffee (FAO, 2012). 68

Nevertheless, most researchers claim that sesame was first cultivated in Africa and later taken to



69	The average yield of sesame is low in Ethiopia. For the years 2005-2012; the highest average
70	productivity of sesame for Tigray was about 9 quintals/hectare, followed by Amhara region
71	about 8 quintals/hectare. This is lower almost by half than the potential yield of the crop
72	estimated by FAO, which is 16 quintals/ha (FAO, 2015). According to Gelalcha (2009), the low
73	sesame productivity is attributed to a combination of various factors. The major constraints
74	include lack of improved seeds (Teklu et al., 2021), drought stress, low fertilizer input, biotic
75	stress, heat, indeterminate flowering nature and shattering of capsules at maturity and paucity of
76	knowledge on postharvest crop management practices (Endale, 2017). As a rainfed crop
77	commonly cultivated in arid and semiarid tropics, sesame is frequently exposed to terminal
78	drought (Pandey et al., 2021) and such exposure has reduced grain yield by 52% as compared to
79	the non-stressed ones (Kim, Park & Jenks, 2007; Golestani & Pakniyat, 2015). Moreover, the
80	rainfall distribution in the study area is significantly erratic that impedes the productivity of
81	sesame. The crop is sensitive to drought, especially at the vegetative stage (Boureima et al.,
82	2011). This is reflected in the changes that occur subsequently in plant metabolism, growth,
83	development and yield. However, the effect of drought is more severe on seed yield than other
84	morphological characters. According to Kim, Park & Jenks, (2007) sesame yield reduction owes
85	to the decreased number of seeds under drought stress. Research reports have revealed that
86	various sesame varieties show variable responses to drought, with some varieties being highly
87	tolerant and others more susceptible (Boureima et al., 2011).
88	Despite the high genetic diversity of sesame in the country; more than 870 accessions (Teshome,
89	Tesfaye & Bekele, 2015), studies focusing on evaluation of sesame genotypes under drought
90	stress conditions using agromorphological traits and drought tolerance indices are scarce.
91	Selecting genotypes with optimal performance under both stress and non-stress conditions from



97

- other groups is the best tool for identifying genotypes for drought tolerance (Fernandez, 1992).
- 93 Several such criteria have been proposed, most of which were used in this study to select the best
- 94 sesame varieties for drought tolerance and recommend suitable sesame varieties in drought
- 95 prone areas in the country.

MATERIAL and METHODS

Description of the experimental area

- 98 The experiment was conducted at Liben senior secondary and preparatory school found in
- 99 Amhara Region, western Gojjam zone, North Achefer district (Fig. 1). The school is located at
- 100 11°41'51"N latitude and 36°56' 35' east longitude. The area has arid and semi-arid climatic
- conditions with soil suitable for irrigation. The altitude of the district ranges 1500 to 1800 meter
- above sea level (m.a.s.l). It is also characterized by unimodal rainfall with an average annual
- rainfall ranging from 1000 to 1500mm. The minimum and maximum daily temperatures are
- 104 25°C and 30°C, respectively (NADoA, 2013 cited in *Demeke, Mekuriaw & Asmare, 2017*).

105 Study plant materials

- A total of four sesame varieties were used in this study. All of them were released varieties; T-
- 85, Kelafo-74, Mehado-80 and Abasena. The former two were released in 1976 and the later in
- 108 1989. They are the main export varieties and well known by their market names as humera-1,
- gondar-1, wollega and abasena, respectively. Then after, the market names are used. Humera-1 is
- characterized by whitish, large & sweet taste seeds, high productivity, high shattering property,
- 45-50% oil content, maturity time of 110-115 and adapted to the Humera plains (*Jelata, 2012*).
- Gondar-1 has light brown and good uniformity of seed, a maturity date ranging from 110-120
- days. It is released and adapted to the Gode region (*Endale, 2017*). Similarly, wollega is
- characterized by its small grey seeds, high oil content (49-56%), low sweetness, maturity date of



100-110 days and released and adapted to irrigated areas of Awash valley (*Jelata*, 2012; Endale, 115 2017). Abasena on the other hand is characterized by gray, large and sweet taste seeds, high 116 productivity, high shattering property, 44-48% oil content, maturity time of 110-120 days and 117 adapted to high rainfall (Bekora, 2013). Seeds of humera-1, gondar-1 and abasena were collected 118 from Amhara Agricultural Research Institute, Gondar Agricultural Center, and seed of wollega 119 120 variety was collected from local farmers around the study area. **Experimental Design and Treatments** 121 The experimental field was tilled using human power. All the weeds and large plant debris were 122 removed manually from the experimental field. Then, it was exposed to sunshine for 6 days 123 prior to the next tilling. The land ploughed again and again until smooth soil particles were 124 obtained as this is necessary for the better growth of the crop. The field was ploughed, leveled, 125 ridged and divided into plots before sowing. 126 127 The field experiment was carried out using factorial design laid down in a randomized complete 128 block design (RCBD) with three replications. The size of the experimental field was 39m² 129 (9.75m x 4m) consisting of 24 plots each with 1m² area, distributed in three replications that are 130 131 0.5m far apart. Each replication consisted of eight plots with 0.25m spacing to prevent lateral penetration of water. Each plot had three rows with 37cm spacing. There was 18.5cm spacing 132 between sesame plants. Around each plot, 13cm area was left to avoid edge effect. Four holes 133 134 were prepared along each row and three seeds were hand sown in each. The sesame varieties were assigned to plots randomly by lottery method. Thinning to one plant/hole was carried out 135 after complete emergence. N and P inorganic fertilizers were applied at the rate of 100kg/ha 136 137 (Zenawi & Mizan, 2019) and 46kg/ha (Gebremariam, 2015), respectively. All the P and half of the N fertilizers were applied at the time of sowing and the remaining half N at branching stage. 138



140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

Data collection

The common management practices such as weeding, pesticide and other parasites monitoring and control were carried out as needed. Treatments consisted of four sesame varieties grown under two water levels. The water levels were determined gravimetrically as described by *Turner* (2018). Bottom perforated pots filled with dry soil were weighed and then flooded with water for adequate period of time. The saturated soil was left overnight to drain the excess water by the force of gravity and then weighed. At this condition, the soil water is at its field capacity equivalent to pot capacity. The water level at field capacity was considered as well watered and 50% of the field capacity was taken as water stress treatments. The water level of all treatments was maintained to field capacity until the imposition of water stress. Water stress was imposed 30 days after sowing (DAS). Sesame plants grown under well watered treatment in each plot were irrigated with 333ml of water for every three days and those in the water deficit stressed received half of this volume at the same time interval. Soil analysis Soil samples representative of the experimental field were randomly collected from different points and thoroughly mixed together to form a composite soil sample. The sample was dried, ground and passed through a 2mm sieve and used for further soil property analysis. The pH was determined in 1:1 soil/water ratio using pH meter. The total percent nitrogen was determined by Kjeldahl digestion method, organic carbon content and organic matter were measured by volumetric method as described by Walkley & Black, (1934). The analysis result revealed that the soil had a pH of 6.5, total percent nitrogen (0.084), percent organic carbon content (0.49) and organic matter (0.85).





162	Growth and physiological data of the study species were gathered from three randomly selected
163	and tagged plants in each plot. When the tagged plants are lost due to various reasons, data were
164	collected from other plants in the same plot. Data of both types were collected 30 and 60 days
<mark>165</mark>	after water deficit stress imposition.
166	Growth parameters
167	Plant height of tagged plants was measured from the ground level to the tip of the youngest leaf
168	and expressed in centimeter (cm). Number of leaves and branches of the same plants were
169	counted and recorded. The length of leaves was measured from the leaf base to the tip and width
170	at the maximum area of the blade. Relative growth rate (RGR) was assessed based on height and
171	determined according to <i>Hunt (1990)</i> as $RGR_H = (logH_{t2} - logH_{t1})/t_2 - t_1$ where, H_{t1} is height of
172	the plant measured at t_1 and H_{t2} is height of the plant measured at t_2 , t_1 and t_2 is time of the first
173	and second height measurements, respectively.
174	Physiological parameters
175	Relative water content (RWC)
176	Relative water content was estimated according to Barrs & Weatherly (1962b). Fully expanded
<u>177</u>	leaves from tagged plants were collected and weighed, producing fresh weight (FW). These
178	leaves were preserved in plastic bags and transported to laboratory. Each leaf was floated on
179	distilled water in a Petri dish for 24 hrs. The leaves were then blotted gently with tissue paper
180	and weighed again to produce turgid weight (TW). The leaves were then oven dried at 70°Cs for
181	24 hrs and dry weight (DW) were recorded. Finally, leaf relative water content was calculated as
182	(FW-DW) / (TW-DW)*100.
183	
184	



Yield and Biomass parameters

The number of days required to 50% flowering was noted when 50% of the plants in each plot flowered and expressed in days. The average number was recorded and used for description. The numbers of capsules/pods per plant in each plot were counted from three tagged plants and recorded during data collection periods. All the pods from each plant and plot were collected and dried in an oven at 80°C to constant weight and threshed plot wise. Grain yield was determined for each plot (1m²) and reported in terms of kg/ha as described by *Nadeem et al.*, (2015).

After harvesting the pods, each plant was uprooted and parts were separated into shoot (stem and root) and root. The roots were carefully detached from soils and thoroughly washed with tap water. The shoot and root parts were dried in an oven at 80°C to constant weight and weighed

separately. Harvest index was calculated from biological yield (shoot and root dry weight) and

Harvest index =
$$\frac{\text{Seed yield}}{\text{Biological yield}}$$
 X 100

grain yield in kg/ha according to *Nadeem et al.*, (2015) as follows:

Drought tolerance indices

Drought tolerance indices are important tools that provide better opportunities to select genotypes with good performance under normal and stress conditions. Several of the tolerance indices were calculated as follow:





```
209
Yield stability index (YSI) = Ys/Yp ......Bouslama & Schapaugh, (1984)
           Stress susceptibility index (SSI) = 1-(Ys/Yp)/1- (\bar{Y}s/\bar{Y}p) ... Fischer & Maurer, (1978)
     212
            213
                     Yield potential score index (YPSI) = 0.5 \text{ (MP + STI)} - 0.5 \text{ (SSI + TOL)} \dots Malinowska et al.
           Where,
     214
                     2020)
           Ys is the yield of each genotype under stress; Yp is the yield of each genotype under non-stress
     215
           condition; \bar{Y}s is the mean yield of each genotype under stress and \bar{Y}p is the mean
     216
           yield each genotype under non-stress condition.
     217
     218
           Ranking of sesame varieties
     219
     220
           Different drought tolerance indices discriminate different sesame varieties as drought resistant;
     221
           hence identifying drought tolerant genotypes based on a single index does not produce clear
     222
     223
           results. To identify desirable drought tolerant varieties, the mean rank, standard deviation of
           ranks, and rank sum of all indices were calculated. For screening drought tolerant varieties a rank
     224
           sum (RS) was calculated by using the following relationship formula:
     225
     226
     227
```





Rank Sum = Mean ranks (MR) + standard deviation of ranks (SDR)...Noorifarjam, Farshadfar,

Saeidi, (2013)

228 Data analysis

All the data were analyzed using Statistical Package for Social Sciences (SPSS, version 23).

Multiple comparisons of means were carried out with Tukey HSD test to see variations between

treatments at 30 and 60 days after drought imposition. Factorial Analysis of variance (ANOVA)

was conducted for each trait under WW and WS conditions and results were considered

significant at p < 0.05. Furthermore, principal component analysis (PCA) was used to

characterize trait variation and correlation analysis between grain yield and drought tolerance

indices were analyzed with R software. The biplot was generated using factor analysis and data

processing with R package.

237238

239

240

241

242

243

244

245

246

247

248

230

231

232

233

234

235

236

RESULTS

The sesame varieties showed notable variation among most of the traits between WW and WS conditions. The analysis of variance of the data from the field experiment showed that the differences between treatments and varieties were statistically significant at P < 0.05, 60 days after water stress imposition (Table 1). In this period, plant height was reduced under WS by 23.2, 27.6, 28.9 and 27.4% in abasena, gondar-1, humera-1 and wollega sesame varieties, respectively. The reduction in RGR was nearly similar to the reduction in plant height for all the varieties as it was derived from plant height. Greater reduction was observed in number of leaves, leaf length and leaf width. Accordingly, the number of leaves was reduced by 48.2, 51.2, 48.3 and 50.8 under WS in abasena, gondar-1, humera-1 and wollega varieties. Almost a similar pattern was recorded for leaf length and leaf width (Table 1). The growth of branches was





leaf RWC was observed between plants under WW and WS conditions in all varieties, 250 251 indicating that leaf RWC is not an important parameter for screening sesame varieties for drought tolerance. 252 253 The results also demonstrated a highly significant difference among varieties under the different 254 255 water levels, 60 days after water stress imposition at P < 0.05. Until 30 days, the effect of WS on some growth parameters was not visible. A significant difference was recorded in number of 256 leaves between abasena (6.89) and humera-1 (8.00) varieties under WS condition at 30 days after 257 258 water stress imposition (Table 1). Plants of gondar-1 variety performed the highest in terms of number of leaves, leaf width, and plant height under WW condition at 30 and 60 days after water 259 stress imposition. This difference was statistically significant at P < 0.05 (Table 1). Although 260 261 insignificant, higher numbers of branches were recorded in wollega variety under WW conditions, 60 days after stress imposition (Table 1), which might have implication to yield. 262 Plants of gondar-1 had also significantly higher number of leaves than abasena under WS 263 264 condition. It also showed considerably faster RGR than abasena under WW condition. The average reduction in plant height, number of leaves, leaf length, leaf width and RGR under WS 265 was found to be 21.8, 49.6, 48.4, 47.9 and 21.7%, respectively, 60 days after water stress 266 imposition. In general, gondar-1 and abasena sesame varieties performed the best and least, 267 respectively under the present study experimental conditions (WW and WS). Plants of humera-1 268 and wollega showed moderate growth performance under both water levels (Table 1). 269 270 Plant height showed positive and significant correlation with RGR in all sesame varieties both 271 under WW and WS conditions at P < 0.01. Number of leaves and RWC had higher positive 272

completely suppressed under WS condition in all sesame varieties. An insignificant difference in



274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

correlation with yield in abasena variety under WW conditions (Data not shown). In contrast, almost all the parameters had negative correlation with yield under WS condition. In gondar-1 and humera-1 sesame varieties, almost all growth and physiological parameters had negative correlation with yield both under WW and WS conditions except RWC under WS conditions in humera-1 variety. On the other hand, plant height and RGR had a significantly positive correlation with yield in wollega variety under WS condition (Data not shown). A significant variation in plant height, number of branches, number of leaves and RGR was recorded due to the separate effect of variety and water levels, 60 days after water stress imposition (Table 2). According to the analysis of the result, variety had no significant influence on the leaf length and width of the studied varieties. The water levels had greater and significant effect size than variety. In this regard, the highest reduction in plant height and RGR under WS was recorded by humera-1 variety followed by gondar-1 and wollega, while the lowest reduction was recorded in abasena variety. The difference in reduction due to WS in plant height, number of leaves, leaf length and width ranged from 23.2-27.4, 48.2-51.2, 43.9-53.9 and 45.4-49.8%, respectively (Table 1). The biggest reduction in number of leaves and leaf length was observed in gondar-1 variety followed by wollega. Similarly, humera-1 and wollega varieties showed the biggest reduction in plant height and leaf width under WS condition (Table 1). In the same period, the interactive effect of variety and water levels on growth parameters was insignificant except number of branches where the synergistic effect of the two independent variables was greater (Table 2).

Mean performance of sesame varieties

The sesame varieties demonstrated a statistically significant variation in biomass production, yield and yield related traits under the two water levels (WW and WS) at P < 0.05. A significant





297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

yield was recorded under WS condition among sesame varieties. These traits showed an average reduction of 52.2, 72.5, 54.0, 51.9, and 52.8%, respectively as compared to WW condition. The highest reduction in yield and yield related traits between WW and WS conditions was found in wollega followed by humera-1, while the lowest was found in abasena followed by gondar-1 (Table 3). There was insignificant difference in shoot biomass, harvest index and seed yield between water levels in abasena variety (Table 3). Significant differences were also recorded among varieties both under similar and different water levels (Table 3). Accordingly, wollega, gondar-1 and humera-1 varieties produced considerably greater shoot biomass and biological yield than abasena under WW condition. The variation in shoot biomass between the varieties under WS condition was insignificant at P < 0.05. All varieties under WS had significantly lower root biomass than those in WW condition, implying that enhanced root growth and the consequent increased root/shoot ratio is not a drought tolerance strategy in the study sesame varieties. In this regard, abasena variety had a relatively greater root biomass than the others. Sesame plants of wollega variety produced a relatively higher biological yield followed by gondar-1 both under WW and WS conditions than others (Table 3 and 4). In terms of yield and related traits, gondar-1 variety had greater number of pods per plant followed by humera-1 and wollega varieties under WW and WS conditions (Table 3 and 4). Wollega sesame variety produced a significantly greater seed yield per cm² under WW condition followed by humera-1 and gondar-1 varieties. Correspondingly, variety gondar-1 had better seed

yield than others followed by humera-1 under WS condition. However, the variation among the

varieties in seed yield/cm² under WS condition was insignificant. Wollega sesame variety had

higher harvest index under WW condition attributed to the highest seed yield/cm², while a

reduction in shoot biomass, root biomass, biological yield, number of pods per plant and seed



relatively higher harvest index was recorded under WS condition in abasena and gondar-1 varieties which might be due to the lower biological yield caused by the WS condition. Plants of abasena performed the least in most parameters under both water levels (Table 3 and 4).

When variety, water levels and their interactive effects are analyzed, they produced a statistically significant difference in biological yield, number of pods per plant, Ys, Yp, shoot and root biomass at P < 0.01, 60 days after water stress imposition (Table 4). Harvest index was not significantly affected by variety and water levels. Similarly, the interactive effect (variety * water levels) had no remarkable effect on number of pods per plant. The partial eta squared value revealed that water levels had greater effect on Ys, Yp, biological yield, number of pods per plant, shoot and root biomass followed by variety (Table 4).

Drought tolerance indices

Analysis of variance of Ys, Yp and drought tolerance indices showed a highly significant difference among sesame varieties under the different water levels, indicating the presence of high genetic variability. Drought tolerance indices were calculated on the basis of seed yield ha⁻¹ of 4 sesame varieties from Ys and Yp. The mean seed yield under WW conditions ranged from 2144.44 to 5177.78 kg/ha, while 1344.44 to 1811.11 kg/ha was obtained under water stressed condition. Wollega sesame variety had the highest and the second lowest mean seed yield/ha⁻¹ under WW and WS conditions, respectively (Fig. 2). Gondar-1 and humera-1 varieties achieved the highest seed yield/ha under WS condition. In both water levels, plants of abasena performed the least and the variation was found to be significantly lower than other sesame varieties (Fig. 2).



344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

The study sesame varieties showed a significant difference in drought tolerance indices (Table 5). Wollega sesame variety had significantly higher MP, GMP and TOL than others followed by humera-1 variety. According to these indices, wollega and humera-1 varieties were found more desirable and comparatively drought tolerant. Unlike this, wollega variety had lower STI than gondar-1 and humera-1 varieties (Table 5). The mean STI values demonstrated that gondar-1 and humera-1 have better drought tolerance characteristics. In the present study, abasena sesame variety had the lowest TOL, SSI and the highest YI and YSI of all the study varieties (Table 5). The difference between Ys and Yp values in abasena variety was the lowest indicating the lower sensitivity of the variety to drought stress. Varieties with lower TOL, SSI, higher YI and YSI values are more stable and tolerant, respectively under drought conditions. Based on these indices thus, abasena could be identified as stable and tolerant variety. Plants of wollega variety demonstrated a significantly higher percent reduction than abasena that showed the lowest reduction. The studied varieties revealed a significantly different stress intensity of which the weakest was seen in abasena variety. Similarly, plants of gondar-1 and humera-1 had higher YSSI and YPSI values as compared to others (Table 5). Correlation A strong association between Yp, Ys and drought tolerance indices is an important indicator to select the most desirable drought tolerance indices. Ys had positive and significant correlation with Yp. Moreover, Ys revealed a significantly positive correlation with MP, STI, GMP, YSSI and YPSI at P < 0.01 (Table 6). Similarly, Yp had significantly positive association with MP, STI, GMP, TOL, PR, SI and YSSI at P < 0.01 probability level (Table 6). This showed that MP, STI, GMP and YSSI had significantly higher correlation with yield under WW and WS

conditions so that these indices can be used as criteria to select varieties for drought tolerance.



According to MP and GMP values, wollega and humera-1 were found to be the most desirable 366 and drought tolerant varieties, respectively, while gondar-1 and humera-1 had higher STI and 367 YSSI indicating higher yield performance under drought conditions. Furthermore, TOL had 368 positive and significant correlation with MP, GMP, PR, SSI and SI at P < 0.01. Its correlation 369 with YI, YSI and YPSI was negative and significant (Table 6). 370 371 **Principal Component Analysis** The minimum number of principal components that account for most of the variations in drought 372 tolerance indices was determined based on eigen value. Principal components with the highest 373 eigen values greater than 1 were selected. Accordingly, two principal components fulfilled the 374 acceptable level/variance of the dataset. The first and second principal components (PC1) and 375 (PC2) explained 86.2 % and 12.7 % of the variation, respectively. The PC1 and PC2 together 376 cover 98.9 % of the variation in the dataset (Table 7). The results of the analysis showed that 377 PC1 had large positive association with Yp, TOL, MP and GMP. Others such as YI, YSI and 378 YPSI had negative correlation with PC1. PC2 had large positive association with YPSI, Ys, STI, 379 GMP, MP and YSSI, while negatively correlated with TOL (Table 7). The large positive and 380 negative values of drought tolerance indices indicate the strong effect of each index on each 381 382 principal component. The biplot analysis showed that Yp, TOL, GMP, MP, Ys and YSSI orient in similar direction 383 and had tight angles indicating a positive correlation and strong effect on PC1. Similarly, YPSI, 384 385 Ys, STI and GMP contributed higher proportion to PC2 (Fig. 3). The biplot analysis clustered the study varieties roughly into two groups as low yielding (abasena) and relatively above average 386 387 performing varieties (humera-1, gondar-1 and wollega) under well watered and stressed 388 conditions (Fig. 3). This was corroborated with the values of STI, MP and GMP where the



varieties in the second group had higher average yield values under well watered and water stressed conditions (Table 7 and Fig. 3).

Ranking of sesame varieties

Rank sum was calculated due to the fact that identification of genotypes for drought tolerance using individual indices is difficult. Different indices identified different sesame varieties for drought tolerance. Rank sum for all indices was used as an indicator to select the best varieties. Lower and higher rank sum values indicate high drought tolerant and susceptible genotypes, respectively. Accordingly, humera-1 sesame variety had the lowest rank sum value followed by gondar-1 variety. These varieties represent the best varieties with the highest performance under well watered and water deficit stressed conditions (Table 8). The rank sum result identified abasena sesame variety as the most susceptible and low yielding variety (Table 8).

DISCUSSION

The results of the present study showed that the different agromorphological traits significantly vary among varieties due to water stress and genetic variability. The results revealed that water stress had a remarkable influence on growth parameters specifically plant height, leaf number and number of branches unlike physiological processes. The growth of branches was completely suppressed under WS condition in all varieties. Humera-1 and wollega varieties showed the highest reduction in growth parammeters. Of the sesame varieties, gondar-1 outperformed in agromorphological traits, while abasena was the lowest. This is in line with the findings of *Hassen, (2022), Mewcha et al., (2020), Mekonnen and Sintayehu, (2020)* and *Gebremichael & Parzies,* (2011). *Hassen (2022)* has reported the high heritability of plant height and harvest index of 100 sesame genotypes at Amibara, Ethiopia. Furthermore, nearly similar results of leaf





length and number of branches have been found in gondar-1, humera-1, wollega and argene sesame varieties (*Mewcha et al., 2020*). However, these and other varieties have shown a significantly higher plant height and number of leaves (*Mekonnen & Sintayehu, 2020; Mewcha et al., 2020*). This deviation might be caused by the difference in the type of irrigation system, time of drought imposition, agroecology of the experimental sites that vary in soil fertility and other environmental factors. Furthermore, findings have indicated that the agromorphological performance of sesame varieties varies across seasons (*Hailu et al., 2018; Hassen, 2020; Baraki et al., 2020*).

The data in the present study demonstrated a significant reduction in yield and yield related traits under water stress. The highest reduction was recorded from wollega variety followed by humera-1. Gondar-1 had greater number of pods per plant and yield under WS condition followed by humera-1. Wollega variety had greater seed yield under WW condition. Almost similar, lower and higher yield performances of sesame varieties have been reported so far. According to *Mawcha et al.* (2020), gondar-1, humera-1 and wollega sesame varieties have produced seed yield of 3432.09, 2194.44 and 2377.78kg/ha, respectively, at Humera under supplementary irrigation; a common sesame production area, which is almost similar to the present study (Fig. 2). On the other hand, gondar-1 and humera-1 varieties produced significantly higher yield in the present study (Fig 2) than the same varieties at Dansha grown under normal moisture conditions (588 kg/ha and 542 kg/ha, respectively). This higher yield deviation might be attributed to the high prevalence of bacterial blight disease, as the area is a hotspot for the disease (Golla, Kebede & kindeya, 2020). A higher seed yield of abasena variety was obtained under WW condition (Figure 2) in the present study as compared to Mekonnen and



435	Sintayehu, (2020). The authors have reported 1840 kg/ha and 670 kg/ha seed yield under uniform
436	optimum irrigation and 50% uniform deficit irrigation, respectively at Metema; another hub of
437	sesame production. The yield of the variety under WS condition (Fig. 2) has been lower than
438	75% water deficit imposed at the development stage, 1785 kg/ha (Mekonnen and Sintayehu,
439	2020). Comparatively, close yield performance as to the present study has been found in adi
440	sesame variety treated with 100, 75 and 50% of the evapotranspiration of the crop applied
441	through convention furrow irrigation method in 2015, at Werer research center (Hailu et al.,
442	2018). On the other hand, the study varieties produced significantly greater seed yield than
443	several other varieties and accessions such as serkamo white, adi, acc-00048, acc-00016, acc-
444	00025, acc-00049 and others that have bee evaluated at Werer (Afar Region), Bonta (Afar
445	Region) and Miesso (Oromia) under normal growth conditions (Hassen, 2022). This is
446	corroborated by Bakari et al., (2020) and found to be due to environmental and genetic variation
447	among the sesame varieties. The authors have pointed out that about 42.62, 6.22 and 25.09% of
448	sesame agronomic performance are determined by the environment, genotype and their
449	interactive effect, respectively (Baraki et al., 2020).
450	Selection of crop varieties for drought tolerance drought tolerance indices based on yield under
451	normal and stressed conditions is one of the tools widely used in agriculture. In this study,
452	wollega sesame variety had the highest MP, GMP and TOL values, which agrees with the
453	findings of Fernandez (1992) and Farshadfar, Jamshidi & Aghaee (2012). The highest TOL
454	value indicates the high sensitivity of the variety to water stress. This was evident from the
455	highest percent reduction under water stressed condition (Table 2). On the other hand, gondar-1
456	and humera-1 varieties had higher STI values showing better performance than other varieties.
457	Several studies such as Fernandez (1992), Pireilvatlou, Masjedlou & Aliyou, (2010), Farshadfar,



Jamshidi & Aghaee (2012), Zare (2012), Noorifarjam, Farshadfar, Saeidi, (2013) and Baghery 458 et al. (2022) have identified STI as a reliable index in mungbean, wheat, wheat, barley, wheat 459 and sesame, respectively, for selecting varieties with high drought tolerance and yield under 460 normal and water stressed conditions. This might due to the fact that STI considers potential 461 yield under normal condition, yield under stressful environments and stress intensity (Fernandez, 462 463 1992). Variety abasena had the lowest TOL and SSI and the highest YI and YSI as compared to the other varieties. The TOL and SSI values for the variety showed the lower yield potential 464 under non-stress condition and higher yield under stressed condition, evidently proved by the 465 lowest percent reduction under the two contrasting conditions (Table 2). The variety also 466 characterized by a closer Yp and Ys, implying the lower sensitivity of the variety to water stress 467 which subsequently results in smaller SSI. This is consistent with the findings of Fernandez, 468 (1992) in mungbean, Zare, (2012) in barley, and Baghery et al. (2022) in sesame varieties. 469 The best drought tolerance index is the one that has discernable association with yield under 470 normal and water stress conditions. In the present study, MP, GMP and STI had a significantly 471 higher positive correlation with Yp and Ys. This shows that these indices are effective in 472 identifying varieties under different water stress conditions as supported by the findings of 473 Siahsar, Ganjali & Allahdo, (2010) in lentil. Moreover, Yp had significantly higher positive 474 correlation with TOL, SI and YSSI and, Ys with YSSI and YPSI. These findings are in line with 475 the reports of Zare, (2012), Farshadfar, Jamshidi & Aghaee, (2012), Noorifarjam, Farshadfar, 476 Saedid, (2013), Baghery et al. (2022) and Sun et al. (2023). Drought tolerance Indices correlated 477 with both Yp and Ys have been found suitable for the selection of varieties for water stress 478 (Baghery et al., 2022), indicating increase in yield under water stressed and normal conditions 479 Farshadfar, Jamshidi & Aghaee, (2012). The result of the PCA also revealed that only PC1 and 480



482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

PC2 with eigen value greater than 1 explained 98.9% of the variation, which agrees with the reports of Zare, (2012) in barley and Baghery et al. (2022). According to the analysis, Yp, TOL, MP and GMP had a relatively strong effect on PC1 indicating the high yield potential of indices and PC2 strongly associated to YPSI, Ys, STI and GMP, predictors of drought tolerance. Similar results have been reported in barley (Zare, 2012), sesame (Baghery et al., 2022) and cotton (Sun et al., 2023). Furthermore, the biplot analysis identified two categories of sesame varieties; low yielding (abasena) and above average performing varieties such gondar-1, humera-1 and Wollega (Figure 3). This corresponds with the categorization of Fernadez, (1992) and Sofi et al. (2018) where abasena roughly belongs to group D (poor yield performance under WW and WS conditions), Wollega to group B (good performance under WW, not under WS conditions) and gondar-1 and humera-1 varieties to group A (relatively higher performance in both WW and WS conditions. This might be related to the agroecology where the varieties are released that is characterized by high temperature and transpiration that lead to water stress in the plant. Since identifying the most drought tolerant variety using the indices is difficult, the rank sum and standard deviation of ranks of all indices were calculated. Accordingly, humera-1 followed by gondar-1 were identified as drought tolerant and high yielding, while abasena as the most susceptible and low yielding variety. Similar findings for other species and varieties have been reported by Noorifarjam, Farshadfar, Saeidi, (2013) and Anter & Ashraf, (2018).

CONCLUSION

The present study aimed to identify drought tolerant sesame varieties using agromorphological and drought tolerance indices. Our findings showed a significant variation in agromorphological, seed yield and drought tolerance indices due to water levels, variety and their interaction.

However, the effect of water levels was stronger than others. In support of this, a significant



504	reduction was observed in growth and yield parameters under water stressed conditions. The
505	highest reduction was recorded in wollega. Abasena performed the lowest under both water
506	levels. Humera-1 and gondar-1 varieties performed better both under water levels. They also had
507	higher stress tolerance index, yield stress score index, yield potential score index, geometric
508	mean productivity and mean productivity, implying better tolerance to water stress. In general,
509	humera-1 followed by gondar-1 are identified as high yielding and drought tolerant varieties
510	based on drought tolerance indices, biplot analysis and rank sum of all drought tolerance indices.
511	REFERENCES
512 513	Alegbejo MD, Iwo GA, Abo ME, Idowu AA. 2003. Sesame: A potential industrial and export oilseed crop in Nigeria. <i>J. Sustain Agric</i> . 23:59-76.
514 515	Anter AS, Ashraf AA. 2018. Evaluation performance of new sesame (Sesame indicum L.) lines under normal and drought conditions. <i>Middle East J. Agric. Res.</i> 7(4):1411-1418.
516	Baghery MA, Kazemitabar SK, Dehestani A, Mehrabanjoubani P, Zarrini HN. 2022. Assessment
517	of agro-morphological traits and yield-based tolerance indices in sesame (Sesamum indicum
518	L.) genotypes under drought stress. Indian J. Genet. 82(3): 324-332doi:
519	10.31742/ISGPB.82.3.7.
520	Baraki F, Gebregergis Z, Belay Y, Berhe M, Teame G, Hassen M, Gebremedhin Z, Abadi A,
521	Negash W, Araya G. 2020. Multivariate analysis for yield and yield-related traits of sesame
522	(Sesamum indicum L.) genotypes. Heliyon. https://doi.org/10.1016/j.heliyon.2020.e05295 .
523	Barrs H, Weatherly P. 1962b. Physiological indices for high yield potential in Wheat. J. P.
524	Physio. 25:352-35.
525	Bedigian D. 2015. Systematics and evolution in Sesamum L. (Pedaliaceae), part 1: Evidence
526	regarding the origin of sesame and its closest relatives. Journal of Plant Taxonomy and
527	Geography 70: 1–42.
528	Bekora G. 2013. Oil Seed Production and Marketing Situation in Ethiopia. Ethiopian Ministry of
529	Trade, Addis Abeba.
530	Borchani C, Besbes S, Blecker CH, Attia H. 2010. Chemical characteristics and oxidative
531	stability of sesame seed sesame paste, and olive oils. I. Agri. Sci. Tech. 12: 585–596



- Boureima S, Eyletters M, Diouf M, Diop TA, Van Damme P. 2011. Sensitivity of seed
- germination and seedling radicle growth to drought stress in sesame (*Sesamum indicum* L.).
- 534 *Res. J. Environ. Sci.* 5: 557–564.
- Bouslama M, Schapaugh WT. 1984. Stress tolerance in soybean. Part 1: Evaluation of three
- screening techniques for heat and drought tolerance. *Crop Sci.* 24: 933-937.
- 537 Choukan R, Taherkhani T, Ghannadha MR, Khodarahmi M. 2006. Evaluation of drought
- tolerance in grain maize inbred lines using drought tolerance indices. *Iran J. Agric. Sci.*
- 539 8(1):79-89.
- Coates M, Kitchen R, Kebbell G, Vignon C, Guillemain R, Hofmeister R. 2011. Financing
- agricultural value chains in Africa. Focus on coffee and sesame in Ethiopia. GIZ Bonn and
- Eschborn, Germany.
- 543 CSA. 2011. Agricultural Sample Survey 2010/2011. Area and production of crops. Central
- Statistical Agency of Ethiopia, Addis Ababa.
- 545 CSA. 2013. Agricultural Sample Survey 2010/2011. Area and production of crops. Central
- Statistical Agency of Ethiopia, Addis Ababa.
- Demeke S, Mekuriaw Y, Asmare B. (2017). Assessment of livestock production system and
- feed balance in watersheds of North Achefer District, Ethiopia. *Journal of Agriculture and*
- *Environment for International Development.* 111 (1): 175-190. DOI:
- 550 10.12895/jaeid.20171.574.
- Endale D. 2017. Sesame (Sesamum indicum L.) breeding in Ethiopia. International Journal of
- Novel Res. in Life Sci. 4(1): 1-11. www.noveltyjournals.com
- 553 FAO. 2012. Food and Agriculture Organization. FAOSTAT Database. Available at
- 554 http://faostat.fao.org/site/567/ default.
- FAO. 2015. Analysis of price incentives for Sesame seed in Ethiopia, 2005-2012. Technical
- notes series, MAFAP, by Kuma Worako, T., MasAparisi, A., Lanos, B., Rome.
- 557 Farshadfar E, Jamshidi B, Aghaee M. 2012. Biplot analysis of drought tolerance indicators in
- bread wheat lanraces of Iran. *Int. J. Agri. Crop Sci.* 4(5):226-233.
- Fernandez GCJ. 1992. Effective selection criteria for assessing plant stress tolerance. In:
- Proceeding of the international symposium on adaptation of vegetables and other food crops
- to temperature and water stress. AVRDC Shanhua publication, Taiwan, Pp. 257-270.



- Fischer RA, Maurer R. 1978. Drought resistance in spring wheat cultivars.I: Grain yield
- response. *Austrian Journal of Agricultural Research* 29:897-912.
- Francom MG. 2016. Ethiopia's oilseed production forecast to increase despite drought. GAIN
- Report; Global Agricultural Information Network, USDA Agricultural service, Addis Ababa.
- Gavuzzi P, Rizza F, Palumbo M, Campaline RG, Ricciardi GL, Borghi B. 1997. Evaluation of
- field and laboratory predictors of drought and heat tolerance in winter cereals. Can J. Plant
- 568 *Sci.* 77: 523-531.
- Gebremariam G. 2015. Growth, yield and yield component of Sesame (Sesamum indicum L.) as
- affected by timing of nitrogen application. J. Biol. Agric. and Healthcare. 5(5): 165-169.
- 571 Gebremichael DE, Parzies HK. 2011. Genetic variability among landraces of sesame in
- 572 Ethiopia. *Afri. Crop Sci. J.* 19(1):1-13.
- 573 Gelalcha SD. 2009. Sesame trade arrangements, costs and risks in Ethiopia. Ministry of Foreign
- 574 Affairs, The Haque.
- 575 Golestani M, Pakniyat H. 2015. evaluation of traits related to drought stress in sesame (Sesamum
- 576 *indicum* L.) genotypes. *J. Asian Sci Res.* 5(9): 465-472.
- 577 Golla WN, Kebede AA, Kindeya YB. 2020. Evaluation of sesame genotypes for seed yield and
- bacterial blight (*Xanthomonas campestris pv. sesami*) disease resistance in optimum moisture
- areas of Western Tigray, Ethiopia. Cogent Food & Agriculture 6:1, 1771114, DOI:
- 580 10.1080/23311932.2020.1771114.
- Hailu EK, Urga YD, Sori NA, Borona FR, Tufa KN. 2018. Sesame yield responses to deficit
- irrigation and water application techniques in irrigated agriculture, Ethiopia. *International*
- Journal of Agronomy. https://doi.org/10.1155/2018/5084056.
- Hassen M. 2022. Morphological characteristics of sesame (Sesamum indicum L.) genotypes via
- genetic diversity and characters association in Amibara, Ethiopia. *Ecology and Evolutionary*
- 586 *Biology*. 7(3):30-45. doi: 10.11648/j.eeb.20220703.11.
- 587 Hassen M. 2022. Performance evaluation and GGE biplot analysis of Sesame (Sesamum indicum
- L.) genotypes in Eastern Ethiopia. https://doi.org/10.21203/rs.3.rs-1523137/v1.
- Hunt R. 1990. Basic Growth Analysis: Plant Growth Analysis for Beginners. London, Unwin
- 590 Hyman.
- Jaleta Z. 2012. Sesame (Sesame indicum L.) crop production in Ethiopia: Trends, challenges and
- future prospect. *Science, Technology and Arts Res. J.* 1(3): 1-7.



- Kim KS, Park SH, Jenks MA. 2007. Changes in leaf cuticular waxes of sesame (Sesamum
- indicum L.) plants exposed to water deficit. J. Plant Physiol. 164: 1134–1143.
- 595 Malinowska M, Donnison I, Robson P. 2020. Morphological and physiological traits that explain
- yield response to drought stress in Miscanthus. *Agronomy* 10: DOI:
- 597 10.3390/agronomy10081194.
- 598 Mawcha KT, Gebru MM, Mewael K, Assefa MK, Gebre GG. 2020. Morphological
- characterization and genetic diversity of sesame (Sesamum indicum L.) varieties
- cultivated in Ethiopia. *The Open Agric. J.* 14: 117-129. DOI:
- 601 10.2174/1874331501014010117.
- Mekonnen SA, Sintayehu A. 2020. Performance evaluation of sesame under regulated deficit
- 603 irrigation application in the low land of Western Gondar, Ethiopia. *Int. J. Agronomy*.
- 604 https://doi.org/10.1155/2020/3760349.
- Nadeem A, Kashani S, Ahmed N, Buriro M, Saeed Z, Mohammad F, Ahmed S. 2015. Growth
- and yield of Sesame (Sesamum indicum L.) under the influence of planting geometry and
- irrigation regimes. *American Journal of Plant Sciences*. 6:980-986.
- 608 http://dx.doi.org/10.4236/ajps.2015.67104
- Noorifarjam S, Farshadfar E, Saeidi M. 2013. Evaluation of drought tolerant genotypes in bread
- wheat using yield based screening techniques. Euro. J. Exp. Bio., 2013, 3(1):138-143.
- Pandey BB, Ratnakumar P, Kiran BU, Dudhe, MY, Lakshmi GS, Ramesh K, Guhey A. 2021.
- Identifying Traits Associated With Terminal Drought Tolerance in Sesame (Sesamum
- 613 indicum L.) Genotypes. *Front. Plant Sci.* DOI:10.3389/fpls.2021.739896
- 614 Pireivatlou AS, Masjedlou BD, Aliyev RT. 2010. Evaluation of yield potential and stress
- adaptive trait in wheat genotypes under post anthesis drought stress conditions. Afric J of
- 616 Agric Res. 5: 2829-2836.
- Rosielle AA, Hamblin J. 1981. Theoretical aspects of selection for yield in stress and nonstress
- environment. *Crop Sci.* 21: 493.
- 619 Siahsar BA, Ganjali BAS, Allahdoo M. 2010. Evaluation of drought tolerance indices and their
- relationship with grain yield of lentil lines in drought stressed and irrigated environments.
- 621 Aust. J. Basic & Appl. Sci. 4(9): 4336-4346.



- Sofi PA, Rehman K, Ara A, Gull M. 2018. Stress tolerance indices based on yield, phenology
- and biomass partitioning: A review. *Agricultural Reviews* 39(4): 292-299. DOI:
- 624 10.18805/ag.R-1822.
- 625 Sun F, Chen Q, Chen Q, Jiang M, Qu Y. 2022. Yield-based drought tolerance index evaluates the
- drought tolerance of cotton germplasm lines in the interaction of genotype-by-environment.
- 627 *PeerJ.* DOI 10.7717/peerj.14367.
- Teklu DH, Shimelis H, Tesfaye A, Abady S. 2021. Appraisal of the sesame production
- opportunities and constraints, and farmer-preferred varieties and traits, in Eastern and
- 630 Southwestern Ethiopia. Sustainability 13: https://doi.org/10.3390/su132011202.
- 631 Terefe G., Wakjira A, Berhe M, Tadesse H. 2012. Sesame Production Manual. Ethiopian
- Institute of Agricultural Research, Addis Ababa, Ethiopia.
- 633 Teshome D, Tesfaye K, Bekele E. 2015. Genetic diversity of sesame germplasm collection
- 634 (Sesamum indicum L.): Implication for conservation, improvement and use. *Int. J.*
- 635 *Biotechnol. Mol. Biol. Res.* 6(2): 7-18.
- Turner NC. 2018. Imposing and maintaining soil water deficits in drought studies in pots. Plant
- 637 Soil. https://doi.org/10.1007/s11104-018-3893-1
- Walkley AJ, Black IA. 1934. Estimation of soil organic carbon by the chromic nacid titration
- 639 method. Soil Sci. 37: 29-38.
- Weiss EA. 1983. Sesame. In Oilseed Crops. Longman, London. Pp. 282-340.
- Zare M. 2012. Evaluation of drought tolerance indices for the selection of Iranian barley
- (Hordeum vulgare) cultivars. Afri. J. Biotechnol. 11(93):15975-15981. DOI:
- 643 10.5897/AJB12.2127.
- Zenawi G, Mizan A. 2019. Effect of nitrogen fertilization on the growth and seed yield of
- Sesame (Sesamum indicum L.). Int. J. Agron. https://doi.org/10.1155/2019/5027254



Table 1(on next page)

Growth and relative water content responses of sesame varieties under well watered (WW) and water stressed (WS) conditions. Mean values \pm SE (n = 9)

- 1 Table 1:
- 2 Growth and relative water content responses of sesame varieties under well watered (WW) and water stressed (WS) conditions. Mean
- 3 values $\pm SE (n = 9)$

	30 days after water stress imposition				60 days after water stress imposition				
		Sesame v	arieties			Sesar	ne varieties		
Plant parameters	Treatments	Abasena	Gondar-1	Humera-1	Wollega	Abasena	Gondar-1	Humera-1	Wollega
Dlant haight (ann)	WW	42.17±0.30 ^{aA}	40.67±0.67aA	41.89±0.45 ^{aA}	41.83±0.41 ^{aA}	51.33±0.55aB	56.33±1.83 ^{aA}	54.67±0.44a ^{AB}	53.39±0.60 ^{aAB}
Plant height (cm)	WS	35.89 ± 0.81^{bA}	34.67 ± 0.73^{bA}	35.61 ± 0.44^{bA}	34.78 ± 0.92^{bA}	39.44 ± 0.29^{bA}	$40.78{\pm}1.23^{bA}$	38.89 ± 0.65^{bA}	$38.78{\pm}0.32^{\rm bA}$
NIl Cl	WW	$0.00{\pm}0.00^{aA}$	0.67 ± 0.33^{aA}	$0.67 \pm 0.47 ^{aA}$	1.33 ± 0.75^{aA}	0.00 ± 0.00^{aA}	$2.00{\pm}0.00^{aB}$	$2.00{\pm}0.00^{aB}$	$2.44{\pm}0.44^{aB}$
Number of branches	WS	0.00 ± 0.00^{aA}	0.00 ± 0.00^{aA}	0.00 ± 0.00^{aA}	0.00 ± 0.00^{aA}	0.00 ± 0.00^{aA}	0.00 ± 0.00^{bA}	0.00 ± 0.00^{bA}	0.00 ± 0.00^{bA}
N. 1 01	WW	9.56±0.75aA	12.67±1.37aA	12.67 ± 1.37^{aA} 9.22 ± 0.57^{aA} 10.33 ± 0.82^{aA} 12.44 ± 0.65^{aB}	17.33±1.41 ^{aA}	13.33 ± 0.67^{aB}	14.44±1.04aAB		
Number of leaves	WS	6.89 ± 0.39^{aB}	$8.67 \pm 0.83^{\text{bAB}}$	7.44 ± 0.38^{aA}	7.56 ± 0.53^{aAB}	6.44 ± 0.44^{bA}	8.44±1.09bA	6.89 ± 0.59^{bA}	7.11 ± 0.75^{bA}
T (1 1 ()	WW	8.72 ± 0.35^{aA}	9.72 ± 0.19^{aA}	8.00 ± 0.49^{aA}	8.89 ± 0.49^{aA}	11.39±0.73 ^{aA}	12.89±0.78 ^{aA}	11.00±0.52 ^{aA}	11.56±0.48 ^{aA}
Leaf length (cm)	WS	5.11±0.31bA	4.67±0.41 ^{bA}	4.67 ± 0.29^{bA}	4.33 ± 0.24^{bA}	6.39 ± 0.37^{bA}	5.94±0.13bA	5.89±0.11bA	5.89 ± 0.26^{bA}
	WW	4.39 ± 0.14^{aAB}	5.39 ± 0.20^{aA}	4.11 ± 0.42^{aB}	$4.67{\pm}0.34^{aAB}$	6.61 ± 0.22^{aA}	6.56 ± 0.24^{aA}	6.11±0.30 ^{aA}	6.61 ± 0.26^{aA}
Leaf width (cm)	WS	2.78 ± 0.15^{bA}	2.93±0.13bA	2.53±0.17bA	2.82±0.11bA	3.43±0.15bA	3.58±0.15bA	3.17 ± 0.12^{bA}	3.32 ± 0.14^{bA}
	WW	61.40±4.24 ^{aA}	61.61±2.64 ^{aA}	65.31±2.93 ^{aA}	73.77±9.01aA	46.42±3.79aA	42.85±0.82aA	62.24±14.91 ^{aA}	58.79±7.67 ^a A
Relative water content (%)	WS	50.42±3.10 ^{aA}	62.34 ± 2.86^{aA}	58.50±4.80 ^{aA}	81.14±15.68 ^{aA}	47.79±24.26 ^{aA}	51.91±4.05aA	38.49 ± 4.04^{aA}	21.43±14.87 ^{aA}
Relative growth rate	WW	0.86 ± 0.01 aA	$0.94{\pm}0.03^{aB}$	0.91 ± 0.01^{aAB}	0.89 ± 0.01^{aAB}				
(cm/day)	WS	0.66 ± 0.01^{bA}	0.68 ± 0.02^{bA}	0.65 ± 0.01^{bA}	0.65 ± 0.01^{bA}				

- Mean values in a column followed by different small case letters within the same parameter and mean values in a row followed by
- different upper case letters are significantly different at P < 0.05.



Table 2(on next page)

Two ways ANOVA tests of between subjects effects



1 Table 2:

2 Two ways ANOVA tests of between subjects effects

Source	Dependent Variable	df	Mean	F	Sig.	Partial Eta
			Square			Squared
	Plant height	3	33.3	4.7	0.005	0.2
	Number of branches	3	5.4	24.3	0.000	0.5
	Number of leaves	3	40.1	5.7	0.002	0.2
Variety	Leaf Length	3	3.0	1.5	0.242	0.1
	Leaf width	3	0.7	1.8	0.162	0.1
	RWC	3	121.6	0.3	0.864	0.0
	RGR	3	0.01	4.6	0.006	0.2
	Plant height	1	3762.8	526.9	0.000	0.9
	Number of branches	1	46.7	210.3	0.000	0.8
	Number of leaves	1	924.5	131.3	0.000	0.7
Water levels	Leaf Length	1	580.9	275.6	0.000	0.8
	Leaf width	1	172.7	450.3	0.000	0.9
	RWC	1	297.0	0.6	0.442	0.0
	RGR	1	1.1	520.9	0.000	0.9
	Plant height	3	14.4	2.0	0.121	0.1
	Number of branches	3	5.4	24.3	0.000	0.5
Variety *	Number of leaves	3	7.3	1.0	0.382	0.1
Water levels	Leaf Length	3	3.6	1.7	0.176	0.1
water revers	Leaf width	3	0.12	0.3	0.814	0.0
	RWC	3	157.0	0.3	0.813	0.0
	RGR	3	0.0	2.2	0.103	0.1

Notes: RWC = Relative water content; RGR = Relative growth rate

4



Table 3(on next page)

Mean biomass and yield responses of sesame varieties grown under well watered (WW) and water stressed (WS) conditions. Mean \pm SE (n= 9)

1 Table 3:

2 Mean biomass and yield responses of sesame varieties grown under well watered (WW) and water stressed (WS) conditions. Mean

 $3 \pm SE (n=9)$

Sesame	Water	Shoot biomass	Root biomass Biological yield		Number of	Seed	Harvest Index	
varieties	levels	(g)	(g)	(g)	pods/plant	yield/cm ² (g)	(%)	
Abasena	WW	3.11±0.15 ^b	0.70±0.04a	3.81±0.15 ^b	6.00±0.33b	2.14±0.15°	56.65±3.82a	
	WS	1.68±0.14b	0.39 ± 0.08^{b}	2.07 ± 0.17^{d}	3.11±0.35°	1.42±0.17°	68.40±5.22ª	
Gondar-1	WW	6.11±0.51a	0.78 ± 0.05^{a}	6.89±0.53a	8.67±0.58a	3.67±0.14 ^b	55.84±4.97a	
	WS	2.56±0.24b	0.11±0.01°	2.67 ± 0.25^{c}	4.44±0.44°	1.81±0.08°	70.63±4.28a	
Humera-1	WW	6.00 ± 0.67^{a}	0.44 ± 0.06^{b}	6.54±0.69a	8.22±0.62ab	4.14±0.36 ^b	68.27±6.73a	
	WS	2.78 ± 0.28^{b}	0.04±0.01°	2.82 ± 0.28^{c}	3.78±0.52°	1.71±0.15°	65.13±7.89a	
Wollega	WW	6.56±0.44a	0.45 ± 0.07^{b}	7.11±0.42a	8.22±0.70ab	5.18±0.23a	75.02±5.51a	
	WS	3.23±0.26b	0.14±0.08°	3.41±0.22°	3.56±0.44°	1.66±0.12°	50.66±5.68a	

Mean values in each column that don't share similar letters are significantly different at P < 0.05.



Table 4(on next page)

Two ways ANOVA tests on the effects of variety, water levels and their interaction on yield and yield components of sesame varieties



1 Table 4:

- 2 Two ways ANOVA tests on the effects of variety, water levels and their interaction on yield and
- 3 yield components of sesame varieties

Source	Dependent Variable	df	Mean Square	F	Sig.	Partial Eta
						Squared
	Biological yield	3	18.5	14.0	.000	0.4
	Harvest index	3	67.7	0.236	.871	0.0
	Number of pods	2	12.0	<i>5.5</i>	002	0.2
	per plant	3	12.9	5.5	.002	0.2
Variety	Shoot biomass	3	21.8	16.8	.000	0.4
	Root biomass	3	0.4	12.3	.000	0.4
	Ys	3	217777.8	4.8	0.005	0.2
	Yp	3	7161666.7	28.8	0.000	0.6
	Biological yield	1	201.1	152.6	.000	0.7
	Harvest index	1	1.1	0.0	.952	0.0
	Number of pods	1	207.1	1047	000	0.7
	per plant	1	296.1	124.7	.000	0.7
Water levels	Shoot biomass	1	149.5	115.5	.000	0.6
	Root biomass	1	3.2	109.8	.000	0.6
	Ys	1	49335555.6	1078.7	0.000	0.9
	Yp	1	257645000.0	1034.5	0.000	0.9
	Biological yield	3	5.4	4.1	.010	0.2
	Harvest index	3	1439.6	5.0	.003	0.2
	Number of pods	2	2.0	1.0	214	0.1
Variety * Water	per plant	3	2.9	1.2	.314	0.1
levels	Shoot biomass	3	4.317	3.3	.025	0.1
	Root biomass	3	0.1	4.3	.007	0.2
	Ys	3	217777.8	4.8	0.005	0.2
	Yp	3	7161666.7	28.8	0.000	0.6

Note: Ys = Yield under water stress condition; Yp = Yield under well watered condition





Table 5(on next page)

Drought tolerance index values of sesame varieties grown under well watered (WW) and water stressed (WS) conditions



1 Table 5:

2 Drought tolerance index values of sesame varieties grown under well watered (WW) and water

3 stressed (WS) conditions

Drought		Sesame varieties										
indices	Abasena	Gondar-1	Humera-1	Wollega								
MP	1744.44±67.41°	2738.89±78.07 ^b	2977.78±181.07 ^{ab}	3416.67±136.49a								
STI	1310.91±96.59a	1810.89±100.28a	1809.94±180.26a	1659.86±152.93a								
GMP	1667.12±63.07°	2568.40±73.58 ^b	2709.39±141.57ab	2906.88±134.41a								
TOL	800.00±232.74°	1855.56±157.33 ^b	2333.33±365.53b	3522.22±251.68a								
YI	0.63±0.05 ^a	0.49 ± 0.02^{b}	0.44 ± 0.02^{bc}	0.32 ± 0.02^{c}								
YSI	0.66 ± 0.08^{a}	0.50 ± 0.03^{ab}	0.48 ± 0.07^{ab}	0.32 ± 0.03^{b}								
% Reduction	33.84±7.95 ^a	50.00±2.94ab	52.07±6.95ab	67.52±2.93 ^b								
SSI	0.91±0.21 ^a	0.99 ± 0.06^{a}	0.93 ± 0.12^{a}	0.99±0.04a								
SI	0.37 ± 0.00^d	0.51 ± 0.00^{c}	0.56 ± 0.00^{b}	0.68 ± 0.00^{a}								
YSSI	655.91±46.25 ^a	905.94±50.14 ^a	905.43±90.17 ^a	830.43±76.45a								
YPSI	1127.22±142.67 ^{ab}	1346.62±96.11a	1226.73±98.94 ^{ab}	776.66±152.20 ^b								

⁴ Mean values in a row that share different letters are significantly different at P < 0.05. (Mean \pm

⁵ SE and n = 9). MP = Mean productivity; STI = Stress tolerance index; GMP = Geometric mean

⁶ productivity; TOL = Drought tolerance; YI = Yield index; YSI = Yield stability index; SSI =

⁷ Stress susceptibility index; SI = Stress intensity; YSSI = Yield stress score index and YPSI =

⁸ Yield potential index.



Table 6(on next page)

Correlation coefficient between drought tolerance indices with seed yield under well watered (WW) and water stressed (WS) growing conditions

Table 6:
 Correlation coefficient between drought tolerance indices with seed yield under well watered (WW) and water stressed (WS) growing
 conditions

	Ys	Yp	MP	STI	GMP	TOL	YI	YSI	%Reduction	SSI	SI	YSSI	YPSI
Ys	1	0.28	0.50**	0.74**	0.68**	0.01	0.24	0.25	-0.25	-0.52**	0.34*	0.74**	0.80**
Yp	0.28	1	0.97**	0.59**	0.89**	0.96**	-0.70**	-0.80**	0.80^{**}	0.38^{*}	0.85**	0.59**	-0.29
MP	0.50**	0.97**	1	0.72**	0.97**	0.88**	-0.57**	-0.66**	0.66**	0.22	0.85**	0.72**	-0.06
STI	0.74**	0.59**	0.72**	1	0.81**	0.41^{*}	0.09	-0.24	0.24	0.06	0.29	1.00**	0.53**
GMP	0.68**	0.89**	0.97**	0.81**	1	0.74**	-0.43**	-0.52**	0.52**	0.10	0.79**	0.81**	0.16
TOL	0.01	0.96**	0.88**	0.41*	0.74**	1	-0.79**	-0.90**	0.90^{**}	0.54**	0.79**	0.41^{*}	-0.52*
YI	0.24	-0.70**	-0.57**	0.09	-0.43**	-0.79**	1	0.84**	-0.84**	-0.56**	-0.77**	0.09	0.74*
YSI	0.247	-0.80**	-0.66**	-0.24	-0.52**	-0.90**	0.84**	1	-1.00**	-0.83**	-0.60**	-0.24	0.65*
%Reduction	-0.25	0.80^{**}	0.66**	0.24	0.52**	0.90**	-0.84**	-1.00**	1	0.83**	0.59**	0.24	-0.65
SSI	-0.52**	0.38^{*}	0.22	0.06	0.10	0.54**	-0.56**	-0.83**	0.83**	1	0.10	0.06	-0.59*
SI	0.34^{*}	0.85**	0.85**	0.29	0.79**	0.79**	-0.77**	-0.60**	0.59**	0.10	1	0.29	-0.29
YSSI	0.74**	0.59**	0.75**	1.00**	0.81**	0.41^{*}	0.09	-0.24	0.24	0.10	0.29	1	0.53*
YPSI	0.80^{**}	-0.29	-0.06	0.53**	0.16	-0.52**	0.74**	0.65**	-0.65**	-0.59**	-0.29	0.53**	1

^{**} and * indicate significant differences at P < 0.01 and P < 0.05 probability levels, respectively. MP = Mean productivity; STI = Stress tolerance index; GMP = Geometric mean productivity; TOL = Drought tolerance; YI = Yield index; YSI = Yield stability index; SSI = Stress susceptibility index; SI = Stress intensity; YSSI = Yield stress score index and YPSI = Yield potential index.

13

14



Table 7(on next page)

Eigen analysis of the correlation matrix and weight of each parameter

Table 7:

2 Eigen analysis of the correlation matrix and weight of each parameter

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9	PC10	PC11	PC12	PC13
Eigen value	2.88	1.68	0.76	0.50	0.20	0.09	0.06	0.02	0.00	0.00	0.00	0.00	0.00
Proportion of variance	0.862	0.127	0.011	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Cumulative proportion	0.862	0.989	0.999	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

	Cum.(%)	Ys	Yp	MP	STI	GMP	TOL	YI	YSI	% Reduction	SSI	SI	YSSI	YPSI
Principal component 1	86.24	0.05	0.65	0.35	0.13	0.25	0.60	-0.00	-0.00	0.01	0.00	-0.00	0.07	-0.07
Principal component 2	98.86	0.42	0.03	0.23	0.42	0.34	-0.40	0.00	0.00	-0.01	-0.00	-0.00	0.21	0.52

MP = Mean productivity; STI = Stress tolerance index; GMP = Geometric mean productivity; TOL = Drought tolerance; YI = Yield

⁴ index; YSI = Yield stability index; SSI = Stress susceptibility index; SI = Stress intensity; YSSI = Yield stress score index and YPSI =

⁵ Yield potential index.



Table 8(on next page)

Rank mean, standard deviation of rank and rank sum derived from all drought tolerance indices

1 Table 8:

2 Rank mean, standard deviation of rank and rank sum derived from all drought tolerance indices

Varieties	Rank mean	Standard Deviation of Rank	Rank Sum (RS)		
		(SDR)			
Abasena	3.46	1.13	4.59		
Gondar-1	2.15	0.90	3.05		
Humera-1	2.08	0.64	2.72		
Wollega	2.15	1.35	3.5		

3

Figure 1

Map of the study area showing the experimental site




Figure 2

Mean seed yield of sesame varieties grown under water deficit stress (Ys) and well watered conditions (Yp). Mean values followed by different English alphabets are statistically significant at P < 0.05. Mean \pm SE (n = 9).






Figure 3

Biplot of principal component analysis of sesame genotypes for Ys, Yp and drought tolerance indices.

