

First record of *Apanteles hemara* (N.) on *Leucinodes orbonalis* Guenée and biodiversity of Hymenoptera parasitoids on Brinjal (#88540)

1

First submission

Guidance from your Editor

Please submit by **13 Sep 2023** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

11 Figure file(s)

1 Table file(s)

2 Raw data file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

First record of *Apanteles hemara* (N.) on *Leucinodes orbonalis* Guenée and biodiversity of Hymenoptera parasitoids on Brinjal

Hager M. M. Saleh^{Corresp., 1, 2}, Areej A. Al-Khalaf³, Maha Abdullah Alwaili³, Dey Dey^{Corresp. 1}

¹ National Pusa Collection, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi, 110012, India

² Plant Protection Department, Faculty of Agriculture, Fayoum University, Fayoum, 63514, Egypt

³ Biology Department, Faculty of Science, Princess Nourah Bint Abdulrahman University, Riyadh, 11671, Saudi Arabia

Corresponding Authors: Hager M. M. Saleh, Dey Dey

Email address: hms03@fayoum.edu.eg, ddeyiari@hotmail.com

The brinjal fruit and shoot borer (BFSB), *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae), is a severe pest and causes economic losses for the brinjal crop worldwide. Infested brinjal fruits were collected from vegetable fields at ICAR-Indian Agricultural Research Institute (ICAR-IARI), New Delhi, India, during two consecutive seasons (2021-2022). The larvae of the pest were brought to the lab and reared for the emergence of parasitoids at $25 \pm 0.5^\circ\text{C}$ and relative humidity of $70 \pm 5\%$. Further, surveying Hymenoptera parasitoids in brinjal was done using a sweep net and yellow pan trap during two seasons. Findings indicate that five parasitoid species emerged from *L. orbonalis* viz., *Apanteles hemara* Nixon, 1965, *Bracon greeni* Ashmead 1896, *Goryphus nursei* (Cameron, 1907), *Trathala flavoorbitalis* (Cameron, 1907) and *Spalangia gemina* Boucek 1963. Out of these, *A. hemara* and *S. gemina* were reported for the first time in Delhi, and *A. hemara* was recorded for the first time as a parasite on *L. orbonalis*. Among them, *T. flavoorbitalis* was observed during both seasons and showed higher parasitism of 15.55 and 18.46% during July and August 2022, respectively. However, the average parasitism (%) of *A. hemara*, *B. greeni*, *G. nursei*, *T. flavoorbitalis* and *S. gemina* was 1.20, 1.76, 1.10, 9.28 and 3.10% respectively. In addition, the results showed a significant ($p \leq 0.01$) strongly positive correlation between fruit infestation (%) by *L. orbonalis* and parasitism (%). The survey revealed the presence of a broad group (19 families and 60 species) of Hymenoptera parasitoids in the brinjal crop ecosystem in Delhi which could be valuable in biological control. In light of these results, this study revealed that *A. hemara* and other parasitoids recorded in this study alongside *T. flavoorbitalis* would be ideal biocontrol agents in the BFSB IPM program in Delhi.

First record of *Apanteles hemara* (N.) on *Leucinodes orbonalis* Guenée and biodiversity of Hymenoptera parasitoids on Brinjal

Hager M.M. Saleh^{1,2*}, Areej A. Al-Khalaf³, Maha Abdullah Alwaili³ and Debjani Dey^{1*}

¹ National Pusa Collection, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India; hms03@fayoum.edu.eg (H.M.M.S.).

² Plant Protection Department, Faculty of Agriculture, Fayoum University, Fayoum 63514, Egypt; hms03@fayoum.edu.eg (H.M.M.S.).

³ Biology Department, Faculty of Science, Princess Nourah Bint Abdulrahman University, Riyadh 11671, Saudi Arabia; aaalkhalaf@pnu.edu.sa (A.A.A.).

³ Biology Department, Faculty of Science, Princess Nourah Bint Abdulrahman University, Riyadh 11671, Saudi Arabia; maalwaele@pnu.edu.sa (M.A.A.).

¹ National Pusa Collection, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India; ddeyiari@hotmail.com (D.D.).

Corresponding Authors:

Hager M.M. Saleh

National Pusa Collection, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India.

Email address: hms03@fayoum.edu.eg

Debjani Dey

National Pusa Collection, Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India.

Email address: ddeyiari@hotmail.com

Abstract

The brinjal fruit and shoot borer (BFSB), *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae), is a severe pest and causes economic losses for the brinjal crop worldwide. Infested brinjal fruits were collected from vegetable fields at ICAR-Indian Agricultural Research Institute (ICAR-IARI), New Delhi, India, during two consecutive seasons (2021-2022). The larvae of the pest were brought to the lab and reared for the emergence of parasitoids at $25 \pm 0.5^\circ\text{C}$ and relative humidity of $70 \pm 5\%$. Further, surveying Hymenoptera parasitoids in brinjal was done using a sweep net and yellow pan trap during two seasons. Findings indicate that five parasitoid species emerged from *L. orbonalis* viz., *Apanteles hemara* Nixon, 1965, *Bracon greeni* Ashmead 1896, *Goryphus nursei* (Cameron, 1907), *Trathala flavoorbitalis* (Cameron, 1907) and *Spalangia gemina* Boucek 1963. Out of these, *A. hemara* and *S. gemina* were reported for the first time in Delhi, and *A. hemara* was recorded for the first time as a parasite on *L. orbonalis*. Among them, *T. flavoorbitalis* was observed during both seasons and showed higher parasitism of 15.55 and 18.46% during July and

August 2022, respectively. However, the average parasitism (%) of *A. hemara*, *B. greeni*, *G. nursei*, *T. flavaorbitalis* and *S. gemina* was 1.20, 1.76, 1.10, 9.28 and 3.10% respectively. In addition, the results showed a significant ($p \leq 0.01$) strongly positive correlation between fruit infestation (%) by *L. orbonalis* and parasitism (%). The survey revealed the presence of a broad group (19 families and 60 species) of Hymenoptera parasitoids in the brinjal crop ecosystem in Delhi which could be valuable in biological control. In light of these results, this study revealed that *A. hemara* and other parasitoids recorded in this study alongside *T. flavaorbitalis* would be ideal biocontrol agents in the BFSB IPM program in Delhi.

Keywords: Parasitoids, *Apanteles hemara*, *Trathala flavaorbitalis*, *Leucinodes orbonalis*, survey, Brinjal, Delhi.

Introduction

Brinjal (*Solanum melongena* L.) holds a significant position as the third most widely consumed and economically valuable vegetable in Asia. India is the second largest producer of brinjal in the world, next to China. Despite the overall growth in global eggplant production, its productivity is constrained by challenges posed by insects and diseases (Alam & Salimullah, 2021). The primary insect pest responsible for causing damage to brinjal is the lepidopteran fruit and shoot borer, which seriously threatens its cultivation.

The Brinjal fruit and shoot borer (BFSB) *Leucinodes orbonalis* Guenée is a highly important insect pest that affects brinjal plants worldwide (Nusra et al., 2021). The larvae of BFSB cause damage to eggplant plants by tunneling into the leaf petioles, midribs, and tender shoots, which subsequently leads to the wilting and desiccation of the stems. Furthermore, the larvae feed on the flowers, causing them to fall prematurely or resulting in malformed fruits.

The most significant economic damage caused by BFSB occurs to the fruit itself. The fruit becomes unsuitable for human consumption and sale due to the presence of holes, feeding tunnels, and excrement from larvae. BFSB has a high ability to reproduce, quickly cycles through generations, and causes extensive damage in both wet and dry seasons, posing a significant challenge to plants (Prodhan et al., 2018).

Infestation levels may exceed 90%, resulting in substantial worldwide economic losses estimated at 86-90% (Ghosh et al., 2003). In India, brinjal is sprayed with chemicals 15 to 40 times per season (Watkins et al., 2012). However, this approach is environmentally hazardous, poses health risks to consumers and farmers, and incurs significant costs (Prodhan et al., 2018).

To adopt an environmentally friendly strategy for controlling pests, it is essential to protect and preserve the natural predators and parasitoids that naturally keep pest populations in check. Among the 21 parasitoids reported in relation to BSFB, one of the most prominent parasitoids is *T. flavaorbitalis* (Ranjith et al., 2020). This parasitoid has a high parasitism rate of 61.7% (Srinivasan, 2008). In addition to *T. flavaorbitalis*, *Goryphus nursei* was recorded in Uttar Pradesh and proved to be an active parasitoid, particularly during the winter season, exhibiting a maximum parasitism rate of 7% (Alam et al., 2003).

Trathala flavoorbitalis, a widely recognized parasitoid of *L. orbonalis*, has been observed in different regions of India, especially Bihar, Tamil Nadu, Manipur and Karnataka (Mallik et al., 1989; Yasodha & Natarajan, 2009; Murali et al., 2017; Ranjith et al., 2020; Thokchom et al., 2022). This parasitoid species is globally distributed and can be found in regions such as Afrotropical, Australasian, Eastern Palaearctic, Nearctic, Oceanic, and Oriental. It is particularly prevalent in the Indo-Pacific and Eastern Oriental regions (Rousse & Villemant, 2012).

In Tamil Nadu, (Yasodha & Natarajan, 2009) documented the emergence of 12 parasitoid species from BFSB, belonging to the superfamilies Ichneumonoidea and Chalcidoidea. Furthermore, (Murali et al., 2017) reported the presence of *Spalangia gemina* on BSFB, while *Bracon greeni* was documented by (Venkatraman et al., 1948).

Consequently, this research aims to evaluate the parasitism rate of parasites associated with BFSB during two seasons, which is essential for the success of its biological control. Furthermore, a survey of Hymenoptera parasitoids families in the brinjal crop ecosystem in New Delhi.

Materials & Methods

Insect sampling

This study was carried out to record and assess the potential of the natural enemies associated with the *L. orbonalis* Guenée during two consecutive seasons (November 2021 to October 2022) in vegetable fields at ICAR-Indian Agricultural Research Institute (ICAR-IARI), New Delhi, India.

Weather parameters during the study period such as rainfall, minimum and maximum temperature and relative humidity were recorded from an agrometeorological observatory, Division of Agricultural Physics, IARI, New Delhi and are provided in (Table S1).

To calculate the pest infestation rate (%) in brinjal, the field was divided into four quarters, and 15 plants per quarter were randomly checked. The pest incidence was observed at 7 days' intervals, and infestation (%) was calculated according to the following equation.

$$\text{Fruit infestation (\%)} = \frac{\text{No. of infested fruits}}{\text{Total fruits observed}} \times 100 \quad (1)$$

Infested brinjal fruits were collected, and the larvae were reared for the emergence of parasitoids under laboratory conditions of $25 \pm 0.5^\circ\text{C}$, relative humidity $65 \pm 5\%$ RH and a photoperiod of 12 light: 12 Dark h. The larvae, after pupation were separated and kept in plastic containers until parasitoid emergence. The emerged parasitoids were preserved in 70% alcohol and card mounted for taxonomic studies, and the parasitism percentage during each month was calculated according to the following equation (Van Driesche, 1983).

$$\text{Parasitism (\%)} = \frac{\text{Total parasitoids emerged from larvae or pupa}}{\text{Total No. of larvae or pupa collected from field}} \times 100 \quad (2)$$

In addition, a surveying study for Hymenoptera parasitoids in brinjal was randomly done using a sweep net and yellow pan trap weekly during two seasons (2021-2022).

Identification of parasites

From the collected parasites, big parasitoids were pinned and dry-preserved, while small ones were preserved in 70% ethanol. Parasitoids were identified with the help of (*Bouček, 1951; Habu, 1960; Stary 1975; Husain & Agarwal, 1982; Greco, 1997; Narendran et al., 2001; Belokobylskij, 2003; Jonathan, 2006; Gibson, 2009; Rousse et al., 2011; Rousse & Villemant, 2012; Sheeba & Narendran, 2013; Xu et al., 2013; Akhtar et al., 2014; Ghafouri-Moghaddam et al., 2014; Amer et al., 2016; Cao et al., 2017; Fernandez-Triana et al., 2017; Khalaim, 2018; Ahmed et al., 2020; Zerova & Fursov, 2020; Gull-e-Fareen et al., 2021; Talamas et al., 2021*). The morphological studies were carried out using a Leica S8AP0 stereo microscope and a LEICA M205 C stereozoom automontage microscope. Multi-focused montage images were taken using a LEICA MC190 HD digital camera attached to the LEICA M205 C stereozoom automontage microscope. The photographs and illustrations were processed with Adobe Photoshop CS5 and plates were prepared. The morphological terminology and wing venation are based on (*Nixon 1965; Jonathan, 2006; Gibson, 2009; Sheeba & Narendran, 2013*). All the specimens are deposited in the National Pusa Collection (NPC), ICAR-IARI, New Delhi, India.

Statistical analysis

The parasitism (%) was analyzed statistically, and the extent of parasitism (%) was also subjected to correlation analysis with brinjal fruit infestation (%), also with weather factors using Minitab® Statistical Software (17.0).

Results

The results of this study showed that five parasitoid species viz., *A. hemara*, *B. greeni* (Braconidae), *G. nursei*, *T. flavoorbitalis* (Ichneumonidae) and *S. gemina* (Pteromalidae) were recorded on *L. orbonalis*.

I. Systematic study

The main diagnostic characteristics of five parasitic wasps that emerged from Brinjal fruit and shoot borer (*L. orbonalis* Guenée), as well as their hosts and distribution details, were highlighted as follows.

1. *Apanteles hemara* Nixon, 1965 (Fig. 1)

Diagnosis: Body black. Mandible and labrum orange testaceous, palpi pale yellow and antennae dark brown to black. Fore and mid legs yellowish orange, entire hind coxa densely punctate rugose; hind femur brown, yellow trochanter, hind tibia most often strikingly bicolor, yellow and brown on posterior with spurs white, hind tarsus infusate. Wing hyaline, venation brown; pterostigma mostly brown. Head entirely densely setose. Clypeus ventral margin slightly concave. Face and clypeus moderately and shallowly punctate; antenna slightly shorter than body length, with 14 flagellomere. Mesosoma, mesoscutum and scutellum with relatively coarse and dense punctures (distance between punctures smaller than diameter of puncture). Propodeum with areola complete, propodeal areola strong, centrally smooth, apically pointed and basally petiolate to base of propodeum by two sub-median irregular carinas. Forewing with 2Rs more than twice shorter than

r, R1 very long, pterostigma enlarged. The areolet open. Legs hind coxae entirely punctate rugose. Hind femur slightly swollen. Metasoma T1 of metasoma much longer than wide with strong, longitudinal striation, its margins sub-parallel to strongly converging apically. Tergum 2 wider than long, short, transverse and apically widened. Tergum 3 longer than tergum 2. Ovipositor sheath slightly shorter than tibia. Ovipositor large, usually slightly decurved and gradually tapered. **Material examined:** 1♀ June; 2♀ August; 1♀ September 2022 emerged from *L. orbonalis* and 1♀ August; 2♀ September 2022 brinjal, yellow pan trap, ICAR-IARI, New Delhi. **Host records:** *Tebenna micalis* (Choreutidae); *Cnaphalocrocis trapezalis*, *Herpetogramma stultalis*, *Hydriris ornatalis*, *Omiodes indicatae*, *Spoladea recurvalis* and *Udea ferrugalis* (Crambidae) (Fernandez-Triana et al., 2017) *Hymenia fascialis*, *Pachyzancla stultalis* (Pyralidae) (Nixon 1965). **Distribution:** Kenya, Madagascar, Cape Verde Islands, Mauritius, Saudi Arabia, Senegal, Republic of Congo, South Africa, Yemen, Australia and India (new record established at New Delhi) (Fernandez-Triana et al., 2017).

2. *Bracon greeni* Ashmead 1896 (Fig. 2)

Diagnosis: Body brownish-yellow. Disc of metasoma, extreme apex of second tergite and large dorsal blotches on third and fourth tergites black; wings hyaline, stigma and veins brown. Head and thorax smooth; wings hyaline; stigma and veins brown antenna 24 segmented; vertex rugose, anteriorly smooth, shiny posteriorly and hairy; eyes glabrous, slightly emarginate; temple smooth, shiny and hairy; width of face 1.65x its height; clypeus smooth. Antenna nearly as long as the body. Mesosoma pronotum smooth and shiny; mesoscutum smooth, shiny hairy; notauli weakly impressed; posteriorly, smooth; scutellum smooth, shiny and hairy; metanotum with anterior median carina; propleuron smooth, shiny and hairy; propodeum smooth, shiny with a few hairs anteriorly and posterior median longitudinal carina extending upto its middle; propodeal spiracle round, small and medially placed; fore wing vein 3-SR about 3.0x r. Metasoma broadly oval and shagreen; second to fourth metasomal tergites subequal, remaining a little shorter. Large dorsal blotches on third and fourth tergites black; ovipositor nearly as long as metasoma.

Material examined: 2♀ November; 1♀ December 2021; 1♀ February 2022 emerged from *L. orbonalis* and 1♀ November; 2♀ December 2021 brinjal, yellow pan trap, ICAR-IARI, New Delhi.

Host records: *Adisura atkinsoni*, *Alcides affaber*, *Earias fabia*, *Heliothis obsolete*, *Rabila frontalis* (Sheeba & Narendran, 2013) and *L. orbonalis* (Venkatraman et al., 1948). In the present study, it emerged from *L. orbonalis*.

Distribution: India (Kerala, Uttar Pradesh), Bangladesh, China, and Sri Lanka.

3. *Goryphus nursei* (Cameron, 1907) (Fig. 3)

Diagnosis: Body dark orange except eighth to tenth flagellar segments white above; face and frons along the eye margin, apex of metasoma T1 and T5-T8 light yellow to white; T2-T4 entirely black. Legs in general reddish, except fore and middle tibia and tarsus tending to be brownish; hind femur broadly at apex, whole of tibia and tarsus blackish; hind tibia with a subbasal white band. Head

face and clypeus strongly punctate, clypeus tending to be smooth towards the apex. Antennal scrobe shallow, smooth and shiny. Temple broadly smooth and shiny. Malar space weakly granulose, about 0.7x the basal width of mandible. Mandible teeth subequal. Mesosoma pronotum coarsely striate, Mesoscutum strongly punctate, striate along margins of each lobe. Scutellum smooth and shiny, Metascutellum smooth and shiny. Mesopleuron rugose, tending to be wrinkled above, Propodeum largely reticulate, based on basal carina transwrinkled, both transverse carinae strong and complete, basal carina more or less straight, apical carina evenly and strongly arched, bow-shaped. Areolet small, squares, about 3x as high as the width of bordering veins. Metasoma T1 short, about 1.3x as long as wide at apex, 2nd and 3rd tergite closely punctate; 4th tergite less punctate. Ovipositor tip with distinct teeth ventrally.

Material examined: 1♀ December 2021; 2♀ February 2022 emerged from *L. orbonalis* and 1♀ November 2021; 1♀ January 2022 brinjal, yellow pan trap, ICAR-IARI, New Delhi.

Host records: BFSB *L. orbonalis* (Alam et al., 2003).

Distribution: India (Bihar, Delhi, Gujarat, Haryana, Jharkhand, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Uttarakhand, and Uttar Pradesh), China and Pakistan (Jonathan, 2006; Gupta 1985).

4. *Trathala flavoorbitalis* (Cameron, 1907) (Fig. 4)

Diagnosis: Body brownish-yellow, Antenna scape and pedicel yellowish and flagellomere brownish, vertex (except ocelli area black) and around eye light yellow; Notauli often well marked with yellow. Tegula and scutellum yellow. Metasoma orange, tergites 1-2 and basal triangle on tergite III dark brown. Legs yellow, hind tibia slightly infuscate basally and apically; wings hyaline, pterostigma brown, its anterior half yellow. Sheath and ovipositor dark brown. Head strongly constricted behind eyes. Face densely punctate, clypeus smooth, vertex and frons granulate, center of frons smooth. Temples short and slightly rounded; malar space 0.65 times as long as basal mandible width, occipital carina complete; mandible with equal size teeth; antenna filiform longer than head and mesosoma with 33 flagellomeres; scape length 1.31x pedicel length. Mesonotum densely punctate-shagreened, scutellum and metanotum more sparsely punctate. Scutellum rounded without dorsal lateral carinae. Propodeum densely punctate-shagreened dorsally than laterally. Propodeal carination complete, area basalis small but distinct. Metasoma tergite I a little longer than tergite II and twice longer than tergite III. Tergite I almost smooth, slightly longitudinally strigose at apex. Tergite II three times longer than apically wide and longitudinally striated. Ovipositor shorter than abdomen.

Material examined: 3♀, 1♂ November; 2♀ December 2021; 1♀, 1♂ January; 2♀, 1♂ February; 3♀, 2♂ March; 2♀, 1♂ June; 4♀, 2♂ July; 6♀, 2♂ August; 3♀, 1♂ September; 1♀, 1♂ October 2022 emerged from *L. orbonalis* and 2♀ July; 3♀ August 2022 brinjal, sweeping net, ICAR-IARI, New Delhi.

Host records: Seventy-eight host records, all Lepidoptera (Gelechoidea, Noctuoidea, Pyraloidea, Tineoidea and Tortricidae) (Rousse et al., 2011) brinjal shoot and fruit borer (Alam et al., 2003; Ranjith et al., 2020). In the present study, it emerged from *L. orbonalis*.

Distribution: Reunion, Madagascar ([Rousse et al., 2011](#)). Widespread through Indo-Pacific and Eastern Oriental Areas ([Rousse & Villemant, 2012](#)).

5. *Spalangia gemina* Boucek, 1963 (Fig. 5)

Diagnosis: Body dark. Legs dark except basal four tarsal segments yellow. Head with dense circular setiferous punctures; temple distinctly circular punctures. Gena with malar space distinctly shorter than eye height and malar sulcus absent. Antenna with scape about 6.8 as long as greatest width, the inner and outer surfaces uniformly setose and strongly. Mesosoma pronotal collar in lateral view convex behind neck, with coarsely reticulate-rugose, except for a nearly triangular area close to the crenulate cross-line posteriorly; axillae smooth and shiny except for a few pinprick-like setiferous punctures. Scutellum smooth and shiny except for a few pinprick-like setiferous punctures laterally; mesopleuron with longitudinal carinae extending from subalar area ventrally over almost all of upper mesepimeron, subalar scrobe extending posteroventrally along transepisternal line, and epistern. Fore wing hyaline; bare behind submarginal vein. Propodeum with postspiracular sulcus; dorsal surface punctate-rugose anteriorly and posteriorly, sculptured, sometimes almost smooth; supracoxal; propodeal sides smooth and shiny. Metasoma smooth and shiny. Petiole about 1.7-1.8x as long as medial width; almost smooth to finely, transversely carinate between longitudinal carinae.

Material examined: 1♀ June; 2♀ July; 2♀, 1♂ August; 1♀ September 2022 emerged from *L. orbonalis* and 2♀ July; 1♀ August 2022 brinjal, sweeping net trap, ICAR-IARI, New Delhi.

Host records: Micropezidae, Noctuidae and Tortricidae (Lepidoptera) as primary hosts ([Noyes, 2019](#)) recorded on *L. orbonalis* ([Murali et al., 2017](#)).

Distribution: Afrotropical, Australasian, Oriental, and Neotropical region ([Noyes, 2019](#)).

II. Evaluation of parasitism extent by *T. flavoorbitalis* and other parasitoids on BFSB

During July and August 2022, *T. flavoorbitalis* showed higher parasitism of 15.55 and 18.46%, respectively. However, the average parasitism (%) of *T. flavoorbitalis*, *S. gemina*, *B. greeni*, *A. hemara* and *G. nursei* was 9.28, 3.10, 1.76, 1.2 and 1.1%, respectively (Fig. S1). Throughout the study period, *T. flavoorbitalis* was the dominant parasitoid. Its parasitism in 1st season peaked at 12% in November 2021, while it decreased in December 2021 and again increased during February and March 2022. In the 2nd season, its parasitism peaked at 18.5% in August 2022, and higher fruit infestation (%) was observed (Fig. S2). The parasitism (%) of *B. greeni* was 6.7 and 3.3% during November and December 2021, respectively but was not recorded during the 2nd season. The parasitism (%) of *G. nursei* was 3.3 and 2.9% during December and January, respectively but not recorded in the 2nd season (Fig. 6).

The parasitism (%) of *S. gemina* was 6.12% in August 2022. *A. hemara*, as a new parasitoid (Hymenoptera: Braconidae), was found to parasitizes on the BFSB larvae. This species was recorded in the 2nd season, and its parasitism (%) was 3.3 and 2.4% during June and September, respectively (Fig. 6). The total parasitism (%) of all parasitoids, as well as the fruit infestation (%), reach its peak during August (Fig. 7).

III. Correlation analysis of fruit infestation (%) and parasitism (%) under study conditions

Correlation analysis was performed on the data obtained under this study to explore the interrelationship among the infestation (%) and parasitism (%) with the environmental parameters during the study period. A significant ($p \leq 0.01$) strongly positive correlation ($r = 0.7$) was shown between fruit infestation (%) and total parasitism (%) of parasitoid species. Also, a significant ($p \leq 0.01$) strongly positive correlation ($r = 0.7$) was recorded between infestation (%) and parasitism (%) of *T. flavaorbitalis*. Parasitism (%) of *T. flavaorbitalis* showed a significant ($p \leq 0.5$) moderately negative correlation with rainfall ($r = -0.4$), moderately positive correlation with *Tmin* ($r = 0.3$), RH ($r = 0.4$) and not correlated with *Tmax* ($r = -0.02$). Fruit infestation (%) showed a significantly ($p \leq 0.1-0.5$) moderately positive correlation with *Tmax*, ($r = 0.3$), *Tmin* ($r = 0.5$), RH ($r = 0.4$), and a moderate negative correlation with rainfall ($r = -0.5$). Total parasitism (%) showed a significant ($p \leq 0.5$) moderately negative correlation with rainfall ($r = -0.3$), weakly positive correlation with *Tmin* ($r = 0.2$), moderately positive correlation with RH ($r = 0.4$) and not correlated with *Tmax* ($r = -0.05$).

IV. Survey for study of associated Hymenoptera parasitoids

Parasitoids were collected weekly from brinjal during two seasons (November 2021 to October 2022), using a sweep net and yellow pan trap. The monthly distribution of different Hymenoptera parasitoid families (19 families) collected from brinjal is shown in (Fig. S3). About 60 species were recorded, from which 46 were identified up to the species level, and 14 were identified up to the genus level. Superfamily Chalcidoidea was the most dominant, followed by Ichneumonoidea, Platygastroidae and Ceraphronoidea. Among 60 species recorded, eleven species (Braconidae), ten species (Ichneumonidae), eight species (Chalcididae), eight species (Scelionidae), three species (Eulophidae), four species (Pteromalidae), three species (Dryinidae), three species (Diapriidae), two species (Mymaridae), two species (Figitidae) and one species were recorded in (Platygastriidae, Agonidae, Bethylinidae, Ceraphronidae, Eurytomidae and Aphelinidae) (Table 1 and Fig. 8) families.

Discussion

Several studies carried out in different time-lines suggest the occurrence of different parasitoids on *L. orbonalis*. Along with *T. flavaorbitalis*, a few other Ichneumonids recorded were *Pristomerus testaceus* (Ayyar, 1927), *Eriborus argentiopilosus* (Tewari & Sardana, 1987), *Xanthopimpla punctata* (Navasero, 1983; Navasero & Calilung, 1990), *E. sinicus* (Talekar, 1995) and *Diadegma apostate* (Krishnamoorthy & Mani, 1998). Previous studies suggest the occurrence of braconids viz., *Chelonus* sp. (Sandanayake & Edirisinghe, 1992), *Bracon* sp. (Tewari & Sardana, 1987), *B. greeni* (Venkatraman et al., 1948), and *Phanerotoma* sp. (Tewari & Moorthy, 1984; Sandanayake & Edirisinghe, 1992) *Apanteles* sp. (Navasero, 1983). Among various larval parasitoids, *T. flavaorbitalis* was recorded as the most critical species constituting about 60% of larval parasitoids. It has been recorded as a major parasitoid of *L. orbonalis* in Sri Lanka, Gujarat (India) and Bangladesh, with maximum parasitism of 61.7% (Alam et al., 2003). The superfamily

Chalcidoidea species identified in relation to *L. orbonalis* viz., *Brachymeria* sp., *B. lasus*, *B. obscurata*, *Antrocephalus mitys* (Chalcididae), *S. irregularis*, *S. gemina*, *S. endius* (Pteromalidae) and *Trichogramma* sp. (Trichogrammatidae) (Navasero & Calilung, 1990).

In the present study, five species belong to five genera under three families (Braconidae, Ichneumonidae, and Pteromalidae). *T. flavaorbitalis* had the highest parasitism rate compared to other parasitoids, with an average parasitism rate of 9.28%, while the parasitism rate of *B. greeni*, *A. hemara*, *S. gemina* and *G. nursei* was 1.76, 1.21, 3.11, and 1.06%, respectively. These findings align with the other reports (Alam et al., 2003; Nagalingam, 2006; Ranjith et al., 2020). *T. flavaorbitalis* has also been recorded as an important parasitoid in different countries, such as Hawaii and several places in the USA (Swezey, 1926), Bangladesh (Alam & Sana, 1962), Sri Lanka (Sandanayake & Edirisinghe, 1992), India (Bihar (Mallik et al. 1989), Karnataka (Ranjith et al., 2020); Manipur (Thokchom et al., 2022); Gujarat, Uttar Pradesh (Alam et al., 2003) and Nepal (Kafle, 1970). The current study revealed a higher maximum parasitism of 18.45% on larvae of *L. orbonalis* in August during 2nd season, and our findings agree with (Srinivasan, 2008; Ranjith et al., 2020) indicating that the parasitoid potentially reduced the population of *L. orbonalis* (Srinivasan, 2008; Kumar & Raghuraman, 2014). In this study, during the summer season *A. hemara* was recorded as a parasitoid on *L. orbonalis*, for the first time. This finding agrees with (Navasero, 1983), who recorded *Apanteles* sp. on BFSB in Philippines. *G. nursei* and *B. greeni* were active during the winter but did not appear in the summer, indicating that their activity decreased under high temperatures, according to (Alam et al., 2003).

The brinjal fruit and shoot borer (BFSB) damage varied considerably, and highest damage was observed during summer, while there was very little fruit infestation during the winter season (Fig. 7). This could be due to high summer temperatures and cooling temperatures from December to January (Fig. 7). At the peak period, the pest damage exceeded 70% of total fruits, whereas in January, barely 27% damage was seen. Generally, the level of parasitism concisely concurs with the population of the pest. However, there was a significant correlation between fruit infestation (%) and parasitism (%), and these findings are in line with (Alam et al., 2003; Ranjith et al., 2020).

Conclusion

In summary, the fruit and shoot borer (BFSB) *L. orbonalis*, is a major pest of brinjal worldwide, causing extensive damage to the fruit and making it unsuitable for human consumption. We have recorded five species of parasitoids on *L. orbonalis*, including the first-ever recorded instance of *Apanteles hemara* acting as a parasite on this pest in New Delhi. The survey revealed the presence of various Hymenoptera parasitoids within the brinjal crop ecosystem. However, the parasitism rate varied significantly depending on the environmental conditions during the survey. We found a significant strongly positive correlation between parasitism (%) of parasitoids with fruit infestation (%) by *L. orbonalis*. This research emphasizes the importance of preserving and protecting the natural enemies as they serve as effective biocontrol agents for *L. orbonalis*, thereby reducing the need for pesticides.

Acknowledgements

The authors express their gratitude to ICAR-Indian Agricultural Research Institute, Division of Entomology, for providing necessary research facilities. The authors also would like to thank Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia for the support through University Researchers Supporting Project number PNURSP2023R37.

Author Contributions

Conceptualization, H.M.M.S., A.A.A-K., M.A.A. and D.D.; methodology, H.M.M.S. and D.D.; software, H.M.M.S., A.A.A-K., M.A.A. and D.D.; validation, H.M.M.S., A.A.A-K., M.A.A. and D.D.; formal analysis, H.M.M.S., A.A.A-K., M.A.A. and D.D.; investigation, H.M.M.S., A.A.A-K., M.A.A. and D.D.; resources, H.M.M.S., A.A.A-K., M.A.A. and D.D.; data curation, H.M.M.S., A.A.A-K., M.A.A., A.E. and D.D.; writing-original draft preparation, H.M.M.S., A.A.A-K., M.A.A. and D.D.; writing-review and editing, H.M.M.S., A.A.A-K., M.A.A. and D.D.; visualization, H.M.M.S., A.A.A-K., M.A.A. and D.D.; supervision, H.M.M.S. and D.D.; project administration, H.M.M.S. and D.D.; funding acquisition, H.M.M.S., A.A.A-K., M.A.A. and D.D. All authors have read and agreed to the published version of the manuscript

Data Availability Statement

The data presented in this study are available on request from the corresponding author.

References

- Ahmed S, Bodlah I, Nasir MF, Nadeem, MS. 2020. Genus *Xanthopimpla* Saussure, 1892 (Ichneumonidae: Pimplinae) from pothwar region (Punjab) of Pakistan. *Japs, Journal of Animal and Plant Sciences* **30(3)**:713-722. <https://doi.org/10.36899/JAPS.2020.3.0084>.
- Akhtar MS, Singh LRK, Ramamurthy VV. 2014. New species of the genus *Macrocentrus* Curtis 1833 (Hymenoptera: Braconidae) from India. *The Pan-Pacific Entomologist* **90(1)**:11-15. <https://doi.org/10.3956/2014-90.1.11>.
- Alam I, Salimullah M. 2021. Genetic Engineering of Eggplant (*Solanum melongena* L.): Progress, Controversy and Potential. *Horticulturae* **7(4)**:78. <https://doi.org/10.3390/horticulturae7040078>.
- Alam MZ, Sana DL. 1962. Biology of the brinjal shoot and fruit borer, *Leucinodes orbonalis* G. (Pyralidae: Lepidoptera) in East Pakistan. *Scientist* **5(1-4)**:13-24.
- Alam SN, Rashid MA, Rouf FMA, Jhala RC, Patel JR, Satpathy S, Shivalingaswamy TM, Rai S, Wahundeniya I, Cork A, Ammaranan C, Talekar NS. 2003. Development of an integrated pest management strategy for eggplant fruit and shoot borer (*Leucinodes orbonalis*) in South Asia. *Technical Bulletin* No. 28. AVRDC Publication No. 03-548, pp. 56. ISBN: 92-9058-126-3. <http://www.oisat.org/downloads/fruitandshootborer.pdf>
- Amer FSK, Zeya SB, Veenakumari KA. 2016. Review of the Indian species of *Mymar* Curtis (Hymenoptera: Mymaridae). *Journal of Insect Systematics* **3(1&2)**:16-37.

- 380 **Ayyar TVR. 1927.** The Parasitic Hymenoptera of economic importance noted from South India.
381 *Bulletin of Entomological Research* **18(1)**:73-78.
382 <https://doi.org/10.1017/S0007485300019714>.
- 383 **Belokobylskij SA. 2003.** The species of the genus *Spathius* Nees, 1818 (Hymenoptera:
384 Braconidae: Doryctinae) not included in the monograph by Nixon (1943). *Annales*
385 *Zoologici* **53(3)**:347-488.
- 386 **Bouček Z. 1951.** The first revision of the European species of the family Chalcididae
387 (Hymenoptera). *Acta Entomologica Musei Nationalis Pragae* **27(1)**:1-108.
- 388 **Cao H-X, LaSalle J, Zhu C-D. 2017.** Chinese species of *Pediobius* walker (Hymenoptera:
389 Eulophidae). *Zootaxa*, **4240(1)**:1-71. <https://doi.org/10.11646/zootaxa.4240.1.1>.
- 390 **Fernandez-Triana J, Beaudin M, van Achterberg K, Agbodzavu MK, Othim ST, Nyamu**
391 **FW, Fiaboe KK. 2017.** DNA barcodes, expanded distribution, and redescription of
392 *Apanteles hemara* Nixon, 1965 (Hymenoptera, Braconidae, Microgastrinae), a potential
393 biocontrol species against amaranth leaf-Webbers in Africa. *Journal of Hymenoptera*
394 *Research* **58**:1-15. <https://doi.org/10.3897/jhr.58.13361>.
- 395 **Ghafouri-Moghaddam M, Lotfalizadeh H, Rakhshani EA. 2014.** Survey on hyperparasitoids
396 of the poplar spiral gall aphid, *Pemphigus spyrothecae* Passerini (Hemiptera: Aphididae)
397 in Northwest Iran. *Journal of Crop Protection* **3(3)**:369-376.
- 398 **Ghosh SK, Laskar N, Senapati SK. 2003.** Estimation of loss in yield of brinjal due to pest
399 complex under Terai region of West Bengal. *Environment and Ecology* **21(4)**:764-769.
- 400 **Gibson GAP. 2009.** Revision of new world Spalanginae (Hymenoptera: Pteromalidae). *Zootaxa*
401 **2259(1)**:1-159. <https://doi.org/10.11646/zootaxa.2259.1.1>.
- 402 **Greco CF. 1997.** Specificity and instar preference of *Diplazon laetatorius* (Hym.: Ichneumonidae)
403 parasitizing aphidophagous syrphids (Dipt.: Syrphidae). *Entomophaga*, **42**:315-318.
404 <https://doi.org/10.1007/BF02769823>.
- 405 **Gull-e-Fareen A, Bodlah I, Bodlah MA, Rasheed MT, Ali H, Asif M. 2021.** Colour and
406 distributional pattern of *Callaspidia notata* (Boyer de Fonscolombe, 1832) (Hymenoptera:
407 Figitidae: Aspicerinae) from Pakistan. *Pakistan Journal of Zoology* **53(3)**:1153.
- 408 **Habu AA. 1960.** Revision of the Chalcididae (Hymenoptera) of Japan, with descriptions of sixteen
409 new species. *Bulletin of the National Institute of Agricultural Sciences (Jpn) Ser. C* **11**:131-
410 363.
- 411 **Husain T, Agarwal MM. 1982.** Taxonomic studies on Haltichellinae of India (Hymenoptera:
412 Chalcididae) pt. I. Haltichellini. *Oriental Insects* **16(3)**:313-336.
413 <https://doi.org/10.1080/00305316.1982.10434319>.
- 414 **Jonathan JK. 2006.** Ichneumonologia Indica (Part-1), An Identification Manual on Subfamily
415 Mesosteninae (Hymenoptera: Ichneumonidae). *Zoological Survey of India, Kolkata*: 1-
416 680. ISBN: 81-8171-107-6.
- 417 **Kafle GP. 1970.** New records of some insects and noninsects parasites of *Leucinodes orbonalis*
418 Guenee in Nepal. *Nepalese Journal of Agricultural Sciences* **5**:99-103.

- 419 **Khalaim AI. 2018.** The genera *Allophrys* Förster and *Aneuclis* Förster (Hymenoptera:
420 Ichneumonidae: Tersilochinae) of Vietnam. *Zootaxa* **4378(3)**:414-428.
421 <https://doi.org/10.11646/zootaxa.4378.3.9>.
- 422 **Krishnamoorthy A, Mani M. 1998.** New record of parasitoid *Diadegma apostata* (G.) on brinjal
423 shoot and fruit borer. *Insect Environment* **4(3)**:87.
- 424 **Kumar D, Raghuraman M. 2014.** An over view of integrated management of *Leucinodes*
425 *orbonalis* in brinjal crop in Eastern Uttar Pradesh, India. *Plant Archives* **14(2)**:649-654.
- 426 **Mallik SN, Kumar M, Sinha AN, Karn BP. 1989.** *Trathala flavoorbitalis* Cameron
427 (Ichneumonidae) a parasite of *Leucinodes orbonalis* Guen. from Bihar. *Current Science*
428 **58(19)**:1098-1099.
- 429 **Murali S, Jalali SK, Kariyanna B, Shylesha AN, Shivalinga Swamy TM, Gandhi Gracy R.**
430 **2017.** Documentation of parasitoid complex on brinjal fruit and shoot borer, *Leucinodes*
431 *orbonalis* Guenee. *Bulletin of Environment, Pharmacology and Life Sciences* **6(2)**:452-
432 459.
- 433 **Nagalingam T. 2006.** Association of parasitoid, *Trathala flavoorbitalis* with brinjal shoot and pod
434 borer *Leucinodes orbonalis* in the mid country region of Sri Lanka. *Sri Lankan Journal of*
435 *Agricultural Sciences* **43**:124-132.
- 436 **Narendran TC, Galande SM, Mote UN. 2001.** Two new species of Eulophidae (Hymenoptera:
437 Chalcidoidea) from Maharashtra, India. *Uttar Pradesh Journal of Zoology* **21(2)**:153-157.
- 438 **Navasero MV, Calilung VJ. 1990.** Biology of the eggplant fruit and shoot borer, *Leucinodes*
439 *orbonalis* Guenee (Lepidoptera: Pyralidae) in the Philippines. **Philippine Entomologist**
440 **8(2)**:749-760.
- 441 **Navasero MV. 1983.** Biology and chemical control of the eggplant fruit and shoot borer,
442 *Leucinodes orbonalis* Guenee (Pyraustidae: Lepidoptera). BSc Thesis, University of the
443 Philippines, Los Baños.
- 444 **Nixon GEJ. 1965.** A reclassification of the tribe Microgasterini (Hymenoptera: Braconidae).
445 *Bulletin of the British Museum (Natural History) Entomology* **2**: 1-284.
446 <https://doi.org/10.5962/p.144036>.
- 447 **Noyes JS. 2019.** Universal Chalcidoidea Database. World Wide Web electronic publication.
448 <http://www.nhm.ac.uk/chalcidoids> (accessed on 15 April 2023).
- 449 **Nusra MSF, Udukala DN, Amarasinghe LD, Paranagama PA. 2021.** Volatiles from host plant
450 brinjal attract the brinjal Fruit and Shoot Borer-*Leucinodes orbonalis* Guenee. *Journal of*
451 *Asia-Pacific Entomology* **24(3)**:695-703. <https://doi.org/10.1016/j.aspen.2021.06.002>.
- 452 **Prodhan MZH, Hasan MT, Chowdhury MMI, Alam MS, Rahman ML, Azad AK, Hossain**
453 **MJ, Naranjo SE, Shelton AM. 2018.** Bt eggplant (*Solanum melongena* L.) in Bangladesh:
454 Fruit production and control of eggplant fruit and shoot borer (*Leucinodes orbonalis*
455 Guenee), effects on non-target arthropods and economic returns. *PLoS One*
456 **13(11)**:e0205713. <https://doi.org/10.1371/journal.pone.0205713>.
- 457 **Ranjith M, Nelson SJ, Sithanantham S, Hegde DR. 2020.** Parasitisation of brinjal shoot and
458 fruit borer, *Leucinodes orbonalis* Guenee by *Trathala flavoorbitalis* Cameron. *Pest*

- 459 *Management in Horticultural Ecosystems* **26**(1):104-108. [https://doi.org/10.5958/0974-](https://doi.org/10.5958/0974-4541.2020.00017.X)
- 460 [4541.2020.00017.X](https://doi.org/10.5958/0974-4541.2020.00017.X).
- 461 **Rousse P, Villemant C, Seyrig A. 2011.** Ichneumonid wasps from Madagascar. V.
- 462 Ichneumonidae Cremastinae. *Zootaxa* **3118**(1-5): 1-30.
- 463 <https://doi.org/10.11646/zootaxa.3118.1.1>.
- 464 **Rousse P, Villemant C. 2012.** Ichneumons in Reunion Island: a catalogue of the local
- 465 Ichneumonidae (Hymenoptera) species, including 15 new taxa and a key to species.
- 466 *Zootaxa* **3278**(1):1-57. <https://doi.org/10.11646/zootaxa.3278.1.1>.
- 467 **Sandanayake WRM, Edirisinghe JP. 1992.** *Trathala Flavoorbitalis*: Parasitization and
- 468 development in relation to host stage attacked. *International Journal of Tropical Insect*
- 469 *Science* **13**: 287-292. <https://doi.org/10.1017/S1742758400013539>.
- 470 **Sheeba M, Narendran TCA. 2013.** Review of *Bracon* species of India with a key to Indian
- 471 Species. *Zoological Survey of India, Kolkata* **348**:1-52. ISBN: 978-81-8171-343-8.
- 472 **Srinivasan R. 2008.** Integrated Pest Management for eggplant fruit and shoot borer (*Leucinodes*
- 473 *orbonalis*) in south and southeast Asia: Past, Present and Future. *Journal of Biopesticides*
- 474 **1**(2):105-112. <https://doi.org/10.57182/jbiopestic.1.2.105-112>.
- 475 **Stary P. 1975.** *Aphidius colemani* Viereck: its taxonomy, distribution and host range
- 476 (Hymenoptera, Aphidiidae). *Acta entomologica bohemoslovaca* **72**(3):156-163.
- 477 **Swezey OH. 1926.** Recent introductions of beneficial insects in Hawaii. *Journal of Economic*
- 478 *Entomology* **19**(5):714-720. <https://doi.org/10.1093/jee/19.5.714>.
- 479 **Talamas EJ, Bremer JS, Moore MR, Bon MC, Lahey Z, Roberts CG, Combee LA,**
- 480 **McGathey N, Van Noort S, Timokhov AV, Hougardy E. 2021.** A maximalist approach
- 481 to the systematics of a biological control agent: *Gryon aetherium* Talamas, sp. nov.
- 482 (Hymenoptera, Scelionidae). *Journal of Hymenoptera Research* **87**:323-480.
- 483 <https://doi.org/10.3897/jhr.87.72842>.
- 484 **Talekar NS. 1995.** Integrated insect and disease management (IPM) for environment-friendly
- 485 production of safe vegetables. Publication No.00-505 Shanhua, Taiwan: *Asian Vegetable*
- 486 *Research and Development Center*:16.
- 487 **Tewari GC, Moorthy PNK. 1984.** New record of two parasites of brinjal shoots and fruit borer,
- 488 *Leucinodes orbonalis* Guenee. *Entomon* **9**:63-64.
- 489 **Tewari GC, Sardana HR. 1987.** *Eriborus argenteopilosus* (Cameron)-a new parasite of
- 490 *Leucinodes orbonalis* Guen. *Entomon* **12**(3):227-228.
- 491 **Thokchom S, Gupta A, Akoijam R, Devesh A. 2022.** Parasites of *Leucinodes orbonalis* Guenee
- 492 from Manipur. *Indian Journal of Entomology* **85**(1):188-189.
- 493 <https://doi.org/10.55446/IJE.2022.524>.
- 494 **Van Driesche RG. 1983.** Meaning of “percent parasitism” in studies of insect parasitoids.
- 495 *Environmental Entomology* **12**(6):1611-1622. <https://doi.org/10.1093/ee/12.6.1611>.
- 496 **Venkatraman TV, Gupta SN, Negi PS. 1948.** Preliminary trials in the propagation of *Bracon*
- 497 (*Microbracon*) *greeni* Ashmead on unnatural hosts. In **Proceedings/Indian Academy of**
- 498 **Sciences**, Springer India: New Delhi, India **27**(4):92-104.

- 499 **Watkins PR, Huesing JE, Margam V, Murdock LL, Higgins TJV. 2012.** Insects, nematodes,
500 and other pests. In **Plant Bio-technology and Agriculture**, Academic Press, pp. 353-370.
501 <https://doi.org/10.1016/B978-0-12-381466-1.00023-7>.
- 502 **Xu Z, Olmi M, He J. 2013.** Dryinidae of the oriental region (Hymenoptera: Chrysidoidea).
503 *Zootaxa* **3614(1)**:1-460. <https://doi.org/10.11646/zootaxa.3614.1.1>.
- 504 **Yasodha P, Natarajan N. 2009.** Record of parasitoids and pathogens on *Leucinodes orbonalis*.
505 *Madras Agricultural Journal* **96(1-6)**:194-199.
- 506 **Zerova MD, Fursov VN. 2020.** A Review of the Palaearctic Species of the Genus *Eurytoma* Ill.
507 of the robusta Group, with Description of a New Species from Uzbekistan (Hymenoptera,
508 Chalcidoidea: Eurytomidae). *Entomological Review* **100**:563-576.

Table 1(on next page)

Table 1 Surveying of the Hymenoptera parasitoids in brinjal during two seasons (November 2021 to October 2022).

* ^{1st}, first season; 2nd, second season; +, Presence; -, Absence

**1 Table 1 Surveying of the Hymenoptera parasitoids in brinjal during two seasons (November
2 2021 to October 2022).**

S.No.	Families/Parasitoids *	1 st	2 nd	S.No	Families/Parasitoids	1 st	2 nd
Family Ichneumonidae				Family Platygasteridae			
1	<i>Trathala flavoorbitalis</i>	+	+	39	<i>Macroteleia livingstoni</i>	-	+
2	<i>Xanthopimpla punctate</i>	-	+	Family Eulophidae			
3	<i>Xanthopimpla flavolineata</i>	+	+	40	<i>Tetrastichus schoenobii</i>	+	-
4	<i>Ichneumon</i> sp.	+	-	41	<i>Pediobius foveolatus</i>	+	+
5	<i>Diplazon laetatorius</i>	+	+	42	<i>Quadrastichus</i> sp.	+	+
6	<i>Goryphus nursei</i>	+	+	Family Agonidae			
7	<i>Goryphus apollonis</i>	-	+	43	<i>Eupristina saundersi</i>	+	+
8	<i>Aneucelis</i> sp.	+	-	Family Aphelinidae			
9	<i>Temelucha</i> sp.	-	+	44	<i>Aphelinus asychis</i>	+	+
10	<i>Amyosoma chinese</i>	-	+	Family Mymaridae			
Family Braconidae				45	<i>Anagrus atomus</i>	+	-
11	<i>Macrocentrus delhiensis</i>	+	-	46	<i>Mymar taprobanicum</i>	+	+
12	<i>Cotesia</i> sp.			Family Dryinidae			
13	<i>Aphidius colemani</i>	+	+	47	<i>Aphelopus</i> sp.	+	+
14	<i>Microplates</i> sp.	+	+	48	<i>Anteon achterbergi</i>	-	+
15	<i>Chelonus blackburni</i>	-	+	49	<i>Anteon yasumatsui</i>	-	+
16	<i>Bracon carpomyiae</i>	+	+	Family Diapriidae			
17	<i>Apanteles hemara</i>	-	+	50	<i>Trichopria</i> sp.	+	+
18	<i>Bracon greeni</i>	+	-	51	<i>Trichopria keralensis</i>	+	-
19	<i>Phaenocarpa</i> sp.	+	-	52	<i>Psillus</i> sp.	+	+
20	<i>Spathius helle Nixon</i>	+	-	Family Bethyridae			
21	<i>Phanerotoma</i> sp.			53	<i>Goniozus indicus</i>	+	+
Family Chalcididae				Family Ceraphronidae			
22	<i>Brachymeria bengalensis</i>	+	+	54	<i>Aphanogmus fijiensis</i>	+	-
23	<i>Brachymeria lasus</i>	+	-	Family Pteromalidae			
24	<i>Brachymeria hime</i>	+	+	55	<i>Pachyneuron Solitarium</i>	+	-
25	<i>Brachymeria podagrica</i>	+	-	56	<i>Spalangia gemina</i>	+	+
26	<i>Dirhinus auratus</i>	+	-	57	<i>Pteromalus puparum</i>	+	+
27	<i>Antrocephalus sepyra</i>	+	+	58	<i>Sphegigaster brunneicornis</i>	+	+
28	<i>Kriechbaumerella kraussi</i>	+	+	Family Figitidae			
29	<i>Kriechbaumerella pulvinatus</i>	+	+	59	<i>Alloxysta rubidus</i>	+	-
Family Eurytomidae				60	<i>Callaspidia notate</i>	+	-
30	<i>Eurytoma</i> sp.	+	-				
Family Scelionidae							
31	<i>Trissolcus basalis</i>	-	+				
32	<i>Baryconus europaeus</i>	+	+				
33	<i>Scelio</i> sp.	-	+				
34	<i>Hadronotus fulviventris</i>	+	+				
35	<i>Hadronotus hogenakalense</i>	+	+				
36	<i>Gryonoides pulcherllus</i>	-	+				
37	<i>Protelenomus</i> sp.	-	+				
38	<i>Telenomus dignus</i>	-	+				

3 * 1st, first season; 2nd, second season; +, Presence; -, Absence

Figure 1

Figure 1 *Apanteles hemara* Nixon, 1965 (female)

A. Dorsal habitus, **B.** Fore and hind wings, **C.** Dorsal view of mesosoma, **D.** Dorsal view of metasoma and **E.** Emerged parasite from host.




Figure 1 *Apanteles hemara* Nixon, 1965 (female); **A.** Dorsal habitus, **B.** Fore and hind wings, **C.** Dorsal view of mesosoma, **D.** Dorsal view of metasoma and **E.** Emerged parasite from host.

1

Figure 2

Figure 2 *Bracon greeni* Ashmead 1896 (female)

A. Lateral habitus, **B.** Frontal view of head, **C.** Dorsal view of head, **D.** Dorsal view of metasoma, **E.** Fore wing and **F.** Emerged parasite from host.

1




Figure 2 *Bracon greeni* Ashmead 1896 (female); **A.** Lateral habitus, **B.** Frontal view of head, **C.** Dorsal view of head, **D.** Dorsal view of metasoma, **E.** Fore wing and **F.** Emerged parasite from host

2



1

Figure 3

Figure 3 *Goryphus nursei* (Cameron, 1907) (female)

A. Dorsal habitus, **B.** Frontal view of head, **C.** Dorsal view of mesosoma, **D.** Fore and hind wings and **E.** Dorso-lateral view of metasoma.

1




Figure 3 *Goryphus nursei* (Cameron, 1907) (female); **A.** Dorsal habitus, **B.** Frontal view of head, **C.** Dorsal view of mesosoma, **D.** Fore and hind wings and **E.** Dorso-lateral view of metasoma.

2
3
4
5
6
7
8
9

Figure 4

Figure 4 *Trathala flavoorbitalis* (Cameron, 1907) (female)

A. Dorso-lateral habitus, **B.** Dorsal view of head, **C.** Frontal view of head, **D.** Dorsal view of mesosoma, **E.** Fore and hind wings and **F.** Lateral view of metasoma.




Figure 4 *Trathala flavoorbitalis* (Cameron, 1907) (female); **A.** Dorso-lateral habitus, **B.** Dorsal view of head, **C.** Frontal view of head, **D.** Dorsal view of mesosoma, **E.** Fore and hind wings and **F.** Lateral view of metasoma.

Figure 5

Figure 5 *Spalangia gemina* Boucek, 1963 (female)

A. Lateral habitus, **B.** Frontal view of head, **C.** Mesopleuron, **D.** Dorsal view of mesosoma and **E.** Fore wing.




Figure 5 *Spalangia gemina* Boucek, 1963 (female); **A.** Lateral habitus, **B.** Frontal view of head, **C.** Mesopleuron, **D.** Dorsal view of mesosoma and **E.** Fore wing.

Figure 6

Figure 6 Parasitism (%) of all parasitoid species on *L. orbonalis* during two seasons (November 2021 to October 2022)

No Legend




Figure 6 Parasitism (%) of all parasitoid species on *L. orbonalis* during two seasons (November 2021 to October 2022)

Figure 7

Figure 7 Monthly fruit infestation (%) by *L. orbonalis* and parasitism (%) of parasitoids during the study period (November 2021 to October 2022)

No Legend




Figure 7 Monthly fruit infestation (%) by *L. orbonalis* and parasitism (%) of parasitoids during the study period (November 2021 to October 2022)

Figure 8(on next page)

Figure 8 Some Hymenoptera parasitoid species collected during the study period (November 2021 to October 2022)

No Legend



Hadronotus hogenakalense



Telenomus dignus



Hadronotus fulviventris



Macroteleia livingstoni



Baryconus europaeus



Gryonoides pulcherillus



Pediobius foveolatus ♂



Aphanogmus fijiensis



Eurytoma sp



Sphegigaster brunneicornis



Aphelopus sp



Callaspidia notate




Figure 8 Some Hymenoptera parasitoid species collected during the study period (November 2021 to October 2022)