

Population structure and fecundity of a Xanthid crab *Leptodius exaratus* (H. Milne Edwards, 1834) on the rocky shore of Gujarat state, India

Krupal Patel ^{Equal first author, 1}, Heris Patel ^{Equal first author, 1}, Swapnil Gosavi ², Kauresh Vachhrajani ², Jigneshkumar Trivedi ^{Corresp. 1}

¹ Department of Life Sciences, Hemchandracharya North Gujarat University, Patan, Gujarat, India

² Department of Zoology, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India

Corresponding Author: Jigneshkumar Trivedi
Email address: jntrivedi26@yahoo.co.in

Background A Xanthid crab, *Leptodius exaratus* (H. Milne Edwards, 1834), was examined for its population structure and breeding biology on the rocky intertidal region of Shivrajpur village in Saurashtra coast, Gujarat state.

Method Month wise sampling was conducted from March 2021 to February 2022 during low tide using catch per unit effort in the 500 m² area. The sampled specimens were categorised into male, non-ovigerous females or ovigerous females. In order to estimate fecundity, the morphology of the crab specimens (carapace width and body weight) as well as the size of egg, number of eggs and weight of egg mass were recorded.

Results A total 1215 individuals were sampled of which 558 individuals were males and 657 individuals were females. The size (carapace width) of males ranges from 5.15 mm to 29.98 mm, while females ranges from 5.26 mm to 28.63 mm which shows that the average size of male and female individuals did not differ significantly. The overall as well as monthly sex ratio was skewed towards males with a bimodal distribution while unimodal in females. The population breeds year-round, which was indicated by the occurrence of ovigerous females throughout the year. However, the maximum percentage occurrence of ovigerous females was observed from December to April which indicates the peak breeding season. The size of egg, number of eggs and weight of egg mass were shown to positively correlate with the morphology of ovigerous females (carapace width and wet weight).

Population structure and fecundity of a Xanthid crab *Leptodius exaratus* (H. Milne Edwards, 1834) on the rocky shore of Gujarat state, India

Krupal Patel^{1¶}, Heris Patel^{1¶}, Swapnil Gosavi², Kauresh Vachhrajani², and Jigneshkumar
Trivedi^{1*}

¹ Animal Taxonomy and Ecology Laboratory, Department of Life Science, Hemchandracharya
North Gujarat University, Patan–384265, Gujarat, India.

² Marine Biodiversity and Ecology Laboratory, Department of Zoology, The Maharaja Sayajirao
University of Baroda, Vadodara–390002, Gujarat, India

*Corresponding Author: Jigneshkumar Trivedi¹

Animal Taxonomy and Ecology Laboratory, Department of Life Science, Hemchandracharya
North Gujarat University, Patan–384265, Gujarat, India

Email address: jntrivedi26@yahoo.co.in

[¶] These two authors contributed equally to this work

Abstract

Background A Xanthid crab, *Leptodius exaratus* (H. Milne Edwards, 1834), was examined for its population structure and breeding biology on the rocky intertidal region of Shivrajpur village in Saurashtra coast, Gujarat state.

Method Month wise sampling was conducted from March 2021 to February 2022 during low tide using catch per unit effort in the 500 m² area. The sampled specimens were categorised into male, non-ovigerous females or ovigerous females. In order to estimate fecundity, the morphology of the crab specimens (carapace width and body weight) as well as the size of egg, number of eggs and weight of egg mass were recorded.

Results A total 1215 individuals were sampled of which 558 individuals were males and 657 individuals were females. The size (carapace width) of males ranges from 5.15 mm to 29.98 mm, while females ranges from 5.26 mm to 28.63 mm which shows that the average size of male and female individuals did not differ significantly. The overall as well as monthly sex ratio was skewed towards males with a bimodal distribution while unimodal in females. The population breeds year-round, which was indicated by the occurrence of ovigerous females throughout the year. However, the maximum percentage occurrence of ovigerous females was observed from December to April which indicates the peak breeding season. The size of egg, number of eggs and weight of egg mass were shown to positively correlate with the morphology of ovigerous females (carapace width and wet weight).

Keywords: Breeding, Population structure, Rocky shore, Saurashtra coast, Sex ratio, Xanthidae

40 Introduction

41 Investigation on the population structure of intertidal crabs started in early 1940s (Flores and
 42 Paula, 2002), which can reveal the patterns of species interactions and their roles within
 43 ecosystems. Accounts on population structure and breeding biology majorly try to understand the
 44 genetic diversity, age, spatial distribution, abundance, sex ratio, variation in year-round
 45 composition, fecundity of the species, as well as juvenile recruitment (Litulo, 2005; Saher and
 46 Qureshi, 2010; Manzoor et al., 2016; Hu et al., 2015). A species' life history might differ by
 47 habitat or even by location. For example, a slight variation in the latitude leading to climatic
 48 variation can cause differences between the populations. Variations in the population trends are
 49 also due to the effects of several biotic and abiotic factors affecting populations differently
 50 (Lycett et al., 2020). Studies on the population structure and breeding biology of a species can
 51 help determine its ecological stability in a given habitat and also contribute to our understanding
 52 of the species' biology. (Santos et al., 1995; Takween and Qureshi, 2005). This knowledge helps
 53 ecologists understand how different species coexist, compete, and interact, influencing
 54 ecosystem dynamics. However, such studies have not been carried out so far on some of the
 55 commonly occurring brachyuran crabs on Gujarat state.

56 The Saurashtra coast of Gujarat state, India, is characterised by its rocky intertidal coasts, which
 57 support a great diversity of marine organisms, including intertidal crustaceans, especially crab
 58 population. With its major inhabiting marine intertidal species, majority of the crab studies have
 59 focused on the diversity (Trivedi and Vachhrajani, 2013a, b; 2015; 2016a; 2018; Trivedi et al.,
 60 2015a, b;; 2021; Gosavi et al., 2017a, b; 2021; Patel et al., 2020a; 2021a; Bhat and Trivedi, 2021;
 61 Padate et al., 2022). Very less is known about the population structure of these important
 62 intertidal organisms (Trivedi and Vachhrajani, 2016b; 2017; Patel et al., 2020b, 2022). *Leptodius*

exaratus (H. Milne Edwards, 1834) is a xanthid crab that is commonly found in the rocky shores of Saurashtra coast (Patel et al., 2021b). This crab species has been commonly reported from the rocky intertidal regions of Indo-Pacific region (Kneib and Weeks, 1990; Naderloo, 2017). It is an omnivorous species that prefers benthic fauna over algae to feed upon and is expected to have a considerable impact on how the benthic ecosystem is structured (Al-Wazzan et al., 2020).

In the Indian subcontinent *L. exaratus* is recorded from Andaman and Nicobar Islands, Maharashtra, Tamil Nadu, Lakshadweep Islands, Goa, Karnataka, and Gujarat (Trivedi et al., 2018). Though the species is very commonly found on the Saurashtra coast of Gujarat, studies only on its taxonomy (Chopra and Das, 1937; Trivedi and Vachhrajani, 2015) and colour variation (Patel et al., 2021b) have been carried out so far. Hence the current investigation was aimed to 1) understand the population structure and 2) study the breeding biology of *L. exaratus* occurring on the rocky shore of Saurashtra coast Gujarat state, India in order to obtain knowledge about the ecology of rocky intertidal habitats. Studying the population structure and breeding biology of *L. exaratus* which is commonly found on the Saurashtra coast, would provide a baseline data that plays a pivotal role in understanding the effects of changing environment, habitat, or anthropogenic pressure. The present study will help in elucidating the coastal health of study area.

Materials & Methods

Study area

The investigation was conducted on the rocky shore of Shivrajpur (22°19'55" N 68°57'03" E) which is located on the Saurashtra coast of Gujarat state, India (Figure 1). *Leptodius exaratus* is a

small crab found abundantly in the rocky shore in this study area (Figure 2). During the low tide, the exposed area of rocky intertidal region varies from 60 to 150 m.

Field methods

Monthly field work was conducted for 12 consecutive months from March 2021 to February 2022. The month wise data was compiled into winter season (November to February months), summer season (March to June months), and monsoon seasons (July to October months), following Rao and Rama-Sharma (1990), to observe the seasonal variation. Catch-per-unit effort using the hand-picking method was used for the collection of specimens for a time period of 4 hours at the time of low tide. When the water receded, a 500 m² area in the intertidal region was marked off and thoroughly examined for the presence of *L. exaratus*. Small rocks were also upturned for the presence of *L. exaratus*, which they prefer to occupy. Whenever an individual crab was encountered, the crab was collected and preserved in 10% formalin pending additional examination.

Laboratory analysis

The crabs were identified on the basis of their morphological characters as follows using standard identification key provided by Lee et al. (2013): Carapace is transversely sub ovate, and lightly granular. There are four large, triangular teeth on the anterolateral border, behind the exorbital angle. Male abdomen tapered, somites 3–5 fused, somite 6 elongated, 1.6 times longer than the telson. Chelipeds are unequal in size; fingers are stout with dark pigmentation excluding the tips, which are white in colour. Further, the individuals were categorised as male, non-ovigerous female, or ovigerous female (Figure 2). For morphological character, carapace width (CW) was measured by digital vernier callipers (Mitutoyo 500-197-20) (0.01 mm accuracy) and

wet weight of crabs was measured using weighing balance (Sartorius–BSA224S–CW) (0.001 g accuracy).

The following method for fecundity study was adopted from Patel et al. (2023). Fecundity estimation conducted by cautiously taking out the mass of eggs present on the pleopods of ovigerous females (n=34) and measuring three parameters i.e., total number of eggs, weight of egg mass and size of eggs (diameter). For the total number of eggs, the egg mass was transferred into 20 ml of sea water and mixed gently so that the eggs got distributed evenly in the water. From this solution, three samples of 2 ml each were taken in a petri dish and observed under a stereo zoom microscope (Matlab–PST–901) to count the total number of eggs. The total number of eggs in each sample was divided by 3 and multiplied by the dilution factor (10) to obtain the total number of eggs (Litulo, 2004). Ovigerous females were weighed both with and without egg mass, and the difference in their weight was considered as the weight of egg mass. Eggs (n=10) from each ovigerous female were measured by means of an ocular micrometre under a microscope for the size range (Saher and Qureshi, 2010).

Data analysis

Population structure

The specimens were grouped in 2 mm size class intervals from 4 mm to 30 mm CW in order to get the overall size frequency distribution. Shapiro Wilk test was conducted to analyse the normality of the collected data, which suggests that the data distribution was not normal ($p < 0.001$). Hence, non-parametric analysis was carried out. To investigate the difference in the variance of mean values of the carapace width of male, non-ovigerous, and ovigerous individuals Kruskal-Wallis (KW) test was conducted. On getting a significant difference ($p < 0.05$) in the CW between the sexes, a multiple comparison analysis using Dunn's post hoc test was used to do a

multiple comparison study. Monthly variations in the size (CW) and sex composition of *L. exaratus* individuals were obtained by plotting the data on individuals' carapace width and sex. The ratio of males and females (ovigerous and non-ovigerous females) was evaluated by the means of chi-square test (χ^2). The size at first maturity was determined by calculating the percentage of ovigerous females across various size classes from the total number of samples collected. Juveniles were defined as individuals that were smaller than the smallest ovigerous female (Baeza et al., 2013). The effect of temperature on *L. exaratus* breeding and juvenile settling was examined by plotting monthly data on the incidence of juvenile and ovigerous females against ambient temperature. The relationship between the mean ambient temperature and the relative juvenile frequency was examined using Pearson's correlation analysis.

Fecundity

To investigate the relationship between the morphological features of eggs (total number of eggs, egg mass weight and size of eggs) and crabs' morphology (CW and weight) regression analysis was performed. At $p < 0.05$, the statistical significance was deemed significant. Microsoft Excel and PAST software, version 4.03 (Hammer et al., 2001), were used to carry out statistical analyses.

Results

During the study period, 1215 individuals were investigated in total; 558 of them were male (45.93%) and 657 of them were females (54.07%) (Table 1). The carapace width of *L. exaratus* males ranged from 5.15 mm to 29.98 mm, while in case of females it ranged from 5.26 mm to 28.63 mm. The size differences between the male and female individuals were not statistically significant (Kruskal-Wallis, $H = 0.209$, $p = 0.646$). (Table 1).

Table 2 shows that the year-round average total sex ratio (1:1.2) for *L. exaratus* was significantly different from the predicted 1:1 proportion ($\chi^2 = 4.1219$, $p = 0.042$) and biased towards females. Month wise, female biased sex ratio was observed in almost all the months except September (1:0.8), October (1:0.9), and November (1:0.7) (Table 2). November had the highest percentage of male occurrences (57.89%), while April had the lowest rate (37.41%). In terms of females, the highest percentages of non-ovigerous female occurrences were observed in June (50%) and August (50%), while the lowest percentages were observed in January (27.18%). Ovigerous females were collected all year, which shows the species is breeding all year-round. However, from December to April, the greatest percentage of occurrence was recorded of ovigerous females, suggesting a peak in the breeding season.

The individuals of *L. exaratus* occurred in all the size classes between 4 mm to 30 mm. It was observed that males exhibited bimodal pattern of distribution having maximum occurrence in 6–8 mm CW size class and 24–26 mm CW size class. On the other hand, females exhibited unimodal pattern of size frequency distribution, with maximum occurrence recorded in 14–16 mm CW size class (Figure 3).

Moreover, there was a considerable variation in the occurrence of adults, ovigerous females and juveniles (< 12 mm) during different months of the year (Figure 4). It was found that in April, May, June, and July (summer and early monsoon season) the population of juveniles was least as compared to the adult population. Moderately, less number of juveniles were also observed during December to March (winter and early summer season) than August to November (Monsoon and early winter season) as compared to adult male and female (Figure 4).

Males had a bimodal distribution during most of the months, while non-ovigerous females showed a unimodal distribution pattern, as was also observed in ovigerous females. Furthermore, it was also observed that juveniles were present all year round (Figure 5–7). A negative correlation (Pearson's correlation, $r = -0.39$) was observed between the mean ambient temperature and relative frequency of juveniles (Figure 8).

The results of fecundity revealed that the CW of ovigerous females was between 10.38 mm and 24.02 mm, with their average size being 17.95 ± 3.81 mm ($n = 34$). The wet body weight of the ovigerous females was recorded between 0.41 and 4.64 g, with the mean weight being 2.01 ± 1.1 g ($n = 34$). The average number of eggs observed was $4,529 \pm 2,003$ ($n = 34$), with the minimum and maximum reported being 920 and 8,730 eggs, respectively. The average egg size ($n = 34$) was 0.36 ± 0.07 mm, with the minimum and maximum observed sizes being 0.19 and 0.54 mm, respectively. The average egg mass weight ($n = 34$) was 0.29 ± 0.18 g, with the minimum and maximum observed egg mass weights being 0.04 and 0.88 g, respectively (Table 3). The ovigerous females' carapace width and body weight were shown to be significantly correlated with both the egg weight and total number of eggs (Figure 9).

Discussion

A Significant variation was observed in the average carapace width of different sexes of *Dotilla blanfordi*, where it was found that male individuals were significantly larger than females. Studies conducted on the population structure of *Matuta planipes* and *Ashtoret lunaris* (Saher et al., 2017), *Uca bengali* (Tina et al., 2015), *Scylla olivacea*, *S. tranquebarica*, and *S. paramamosain* (Waiho et al., 2021), *Clibanarius ransonii* (Patel et al., 2020a; 2022), *C. rhabdodactylus* (Patel et al., 2023), and *Diogenes custos* (Patel et al., 2020b) also exhibited

similar results. It has been observed that the growth rate of female individuals is generally reduced as a result of greater energy investment in gonadal development, which leads to decreased somatic growth in comparison to male individuals (Mantelatto et al., 2010). Another hypothesis suggests that the chances of attracting and obtaining females for the purpose of mating increases with increased size of male individuals (Wada et al., 2000), while the difference in size also reduces intraspecific competition among different sexes for available resources (Abram, 1988).

Overall sex ratio (1:1.2) was found to be female-biased, while month-wise also female biased sex ratio was observed except September, October and November months. In general, natural selection promotes a sex ratio of 1:1 parental expenditure on offspring (Taylor, 1996); however, deviation from the ideal sex ratios is common in marine crustaceans, as observed in *Calcinus tibicen* (Fransozo and Mantelatto, 1998), *Limulus polyphemus* (Smith et al., 2002), *Crangon crangon* (Siegel et al., 2008), *Opusia indica* (Saher and Qureshi, 2011), and *Macrophthalmus (Venitus) dentipes* (Qureshi and Saher, 2012). The sex ratio also differed during different growth stages, with an ideal sex ratio (1:1) in smaller size classes (1–3 mm CW), female biased in intermediate size classes (3–6 mm CW) and exclusively male biased in larger individuals (6–8 mm CW). Certain other studies have found similarly results (Gherardi and Nardone, 1997; Bezerra and Matthews-Cascon, 2007; Mishima and Henmi 2008; Manzoor et al., 2016). Numerous factors can be responsible for such deviation in the sex ratio including competition in local mate (Hamilton, 1967), differences in the efficiency of utilizing local resources that biases sex ratios (Silk, 1983), difference in the investment in male and female offspring (Kobayashi et al., 2018), and sexual selection (Swanson et al., 2013). Sexual dimorphism in size could be one of the reasons for the different sex ratio from the ideal 1:1 in different size classes. Higher male

217 mortality in the intermediate-size classes often leads to a female biased sex ratio (Asakura,
 218 1995). Moreover, males grow to bigger sizes quickly than females, leading to male biased sex
 219 ratio on the larger size classes (Wenner, 1972). Disparities in sexual mortality and dispersion
 220 may potentially contribute to unbalanced sex ratios in crab populations (Johnson, 2003).
 221 Present investigation found that the size frequency distribution of *L. exaratus* males had a
 222 bimodal distribution, while the females had a unimodal distribution. Also, there was a
 223 considerable difference in the seasonal size frequency distribution. Similar results have been
 224 observed in *Paguristes tortugae* (Mantelatto and Sousa, 2000), *Chaceon affinis* (López Abellán
 225 et al., 2002), *Pilumnus vespertilio* (Litulo, 2005), *Dilocarcinus pagei* (Taddei et al., 2015) *Aegla*
 226 *georginae* (Copatti et al., 2016) and *Clibanarius rhabdodactylus* (Patel et al., 2023). Over time,
 227 the population size and frequency of dispersion may be significantly changed by the rapid
 228 recruitment of larvae and reproductive rate (Thurman, 1985). Such distributions have been
 229 explained by a variety of theories, such as differential patterns of migration (Flores and
 230 Negreiros-Fransozo, 1999), **growth rate** (Negreiros-Fransozo et al., 2003), and differential death
 231 rate (**Díaz** and Conde, 1989). It is commonly found in organisms that undergo several rounds of
 232 reproduction and generate a large number of clutches every season (Zimmerman and Felder,
 233 1991). Unimodality is often seen in stable populations that have approximately equal numbers of
 234 new members and emigrants, consistent recruitment and mortality rates throughout the course of
 235 the life cycle, and steady demographics (Thurman, 1985; Díaz and Conde, 1989) whereas,
 236 bimodality could be an indication of the general tendencies in population increase.
 237 The ambient temperature of the study site ranged from 22.1 to 32.5 °C, which is within the range
 238 of a tropical environment that may support continuous reproduction. Hence, there was **year-**
 239 **round** occurrence of ovigerous females suggesting that *L. exaratus* is a continuously breeding

species that has maximum recorded frequency from December to April. Similarly, studies carried out on *L. exaratus* (Al-Wazzan et al., 2020), *Scylla olivacea* (Ali et al., 2020), *Opusia indica* (Saher and Qureshi, 2011), *Emerita portoricensis* and *E. asiatica* (Goodbody, 1965), *Ilyoplax frater* (Saher and Qureshi, 2010), *Diogenes brevirostris* (Litulo, 2004) and *Petrochirus diogenes* (Bertini and Fransozo, 2002) did not find any association between the frequency occurrence of ovigerous females and ambient temperature. As *L. exaratus* is common inhabitant of upper intertidal region where higher temperature during summer season can greatly increase the desiccation risk, leading to migration of ovigerous females in deeper water (Allen, 1966; Asakura, 1987; Al-Wazzan et al., 2020) resulting in decreased abundance in the intertidal region. As a result, seasonal fluctuations in abundance reflect both migration and mortality, while summer abundance estimates may underestimate the size of the local population. However, it was found that juvenile percentage occurrence increased with a decline in ovigerous female percentage occurrence, whereas juvenile percentage occurrence declined when ovigerous female percentage occurrence increased. Such outcomes demonstrates that the species may recruit juveniles throughout the year as a result of rapid reproduction and a short incubation time. Similar outcomes have been reported in several other studies including *Deiratonotus japonicus* (Oh and Lee, 2020), *Scylla olivacea* (Rouf et al., 2021), *Clibanarius rhabdodactylus* (Patel et al., 2023), *Dardanus deformis* (Litulo, 2005), and *Menippe nodifrons* (Fransozo et al., 2000). There are a number of variables, including the availability of food for adults (Goodbody, 1965), the ecology of larvae (Reese, 1968), the amount of time to attain sexual maturity, the timing of mating and gonadal development, as well as the length of the incubation period (Sastry, 1983), which can lead to periodicity in the reproductive season. A variety of abiotic and biotic variables, including water temperature (Chou et al., 2019), salinity (Huang et al., 2022), the nutritional

quality of the females (Matias et al., 2016), variations in photoperiod (Zhang et al., 2023), the amount and availability of nutrition (Viña-Trillos et al., 2023), and the threat of predation (Touchon et al., 2006), may affect the reproductive maxima among populations.

It was found that the CW and wet body weight of *L. exaratus* were having positive correlation with total number of eggs and egg mass weight. Several other studies have also found similar results (Patel et al., 2023; Crowley et al., 2019; Hamasaki et al., 2021; Aviz et al., 2022; Mustaquim et al., 2022). Additionally, it has been demonstrated that ovigerous females with the same CW had variations in the number of eggs, egg mass weight, and egg size resulting from variations in the food supply, variation in egg production, and egg loss (Hines, 1982).

Conclusions

The goal of the current study was to better understand the **population structure and** breeding biology of *L. exaratus*. Significant sexual dimorphism was found, with males being larger than females, most likely as a result of the size of gamete formation differing between the sexes and females investing more in egg production. **Total sex ratio of species was 1:1.2** and monthly populations may be a result of differential biology and behaviour as well as the impact of biotic and abiotic variables on male and female individuals. The year-round occurrence of ovigerous females suggests continuous breeding of the population and an inverse relationship between the peak in juvenile recruitment and the occurrence of ovigerous females which is a common phenomenon of tropical brachyuran crabs. There was a positive correlation between the egg parameters (weight of egg mass and number of eggs) and the morphology of ovigerous females (carapace width and body weight). Fecundity may be impacted by a variety of internal and external variables, such as the amount of energy used for somatic development and egg

production. The present study was conducted at Shivrajpur village, a renowned tourist site with a blue-flag beach where various water sports activities take place. These activities along with higher tourist rush at the study site that may impact the habitat composition of the coast and also potentially influencing the ecology of *L. exaratus*. Furthermore, our findings will contribute to understanding the species' response to environmental changes, as both population structure and fecundity are closely tied to environmental variables.

Acknowledgements

All the authors are thankful to Dhruva Trivedi for technical assistance in field work.

References

- Abrams PA. 1988. Sexual difference in resource use in hermit crabs; consequences and causes. In: Chelazzi G, Vannini M eds. *Behavioral adaptation to intertidal life*. MA: Springer, 283–296.
- Ali MY, Hossain MdB, Sana S, Rouf MA, Yasmin S, Sarower MdG. 2020. Identifying peak breeding season and estimating size at first maturity of mud crab (*Scylla olivacea*) from a coastal region of Bangladesh. *Heliyon* 6:e04318. DOI: <https://doi.org/10.1016/j.heliyon.2020.e04318>.
- Allen JA. 1966. The rhythms and population dynamics of decapod Crustacea. *Oceanography and Marine Biology* 4:247-265.

- 304 Al-Wazzan Z, Vay LL, Behbehani M, Giménez L. 2020. Scale-dependent spatial and temporal
305 patterns of abundance and population structure of the xanthid crab *Leptodius exaratus* on
306 rocky shores in Kuwait. *Regional Studies in Marine Science* 37:101325. DOI:
307 <https://doi.org/10.1016/j.rsma.2020.101325>.
- 308 Asakura A. 1987. Preliminary observations on the offshore mass migration of the sand-dwelling
309 hermit crab, *Diogenes nitidimanus* terao (Crustacea decapoda: Anomura). *Journal of*
310 *Ethology* 5(2):207-209.
- 311 Asakura A. 1995. Sexual differences in life history and resource utilization by the hermit crab.
312 *Ecology* 76:2295–2313. DOI: <https://doi.org/10.2307/1941703>.
- 313 Aviz D, Carmona PA, Barbosa AC de C, Santos CRM dos. 2022. Fecundity and reproductive
314 patterns of the fiddler crab *Uca maracoani* Latreille 1802-1803 in an Amazonian estuary
315 in northern Brazil. *Invertebrate Reproduction & Development* 66:197–207. DOI:
316 <https://doi.org/10.1080/07924259.2022.2125353>.
- 317 Baeza JA, Furlan M, Almeida AC, Barros-Alves S de P, Alves DFR, Fransozo V. 2013.
318 Population dynamics and reproductive traits of the ornamental crab *Porcellana sayana*:
319 implications for fishery management and aquaculture. *Sexuality and Early Development*
320 *in Aquatic Organisms* 1:1–12. DOI: <https://doi.org/10.3354/sedao00002>.
- 321 Bertini G, Fransozo A. 2002. Breeding season of the hermit crab *Petrochirus diogenes*
322 (Anomura: Diogenidae) in the north coast of São Paulo State, Brazil. In: Escobar-Briones
323 E, Alvarez F eds. *Modern Approaches to the Study of Crustacea*. Boston, MA: Springer,
324 145–152.

- 325 Bezerra LEA, Matthews-Cascon H. 2007. Population and reproductive biology of the fiddler
326 crab *Uca thayeri* Rathbun, 1900 (Crustacea: Ocypodidae) in a tropical mangrove from
327 Northeast Brazil. *Acta Oecologica* 31:251–258. DOI:
328 <https://doi.org/10.1016/j.actao.2006.10.003>.

- 329 Bhat M, Trivedi J. 2021. New record of the Sesarmid crab *Episesarma versicolor* (Tweedie,
330 1940) (Crustacea: Decapoda: Sesarmidae) from the West Coast of India. *Journal of*
331 *Biological Studies* 3:165–170. <https://onlinejbs.com/index.php/jbs/article/view/5447>.

- 332 Chopra BN, Das KN. 1937. Further notes on Crustacea Decapoda in the Indian Museum. IX. On
333 three collections of crabs from Tavoy and Mergui Archipelago. *Records of the Indian*
334 *Museum* 39:377–434.

- 335 Chou C-C, Head ML, Backwell PRY. 2019. Effects of temperature on reproductive timing and
336 hatching success in a tropical fiddler crab. *Biological Journal of the Linnean Society* 128.
337 DOI: <https://doi.org/10.1093/biolinnean/blz157>.

- 338 Copatti CE, Legramanti RP, Trevisan A, Santos S. 2016. Growth, sexual maturity and sexual
339 dimorphism of *Aegla georginae* (Decapoda: Anomura: Aeglidae) in a tributary of the
340 Ibicuí River in southern Brazil. *Zoologia (Curitiba)* 33. DOI:
341 <https://doi.org/10.1590/s1984-4689zool-20160010>.

- 342 Crowley CE, Shea CP, Gandy RL, Daly KL. 2019. Fecundity **assessment of stone crabs** in the
343 Eastern Gulf of Mexico. *Marine and Coastal Fisheries* 11:32–47. DOI:
344 <https://doi.org/10.1002/mcf2.10059>.

- 345 Díaz H, Conde JE. 1989. Population dynamics and life history of the mangrove crab *Aratus*
346 *pisonii* (Brachyura, Grapsidae) in a marine environment. *Bulletin of Marine Science*
347 45:148–163.
- 348 Flores A, Negreiros-Fransozo ML. 1999. Allometry of the secondary sexual characters of the
349 shore crab *Pachygrapsus transversus* (Gibbes, 1850) (Brachyura, Grapsidae).
350 *Crustaceana* 72:1051–1066. DOI: <https://doi.org/10.1163/156854099504013>.
- 351 Flores AAV, Paula J. 2002. Population dynamics of the shore crab *Pachygrapsus marmoratus*
352 (Brachyura: Grapsidae) in the central Portuguese coast. *Journal of the Marine Biological*
353 *Association of the United Kingdom* 82:229–241. DOI:
354 <https://doi.org/10.1017/s0025315402005404>.
- 355 Fransozo A, Bertini G, Correa M. 2000. Population biology and habitat utilization of the stone
356 crab *Menippe nodifrons* in the Ubatuba region, São Paulo, Brazil. *Biodiversity Crisis and*
357 *Crustacea* 12:275–281.
- 358 Fransozo A, Mantelatto FLM. 1998. Population structure and reproductive period of the tropical
359 hermit crab *Calcinus tibicen* (Decapoda: Diogenidae) in the region of Ubatuba, São
360 Paulo, Brazil. *Journal of Crustacean Biology* 18:738–745.
- 361 Gherardi F, Nardone F. 1997. The question of coexistence in hermit crabs: population ecology of
362 a tropical intertidal assemblage. *Crustaceana* 70:608–629.
- 363 Goodbody I. 1965. Continuous breeding in populations of two tropical crustaceans, *Mysidium*
364 *columbiae* (Zimmer) and *Emerita portoricensis* Schmidt. *Ecology* 46:195–197. DOI:
365 <https://doi.org/10.2307/1935274>.

- 366 Gosavi S, Purohit B, Mitra S, Patel K, Vachhrajani K, Trivedi J. 2021. Annotated checklist of
367 marine decapods (Crustacea: Decapoda) of Gujarat state with three new records.
368 *Proceedings of the Marine Biology Research Symposium – MBRS 2021*:45–66.
- 369 Gosavi S, Trivedi J, Trivedi D, Vachhrajani K. 2017a. First record of *Leucisca squalina*
370 MacLeay, 1838 (Decapoda: Leucosiidae) from Gujarat state, India. *Journal of*
371 *Entomology and Zoology Studies* 5:400–402.
- 372 Gosavi S, Trivedi J, Trivedi D, Vachhrajani K. 2017b. New records of Anomuran crabs
373 (Crustacea: Decapoda: Anomura) from Gujarat, India. *Journal of Entomology and*
374 *Zoology Studies* 5:658–662.
- 375 Hamasaki K, Ishii M, Dan S. 2021. Seasonal variability in fecundity and egg size in the
376 porcellanid crab *Petrolisthes japonicus* (Decapoda: Anomura: Porcellanidae). *Crustacean*
377 *Research* 50:17–27. DOI: https://doi.org/10.18353/crustacea.50.0_17.
- 378 Hamilton WD. 1967. Extraordinary sex ratios. *Science*. 156:477–488.
- 379 Hammer Ø, Harper DAT, Ryan PD. 2001. PAST-palaeontological statistics. *Palaeontol.*
380 *Electron* 4:1–9.
- 381 Hines AH. 1982. Allometric constraints and variables of reproductive effort in brachyuran crabs.
382 *Marine Biology* 69:309–320. DOI: <https://doi.org/10.1007/bf00397496>.
- 383 Hu MPK, Kwan B, Wang Y, Cheung S, Shin P. 2015. Population Structure and Growth of
384 Juvenile Horseshoe Crabs *Tachypleus tridentatus* and *Carcinoscorpius rotundicauda*
385 (Xiphosura) in Southern China. In: Carmichael R, Botton M, Shin P, Cheung S eds.

386 *Changing global perspectives on horseshoe crab biology, conservation and management.*
387 167–180.

388 Huang X, He L, Tan R, Feng G, Geng Z, Zhao F, Zhang T, Zhuang P. 2022. Effects of salinity
389 on reproductive characteristics and embryo quality of *Eriocheir sinensis*. *Aquaculture*
390 *Research* 53:4970–4979. DOI: <https://doi.org/10.1111/are.15983>.

391 Johnson P. 2003. Biased sex ratios in fiddler crabs (Brachyura, Ocypodidae): a review and
392 evaluation of the influence of sampling method, size class, and sex-specific mortality.
393 *Crustaceana* 76:559–580. DOI: <https://doi.org/10.1163/156854003322316209>.

394 Kneib RT, Weeks CA. 1990. Intertidal distribution and feeding habits of the mud crab *Eurytemora*
395 *limosum*. *Estuaries* 13:462–468.

396 Kobayashi M, Wong YH, OguroOkano M, Dreyer N, Høeg, Jens T, Yoshida R, Okano K. 2018.
397 Identification, characterization, and larval biology of a rhizocephalan barnacle, *Sacculina*
398 *yatsui* Boschma, 1936, from northwestern Japan (Cirripedia: Sacculinidae). *Journal of*
399 *Crustacean Biology* 38:329–340.

400 Lee SK, Mendoza JCE, Ng PKL, Kim W. 2013. On the identity of the indo-west pacific littoral
401 xanthid crab, *Leptodius exaratus* (H. Milne Edwards, 1834) (crustacea: decapoda:
402 brachyura: xanthidae). *Raffles Bulletin of Zoology* 61:189–204.

403 Litulo C. 2004. Fecundity of the pantropical fiddler crab *Uca annulipes* (H. Milne Edwards,
404 1837) (Brachyura: Ocypodidae) at Costa do Sol mangrove, Maputo bay, Southern
405 Mozambique. *Western Indian Ocean Journal of Marine Science* 3:87–91.

- 406 Litulo C. 2005. Population structure and breeding biology of the hairy crab *Pilumnus vesperilio*
407 (Fabricius, 1793) (Crustacea: Brachyura: Pilumnidae) in southern Mozambique. *Journal*
408 *of Natural History* 39:1359–1366. DOI: <https://doi.org/10.1080/00222930400010070>.
- 409 López Abellán LJ, BalgueríasE, Fernández-Vergaz V. 2002. Life history characteristics of the
410 deep-sea crab *Chaceon affinis* population off Tenerife (Canary Islands). *Fisheries*
411 *Research* 58:231–239. DOI: [https://doi.org/10.1016/s0165-7836\(01\)00384-8](https://doi.org/10.1016/s0165-7836(01)00384-8).
- 412 Lycett KA, Shields JD, Chung JS, Pitula JS. 2020. Population Structure of the Blue Crab
413 *Callinectes sapidus* in the Maryland Coastal Bays. *Journal of Shellfish Research* 39:699–
414 713. DOI: <https://doi.org/10.2983/035.039.0316>.
- 415 Mantelatto FLM, Sousa LM. 2000. Population biology of the hermit crab *Paguristes tortugae*
416 Schmitt, 1933 (Anomura, Diogenidae) from Anchieta Island. *Nauplius* 8:185–193.
- 417 Mantelatto FL, Fernandes-Góes LC, Fantucci MZ, Biagi R, Pardo LM, Marcos de Góes J. 2010.
418 A comparative study of population traits between two South American populations of the
419 striped-legged hermit crab *Clibanarius vittatus*. *Acta Oecologica* 36:10–15. DOI:
420 <https://doi.org/10.1016/j.actao.2009.09.003>.
- 421 Manzoor R, Haider S, Fatima M, Qari R. 2016. Study on abundance, breeding and growth of the
422 Ocypodide crab *Dotilla blanfordi* in Karachi Coast, Pakistan. *International Journal of*
423 *Marine Science* 6:1–14. DOI: <https://doi.org/10.5376/ijms.2016.06.0022>.
- 424 Matias D, Joaquim S, Matias AM, Leitão A. 2016. Reproductive effort of the European clam
425 *Ruditapes decussatus* (Linnaeus, 1758): influence of different diets and temperatures.

- 426 *Invertebrate Reproduction & Development* 60:49–58. DOI:
427 <https://doi.org/10.1080/07924259.2015.1126537>.
- 428 Mishima S, Henmi Y. 2008. Reproduction and embryonic diapause in the hermit crab *Pagurus*
429 *nigrofascia*. *Crustacean Research* 37:26–34. DOI:
430 https://doi.org/10.18353/crustacea.37.0_26.
- 431 Mustaqim J, Khatoon S, Rashid S. 2022. A note on sex ratio, size at maturity, fecundity and
432 breeding season of the portunid crab, *Thalamita crenata* Rüppell, 1830 from the Pakistani
433 coast. *Crustaceana* 95:127–136. DOI: <https://doi.org/10.1163/15685403-bja10179>.
- 434 Naderloo R. 2017. *Atlas of Crabs of the Persian Gulf*. Switzerland: Springer Cham. DOI:
435 <https://doi.org/10.1007/978-3-319-49374-9>.
- 436 Negreiros-Fransozo ML, Costa TM, Colpo KD. 2003. Allometric growth in the fiddler crab *Uca*
437 *thayeri* (Brachyura, Ocypodidae) from a subtropical mangrove. *Journal of Crustacean*
438 *Biology* 23:273–279. DOI: <https://doi.org/10.1163/20021975-99990337>.
- 439 Oh I-K, Lee S. 2020. Effects of temperature on the survival and larval development of
440 *Deiratonotus japonicus* (Brachyura, Camptandriidae) as a biological indicator. *Journal of*
441 *Marine Science and Engineering* 8:213–213. DOI: <https://doi.org/10.3390/jmse8030213>.
- 442 Padate VP, Patel KJ, Rivonker CU, Trivedi JN. 2022. On Indian species of *Nanosesarma*
443 Tweedie, 1950 (Decapoda: Brachyura: Sesarmidae). *Nauplius* 30:e2022031. DOI:
444 <https://doi.org/10.1590/2358-2936e2022031>.

- 445 Patel P, Patel K, Trivedi J. 2020a. First record of Hermit crab *Clibanarius ransoni* Forest, 1953
446 (Crustacea: Anomura: Diogenidae) from India. *Journal of Biological Studies* 3:19–23.
447 <https://onlinejbs.com/index.php/jbs/article/view/4601>
- 448 Patel D, Patel K, Patel P, Trivedi J. 2020b. Shell utilization pattern by the hermit crab *Diogenes*
449 *custos* (Fabricius, 1798) along Gulf of Kachchh, Gujarat, India. *Journal of Biological*
450 *Studies* 3(2):79–95. <https://onlinejbs.com/index.php/jbs/article/view/5141>
- 451 Patel K, Patel P, Trivedi J. 2021a. First record of *Manningis arabicum* (Jones and Clayton, 1983)
452 (Decapoda, Brachyura, Camptandriidae) from India. *Nauplius* 29:e2021017. DOI:
453 <https://doi.org/10.1590/2358-2936e2021017>.
- 454 Patel H, Patel K, Trivedi J. 2021b. Study of colour variation in intertidal crab *Leptodius exaratus*
455 (H. Milne Edwards, 1834) inhabiting rocky shores of Saurashtra coast, Gujarat, India. In:
456 *Proceedings of the Marine Biology Research Symposium (MBRS 2021)*. 67–80.
- 457 Patel KJ, Vachhrajani KD, Trivedi J. 2022. Study on Shell Utilization Pattern of Two Sympatric
458 Hermit Crab Species on the Rocky Intertidal Region of Veraval, Gujarat, India.
459 *Thalassas: An International Journal of Marine Sciences* 39:125–137. DOI:
460 <https://doi.org/10.1007/s41208-022-00487-5>.
- 461 Patel KJ, Vachhrajani KD, Trivedi J. 2023. Population structure and reproductive biology of
462 *Clibanarius rhabdodactylus* Forest, 1953 (Crustacea: Anomura: Diogenidae) in Gujarat
463 state, India. *Regional Studies in Marine Science* 63:103033. DOI:
464 <https://doi.org/10.1016/j.rsma.2023.103033>.

- 465 Qureshi NA, Saher NU. 2012. Density, Distribution and Population Biology of *Macrophthalmus*
466 (*Venitus*) *dentipes* Lucas, 1836, from Mangrove Areas of Pakistan. *Pakistan Journal of*
467 *Zoology* 44:615–623.
- 468 Rao GS, Rama-Sharma DV. 1990. Meiobenthos of the Gosthani estuary. *Indian Journal of*
469 *Marine Sciences* 19: 171–173.
- 470 Reese ES. 1968. Annual breeding seasons of three sympatric species of tropical intertidal hermit
471 crabs, with a discussion of factors controlling breeding. *Journal of Experimental Marine*
472 *Biology and Ecology* 2:308–318. DOI: [https://doi.org/10.1016/0022-0981\(68\)90022-1](https://doi.org/10.1016/0022-0981(68)90022-1).
- 473 Rouf MA, Shahriar SIM, Antu A-H, Siddiqui MN. 2021. Population parameters of the orange
474 mud crab *Scylla olivacea* (Herbst, 1796) from the Sundarban mangrove forest in
475 Bangladesh. *Heliyon* 7:e06223. DOI: <https://doi.org/10.1016/j.heliyon.2021.e06223>.
- 476 Saher NU, Amanat Z, Gondal MA, Qureshi NA. 2017. Distribution, abundance and population
477 ecology of *Ashtoret lunaris* (Forskel, 1775) and *Matuta planipes* Fabricius, 1798 from the
478 Sonmiani Bay (Lagoon), Pakistan. *Pakistan Journal of Zoology* 49:425–434. DOI:
479 <https://doi.org/10.17582/journal.pjz/2017.49.2.425.434>.
- 480 Saher NU, Qureshi NA. 2010. Zonal distribution and population biology of *Ilyoplax frater*
481 (Brachyura: Ocypodoidea: Dotillidae) in a coastal mudflat of Pakistan. *Current Zoology*
482 56:244–251. DOI: <https://doi.org/10.1093/czoolo/56.2.244>.
- 483 Saher NU, Qureshi NA. 2011. Density, distribution and population structure of *Opusia indica*
484 (Ocypodoidea: Camptandriidae) in a coastal mangrove creek in Pakistan. *Biologia*
485 66:138–145. DOI: <https://doi.org/10.2478/s11756-010-0142-3>.

486 Santos S, Negreiros-Fransozo ML, Padovani CR. 1995. Relação do peso em função da largura da
487 carapaça do siri candeias *Portunus spinimanus* Latreille, 1819 (Decapoda Portunidae).
488 *Arquivos de Biologia e Tecnologia* 38(3):715–724.

489 Sastry A, Vernberg FJ, Vernberg W. 1983. Ecological aspects of reproduction. In: Vernberg FJ
490 ed. *Environmental Adaptations*. New York, UK: Academic Press, 1–410.

491 Siegel V, Damm U, Neudecker T. 2008. Sex-ratio, seasonality and long-term variation in
492 maturation and spawning of the brown shrimp *Crangon crangon* (L.) in the German
493 Bight (North Sea). *Helgoland Marine Research* 62:339–349. DOI:
494 <https://doi.org/10.1007/s10152-008-0121-z>.

495 Silk JB. 1983. Local resource competition and the evolution of male-biased sex ratios. *Journal of*
496 *Theoretical Biology* 108 (2):203–213. DOI: [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-5193(84)80066-1)
497 [5193\(84\)80066-1](https://doi.org/10.1016/S0022-5193(84)80066-1).

498 Smith DR, Pooler PS, Swan BL, Michels SF, Hall WR, Himchak PJ, Millard MJ. 2002. Spatial
499 and temporal distribution of horseshoe crab (*Limulus polyphemus*) spawning in Delaware
500 Bay: Implications for monitoring. *Estuaries* 25:115–125. DOI:
501 <https://doi.org/10.1007/bf02696055>.

502 Swanson BO, George MN, Anderson SP, Christy JH. 2013. Evolutionary variation in the
503 mechanics of fiddler crab claws. *BMC Evolutionary Biology* 13(137):1–11. DOI:
504 <https://doi.org/10.1186/1471-2148-13-137>.

505 Taddei FG, Davanso TM, Castiglioni L, Herrera DR, Fransozo A, Caetano R. 2015. Population
506 structure, recruitment, and mortality of the freshwater crab *Dilocarcinus pagei* Stimpson,

- 507 1861 (Brachyura, Trichodactylidae) in Southeastern Brazil. *Invertebrate Reproduction &*
508 *Development* 59:189–199. DOI: <https://doi.org/10.1080/07924259.2015.1081638>.
- 509 Takween W, Qureshi N. 2005. Population structure and reproductive biology of four species of
510 swimming crabs (Crustacea: Brachyura: Portunidae) from coastal area of Karachi,
511 Pakistan. *Pakistan Journal of Marine Sciences* 14:107–121.
- 512 Taylor DR. 1996. Parental Expenditure and Offspring Sex Ratios in the Dioecious Plant *Silene*
513 *alba* (= *Silene latifolia*). *The American Naturalist* 147:870–879. DOI:
514 <https://doi.org/10.1086/285883>.
- 515 Thurman CL. 1985. Reproductive biology and population structure of the fiddler crab *Uca*
516 *subcylindrical* (stimpson). *The Biological Bulletin* 169:215–229. DOI:
517 <https://doi.org/10.2307/1541399>.
- 518 Tina FW, Jaroensutasinee M, Sutthakiet O, Jaroensutasinee K. 2015. The fiddler crab, *Uca*
519 *bengali* Crane, 1975: population biology and burrow characteristics on a riverbank
520 in southern Thailand. *Crustaceana* 88:791–807. DOI: [https://doi.org/10.1163/15685403-](https://doi.org/10.1163/15685403-00003450)
521 00003450.
- 522 Touchon JC, Gomez-Mestre I, Warkentin KM. 2006. Hatching plasticity in two temperate
523 anurans: responses to a pathogen and predation cues. *Canadian Journal of Zoology*
524 84:556–563. DOI: <https://doi.org/10.1139/z06-058>.
- 525 Trivedi JN, Vachhrajani KD. 2013a. Taxonomic account of genus *Scylla* (de Haan, 1833) from
526 Gujarat State, India with two new records of species. *Arthropods* 2:159–171.

- 527 Trivedi JN, Vachhrajani KD. 2013b. First record of two porcellanid crabs from Gujarat state,
528 India (Crustacea: Decapoda: Porcellanidae). *Journal of the Marine Biological*
529 *Association of India* 55:55–58. DOI: <https://doi.org/10.6024/jmbai.2013.55.1.01756-09>.
- 530 Trivedi JN, Vachhrajani KD. 2015. First record of brachyuran crab *Leptodius affinis* (De Haan,
531 1835) (Crustacea: Decapoda: Xanthidae) from the western coast of India. *Marine*
532 *Biodiversity Records* 8:1–5. DOI: <https://doi.org/10.1017/s175526721400133x>.
- 533 Trivedi J, Trivedi, D, Soni, G, Purohit B, Vachhrajani. KD. 2015a. On new records of hermit
534 crabs (Anomura: Paguroidea: Diogenidae) from Gujarat state of India. *Electronic Journal*
535 *of Environmental Science* 8:33–42.
- 536 Trivedi JN, Soni GM, Trivedi DJ, Vachhrajani KD. 2015b. A new species of *Ilyoplax*
537 (Decapoda, Brachyura, Dotillidae) from Gujarat, India. *Journal of Asia-Pacific*
538 *Biodiversity* 8:173–177. DOI: <https://doi.org/10.1016/j.japb.2015.02.005>.
- 539 Trivedi JN, Osawa M, Vachhrajani KD. 2016. A new species of the genus *Diogenes* Dana, 1851
540 (Crustacea: Decapoda: Anomura: Diogenidae) from Gujarat, northwestern India. *Zootaxa*
541 4208:189–197. DOI: <https://doi.org/10.11646/zootaxa.4208.2.6>.
- 542 Trivedi JN, Soni GM, Vachhrajani KD. 2016. A new species of *Lyphira* Galil, 2009 (Crustacea,
543 Decapoda, Leucosiidae) from Gujarat, India. *Tropical Zoology* 29:148–154. DOI:
544 <https://doi.org/10.1080/03946975.2016.1181954>.
- 545 Trivedi JN, Vachhrajani KD. 2016a. On burrow morphology of the ghost crab, *Ocypode*
546 *ceratophthalmus* (Decapoda: Brachyura: Ocypodidae) from sandy shore of Gujarat,

- 547 India. *International Journal of Marine Science* 6:1–10. DOI:
548 <https://doi.org/10.5376/ijms.2016.06.0015>.
- 549 Trivedi J, Vachhrajani K. 2016b. On new record of *Pagurus kulkarni* Sankolli, 1962 (Crustacea:
550 Anomura: Paguridae) from Gujarat, India. *International Journal of fisheries and Aquatic*
551 *Studies* 4:183–185.
- 552 Trivedi JN, Vachhrajani KD. 2017. Seasonal Burrow Distribution of the Ghost Crab, *Ocypode*
553 *ceratophthalmus* (Pallas, 1772), on Sandy Shores of Gujarat, India. *International Journal*
554 *of Marine Science* 7:344–352. DOI: <https://doi.org/10.5376/ijms.2017.07.0034>.
- 555 Trivedi JN, Trivedi DJ, Vachhrajani KD, Ng PKL. 2018. An annotated checklist of the marine
556 brachyuran crabs (Crustacea: Decapoda: Brachyura) of India. *Zootaxa* 4502:1–83. DOI:
557 <https://doi.org/10.11646/zootaxa.4502.1.1>.
- 558 Trivedi JN, Vachhrajani KD. 2018. On new record of brachyuran crab *Scopimera crabicauda*
559 Alcock, 1900 (Crustacea: Decapoda) from India. *Journal of the Marine Biological*
560 *Association of India* 60:105–107. DOI: <https://doi.org/10.6024/jmbai.2018.60.1.1945-17>.
- 561 Trivedi J, Gosavi S, Vachhrajani K. 2020. First record of three species of spider crabs from west
562 coast of India (Crustacea: Decapoda: Brachyura). *Journal of Biological Studies* 3:24–29.
- 563 Trivedi J, Mitra S, Patel P, Gosavi S, Vachhrajani K. 2020. Four New Records of Brachyuran
564 Crabs (Crustacea: Decapoda: Brachyura) from India. *Thalassas: An International Journal*
565 *of Marine Sciences* 37:235–242. DOI: <https://doi.org/10.1007/s41208-020-00252-6>.

- 566 Trivedi J, Mitra S, Patel P, Maheta N, Patel K, Ng PKL. 2021. On the Indian species of
567 *Eurycarcinus* A. Milne-Edwards, 1867, *Heteropanope* Stimpson, 1858, and *Pilumnopus*
568 A. Milne-Edwards, 1867 (Decapoda: Brachyura: Pilumnidae). *Nauplius* 29:e2021004.
569 DOI: <https://doi.org/10.1590/2358-2936e2021004>.

- 570 ViñaTrillos N, Brante A, Urzúa Á. 2023. Intraspecific variation in reproductive traits and
571 embryo elemental composition of the crab *Hemigrapsus crenulatus* (Milne Edwards,
572 1837) across fluctuating coastal environments along Chilean coasts. *Marine*
573 *Environmental Research* 188:106023. DOI:
574 <https://doi.org/10.1016/j.marenvres.2023.106023>.

- 575 Wada S, Kitaoka H, Goshima S. 2000. Reproduction of the hermit crab *Pagurus lanuginosus* and
576 comparison of reproductive traits among sympatric species. *Journal of Crustacean*
577 *Biology* 20:474–478.

- 578 Waiho K, Ikhwanuddin M, Abualreesh MH, Shu-Chien AC, Ishak SD, Jalilah M, Azmie G,
579 Fazhan H. 2021. Intra- and Interspecific Variation in Sexual Dimorphism Patterns of
580 Mud Crab Genus *Scylla* Along the Equatorial Region. *Frontiers in Marine Science* 8.
581 DOI: <https://doi.org/10.3389/fmars.2021.690836>.

- 582 Wenner AM. 1972. Sex Ratio as a function of size in marine crustacea. *The American Naturalist*
583 106:321–350. DOI: <https://doi.org/10.1086/282774>.


- 584 Zhang B, Yu C, Xu Y, Huang Z, Cai Y, Li Y. 2023. Hepatopancreas immune response during
585 different photoperiods in the Chinese mitten crab, *Eriocheir sinensis*. *Fish & Shellfish*
586 *Immunology* 132:108482. DOI: <https://doi.org/10.1016/j.fsi.2022.108482>.

587 Zimmerman TL, Felder DL. 1991. Reproductive ecology of an intertidal brachyuran crab,
 588 *Sesarma* sp. (nr. *reticulatum*), from the Gulf of Mexico. *The Biological Bulletin* 181:387–
 589 401. DOI: <https://doi.org/10.2307/1542359>.

Figure 1

Map of study area: a. Gujarat state; b. Shivrajpur village, Gujarat state, India. (Prepared using QGIS version 3.14).

a



b



Figure 2

Morphology of *L. exaratus* from Shivraipur, Gujarat state, India; a. dorsal view; b. ventral view male (CW326.4 mm); c. ventral view ovigerous female (CW318.4 mm), d. ventral view female (CW318.2 mm). Scale bar 5 mm.




Figure 3

Overall size frequency distribution of *L.exaratus* collected from Shivrajpur, Gujarat state, India.




Figure 4

Percentage of different demographic categories of *L.exaratus* from Shivrajpur, Gujarat state, India during the 12 months of study period.




Figure 5

Size frequency distribution of *L. exaratus* in each month from Shivrajpur, Gujarat state, India; a. January, b. February, c. March, d. April.




Figure 6

Size frequency distribution of *L. exaratus* in each month from Shivrajpur, Gujarat state, India; a. May b. June c. July d. August.




Figure 7

Size frequency distribution of *L. exaratus* in each month from Shivrajpur, Gujarat state, India; a. September, b. October, c. November, d. December.




Figure 8

Association between the juveniles (of both sexes) and ovigerous female occurrence of *L.exaratus* with monthly ambient and water temperatures from Shivrajpur, Gujarat state, India.




Figure 9

Relationship of *L. exaratus* carapace width (mm) with a. total number of eggs; b. egg weight; and c. average egg size; and crab weight (g) with d. total number of eggs; e. egg weight; and f. average egg size.




Table 1(on next page)

Carapace width values of male and female individuals of *L. exaratus* from Shivrajpur, Gujarat state, India. (n= total number of individuals, CW= carapace width). *Significant level if $p < 0.05$ (*); $p < 0.01$ (**); $p < 0.001$ (***)

Sex	n	Min. CW (mm)	Max. CW (mm)	Mean \pm SD
Male	558	5.15	29.98	15.967 \pm 5.27*
Female	657	5.26	28.63	15.48 \pm 3.77*

1

2

Table 2 (on next page)

Total number of *L. exaratus* specimens collected from Shivrajpur, Gujarat state, India.

(M: Male; NOF: Non-ovigerous female; OF: Ovigerous female) Chi-square test (χ^2) = 4.1219, P=0.042, the result is significant at (*) $p < 0.05$.

Month	M	%	NOF	%	OF	%	NOF+ OF	%	Sex ratio
January	51	49.51	28	27.18	24	23.30	52	50.49	1:1.02
February	56	43.41	40	31.01	33	25.58	73	56.59	1:1.3
March	55	44.00	42	33.60	28	22.40	70	56.00	1:1.3
April	55	37.41	51	34.69	41	27.89	92	62.59	1:1.7
May	69	40.83	69	40.83	31	18.34	100	59.17	1:1.4
June	20	43.48	23	50.00	3	6.52	26	56.52	1:1.3
July	17	45.95	14	37.84	6	16.22	20	54.05	1:1.2
August	44	44.00	50	50.00	6	6.00	56	56.00	1:1.3
September	54	56.25	37	38.54	5	5.21	42	43.75	1:0.8
October	45	53.57	34	40.48	5	5.95	39	46.43	1:0.9
November	44	57.89	22	28.95	10	13.16	32	42.11	1:0.7
December	48	46.60	32	31.07	23	22.33	55	53.40	1:1.1
Total	558	45.93	442	36.38	215	17.70	657	54.07	1:1.2

1

Table 3(on next page)

. Summary of different morphological parameters of *L. exaratus* ovigerous females and eggs from Shivrajpur, Gujarat state, India. (n= total individuals; SD= standard deviation).

Variables	n	Mean \pm SD	Min.	Max.
Crab weight (g)	34	2.04 \pm 1.2	0.41	4.64
Weight of egg mass (g)	34	0.29 \pm 0.18	0.04	0.88
Carapace length (mm)	34	12.14 \pm 2.4	6.92	15.88
Carapace width (mm)	34	18.1 \pm 3.8	10.38	24.02
Egg number	34	4529 \pm 2003	920	8730
Egg size (mm)	34	0.36 \pm 0.07	0.19	0.54

1