

GPR27 expression correlates with prognosis and tumor progression in gliomas

Changcheng Cai¹, Libo Hu¹, Ke Wu², Yinggang Liu^{Corresp. 1}

¹ Suining Central Hospital, Suining, China

² Xichang People's Hospital, Xichang, China

Corresponding Author: Yinggang Liu

Email address: yinggangliu@suiningcentralhosp.com

Backgrounds: Glioma is a highly aggressive type of brain tumor, and its prognosis is still poor despite recent progress in treatment strategies. G protein-coupled receptor 27 (GPR27) is a member of the G protein-coupled receptor family and has been reported to be involved in various cellular processes, including tumor progression. Nevertheless, the clinical potential and tumor-related role of GPR27 in glioma remain unknown. Here we aimed to explore the function and role of GPR27 in gliomas.

Methods: In the current study, we evaluated the expression and clinical significance of GPR27 in gliomas using data from GTEx and TCGA datasets. We also conducted cellular experiments to evaluate the functional role of GPR27 in glioma cell growth.

Results: We found that glioma tissues had significantly lower GPR27 expression levels than normal brain specimens, and its expression level was closely associated with disease status. Of note, GPR27 was negatively correlated with WHO grade, with grade IV samples showing the lowest GPR27 levels, while grade II samples showed the highest levels. Patients with IDH mutation or 1p/19q co-deletion exhibited higher GPR27 levels. In addition, lower GPR27 levels were correlated with higher death possibilities. In cellular experiments, we confirmed that GPR27 inhibited glioma cell growth.

Conclusions: Our results indicate that GPR27 may function as a potential prognostic biomarker and therapeutic target in gliomas. Further studies are needed to illustrate the signaling mechanism and clinical implications of GPR27 in gliomas.

GPR27 expression correlates with prognosis and tumor progression in gliomas

Changcheng Cai ^{1, *}, Libo Hu ^{1, *}, Ke Wu ², Yinggang Liu ¹

1. Department of Neurosurgery, Suining Central Hospital, Suining 629000, Sichuan, China.

2. Department of Neurosurgery, Xichang People's Hospital, Xichang 615000, Sichuan, China.

* Changcheng Cai and Libo Hu contributed equally to this study.

Corresponding to Yinggang Liu, Suining Central Hospital, Desheng West Road 127#, Suining 629000, Sichuan, China.

Tel: 86-19182577656

Email: lyg302905376@163.com; yinggangliu@suiningcentralhosp.com

ORCID: 0009-0001-8661-9781

Subtitle: GPR27 correlates with glioma progression

Funding: The authors received no funding for this work.

Acknowledgement: None

Conflict of Interest: None.

Data availability: Original data was provided and uploaded.

Ethics

The Ethics Committee of Suining Central Hospital requires no ethic approval and grants an exemption from informed consent for this public-database study.

Abstract

Backgrounds: Glioma is a highly aggressive type of brain tumor, and its prognosis is still poor despite recent progress in treatment strategies. G protein-coupled receptor 27 (GPR27) is a member of the G protein-coupled receptor family and has been reported to be involved in various cellular processes, including tumor progression. Nevertheless, the clinical potential and tumor-related role of GPR27 in glioma remain unknown. Here we aimed to explore the function and role of GPR27 in gliomas.

Methods: In the current study, we evaluated the expression and clinical significance of GPR27 in gliomas using data from GTEx and TCGA datasets. We also conducted cellular experiments to evaluate the functional role of GPR27 in glioma cell growth.

Results: We found that glioma tissues had significantly lower GPR27 expression levels than normal brain specimens, and its expression level was closely associated with disease status. Of note, GPR27 was negatively correlated with WHO grade, with grade IV samples showing the lowest GPR27 levels, while grade II samples showed the highest levels. Patients with IDH mutation or 1p/19q co-deletion exhibited higher GPR27 levels. In addition, lower GPR27 levels were correlated with higher death possibilities. In cellular experiments, we confirmed that GPR27 inhibited glioma cell growth.

Conclusions: Our results indicate that GPR27 may function as a potential prognostic biomarker and therapeutic target in gliomas. Further studies are needed to illustrate the signaling mechanism and clinical implications of GPR27 in gliomas.

Keywords

Biomarkers, Cell proliferation, Glioblastoma, Neuroscience, Molecular biology, Survival analysis.

Introduction

Gliomas and glioblastomas are types of primary brain tumors that originate from glial cell, which are the supportive cells of the central nervous system [1]. Glioma is the most common form of brain malignancies, accounting for about 80% of brain tumors, and they can vary in grade from slow-growing low-grade tumor to aggressive high-grade tumor. Among gliomas, glioblastoma, also known as glioblastoma multiforme (GBM), represents the most aggressive subtype, representing the highest grade of glioma [2]. Glioblastoma is characterized by its rapid growth, infiltrative nature, and resistance to treatment, making it one of the most challenging cancers to manage. Understanding the basic features and biology of gliomas and glioblastomas is crucial for advancing diagnosis, prognosis, and treatment strategies for these complex brain tumors [3].

GPR27, a member of the G protein-coupled receptor family, is a protein-coding gene located on human chromosome 3q25.1 [4]. It encodes a 7-transmembrane receptor that expresses in various tissues and is involved in multiple cellular functions including neurotransmission, immune activation, and cellular growth [5, 6]. Recent research has implied that GPR27 might play critical roles during cancer progression, particularly in hepatocellular carcinoma (HCC) [7] and breast cancer [8]. Abnormal expression of GPR27 has been observed in these cancers, and it has been recognized to contribute to tumor growth and angiogenesis. For example, Wang et al. discussed function of GPR27 in HCC progression and its potential mechanism of action through MAPK-ERK signaling pathway [7]. The study highlights the importance of GPR27 in HCC, a type of liver cancer, and proposes that GPR27 may promote tumor progression by activating the MAPK/ERK pathway. The paper provides insights into the potential molecular mechanisms underlying HCC development and indicates that GPR27 could be a valuable therapeutic target in HCC treatment. However, further study will be needed to fully understand the exact mechanisms by which GPR27 affects cancer development and to explore its potential as a treatment target for cancers.

This study sheds light on expression and clinical relevance of GPR27 in gliomas and provides evidence for its oncogenic role in glioma growth. The findings presented here have the potential

to enhance our understanding of underlying mechanism of glioma progression and could contribute to the development of novel therapeutic strategies for this devastating disease.

Methods

In silico data collection and analysis

RNA expression information in the Fragments Per Kilobase per Million format (FPKM) as well as related clinic-pathological data of glioma samples (n=689) and normal brain samples (n=1157) were downloaded from TCGA (<https://portal.gdc.cancer.gov/projects/TCGA-GBM>) and GTEx (<https://www.gtexportal.org/home/>). The clinic-pathological information of glioma patients was also extracted. The difference of various clinicopathological parameters were compared between the high-GPR27 and low-GPR27 expression groups. Regression analyses were used to assess the relationships between GPR27 expression and clinicopathological variables of glioma cases.

Survival information of glioma cases in TCGA-glioma was analyzed. Kaplan-Meier survival as well as multivariate Cox regression analyses was conducted to identify the patients' prognoses according to GPR27 level as well as other clinic-pathological characteristics.

Establishment of nomogram and calibration curves

We employed the "RMS" package in R to construct the nomogram for predicting possibility of individual survival. The calibration of the nomogram was evaluated through calibration curves.

Immune infiltration analyses

We evaluated the association between GPR27 expression and immune cell infiltration in gliomas using ssGSEA algorithm available in the "GSVA" R package [9]. This algorithm helped us assess infiltration status of 24 kinds of immune cell types [10]. We performed Spearman correlation analysis to clarify correlation between GPR27 expression and immune cell infiltration status.

Cell culture and transfection

U87 and U251 human glioblastoma cells will be obtained from ATCC and cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS and 1% penicillin-streptomycin at 37°C with 5% CO₂. Culturing medium was replaced every two days, and cells were subseeded once they reach 70-80% confluence. Transient transfection of siRNAs targeting GPR27 or scramble control siRNA was achieved with Lipofectamine 3000 based on manufacturer's instructions. Afterwards, cells were incubated to allow for transfection. After the incubation period, the transfection medium will be replaced with fresh complete growth medium to allow for further cell culture or further experimental tests.

Cell proliferation assay

Transfected cells were seeded at 5000 cells/well. Then CCK-8 experiment was done according to the manufacturer's instruction. Briefly, the medium was replaced with CCK-8 reagent, and the cells were incubated for 2 hours. The absorbance of the formazan dye, which is proportional to the number of viable cells, was measured spectrophotometrically at 450 nm. The percentage of cell viability or proliferation was calculated by normalizing to the control group. Results were expressed as mean \pm SD to determine statistical significance.

Statistics

All statistical analyses and plots were performed using R (version 4.1.3). Statistical significance was set at $P < 0.05$. * indicates $P < 0.05$, ** indicates $P < 0.01$, *** indicates $P < 0.001$.

Ethics

The Ethics Committee of Suining Central Hospital requires no ethic approval and grants an exemption from informed consent for this public-database study.

Results

Aberrant GRP27 expression in gliomas

As shown in Figure 1A, RNA expression data from GTEx and TCGA datasets suggested that glioma tissues possessed a significant lower GPR27 level than normal brain tissues ($P < 0.001$). Meanwhile, GPR27 expression was closely correlated with disease status. For example, GPR27 was negatively correlated with WHO grade, on that grade IV samples showed the lowest GPR27 level while grade II samples showed the highest GPR27 level (Figure 1B, $P < 0.001$). In contrast, patients with IDH mutation (Figure 1C) or 1p/19q co-deletion (Figure 1D) exhibited higher GPR27 levels ($P < 0.001$). Moreover, in silico analysis revealed that lower-GPR27 level was correlated with higher death rates (Figure 1E, $P < 0.001$) although its prognostic significance remains further investigation.

Correlations between GPR27 expression and clinic-pathological characteristics of gliomas

By dividing TCGA cohorts into high-GPR27 group ($n=350$) and low-GPR27 group ($n=349$), we further analyzed its correlation with clinic-pathological variables (Table 1). Accordingly, lower GPR27 expression was observed in elder patients, higher-grade gliomas, as well as in the specimens with wild-type IDH status or non-codeletion of 1p/19q (all $P < 0.001$). Considering all the above-mentioned variables had been reported to be correlated with patients' prognosis, we were engaged to further explore whether GPR27 can affect the overall survival of glioma patients.

GPR27 is an independent survival predictor of gliomas

Kaplan-Meier survival analysis showed that patients with lower GPR27 level had significantly worse prognosis (Figure 2A, $P < 0.001$). Briefly, the median overall survival time of low-GPR27 group was 26.3 months (95% CI 23.6-34.1 months), while was up to 99.6 months (95% CI 68.4-135.6 months) of high-GPR27 group. In addition to GPR27, several other variables were identified with prognostic significances in univariate analyses. For example, comparing with WHO grade II patients, WHO grade III patients showed a death hazard ratio as 2.967 (95% CI 1.986 - 4.433, $P < 0.001$) and WHO grade IV patients showed a death hazard ratio as 18.6 (95%

CI 12.448 - 27.794, $P < 0.001$). Consistent with our previous data, mutated IDH status indicated a better survival with a death hazard ratio as 0.116 (95% CI 0.089 - 0.151, $P < 0.001$) comparing to those with wild-type IDH status. Similarly, co-deletion of 1p/19q was a favorable prognostic factor with a death hazard ratio as 0.225 (95% CI 0.147 - 0.346, $P < 0.001$).

To figure out the independent prognostic factors, we further subjected the variables into a Cox multivariate regression model for survival analysis (Table 2). As a result, elder age and higher WHO grade were confirmed as independent unfavorable prognosis factors. In contrast, mutated IDH status was identified as an independent favorable prognostic factor (hazard ratio 0.265, 95% CI 0.176 - 0.399, $P < 0.001$). Importantly, our data, for the first time, showed that higher GPR27 expression was an independent benefit biomarker for glioma prognosis (hazard ratio 0.679, 95% CI 0.486 - 0.947, $P = 0.023$).

Based on the multivariate survival analysis, we also established a nomogram to help predict overall survival of glioma patients (Figure 2B); the variables in the nomogram included patients' age, gender, WHO grade, IDH status, 1p/19q co-deletion, and GPR27 expression level.

GPR27 inhibits proliferation of glioma cells

We next conducted western blot experiments to confirm the efficiency of knocking down GPR27 in U87 and U251 cell lines. The data obtained from the western blot experiments demonstrated a significant reduction in GPR27 expression levels, which indicated that the knockdown was successful (Figure 3A). To further investigate the role of GPR27 in cell proliferation, we conducted CCK-8 experiments. The results showed that silencing GPR27 significantly inhibited the proliferation (Figure 3B- 3C) capacities of both U87 and U251 cell lines. These findings suggested that GPR27 plays a tumor-promoting role in glioma cells. The inhibition of GPR27 expression may be a promising therapeutic approach for treating glioma. Overall, the results of this study provide important insights into the molecular mechanisms underlying glioma development and progression, and may have significant implications for the development of novel therapeutic strategies for this devastating disease.

GPR27 is correlated with the immune cell infiltration in gliomas

The results of GSEA showed that the GPR27 has a negative association with macrophages ($R=-0.551$, $P < 0.001$), neutrophils ($R=-0.473$, $P<0.001$), aDC ($R=-0.405$, $P<0.001$), eosinophils ($R=-0.397$, $P<0.001$), iDC cells ($R=-0.324$, $P<0.001$), cytotoxic cells ($R=-0.286$, $P<0.001$). etc. These findings indicate that GPR27 may play a crucial role in modulating the immune response in the tumor microenvironment. Oppositely, GPR27 shows positive associations with pDC cells ($R=0.325$, $P<0.001$), TFH cells ($R=0.281$, $P=0.003$), NK CD56bright cells ($R=0.251$, $P<0.001$), Tcm cells ($R=0.214$, $P<0.001$), TReg cells ($R=0.190$, $P<0.001$), etc (Figure 4A-C). These results suggest that GPR27 may play a significant role in promoting the differentiation and activation of these immune cells.

Discussions

The results presented in this study suggest that GPR27 expression is significantly reduced in glioma tissues when compared to normal brain tissues and that GPR27 expression levels are closely correlated with the disease status of glioma patients. Specifically, GPR27 expression was negatively correlated with WHO grade, and patients with IDH mutation or 1p/19q co-deletion exhibited higher GPR27 levels. Additionally, in silico analysis showed that lower GPR27 expression was correlated with higher death rates in glioma patients. In line with these findings, the authors performed further analyses and showed that low GPR27 expression was associated with elder patients, higher-grade gliomas, and wild-type IDH status or non-codeletion of 1p/19q. Importantly, Kaplan-Meier survival analysis indicated that lower GPR27 expression was a significant independent predictor of poor overall survival in glioma patients. Taken together, clinical data analyses suggest that GPR27 may serve as a useful prognostic biomarker for glioma and may have implications for the development of novel therapeutic strategies.

Consistently, cellular experiments suggest that GPR27 plays a crucial role in the development and progression of glioma, as silencing of GPR27 expression significantly inhibited the

proliferation capacities of U87 and U251 cell lines. These findings are consistent with previous reports that have demonstrated that GPR27 is overexpressed in several types of cancer and promotes tumor growth and metastasis. For example, Wang et al. confirmed that GPR27 expression was upregulated in HCC tissues and cell lines. They then used cellular experiments to show that knockdown of GPR27 inhibited HCC cell proliferation, migration, and invasion. Further analysis revealed that GPR27 acted through the MAPK/ERK pathway to promote HCC progression [7]. Besides the classical downstream G-protein signaling, GPR27 can also activate beta-arrestin-biased signaling pathways [6]. Meanwhile, beta-arrestins had been well-acknowledged to be involved in malignancies including glioblastoma [11, 12]. Therefore, it is highly likely that GPR27 may promote glioma progression via beta-arrestin-downstream signaling pathways. The successful knockdown of GPR27 in this study confirms the specificity of the effect observed on cell proliferation. Furthermore, the identification of GPR27 as a potential therapeutic target for glioma treatment is of significant clinical relevance, as there is a dire need for novel therapeutic strategies for this devastating disease. Further studies will be necessary to further explore its functional mechanisms.

The present study provides important insights underlying glioma development and progression. The observed negative correlation between GPR27 and various immune cell types suggests that GPR27 may play a role in immune evasion by glioma cells, and further investigations into the mechanism underlying this association could help to develop effective immunotherapeutic strategies [13, 14]. Moreover, the identification of GPR27 as a potential therapeutic target could pave the way for the development of new drugs that selectively inhibit GPR27 expression or function, which could offer a promising avenue for glioma treatment. However, more research is needed to fully understand the mechanisms underlying the role of GPR27 in glioma development and progression, as well as the potential therapeutic implications of targeting GPR27.

Furthermore, our data identified significant correlations between GPR27 and different immune cells. The implications of these findings could be far-reaching in the field of cancer immunotherapy. Given the crucial role that the tumor microenvironment plays in cancer

progression [15], GPR27 could serve as a potential target for the development of novel cancer immunotherapies. By inhibiting the expression of GPR27, it may be possible to modulate the immune response in the tumor microenvironment to promote an anti-tumor immune response. Alternatively, targeting GPR27 could be used to promote the differentiation and activation of immune cells that are positively associated with GPR27, such as pDC cells and TReg cells. Overall, these findings provide important insights into the role of GPR27 in the modulation of the immune response in the tumor microenvironment and could have significant implications for the development of novel cancer immunotherapies.

The study has several limitations, such as a restricted sample size, which may not be representative of the entire population. Moreover, the study's outcomes may not be generalized to other regions due to its geographic confinement. The self-reported data may also be influenced by potential biases. Furthermore, the study's cross-sectional design prevents establishing a cause-and-effect relationship between variables.

Conclusions

Taken together, our data highlights the potential clinical significance of GPR27 in gliomas. The findings indicate that GPR27 expression is significantly lower in glioma tissues than in normal brain tissues and is negatively correlated with WHO grade. Additionally, lower GPR27 expression is associated with elder patients, higher-grade gliomas, and specimens with wild-type IDH status or non-codeletion of 1p/19q, and predicts worse prognosis. These results suggest that GPR27 may be a potential prognostic biomarker for glioma patients. However, further research is needed to fully understand the role of GPR27 in gliomas and to explore its potential as a therapeutic target.

Acknowledgement

251 None.

252

Figure legends

Figure 1. The correlation between GPR27 expression and disease status in glioma.

A) RNA level data from GTEx and TCGA datasets revealed that glioma tissues had significantly lower GPR27 expression levels than normal brain tissues ($P < 0.001$).

B) GPR27 expression was negatively correlated with WHO grade, with grade IV samples exhibiting the lowest GPR27 levels and grade II samples showing the highest levels ($P < 0.001$).

C) Patients with IDH mutation showed higher GPR27 levels ($P < 0.001$).

D) Similarly, patients with 1p/19q co-deletion exhibited higher GPR27 levels ($P < 0.001$).

E) In silico analysis indicated that lower GPR27 expression levels were correlated with higher death rates ($P < 0.001$).

Figure 2. Prognostic significance of GPR27 on glioma patients.

A) Kaplan-Meier survival analysis of glioma patients based on GPR27 expression level. Patients with lower GPR27 level had significantly worse prognosis compared to those with higher GPR27 level ($P < 0.001$).

B) A nomogram established based on the multivariate survival analysis. The variables in the nomogram included patients' age, gender, WHO grade, IDH status, 1p/19q co-deletion, and GPR27 expression level.

Figure 3. GPR27-knockdown inhibit glioma progression.

A) Western blot analysis confirmed the successful knockdown of GPR27 expression in U87 and U251 cell lines.

276 B) CCK-8 assay results indicated that silencing GPR27 significantly inhibited the proliferation of
277 U87 cells compared to the control group.

278 C) CCK-8 assay results indicated that silencing GPR27 significantly inhibited the proliferation of
279 U251 cells compared to the control group.

280

281 **Figure 4. The correlation between immune cell infiltration and GPR27 expression in**
282 **glioma.**

283 A) Spearman analysis result showed the correlation between the infiltration of 24 types of
284 immune cells and GPR27 expression in glioma tissues.

285 B) and C) showed representative infiltration level of pDC and macrophages in cells with different
286 GPR27 level.

287 DC, dendritic cells; pDC, plasmacytoid DC; aDC, activated DC; iDCs, immature DCs; NK, natural
288 killer cells; Tgd, T gamma delta; TReg, regulatory T cells; Tem, T effector memory; Tcm, T
289 central memory; Th1 cells, type 1 Th cells; Th2 cells, type 2 Th cells; Th17 cells, type 17 Th cells;
290 TFH, T follicular helper.

291

292

References

1. Omuro A, DeAngelis LM. Glioblastoma and other malignant gliomas: a clinical review. *Jama*. 2013;310(17):1842-50.
2. Wen PY, Reardon DA. Progress in glioma diagnosis, classification and treatment. *Nature Reviews Neurology*. 2016;12(2):69-70.
3. Melin BS, Barnholtz-Sloan JS, Wrensch MR, Johansen C, Il'yasova D, Kinnersley B, et al. Genome-wide association study of glioma subtypes identifies specific differences in genetic susceptibility to glioblastoma and non-glioblastoma tumors. *Nature genetics*. 2017;49(5):789-94.
4. Pillaiyar T, Rosato F, Wozniak M, Blavier J, Charles M, Laschet C, et al. Structure-activity relationships of agonists for the orphan G protein-coupled receptor GPR27. *European Journal of Medicinal Chemistry*. 2021;225:113777.
5. Chopra DG, Yiv N, Hennings TG, Zhang Y, Ku GM. Deletion of Gpr27 in vivo reduces insulin mRNA but does not result in diabetes. *Scientific reports*. 2020;10(1):1-9.
6. Dupuis N, Laschet C, Franssen D, Szpakowska M, Gilissen J, Geubelle P, et al. Activation of the orphan G protein-coupled receptor GPR27 by surrogate ligands promotes β -arrestin 2 recruitment. *Molecular Pharmacology*. 2017;91(6):595-608.
7. Wang H, Du D, Huang J, Wang S, He X, Yuan S, et al. GPR27 Regulates Hepatocellular Carcinoma Progression via MAPK/ERK Pathway. *Cancer Management and Research*. 2022:1165-77.
8. Milioli HH, Tishchenko I, Riveros C, Berretta R, Moscato P. Basal-like breast cancer: molecular profiles, clinical features and survival outcomes. *BMC medical genomics*. 2017;10:1-17.
9. Hänzelmann S, Castelo R, Guinney J. GSEA: gene set variation analysis for microarray and RNA-seq data. *BMC bioinformatics*. 2013;14:1-15.
10. Bindea G, Mlecnik B, Tosolini M, Kirilovsky A, Waldner M, Obenauf AC, et al. Spatiotemporal dynamics of intratumoral immune cells reveal the immune landscape in human cancer. *Immunity*. 2013;39(4):782-95.
11. Lan T, Wang H, Zhang Z, Zhang M, Qu Y, Zhao Z, et al. Downregulation of β -arrestin 1 suppresses glioblastoma cell malignant progression vis inhibition of Src signaling. *Experimental cell research*. 2017;357(1):51-8.
12. Bae W-Y, Choi J-S, Nam S, Jeong J-W. β -arrestin 2 stimulates degradation of HIF-1 α and modulates tumor progression of glioblastoma. *Cell Death & Differentiation*. 2021;28(11):3092-104.
13. Njonkou R, Jackson CM, Woodworth GF, Hersh DS. Pediatric glioblastoma: mechanisms of immune evasion and potential therapeutic opportunities. *Cancer Immunology, Immunotherapy*. 2022;71(8):1813-22.
14. Razavi S-M, Lee KE, Jin BE, Aujla PS, Gholamin S, Li G. Immune evasion strategies of glioblastoma. *Frontiers in surgery*. 2016;3:11.
15. Barthel L, Hadamitzky M, Dammann P, Schedlowski M, Sure U, Thakur BK, et al. Glioma: molecular signature and crossroads with tumor microenvironment. *Cancer and Metastasis Reviews*. 2022:1-23.

Table 1 (on next page)

Table 1. Basic information of enrolled glioma patients.

Table 1. Basic information of enrolled glioma patients.

Characteristics	Low GPR27	High GPR27	P value
Total cases, n	349	350	
Age, n (%)			< 0.001***
≤ 60 years old	245 (35.1%)	311 (44.5%)	
> 60 years old	104 (14.9%)	39 (5.6%)	
Gender, n (%)			0.134
Female	139 (19.9%)	159 (22.7%)	
Male	210 (30%)	191 (27.3%)	
WHO grade, n (%)			< 0.001***
G2	67 (10.5%)	157 (24.6%)	
G3	119 (18.7%)	126 (19.8%)	
G4	141 (22.1%)	27 (4.2%)	
IDH status, n (%)			< 0.001***
WT	189 (27.4%)	57 (8.3%)	
Mutation	152 (22.1%)	291 (42.2%)	
1p/19q codeletion, n (%)			< 0.001***
Non-codel	328 (47.4%)	192 (27.7%)	
Codel	14 (2%)	158 (22.8%)	
OS event, n (%)			< 0.001***
Alive	162 (23.2%)	265 (37.9%)	
Dead	187 (26.8%)	85 (12.2%)	

Table 2 (on next page)

Table 2. Univariate and multivariate Cox regression analyses of the overall survival of glioma patients.

Table 2. Univariate and multivariate Cox regression analyses of the overall survival of glioma patients.

Characteristics	Total (N)	Univariate analysis		Multivariate analysis	
		Hazard ratio (95% CI)	P value	Hazard ratio (95% CI)	P value
Age	698				
≤ 60 years old	555	Reference		Reference	
> 60 years old	143	4.696 (3.620 - 6.093)	< 0.001***	1.429 (1.046 - 1.952)	0.025*
Gender	698				
Female	297	Reference		Reference	
Male	401	1.250 (0.979 - 1.595)	0.073	1.185 (0.903 - 1.553)	0.220
WHO grade	636				
G2	223	Reference		Reference	
G3	245	2.967 (1.986 - 4.433)	< 0.001***	1.873 (1.223 - 2.868)	0.004**
G4	168	18.600 (12.448 - 27.794)	< 0.001***	4.253 (2.525 - 7.162)	< 0.001***
IDH status	688				
WT	246	Reference		Reference	
Mutation	442	0.116 (0.089 - 0.151)	< 0.001***	0.265 (0.176 - 0.399)	< 0.001***
1p/19q codeletion	691				
Non-code1	520	Reference		Reference	
Code1	171	0.225 (0.147 - 0.346)	< 0.001***	0.843 (0.494 - 1.440)	0.533
GPR27	698				
Low	349	Reference		Reference	
High	349	0.326 (0.251 - 0.423)	< 0.001***	0.679 (0.486 - 0.947)	0.023*

Figure 1

Figure 1. The correlation between GPR27 expression and disease status in glioma.

A) RNAseq data from GTEx and TCGA datasets revealed that glioma tissues had significantly lower GPR27 expression levels than normal brain tissues ($P < 0.001$). B) GPR27 expression was negatively correlated with WHO grade, with grade IV samples exhibiting the lowest GPR27 levels and grade II samples showing the highest levels ($P < 0.001$). C) Patients with IDH mutation showed higher GPR27 levels ($P < 0.001$). D) Similarly, patients with 1p/19q co-deletion exhibited higher GPR27 levels ($P < 0.001$). E) In silico analysis indicated that lower GPR27 expression levels were correlated with higher death rates ($P < 0.001$).









Figure 2

Figure 2. Prognostic significance of GPR27 on glioma patients.

A) Kaplan-Meier survival analysis of glioma patients based on GPR27 expression level. Patients with lower GPR27 level had significantly worse prognosis compared to those with higher GPR27 level ($P < 0.001$). B) A nomogram established based on the multivariate survival analysis. The variables in the nomogram included patients' age, gender, WHO grade, IDH status, 1p/19q co-deletion, and GPR27 expression level.

A



B




Figure 3

Figure 3. GPR27-knockdown inhibit glioma progression.

A) Western blot analysis confirmed the successful knockdown of GPR27 expression in U87 and U251 cell lines. B) CCK-8 assay results indicated that silencing GPR27 significantly inhibited the proliferation of U87 cells compared to the control group. C) CCK-8 assay results indicated that silencing GPR27 significantly inhibited the proliferation of U251 cells compared to the control group.




Figure 4

Figure 4. The correlation between immune cell infiltration and GPR27 expression in glioma.

A) Spearman analysis result showed the correlation between the infiltration of 24 types of immune cells and GPR27 expression in glioma tissues. B) and C) showed representative infiltration level of pDC and macrophages in cells with different GPR27 level. DC, dendritic cells; pDC, plasmacytoid DC; aDC, activated DC; iDCs, immature DCs; NK, natural killer cells; Tgd, T gamma delta; TReg, regulatory T cells; Tem, T effector memory; Tcm, T central memory; Th1 cells, type 1 Th cells; Th2 cells, type 2 Th cells; Th17 cells, type 17 Th cells; TFH, T follicular helper.

