

Health and ecological risk of heavy metals in agricultural soils related to Tungsten mining in Southern Jiangxi Province, China

Jinhu Lai¹, Yan Ni², Jinying Xu^{Corresp., 1}, Daishe Wu^{Corresp. 1, 3}

¹ School of Resources and Environment and Key Laboratory of Poyang Lake Environment and Resource Utilization of Ministry of Education, Nanchang University, Nanchang 330031, China

² College of Ecology and Environment, Yuzhang Normal University, Nanchang 330031, China

³ School of Materials and Chemical Engineering, Pingxiang University, Pingxiang 337055, China

Corresponding Authors: Jinying Xu, Daishe Wu

Email address: xujy2020@ncu.edu.cn, dswncu@126.com

Background. Dayu County, a major tungsten producer in China, experiences severe heavy metal pollution. This study evaluated the pollution status, the accumulation characteristics in paddy rice, and the potential ecological risks of heavy metals in agricultural soils near tungsten mining areas of Dayu County. Furthermore, the impacts of soil properties on the accumulation of heavy metals in soil were explored.

Methods. The geo-accumulation index (I_{geo}), the contamination factor (CF), and the pollution load index (PLI) were used to evaluate the pollution status of metals (As, Cd, Cu, Cr, Pb, Mo, W, and Zn) in soils. The ecological risk factor (RI) was used to assess the potential ecological risks of heavy metals in soil. The health risks and accumulation of heavy metals in paddy rice were evaluated using the health risk index and the translocation factor (TF), respectively. Pearson's correlation coefficient was used to discuss the influence of soil factors on heavy metal contents in soil.

Results. The concentrations of metals exceeded the respective average background values for soils. The levels of As, Cd, Mo, and W exceeded the risk screening values for Chinese agricultural soil contamination and the Dutch standard. Heavy metals, especially Cd, were enriched in paddy rice. The evaluation of I_{geo} , CF, and PLI indicated that soil was polluted by Cd, Mo, and W to varying degree. The RI results indicated that Cd posed the highest risk near tungsten mining areas. The non-carcinogenic and total carcinogenic risks were above the threshold values for As and Cd. Correlation analysis indicated that K_2O , Na_2O , and CaO are main factors affecting the accumulation and migration of heavy metals in soils and plants.

Health and ecological risk of heavy metals in agricultural soils related to Tungsten mining in Southern Jiangxi Province, China

Jinhu Lai¹, Yan Ni², Jinying Xu¹, Daise Wu^{1,3}

¹ School of Resources and Environment and Key Laboratory of Poyang Lake Environment and Resource Utilization of Ministry of Education, Nanchang University, Nanchang 330031, China

² College of Ecology and Environment, Yuzhang Normal University, Nanchang 330031, China

³ School of Materials and Chemical Engineering, Pingxiang University, Pingxiang 337055, China

Corresponding Author:

Jinying Xu

Email address: xujy2020@ncu.edu.cn

Daishe Wu^{1,3}

Email address: dswncu@126.com

Abstract

Background. Dayu County, a major tungsten producer in China, experiences severe heavy metal pollution. This study evaluated the pollution status, the accumulation characteristics in paddy rice, and the potential ecological risks of heavy metals in agricultural soils near tungsten mining areas of Dayu County. Furthermore, the impacts of soil properties on the accumulation of heavy metals in soil were explored.

Methods. The geo-accumulation index (I_{geo}), the contamination factor (CF), and the pollution load index (PLI) were used to evaluate the pollution status of metals (As, Cd, Cu, Cr, Pb, Mo, W, and Zn) in soils. The ecological risk factor (RI) was used to assess the potential ecological risks of heavy metals in soil. The health risks and accumulation of heavy metals in paddy rice were evaluated using the health risk index and the translocation factor (TF), respectively. Pearson's correlation coefficient was used to discuss the influence of soil factors on heavy metal contents in soil.

Results. The concentrations of metals exceeded the respective average background values for soils. The levels of As, Cd, Mo, and W exceeded the risk screening values for Chinese agricultural soil contamination and the Dutch standard. Heavy metals, especially Cd, were enriched in paddy rice. The evaluation of I_{geo} , CF, and PLI indicated that soil was polluted by Cd, Mo, and W to varying degree. The RI results indicated that Cd posed the highest risk near tungsten mining areas. The non-carcinogenic and total carcinogenic risks were above the threshold values for As and Cd. Correlation analysis indicated that K_2O , Na_2O , and CaO are main factors affecting the accumulation and migration of heavy metals in soils and plants.

Introduction

Mining is one of the most economically important activities. However, the heavy metal pollution associated with mining is a major concern (Cheng et al., 2018; Hu et al., 2018; Liu et al., 2005). Mining and smelting operations release large quantities of heavy metals into groundwater, atmospheric dust, and runoff (Han et al., 2021; Hui et al., 2021; Mokhtari et al., 2018; Mugova & Wolkersdorfer, 2022). Through these migration pathways, heavy metals can accumulate in the soil, posing ecological risks and health risks via bioaccumulation and biomagnification. Additionally, agricultural activities such as planting, livestock rearing, and aquaculture have exacerbated the extensive distribution and bioaccumulation of heavy metals in soil (Jiang et al., 2017). This is especially the case for agricultural areas neighboring mining sites irrigated with mining sewage, and heavy metals in soil can have persistent and irreversible toxic effects on various organisms (Chen et al. 2022; Wang et al. 2018). In this context, it is of great significance to gain insight into the heavy metal pollution of soils in close vicinity to mining and agricultural areas and to understand the potential risks.

Soil properties are crucial factors influencing the distribution and environmental behavior of heavy metals. Research indicates that soil pH and organic matter directly impact the speciation of heavy metals, while soil cation exchange capacity (CEC) regulates the adsorption capacity of heavy metals in the soil (Liang et al., 2019). Studies further suggest that soil oxides can influence the transfer and accumulation of heavy metals in crops. For instance, the presence of K_2O as a component in fertilizers affects crop growth, thereby influencing the enrichment of heavy metals (Lai et al., 2023). Additionally, CaO and Na_2O can alter soil pH (Lambers & Barrow, 2020), and Fe_2O_3 can reduce the solubility of metals, thus affecting their accumulation in crops (Xu et al., 2021). Existing research indicates that the impact of soil properties on the distribution and accumulation of heavy metals is region-specific. Therefore, it is essential to conduct specific studies based on regional differences.

Southern Jiangxi, particularly the southern region of Ganzhou, is recognized as one of the world's crucial tungsten mining centers (Liao et al., 2020). The substantial wastewater, emissions, and residues generated during the extraction of mineral resources introduce tungsten and associated heavy metals into the surrounding farmlands (Hu, 2012), causing significant heavy metal pollution in many agricultural soils. For example, existing studies indicate that in the vicinity of the tungsten mining areas in southern Jiangxi, levels of arsenic, cadmium, mercury, and lead in vegetable gardens and rice fields significantly exceed Chinese soil standards (Chen et al., 2016; Zhou et al., 2021), resulting in extremely high ecological risks. This severely impedes the healthy development of local agriculture. Therefore, it is essential to conduct relevant research, assess heavy metal pollution, accumulation, and potential ecological and health risks in farmlands around tungsten mines, laying the theoretical foundation for the sustainable development of agriculture in mining areas.

Dayu County, situated in mountainous terrain north of the Zhangjiang River Basin, Southern Jiangxi Province, China, is a representative region with a long history of tungsten mining, dating back to the early 20th century. The mining activities have led to metal pollution of the

surrounding agricultural soil (Wang et al., 2015; Liu et al., 2017). For example, Wang et al. (2015) studied heavy metal pollution in farmland around the tungsten mine in Dayu County and reported that 67% of the studied area was contaminated by heavy metals. Liu et al. (2017) found that the heavy metals (Pb, Mn, Zn, and Cd) near the tailings mainly occurred in bioavailable acid-soluble forms, posing high ecological risks. However, previous studies mainly focused on evaluating pollution in soil near mining areas or tailing ponds. Up to now, few studies have comprehensively analyzed the pollution and the ecological and health risks of metals in agricultural soils affected by mining activities in Dayu County, which is of great significance to the health of agricultural development and human health. In this context, the present study investigated six sites surrounding the Dayu tungsten mine, with the following aims: (i) to assess metal pollution (As, Cd, Cr, Cu, Mo, Pb, W, and Zn) in soil along the Zhangjiang River Basin; (ii) to evaluate ecological and health risks posed by heavy metals in paddy rice and soil near the tungsten mining area; (iii) to explore the influence of soil properties on the heavy metals. The results provide a basis for science-based strategies for the remediation and protection of soils in the tungsten mining region.

Materials and Methods

Study area

Dayu County is located in the southwest of Jiangxi Province, China (E114°–144.44°, N25.15°–25.37°) (Fig. 1). This region contains the highest concentration of tungsten deposits globally, with considerable tungsten-tin mineral occurrences (Zhang et al., 2021). The climate is a typical subtropical humid monsoon with distinct four seasons and abundant rainfall. The annual average temperature is between 19.1 and 20.8°C, and the annual average precipitation is 1318.9 mm. The topography rises in the northern, western, and southern parts of the county, whereas the central and eastern sections form a hilly basin surrounded by mountains on three sides that opens eastward. The Zhangjiang River crosses the county from west to east. The tungsten mines are distributed across the northern mountainous terrain, and farmland occupies the eastern plain. Long-term mineral development has become an important source of heavy metal contamination in this region.

Soil sampling and chemical analysis

Soil samples were taken at a depth of 0–20 cm at six selected areas in four towns (Fujiang, Huanglong, Qinglong, and Chijiang) along the Zhangjiang River (Fig. 1) on sloping land, sloping plain transition land, and plain land. The soil types in the study area are classified as red soil, yellow soil, and alluvial soil. The six sampling sites were termed FJ-N, FJ-S, HL, QL, CJ-W, and CJ-E; FJ-N and FJ-S are sloping land with artificial terraced fields located along the Fujiang River, a main tributary of the Zhangjiang River, located 4 and 2 km, respectively, from historical tungsten mining areas (Xihuashan, Fig. 1), respectively, HL is a control site far away from mining areas, QL is close to active northern mining areas, and CJ-W and CJ-E are plain sites with traditional paddy rice cultivation. To obtain representative samples, at each site, four to

six subsamples were obtained within a 10×10 -m grid at the center and at the four corners of an ideal square and combined to obtain one composite sample (weight: 1 kg) per site, resulting in a total of 72 soil samples and 26 corresponding rice grain samples. We removed stones, plant debris, and other residues from the soil samples. Due to the influence of terrain, the distance between sampling plots was 50–200 meters. Soil samples were obtained using wooden shovels and stored in polyethylene bags. In the laboratory, all samples were air-dried at room temperature, crushed, homogenized, and sieved through a 100-mesh nylon mesh prior to further analysis. Rice grains were placed in a clean, sunny, and ventilated room on a clean wooden board and frequently turned to prevent mold growth. After drying, grain samples were weighed, sieved through a 100-mesh nylon mesh, and stored in polyethylene bags before digestion.

The concentrations of metals (As, Cd, Cu, Cr, Pb, Mo, W, and Zn) in soils were analyzed using an inductively coupled plasma-mass spectrometer (ICP-MS, iCAP Q, Thermo, USA). Prior to analysis, the soil and rice samples were digested by a H_2SO_4 - HNO_3 -HF mixture using a Microwave Digestion System (Multiwave PRO, Anton Paar, Austria) (Li et al. 2006). A blank control, duplicate samples, and standard reference materials (standard sample of lateritic, GSS-5) were employed to ensure data quality. The concentrations of heavy metals were computed as the average values of duplicate samples. The precision and accuracy of the method is expressed in standard deviation and relative error from GSS-5(n=12), respectively. The standard deviation(SD%) of As, Cd, Cr, Cu, Mo, Pb, W, Zn is 2.47, 0.02, 1.77, 2.07, 0.16, 2.37, 0.25, 2.60, respectively. The relative error(RE%) of As, Cd, Cr, Cu, Mo, Pb, W, Zn is 1.12, 2.56, 0.99, -0.26, 2.79, 0.11, -0.55, 1.15, respectively.

In the soil, pH, total organic carbon (TOC), cation exchange capacity (CEC), and metal oxides (Fe_2O_3 , CaO, Na_2O , and K_2O) were also measured. The soil pH was measured using a pH meter in a 1:2.5 soil: water mixture, and the soil TOC was determined via a Torch TOC combustion analyzer (multi N/C 2100, Jena, Germany). The CEC was determined as described elsewhere (Chen & Huang, 2021), and the levels of Fe_2O_3 , CaO, Na_2O , and K_2O were measured via X-ray fluorescence spectrometry (XRF, EDX-8000, Shimadzu, Japan).

Assessment of soil heavy metal contamination

(1) Geo-accumulation (I_{geo}) index

The geo-accumulation (I_{geo}) index was employed to assess the pollution degree of heavy metals in soil (Muller, 1969), according to Equation (1):

$$I_{geo} = \log 2 \left[\frac{C_i}{1.5B_i} \right], \quad (1)$$

where C_i is the mean concentration of the element in the examined soil, and B_i is the geochemical background value in the soil of Jiangxi Province (As:10.4, Cd:0.10, Cu:20.8, Cr:48.0, Pb:32.1, Mo:0.30, W:4.93, Zn:69.0,mg/kg) (CNEMC 1990). The soil pollution levels can be partitioned into seven classes based on the values of I_{geo} : Class 0 (unpolluted; $I_{geo} \leq 0$), Class 1 (unpolluted to moderately polluted; $0 < I_{geo} \leq 1$), Class 2 (moderately polluted; $1 < I_{geo} \leq 2$), Class 3 (moderately to heavily polluted; $2 < I_{geo} \leq 3$), Class 4 (heavily polluted; $3 < I_{geo} \leq 4$),

Class 5 (heavily to extremely polluted; $4 < I_{geo} \leq 5$), and Class 6 (extremely polluted; $I_{geo} > 5$) (Loska et al., 1997).

(2) Contamination factor (CF)

The level of soil contamination is expressed in terms of a contamination factor (CF), calculated following Equation (2):

$$CF = \frac{C_i}{B_i} \quad (2)$$

where C_i is the mean concentration of the element in the examined soil, and B_i is the background value of metals in the soil of Jiangxi Province (As:10.4, Cd:0.10, Cu:20.8, Cr:48.0, Pb:32.1, Mo:0.30, W:4.93, Zn:69.0, mg/kg) (CNEMC 1990). where the contamination factor $CF < 1$ refers to low contamination, $1 \leq CF < 3$ indicates moderate contamination, $3 \leq CF \leq 6$ indicates considerable contamination, and $CF > 6$ indicates very high contamination.

(3) Pollution load index (PLIs)

The overall metal load in soils from each site was computed using the pollution load index (PLIs) (Tomlinson et al., 1980) according to Equation (3):

$$PLIs = (CF_1 \times CF_2 \times CF_3 \times \dots CF_n)^{\frac{1}{n}} \quad (3)$$

According to Tomlinson, CF_n is the contamination factor of the metal n in the sample. The PLIs provide simple but comparative means for assessing site quality, where a value of $PLIs < 1$ denotes low contamination, $1 < PLIs < 2$ denotes moderate contamination, $2 < PLIs < 3$ denotes heavy pollution, and $PLIs > 3$ denotes extremely heavy pollution.

Ecological risk assessment regarding metals in soils

The ecological risk factor (E_r^i) is a measurement of the potential ecological risk (RI) of a given metal and defined as Equation (4):

$$E_r^i = T_r^i \left(\frac{C_r^i}{B_i} \right), \quad (4)$$

where B_i is the background value of metals in soil in Jiangxi Province (As:10.4, Cd:0.10, Cu:20.8, Cr:48.0, Pb:32.1, Mo:0.30, W:4.93, Zn:69.0, mg/kg) (CNEMC 1990), C_r^i is the heavy metal content in the soil, T_r^i is the toxic-response factor, and the T_r^i values of Cr, Cu, Zn, As, Cd, Pb, and W are 2, 2.5, 10, 30, 5, and 2, respectively (Zheng et al., 2020). Hakanson introduced the potential ecological risk index (RI) as a method to quantitatively evaluate the risk of soil heavy metal pollution. The RI represents the integrated potential ecological risk, calculated as the sum of individual risk factors (E_r^i) for each metal contaminant (Hakanson 1980), and is defined as follows (Equation (5)):

$$RI = \sum E_r^i \quad (5)$$

According to Hakanson, E_r^i and RI represent potential ecological risks through different levels (Table 1).

Health risk assessment of heavy metals in paddy rice

As Dayu County is a major rice production area in China, with rice being a staple food for a large part of the population, heavy metal pollution must be considered. The health risk assessment model developed by the United States Environmental Protection Agency (USEPA) was used to evaluate the carcinogenic and non-carcinogenic risks of heavy metals (USEPA, 2011), using Equations (6), (7), (8), (9), and (10):

$$ADD_i = \frac{C_i \times IR \times EF \times ED}{BW \times AT} \quad (6)$$

$$HQ_i = \sum \frac{ADD_i}{RfD} \quad (7)$$

$$HI = \sum HQ_i \quad (8)$$

$$CR_i = \sum ADD_i \times SF_i \quad (9)$$

$$TCR = \sum CR_i \quad (10)$$

where HQ_i and HI are the non-carcinogenic health risks of single and combined metals, respectively, CR and TCR are the carcinogenic health risks of single and combined metals, respectively (Hui et al., 2021), ADD_i refers to the daily intake of heavy metal i in paddy rice (mg/kg/day), C_i is the mean value of metal i in rice (mg/kg), EF is the exposure frequency (365 days/year), ED is the exposure duration (30 years), IR indicates the daily intake amount of rice (kg/day), which was 410.13 g/day according to MEP (2013), BW is the body weight (60.6 kg) (MEP, 2013), AT refers to the average exposure time ($365 \times ED$ days for non-carcinogenic risk; 365×70 days for carcinogenic risk), RfD is the corresponding reference dose of the metals (As: 0.0003, Cd: 0.001, Cr: 0.003, Cu: 0.04, Mo: 0.005, Pb: 0.004, and Zn: 0.3) (mg/kg/day), SF is the slope factor of the metals (As: 1.5, Cd: 6.1, Cr: 0.5, and Pb: 0.0085) (mg/kg/day).

The translocation factor (TF) was used to assess the bioaccumulation abilities of the heavy metals in rice, as described in Equation (11):

$$TF = C_{\text{rice}}/C_{\text{soil}} \quad (11)$$

Statistical analysis

The differences in the heavy metal contents in soils from different sampling sites were determined via one-way analysis of variance (ANOVA). A Shapiro–Wilk test and Leven’s test were performed to ensure the normality and homogeneity of the data prior to ANOVA ($p < 0.05$ or $p < 0.01$). Pearson’s correlation analysis was applied to evaluate the influences of soil factors on the heavy metals in the soils. Statistical analyses were conducted in SPSS 26.0 (IBM, USA) and Origin 2021b (OriginLab Corporation, USA).

Results and discussion

Concentrations and spatial distribution of heavy metals in soil

Figure 2 and Table S1 show the concentrations and basic statistics of the heavy metals in the tested topsoil samples. The mean concentrations of As, Cd, Cu, Cr, Pb, Mo, W, and Zn exceeded their respective average background values (ABV) of Jiangxi Province soil (CNEMC 1990), accounting for 87.5%, 98.6%, 90.7%, 86.1%, 76.4%, 100%, 97.2%, 70.8%, and 20% in the samples, respectively. The levels of Mo and W surpassed the background values by 34.1 and 19.3 times, respectively, and the Cd concentration exceeded the background level by 6.23 times. The mean values of As, Cu, Cr, Pb, and Zn were 1.33–3.21 times higher than the background values. The coefficient of variation (CV) represents the spatial dispersion and degree of variation of heavy metal concentrations, and the larger its value, the greater the impact of external factors on the pollution level (Pan et al., 2016). The CV values for Cd and Mo indicated strong spatial variation ($CV > 100\%$), likely due to anthropogenic activities (Zhang, 2020). The levels of Cu, Cr and Zn showed a moderate spatial variation ($21\% < CV < 50\%$), and for As, Pb, and W, we observed a high variability ($50\% < CV < 100\%$). The risk screening values for agricultural soil contamination are 30, 0.3, 50, 150, 250, and 200 mg kg^{-1} for As, Cd, Cu, Cr, Pb, and Zn, respectively (MEEC, 2018). For Mo and W, the screening values of 3 mg kg^{-1} refer to the Dutch standard (VROM, 2000). In the present study, the mean As and Cd concentrations were 1.11 and 2.07 times, respectively, their corresponding MEEC limits. However, compared to the Dutch standard, the As and Cu levels indicated mild contamination, whereas the levels for Mo and W suggested strong pollution. Compared with the findings for a northern tungsten mine in China, in this study, the contents of heavy metal were much lower, except for Mo and Cr (Wu et al., 2020). The heavy metal concentrations reported for a southern tungsten mining area were not significantly different to those found in the present study, which could be explained by the soil properties of both sites (Guo et al. 2017; Zheng et al. 2020). In this study, As, Cd, Mo, and W were the major pollutants, similar to the findings for residential areas near a Russian tungsten mine (Timofeev et al., 2018).

The mean concentrations of the eight tested heavy metals followed the order $FJ-S > QL > FJ-N > HL > CJ-E > CJ-W$, with a significant distribution throughout the Zhangjiang River basin (Zheng et al., 2020). The spatial distribution of heavy metal concentrations is shown in Figure 1S to Figure 8S. This indicates that the degree of pollution in the sampling areas was significantly impacted by the distance from the tungsten mining area. Sites FJ-S and FJ-N were only 2 and 4 km, respectively, away from the abandoned mining area (Xihuashan) (Wang et al. 2009), with a history of 100 years of mining, resulting in long-term metal accumulation. Site QL was closer to the current mining area Piaotang (Han et al., 2019), with more active metal accumulation. As a control point in the valley on the southern bank of the Zhangjiang River, HL was not only far from historical tungsten mining areas but also was not impacted by the operating mines. Therefore, the concentrations of heavy metals were close to those detected in traditional agricultural areas.

Correlation between heavy metals and soil properties

Table 2 shows the soil properties for the different sampling sites. Of all soil samples, 73.7% were weakly acidic. The pH values ranged from 3.74 to 6.67, with an average of 5.24 ± 0.46 . The average pH value was similar to the background value determined for Ganzhou area (Chen et al., 2019; Wang et al. 2015, 2019). The highest and lowest mean pH values were observed for FJ-N (5.37) and HL (4.93), respectively. This variation can be attributed to differences in resident population and human activities, which can notably influence the soil properties (Shen et al., 2017; Vega et al., 2004).

The concentration of total organic carbon (TOC) in soil samples from site FJ-S was highest, with a mean value of 2.50%. In comparison, soil samples from sites QL, CJ-W, and CJ-E sites showed lower TOC levels, with mean values of 1.83%, 1.85%, and 2.05% respectively. The fractionation and composition of TOC is of great importance when determining the adsorption capacity of soils. The higher TOC content at site FJ-S suggests a greater capacity to adsorb pollutants, likely due to higher inputs of organic matter, such as vegetation litter, expanding the carbon pool. The lower TOC levels in the other sampling sites indicate a more limited organic matter accumulation, constraining the adsorptive potential of the contaminants (Xiao et al. 2017). However, at site QL, with high heavy metal concentrations, an increased level of human activities could result in a low TOC level, which was the reason for the bioavailable metal supplementation via sewage effluent or mineral fertilizers (Nacke et al., 2013).

The concentrations of alkali metal oxides (K_2O and Na_2O) were below the levels of the continental crustal abundances in China (Li et al., 2014). The content of K_2O was significantly higher at FJ-S, QL, and CJ-W (mean value: 2.54%, 2.43%, and 2.54%, respectively), whereas that of Na_2O was significantly higher at FJ-S (mean value: 0.33%). The CaO concentration was higher at sites FJ-S, QL and CJ-E (mean values: 0.25%, 0.26%, and 0.24%, respectively), but this heavy metal was strongly positively correlated with pH (Table S2). At site FJ-N, levels of the above four metallic oxides were lowest. Sites CJ-W and CJ-E had the lowest Fe_2O_3 concentration (mean values: 2.55% and 2.88%, respectively).

Based on the Pearson's correlation coefficients (Table 3), Mo was significantly positively correlated with pH ($p < 0.05$), whereas Cr and Zn were highly significantly positively correlated with TOC ($p < 0.01$). The Cr was highly significantly positively correlated with CEC ($p < 0.01$) and significantly positively correlated to Fe_2O_3 ($p < 0.05$). These results indicate that higher pH values can facilitate the accumulation of Mo in soil, whereas Cr and Zn are mainly accumulated in the TOC. Consequently, rapid organic degradation leads to rapid Zn cycling (Liu & Han, 2021). The Cr levels slightly fluctuated, and Cr was positively correlated with Fe_2O_3 and TOC, indicating that it was derived from natural sources. The heavy metals As, Cd, Cu, Mo, Pb, W, and Zn were highly significantly positively correlated with K_2O and Na_2O ($p < 0.01$), whereas CaO was significantly positively correlated with Mo and Pb ($p < 0.05$) and highly significantly positively correlated with Cu and Zn ($p < 0.01$). As seen in Table S5, CaO was strongly positively correlated with pH and CEC, thereby affecting the bioavailability of heavy metals such as Mo, as a result of mine wastewater treatment (Chen et al., 2007; Meng et al., 2021). This leads us to infer that K_2O and Na_2O may be principal factors in the accumulation and migration

of heavy metals, such as As, Cd, Cu, Mo, Pb, W, and Zn, in soil, whereas CEC and TOC play an important role in Cr accumulation. According to previous studies, K_2O , Na_2O , and CaO are mainly derived from human activities. Whilst K_2O accumulation is the result of the use of inorganic and organic fertilizers (Lambers & Barrow, 2020), Na_2O and CaO are the main agents in the treatment of tungsten ore mining wastewater (Zhang et al., 2012). This suggests that long-term tungsten ore smelting and agricultural fertilizing have a significant impact on the soil heavy metal content, which has also been reported in previous studies (Hui et al., 2021; Zhang et al., 2017).

Pollution and risk assessment of heavy metals in soil

The geo-accumulation index (I_{geo}) has been widely used to assess the pollution status of metals in sediments and soils (Sawe et al., 2019; Timofeev et al., 2020; Zahra et al., 2014). In the present study, the mean values of I_{geo} in soil followed the order $Mo > W > Cd > As > Pb > Cr > Zn$ (Table S3). Regarding the pollution classes, Mo and W (with mean values of 2.59 and 2.51, respectively) pollution was moderate to heavy, whereas Cd pollution was moderate (class 3), and for As and Cu, the levels indicated no to moderate pollution. The mean contamination factors (CFs) (Rastegari Mehr et al., 2017) for As, Cd, Cu, Cr, Pb, Mo, W, and Zn were 2.98, 5.56, 1.83, 1.31, 1.67, 15.16, 16.55, and 1.41, respectively (Table 4). The contamination factors for Cu, Cr, Pb, and Zn indicated moderate contamination (Inengite et al., 2015) and were significantly lower compared to those for the other tested metals; this leads us to infer that they were derived from natural sources rather than human activities. Although As pollution was moderate, it was close to being considerable. The highest contamination factors were observed for Mo and W, indicating very high contamination (mean values: 15.16 and 16.55, respectively), which was due to the long-term tungsten mining in this area (Wu, 1993). In contrast, Cd pollution was moderate to high. The main heavy metals found across all sampling sites were Mo, W, and Cd. As their levels greatly exceeded the ABV, they need to be taken into consideration regarding environmental pollution (Shi et al., 2022).

Figure 3 shows the I_{geo} spatial distribution of the tested heavy metals. Soil I_{geo} values for As, Cu, Cd, and Zn below 0 were found in sites CJ-W and CJ-E, indicating that agricultural soils were largely unaffected by these metals. For Mo and W, the I_{geo} values were higher than 2 and lower than 4 in sites FJ-N, FJ-S, HL, and QL, classifying these sites as moderately to heavily polluted and heavily polluted, respectively. For As, moderate pollution was observed, whereas for Cu, Pb, and Zn, the sites were categorized as unpolluted to moderately polluted. Additionally, Cd pollution was moderate to heavy at sites FJ-S and QL. Based on the I_{geo} findings, sites FJ-S and QL received varying amounts of heavy metals, except Cr. The PLI findings indicated extremely heavy pollution ($PLI > 3$) at sites FJ-S and QL, mostly with Cd, Mo, and W (Table 4). Sites CJ-W and CJ-E exhibited low pollution levels compared to the other sites. The heavy metal accumulation period of site FJ-S was prolonged, and the tailing pond located in the valley was prone to leakage, resulting in the pollution of both surface water and groundwater (Reutova

et al., 2022). Site QL served as a busy transportation hub for tungsten ore, and the accumulation of heavy metals was worsened by irrigation with sewage (Lu et al., 1997).

Table 5 shows the ecological risk posed by heavy metals. The ecological risk factor (E_r^i) values for Cr, Pb and Zn were below 40 in all sampling areas. However, for As, the E_r^i values were close to 40 in sites FJ-S and QL (mean: 41.32 and 39.25, respectively), indicating a moderate risk. The E_r^i values of W (mean: 35.04, 63.98, 34.99 and 36.72) had reached or close to a moderate risk in sites FJ-N, FJ-S, HL and QL near the Tungsten mining area. For Cd, the risk degree ranged from considerable risk to very high risk, indicating serious contamination and human health risks. The RI values indicated a considerable risk at FJ-S (value: 456) and a low risk at CJ-E (value: 124), whereas for the other sites, a moderate risk was detected. Risk analysis revealed Cd as the primary contaminant of concern, with sites FJ-S and QL exhibiting the highest risk quotients and contamination levels. Regarding other heavy metals such as As and W, some sites were moderately polluted. The overall ecological risk pattern highlights Cd contamination, especially at sites FJ-S and QL, requiring targeted risk management.

Concentrations and transfer of metals in paddy rice

Compared to the concentrations of heavy metals for paddy rice (Table 6) (As, $p < 0.01$; Cr, Pb, Mo, Zn, $p < 0.05$; Cd, Cu, $p > 0.05$), the mean values of Cd, Cu, Mo, Pb, and Zn were the highest in FJ-S, whereas the As concentration was highest in FJ-N. In CJ-W, the heavy metal concentrations for paddy rice were the lowest. The total concentration of heavy metals in paddy rice followed the order FJ-S > FJ-N > QL > CJ-E > HL > CJ-W. For each metal of accumulation in paddy rice followed the order Zn > Cu > Cd > As > Cr > Pb > Mo; the Cr concentration was largely stable. Based on these results, large amounts of heavy metals had accumulated in the farmland next to the historical mining area, which further exacerbated metal accumulation in rice. Active mining areas also release large amounts of heavy metals into the soil. As heavy metals are affected by rainfall and enter irrigation water, they accumulate in rice plants (Wu et al., 2020). Based on the TF values, there were differences in the concentration accumulation values (Table S4). Site QL showed the highest TF values for As, Cd, Cu, Mo, and Zn, whereas for Pb and Cr, the highest values were found for CJ-E. For all tested metals, the values were below 1, except for Cd, indicating that paddy rice had a poor ability to accumulate metals in soil. The TF values for Cd were above 2 in FJ-S, HL, and CJ-E and above 5 in QL, indicating serious accumulation.

Soil properties and composition play crucial roles in metal accumulation (Chen et al. 2016; Tunc & Sahin, 2017). In this study, Pearson correlation analysis revealed a significant association of Na_2O and K_2O with the concentration and transfer of specific metals such as Cd. However, this contradicts the expected positive relationship between Na_2O , K_2O , and Cd (Table S5), which suggests that Na_2O and K_2O might hinder metal accumulation in paddy rice by precipitating metals in soils. Previous research has indicated that sodium and potassium are essential nutrients for plant growth, suggesting that they may induce metal accumulation in

plants by promoting plant growth (Shrivastav et al. 2020; Xu et al. 2020). This could potentially explain the positive influence of K_2O and Na_2O on Cd accumulation observed in this study. Furthermore, Na_2O and K_2O might also enhance Cd accumulation in paddy rice by altering the structure of microorganisms (Yang et al. 2020; Wang et al. 2021; Zheng et al. 2021). Nevertheless, this study did not assess indicators related to rice growth and soil microorganisms. Our future research will continue to investigate the impact of K_2O and Na_2O on Cd accumulation, considering rice growth and soil microorganisms. Additionally, we aim to further explore the primary influencing factors of heavy metal enrichment in rice in mining areas.

Assessment of the health risks posed by heavy metals in paddy rice

Figure 4, Figure 5 and Table S6 show the results of the health risk assessment. The average daily dose (ADD) values for non-carcinogenic risk were highest for Cd, Cu, and Zn at site FJ-S, exceeding 0.001 mg/kg/day. However, As was the only metal with an ADD above the 0.001 mg/kg/day reference limit, specifically at site FJ-N. The hazard quotient (HQ) values for As were above 1 across all study areas, whereas that for Cd was below 1 only at CJ-W. At HQ values above 1, there are potential non-carcinogenic effects on humans (USEPA 2001). The order of As HQ was FJ-N > FJ-S > QL > CJ-E > HL > CJ-W (values: 4.03, 3.32, 2.48, 2.47, 2.14, and 1.63, respectively), accounting for 50.88%, 25.40%, 26.05%, 34.69%, 30.01%, and 44.17% of the hazard index (HI), respectively. The levels of Cd HQ followed the order FJ-S > QL > HL > CJ-E > FJ-N > CJ-W (values: 7.99, 5.61, 3.61, 3.18, 2.41, and 0.77), accounting for 61.13%, 58.93%, 50.63%, 44.66%, 30.43%, and 20.87% of the HI, respectively. The HI value exceeded 10 at site FJ-S, indicating health risks and risk of chronic poisoning. The total cancer risk (TCR) was significant across all sites, with As, Cd and Cr cancer risk values (CR) above $1 \times 10^{-4} a^{-1}$ (Dong et al., 2022), indicating high carcinogenic risk. Further, As and Cd may represent a considerable potential ecological risk at some sites (FJ-S and QL), based on the pollution characteristics of agricultural soil in mining zones (Du et al., 2015). According to the conclusion above, it is crucial to focus on the heavy metal pollution situation at sites FJ-S and QL, reduce the planting of edible crops, and perform soil remediation.

Conclusions

The results of this study indicated that the concentrations of heavy metals were higher at sites near tungsten mines and lower in traditional agricultural areas far from tungsten mines, both for soil and rice samples. In paddy rice, Cd was the most abundant heavy metal. The evaluation of I_{geo} and PLI indicated that As, Cd, Mo and W were the main metals polluted the studied soils. ~~The use of agricultural fertilizers, sewage irrigation, and the purification and disposal of mining wastewater leading to heavy metal contamination of soil. Measurements like restoring metal polluted soil by physical, chemical or biological methods should be taken to reduce the ecological and health risk of metals in agricultural soil.~~ The RI evaluation indicates that Cd and As pose a threat to human health in the study region. Additionally, the correlation analysis indicated that soil K_2O , Na_2O , and CaO are principal factors affecting the accumulation and

migration of heavy metals in soil. This study can provide precious information for managing metal polluted paddy soil near mining area. However, there are limitations for this study: Firstly, the fraction of metals is not considered, which is quite vital for evaluating the pollution and accumulation of metals; Secondly, seasonal change of metal fraction in subtropical area is obvious because of the seasonal rainfall, which is not considered in this study. Therefore, our future research should focus on the pollution and accumulation of metals in soils with considering metal fraction and seasonal rainfall to clarify the potential mechanism.

References

- Chen M, Yang T, Yang Q, Xu H, and Nie J. 2016. Characterization of heavy metal pollution in vegetable field soils and health risk assessment in Dayu County, China. *Agricultural Engineering* 48:95-102.
- Chen M, Li F, Hu L, Yang T, Yang Q, Tao M, and Deng Y. 2019. Heavy Metal Pollution in Topsoil and Vegetables in the Typical Mining Area Near Gannan, Jiangxi Province, China. *Environmental Engineering Science* 36:1307-1314. DOI 10.1089/ees.2019.0013
- Chen M, Zhu Y, and Huang W. 2007. On Treatment of Tailing Dam Wastewater by Using Lime for Elimination of Colloid Stability and Flocculating Sedimentation. *China Tungsten Industry*:43-46.
- Chen M; Zheng X; Tao M; Li F; Hu L; Shi Y; Liu Y. 2020. Pollution characteristics and risk assessment of heavy metals in sediment from Taojiang River Basin. *Environmental Chemistry* 39: 2784-2791
- Chen Y, and Huang C. 2021. Hexammine Cobaltic Chloride Exchange Method Verification for Soil Cation-Exchange Capacity. *Anhui Chemical Industry* 47:128-131.
- Chen Y, Qu X, Zhang B, Li X, Zhu G, and Zha T. 2022. Ecological and Health Risk Assessment of Heavy Metal Pollution in Farmland Soil of Xianghe County. *Environmental Science* 43:5728-5741. DOI 10.13227/j.hjx.202202122
- Cheng X, Danek T, Drozdova J, Huang Q, Qi W, Zou L, Yang S, Zhao X, and Xiang Y. 2018. Soil heavy metal pollution and risk assessment associated with the Zn-Pb mining region in Yunnan, Southwest China. *Environmental Monitoring and Assessment* 190:194. DOI 10.1007/s10661-018-6574-x
- China National Environmental Monitoring Centre(CNEMC). 1990. *The Background Values of Chinese Soils*. Beijing:Environmental Science Press of China.
- Dong J, Gao P, Sun H, Zhou C, Zhang X, Xue P, and Liu W. 2022. Characteristics and Health Risk Assessment of Cadmium, Lead, and Arsenic Accumulation in Leafy Vegetables Planted in a Greenhouse. *Environmental Science* 43:481-489. DOI 10.13227/j.hjx.202106002

466 Du P, Xie Y, Wang S, Zhao H, Zhang Z, Wu B, and Li F. 2015. Potential sources of and
467 ecological risks from heavy metals in agricultural soils, Daye City, China. *Environmental*
468 *Science and Pollution Research* 22:3498-3507. DOI 10.1007/s11356-014-3532-1

469 Guo J, Zhao H, Bian X, and Sun X. 2021. Characteristics and ecological risk of soil heavy metals
470 of a Tungsten mine in Yudu, Jiangxi Province. *Geological Bulletin of China* 40:1195-1202.

471 Guo L, Zhao W, Gu X, Zhao X, Chen J, and Cheng S. 2017. Risk Assessment and Source
472 Identification of 17 Metals and Metalloids on Soils from the Half-Century Old Tungsten Mining
473 Areas in Lianhuashan, Southern China. *International journal of environmental research and*
474 *public health* 14:1475. DOI 10.3390/ijerph14121475

475 Hakanson L. 1980. An ecological risk index for aquatic pollution control. a sedimentological
476 approach. *Water Research* 14:975-1001. DOI 10.1016/0043-1354(80)90143-8

477 Han Y, Zhu T, and Wang T. 2019. Joint Characteristics of Piaotang Tungsten Deposits in
478 Southern Jiangxi and Its Influence on Metallogenesis. *Journal of Hebei North University(natural*
479 *science edition)* 35:39-46.

480 Han Z, Golev A, and Edraki M. 2021. A Review of Tungsten Resources and Potential Extraction
481 from Mine Waste. *Minerals* DOI 11. 10.3390/min11070701

482 Hu Z. 2012. Strategic Strategies and Suggestions for Sustainable Development of Tungsten
483 Mines in Jiangxi Province. *China Nonferrous Metals* 3:38-39

484 Hu Z, Wang C, Li K, and Zhu X. 2018. Distribution characteristics and pollution assessment of
485 soil heavy metals over a typical nonferrous metal mine area in Chifeng, Inner Mongolia, China.
486 *Environmental Earth Sciences* 77:638. DOI 10.1007/s12665-018-7771-1

487 Hui W, Hao Z, Hongyan T, Jiawei W, and Anna L. 2021. Heavy metal pollution characteristics
488 and health risk evaluation of soil around a tungsten-molybdenum mine in Luoyang, China.
489 *Environmental Earth Sciences* 80:293. DOI 10.1007/s12665-021-09539-0

490 Inengite A, Abasi C, and Walter C. 2015. Application of Pollution Indices for the Assessment of
491 Heavy Metal Pollution in Flood Impacted Soil. *International Research Journal of Pure and*
492 *Applied Chemistry* 8:175-189. DOI 10.9734/IRJPAC/2015/17859

493 Jiang Y, Chao S, Liu J, Yang Y, Chen Y, Zhang A, and Cao H. 2017. Source apportionment and
494 health risk assessment of heavy metals in soil for a township in Jiangsu Province, China.
495 *Chemosphere* 168:1658-1668. DOI 10.1016/j.chemosphere.2016.11.088

496 Lai J, Liu J, Wu D, and Xu J. 2023. Pollution and health risk assessment of rare earth elements in
497 Citrus sinensis growing soil in mining area of southern China. *PeerJ* 11:e15470. DOI
498 10.7717/peerj.15470

499 Lambers H, and Barrow NJ. 2020. Pervasive use of P₂O₅, K₂O, CaO, MgO, and basic cations,
500 none of which exist in soil. *Biology and Fertility of Soils* 56:743-745. DOI 10.1007/s00374-020-
501 01486-5

502 Li M, Li D, Liu X, and Lin L. 2006. Determination of Metals in Stream Sediment and Soil by
503 ICP-MS. *Journal of Chinese Mass Spectrometry Society*:99-103+109.

Li M, Xi X, Xiao G, Cheng H, Yang Z, Zhou G, Ye J, and Li Z. 2014. National multi-purpose regional geochemical survey in China. *Journal of Geochemical Exploration* 139:21-30. DOI 10.1016/j.gexplo.2013.06.002

Liang Z, Gao L, Zhao X, Chen J, Xie Z, Li S, Li R, and Yang Z. 2019. Assessment of Metal Pollution, Its Potential Health Risks, and Origin in Different Land Use Types in Zhuhai City, China. *Archives of Environmental Contamination and Toxicology* 76:295-307. DOI 10.1007/s00244-018-00590-w

Liao Z; Zou Z. 2020. Southern Jiangxi geological survey team of Jiangxi Bureau of Geology and mineral resources exploration and development. *World Nonferrous Metals* 19:170-171

Liu D, Zhao Y, Zhou D, and Zhang J. 2017. Ecological risk assessment of heavy metals pollution in a tungsten mine soil in south of Jiangxi Province. *Environmental Chemistry* 36:1556-1567.

Liu H, Probst A, and Liao B. 2005. Metal contamination of soils and crops affected by the Chenzhou lead/zinc mine spill (Hunan, China). *Science of The Total Environment* 339:153-166. <https://doi.org/10.1016/j.scitotenv.2004.07.030>

Liu M, and Han G. 2021. Distribution of soil nutrients and erodibility factor under different soil types in an erosion region of Southeast China. *PeerJ* 9:e11630. DOI 10.7717/peerj.11630

Liu X, Zhang J, Huang F, Cheng J, Lu K, Yang J, Wang M, Wang Y, Qiu J, and Zhang X. 2022. Tungsten deposits in southern Jiangxi Province: Constraints on the origin of wolframite from in-situ U-Pb isotope dating. *Ore Geology Reviews* 143:104774. <https://doi.org/10.1016/j.oregeorev.2022.104774>

Loska K, Cebula J, Pelczar J, Wiechuła D, and Kwapuliński J. 1997. Use of Enrichment, and Contamination Factors Together with Geoaccumulation Indexes to Evaluate the Content of Cd, Cu, and Ni in the Rybnik Water Reservoir in Poland. *Water, Air, and Soil Pollution* 93:347-365. DOI 10.1023/A:1022121615949

Lu W, Zheng J, and Lai B. 1997. Influence of Cadmium Pollution from The Dayu Tungsten Mine on Human being. *Geological Exploration for Non-Ferrous Metals*:53-57.

Lv L, Jiao Z, Ge S, Zhan W, Ruan X, and Wang Y. 2022. Assessment of Cd Pollution in Paddy Soil-Rice System in Silver Mining-Affected Areas: Pollution Status, Transformation and Health Risk Assessment. *Int J Environ Res Public Health* 19. DOI 10.3390/ijerph191912362

MEP. 2013. Exposure Factors Handbook of Chinese Population(Adults). Ministry of Environmental Protection of China. *China Environmental Press*

MEEC. 2018. Soil environment quality risk control standard for soil contamination of of agriculture land. *Ministry of Ecology and Environment of China* GB 15618-2018.

Meng X, Sun W, Gao Z, Han H, Zhang C, Liu R, Chen P, Liu R, Yue T, Wang Y, Kang J, Lin S, Jing G, Wu M, and Yang L. 2021. Treatment and Recycling of Tungsten Dressing and Smelting Wastewater. *Metal Mine*:86-97. DOI 10.19614/j.cnki.jsks.202106009

Mokhtari AR, Feiznia S, Jafari M, Tavili A, Ghaneei-Bafghi M-J, Rahmany F, and Kerry R. 2018. Investigating the Role of Wind in the Dispersion of Heavy Metals Around Mines in Arid

Regions (a Case Study from Kushk Pb–Zn Mine, Bafgh, Iran). *Bulletin of Environmental Contamination and Toxicology* 101:124-130. DOI 10.1007/s00128-018-2319-3

Mugova E, and Wolkersdorfer C. 2022. Identifying potential groundwater contamination by mining influenced water (MIW) using flow measurements in a sub-catchment of the “Cradle of Humankind” Unesco World Heritage Site, South Africa. *Environmental Earth Sciences* 81:104. DOI 10.1007/s12665-022-10224-z

Muller G. 1969. Index of geoaccumulation in sediments of the Rhine River. *Geology Journal*:108–118.

Nacke H, Gonçalves AC, Schwantes D, Nava IA, Strey L, and Coelho GF. 2013. Availability of Heavy Metals (Cd, Pb, and Cr) in Agriculture from Commercial Fertilizers. *Archives of Environmental Contamination and Toxicology* 64:537-544. DOI 10.1007/s00244-012-9867-z

Pan L, Ma J, Hu Y, Su B, Fang G, Wang Y, Wang Z, Wang L, and Xiang B. 2016. Assessments of levels, potential ecological risk, and human health risk of heavy metals in the soils from a typical county in Shanxi Province, China. *Environmental Science and Pollution Research* 23:19330-19340. DOI 10.1007/s11356-016-7044-z

Rastegari Mehr M, Keshavarzi B, Moore F, Sharifi R, Lahijanzadeh A, and Kermani M. 2017. Distribution, source identification and health risk assessment of soil heavy metals in urban areas of Isfahan province, Iran. *Journal of African Earth Sciences* 132:16-26. DOI 10.1016/j.jafrearsci.2017.04.026

Reutova NV, Reutova TV, Dreeva FR, and Shevchenko AA. 2022. Long-term impact of the Tyrnyauz tungsten–molybdenum mining and processing factory waste on environmental pollution and children's population. *Environmental Geochemistry and Health* 44:4557-4568. DOI 10.1007/s10653-022-01221-z

Sawe SF, Shilla DA, and Machiwa JF. 2019. Assessment of enrichment, geo-accumulation and ecological risk of heavy metals in surface sediments of the Msimbazi mangrove ecosystem, coast of Dar es Salaam, Tanzania. *Chemistry and Ecology* 35:835-845. DOI 10.1080/02757540.2019.1663344

Shen F, Liao R, Ali A, Mahar A, Guo D, Li R, Xining S, Awasthi MK, Wang Q, and Zhang Z. 2017. Spatial distribution and risk assessment of heavy metals in soil near a Pb/Zn smelter in Feng County, China. *Ecotoxicology and Environmental Safety* 139:254-262. DOI 10.1016/j.ecoenv.2017.01.044

Shrivastav P, Prasad M, Singh TB, Yadav A, Goyal D, Ali A, and Dantu PK. 2020. Role of Nutrients in Plant Growth and Development. In: Naeem M, Ansari AA, and Gill SS, eds. *Contaminants in Agriculture Sources, Impacts and Management*. Gewerbestrasse 11, 6330 Cham, Switzerland: Springer, 43-54. DOI 10.1007/978-3-030-41552-5

Shi J, Du P, Luo H, Wu H, Zhang Y, Chen J, Wu M, Xu G, and Gao H. 2022. Soil contamination with cadmium and potential risk around various mines in China during 2000–2020. *Journal of Environmental Management* 310:114509. DOI 10.1016/j.jenvman.2022.114509

Timofeev I, Kosheleva N, and Kasimov N. 2018. Contamination of soils by potentially toxic elements in the impact zone of tungsten-molybdenum ore mine in the Baikal region: A survey

and risk assessment. *Science of The Total Environment* 642:63-76. DOI 10.1016/j.scitotenv.2018.06.042

Timofeev I, Shartova N, Kosheleva N, and Kasimov N. 2020. Potentially toxic elements in urban topsoils and health risk assessment for the mining W–Mo center in the Baikal region. *Environmental Geochemistry and Health* 42:221-240. DOI 10.1007/s10653-019-00340-4

Tomlinson DL, Wilson JG, Harris CR, and Jeffrey DW. 1980. Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer Meeresuntersuchungen* 33:566-575. DOI 10.1007/BF02414780

Tunc T, and Sahin U. 2017. Yield and Heavy Metal Content of Wastewater-Irrigated Cauliflower and Soil Chemical Properties. *Communications in Soil Science and Plant Analysis* 48:1194-1211. DOI 10.1080/00103624.2017.1341910

USEPA. 2011. *Exposure Factors Hand-book 2011 Edition (final)*.

Vega FA, Covelo EF, Andrade ML, and Marcet P. 2004. Relationships between heavy metals content and soil properties in minesoils. *Analytica Chimica Acta* 524:141-150. DOI 10.1016/j.aca.2004.06.073

VROM. 2000. *Circular on target values and intervention values for soil remediation*. Dutch Ministry of Housing Spatial Planning and Environment.

Wang F, Huang Y, Wang X, Gao Z, Yu F, Xu F, Bao Q, Hu Y, Qiao M, Jin S, Li J, and Xiang M. 2015. Ecological risk assessment of heavy metals in surrounding soils of tungsten ores: Comparison of different evaluation methods. *Environmental Chemistry* 34:225-233.

Wang L, Zhao S, Li S, Qin L, Sun X, Lei X, Chen S, and Wang M. 2021. Effect of sodium salt with varied anions on Cd fractions and microbial community in soil. *China Environmental Science* 41:4221-4230. DOI 10.19674/j.cnki.issn1000-6923.20210517.008

Wang S, Wu W, Liu F, Zhao M, Qiu J, and Wu J. 2018. Assessment of human health risks of heavy metals in the typical sewage irrigation areas. *China Environmental Science* 38:1550-1560. DOI 10.19674/j.cnki.issn1000-6923.2018.0188

Wang Y, Xiao J, Hong Y, Zhou Y, Han Z, Xie M, Wang D, and Guo J. 2009. Preliminary Study on the Geological Characteristics and Metallogenic Model of the Xihuashan Tungsten Deposit in Dayu, Jiangxi Province. *Acta Mineralogica Sinica* 29:30-31. DOI 10.16461/j.cnki.1000-4734.2009.s1.186

Wang Y, Zou Y, Wang H, and Wang C. 2019. Geochemical characteristics of Se and heavy metal elements in the soil of Youshan, Xinfeng County, Jiangxi Province. *East China Geology* 40:152-160. DOI 10.16788/j.hddz.32-1865/p.2019.02.009

Wu G, Wang C, and Chen H. 2020. Eco-environmental assessment and genetic analysis of heavy metal pollution in the soil around the abandoned tungsten-molybdenum mine area in Inner Mongolia. *Geology in China* 47:1838-1852.

Wu T. 1993. A Brief History of Tungsten, Antimony and Tin Industry Development in Modern China. *Development Research*:56-60. DOI 10.13483/j.cnki.kfyj.1993.03.016

Xiao A, Ouyang Y, Li WC, and Ye Z. 2017. Effect of organic manure on Cd and As accumulation in brown rice and grain yield in Cd-As-contaminated paddy fields. *Environmental Science and Pollution Research* 24:9111-9121. DOI 10.1007/s11356-017-8460-4

Xu J, Wang X, Wang J, Xu L, Zheng X, Zhang Y, and Hu C. 2021. Dominant environmental factors influencing soil metal concentrations of Poyang Lake wetland, China: Soil property, topography, plant species and wetland type. *Catena* 207:105601. DOI 10.1016/j.catena.2021.105601

Xu J, Zheng L, Xu L, and Wang X. 2020. Uptake and allocation of selected metals by dominant vegetation in Poyang Lake wetland: From rhizosphere to plant tissues. *Catena* 189:104477. DOI 10.1016/j.catena.2020.104477

Yang J, Dai Y, Liu Y, Duan S, Li Y, Hu R, Zhou Z, Shi Y, Liu H, and Wang S. 2020. Reduced cadmium accumulation in tobacco by sodium chloride priming. *Environmental Science and Pollution Research* 27:37410-37418. DOI 10.1007/s11356-020-09134-z

Zahra A, Hashmi MZ, Malik RN, and Ahmed Z. 2014. Enrichment and geo-accumulation of heavy metals and risk assessment of sediments of the Kurang Nallah—Feeding tributary of the Rawal Lake Reservoir, Pakistan. *Science of The Total Environment* 470-471:925-933. DOI 10.1016/j.scitotenv.2013.10.017

Zhang B. 2020. Distribution characteristics and evaluation of heavy metals in soils of Fangshan District, Beijing Master of Engineering Master. Thesis, China University of Geosciences (Beijing).

Zhang C, Liu J, He J, and Mao S. 2012. Study on the Beneficiation Technology of a Tungsten-Molybdenum Ore of Jiangxi. *Nonferrous Metals(Mineral Processing Section)*:36-38.

Zhang H, Huang B, Dong L, Hu W, Akhtar MS, and Qu M. 2017. Accumulation, sources and health risks of trace metals in elevated geochemical background soils used for greenhouse vegetable production in southwestern China. *Ecotoxicology and Environmental Safety* 137:233-239. DOI 10.1016/j.ecoenv.2016.12.010

Zhang J, Liu X, Li W, Zeng Z, Hu H, Peng L, Cheng J, Lu K, and Li P. 2021. The metallogenic epoch and geological implications of the tungsten-tin polymetallic deposits in southern Jiangxi Province, China: Constraints from cassiterite U-Pb and molybdenite Re-Os isotopic dating. *Ore Geology Reviews* 134:104159. DOI 10.1016/j.oregeorev.2021.104159

Zheng K, Ma Z, Sun B, and Liang Y. 2021. Effects of Fertilization Strategies on the Cadmium Resistance of Paddy Soil Microorganisms. *Environmental Science* 42:394-402. DOI 10.13227/j.hjxx.202006154

Zheng X-J, Chen M, Wang J-F, Li F-G, Liu Y, and Liu Y-C. 2020. Ecological Risk Assessment of Heavy Metals in the Vicinity of Tungsten Mining Areas, Southern Jiangxi Province. *Soil & Sediment Contamination* 29:665-679. DOI 10.1080/15320383.2020.1763912

Zhou M; Tang Z; Zhang M; Liang X; Zhan L. 2021. Characteristics and health risk assessment of heavy metals in soil-rice system in the Ganzhou area, Jiangxi Province. *Geological Bulletin of China* 12:2149-2158

Figure 1

The study area and sampling sites of soils and paddy rice(site1:FJ-N,site2:FJ-S,site3:HL,site4:QL,site5:CJ-W,site6:CJ-E;Satellite image credit: <http://www.google.com/permissions/geoguidelines/attr-guide.html>)

The little red points are samples of soil. The black triangles are sampling sits in study regin. The red five-pointed stars are tungsten mining aeras. The river is represented by blue curve. Other curves represent administrative areas on the map.




Figure 2

Concentration of heavy metal in soil(mg/kg)

The X-axis represents the sampling sites. The Y-axis represents the concentration of heavy metals. Box charts show the distribution of data. Different letters of "a", "b", "c" indicate the significance of concentration of heavy metals' differences among sites.




Figure 3

Geo-accumulation index (I_{geo}) of heavy metal in soil in sampling sites (site1:FJ-N, site2:FJ-S, site3:HL, site4:QL, site5:CJ-W, site6:CJ-E)

The X-axis represents the sampling sites. The Y-axis represents the I_{geo} values of heavy metals. Box charts show the distribution of data.




Figure 4

ADD of heavy metal in paddy rice for non-carcinogenic in sampling areas (site1:FJ-N, site2:FJ-S, site3:HL, site4:QL, site5:CJ-W, site6:CJ-E)

The X-axis represents the sampling sites. The Y-axis represents the ADD values of heavy metals. Box charts show the distribution of data.




Table 1(on next page)

The categories of E_r^i and RI values

E_r^i and RI represents potential ecological risks through different levels.

Table 1 The categories of E_r^i and RI values2

Table 2 (on next page)

Concentration of components of Soil properties

Compared to concentration of components of Soil properties in sampling sites.

Notes: Different letters of “a”, “b”, “c” indicate the significance of soil factors’ differences among sites.

Table 3(on next page)

The Pearson’s correlation between soil metals and soil properties

Each value indicates the Pearson’s correlation between heavy metal concentration and soil properties.

Table 3 The Pearson's correlation between soil metals and soil properties

	As	Cd	Cr	Cu	Mo	Pb	W	Zn
pH	0.100	0.110	-0.098	0.198	0.289*	0.163	0.130	0.142
TOC	-0.068	0.202	0.556**	0.063	-0.165	0.042	-0.066	0.351**
K ₂ O	0.386**	0.504**	-0.326**	0.520**	0.583**	0.681**	0.490**	0.483**
Na ₂ O	0.372**	0.467**	-0.536**	0.498**	0.718**	0.589**	0.492**	0.410**
CaO	-0.001	0.199	-0.112	0.318**	0.292*	0.289*	0.120	0.324**
Fe ₂ O ₃	0.046	-0.078	0.281*	0.091	0.053	0.021	0.128	0.087
CEC	-0.043	0.015	0.511**	0.087	-0.136	-0.052	-0.142	0.149

Notes: ** Correlation is significant at the 0.01 level(2-tailed). * Correlation is significant at the 0.05 level(2-tailed).

Table 4(on next page)

Heavy metal contamination factors (CFs) and pollution load indexes (PLIs) for metal in soil

The data indicates the degree of contamination and pollution level of metals.

Table 5(on next page)

Ecological risk of heavy metal in soil

The data indicates the ecological risk level of metal in different site.

1

Table 5 Ecological risk of heavy metal in soil

Site	E_r^i							RI
	As	Cd	Cu	Cr	Pb	W	Zn	
FJ-N	36.09	113	8.30	3.15	6.55	35.04	1.31	204
FJ-S	41.32	320	12.72	2.81	12.98	63.98	2.11	456
HL	27.28	161	8.86	2.51	8.13	34.99	1.26	244
QL	39.25	220	11.72	2.60	9.87	36.72	1.71	322
CJ-W	17.80	104	6.88	2.38	7.20	16.99	1.05	157
CJ-E	16.93	81.30	6.50	2.30	5.42	10.91	0.99	124

2

Table 6(on next page)

Concentration of heavy metals in paddy rice

The data indicates concentration of paddy rice at different sampling sites. Notes:Different letters of“a”,“b”,“c”indicates the significant difference of heavy metal concentrations among sites at $p < 0.05$ or $p < 0.01$.

Table 6 Concentration of heavy metals in paddy rice

sites	As ($p<0.01$)	Cd ($p>0.05$)	Cr ($p<0.05$)	Cu ($p>0.05$)	Mo ($p<0.05$)	Pb ($p<0.05$)	Zn ($p<0.05$)
FJ-N	0.18 ^c ±0.0 7	0.36 ^{ab} ±0.01	0.08 ^a ±0.001	4.31 ^a ±0.8 8	0.005 ^a ±0.002	0.05 ^{ab} ±0.03	20.61 ^{bc} ±3.46
FJ-S	0.15 ^{bc} ±0.01	1.18 ^b ±0.18	0.08 ^a ±0.001	5.58 ^b ±0.36	0.009 ^b ±0.005	0.08 ^b ±0.02	21.37 ^c ±1.36
HL	0.09 ^a ±0.02	0.53 ^{ab} ±0.075	0.08 ^a ±0.001	4.57 ^{ab} ±0.68	0.005 ^a ±0.002	0.04 ^a ±0.01	15.55 ^a ±1.53
QL	0.11 ^{ab} ±0.03	0.83 ^{ab} ±0.064	0.08 ^a ±0.001	4.69 ^{ab} ±0.84	0.004 ^a ±0.003	0.04 ^a ±0.01	17.46 ^{ab} ±2.29
CJ-W	0.07 ^a ±0.03	0.11 ^a ±0.09	0.10 ^{ab} ±0.01	3.86 ^a ±0.68	0.001 ^a ±0.004	0.03 ^a ±0.01	15.48 ^a ±1.79
CJ-E	0.11 ^{ab} ±0.02	0.47 ^b ±0.44	0.11 ^b ±0.02	4.17 ^a ±0.98	0.005 ^a ±0.002	0.06 ^{ab} ±0.03	18.19 ^{abc} ±3.45

Notes: Different letters of 'a', 'b', 'c' indicates the significant difference of heavy metal concentrations among sites at $p < 0.05$ or $p < 0.01$.

Figure 5

ADD of heavy metal in paddy rice for carcinogenic in sampling areas (site1:FJ-N, site2:FJ-S, site3:HL, site4:QL, site5:CJ-W, site6:CJ-E)

The X-axis represents the sampling sites. The Y-axis represents the ADD values of heavy metals. Box charts show the distribution of data.

