

Exogenous Hemin enhances the antioxidant defense system of rice by regulating the AsA-GSH cycle under NaCl stress

Fengyan Meng^{1, 2}, Naijie Feng^{Corresp., 1, 2, 3}, Dianfeng Zheng^{Corresp., 1, 2, 3}, Meiling Liu^{1, 2}, Hang Zhou^{2, 3}, Rongjun Zhang^{1, 2}, XiXin Huang^{1, 2}, Anqi Huang^{1, 2}

¹ College of Coastal Agriculture Sciences, Guangdong Ocean University, Zhanjiang, Guangdong, China

² South China Center of National Saline-tolerant Rice Technology Innovation Center, Zhanjiang, Guangdong, China

³ Shenzhen Research Institute of Guangdong Ocean University, Shenzhen, Guangdong, China

Corresponding Authors: Naijie Feng, Dianfeng Zheng

Email address: fengnj@gdou.edu.cn, zhengdf@gdou.edu.cn

Abiotic stress caused by soil salinization remains a major global challenge that threatens and severely impacts worldwide crop growth causing yield reduction. Rice is an important economic crop affected by salt stress. In this study, we aimed to investigate the damage of salt stress on the leaf physiology of two rice varieties (HuanghuaZhan, HHZ and XiangliangYou 900, XLY900) and the regulatory mechanism of Hemin to maintain seedling growth under the stressful environment. Therefore, at the three leaves and one heart, leaves were foliar sprayed with 5 $\mu\text{mol}\cdot\text{L}$ Hemin or 25 $\mu\text{mol}\cdot\text{L}$ ZnPP (Zinc protoporphyrin IX) followed by 50 $\text{mmol}\cdot\text{L}$ NaCl stress. The findings revealed that salt stress increased antioxidant enzyme activity and decreased the contents of nonenzymatic antioxidants such as ascorbate (AsA) and glutathione (GSH). Furthermore, the content of osmoregulatory substances like soluble proteins and proline was raised. Moreover, the increase in reactive oxygen species (ROS) content indicated that the oxidative stress deepened. However, spraying Hemin increased the activities of antioxidants such as superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) and accelerated AsA-GSH cycling to remove excess ROS. In summary, Hemin improved salt stress's effects on rice leaves' physiological characteristics due to improved antioxidant defense mechanisms that impeded lipid peroxidation. Thus, Hemin could somewhat decelerate the damages caused by salt stress.

Exogenous Hemin enhances the antioxidant defense system of rice by regulating the AsA-GSH cycle under NaCl stress

Fengyan Meng^{1,2}, Naijie Feng^{1,2,3}, Dianfeng Zheng^{1,2,3}, Meiling Liu^{1,2}, Hang Zhou^{2,3}, Rongjun Zhang^{1,2}, Xixin Huang^{1,2}, Anqi Huang^{1,2}

¹ College of Coastal Agriculture Sciences, Guangdong Ocean University, Zhanjiang, Guangdong 524088, China

² South China Center of National Saline-tolerant Rice Technology Innovation Center, Zhanjiang, Guangdong 524088, China

³ Shenzhen Research Institute of Guangdong Ocean University, Shenzhen, Guangdong 518108, China

Corresponding Author:

Naijie Feng^{1,2,3} and Dianfeng Zheng^{1,2,3}

College of Coastal Agriculture Sciences, Guangdong Ocean University, Zhanjiang, Guangdong, 524088, China

Email address: fengnj@gdou.edu.cn and zhengdf@gdou.edu.cn

Abstract

Abiotic stress caused by soil salinization remains a major global challenge that threatens and severely impacts worldwide crop growth causing yield reduction. Rice is an important economic crop affected by salt stress. In this study, we aimed to investigate the damage of salt stress on the leaf physiology of two rice varieties (HuanghuaZhan, HHZ and XiangliangYou 900, XLY900) and the regulatory mechanism of Hemin to maintain seedling growth under the stressful environment. Therefore, at the three leaves and one heart, leaves were foliar sprayed with 5 $\mu\text{mol}\cdot\text{L}^{-1}$ Hemin or 25 $\mu\text{mol}\cdot\text{L}^{-1}$ ZnPP (Zinc protoporphyrin IX) followed by 50 $\text{mmol}\cdot\text{L}^{-1}$ NaCl stress. The findings revealed that salt stress increased antioxidant enzyme activity and decreased the contents of nonenzymatic antioxidants such as ascorbate (AsA) and glutathione (GSH). Furthermore, the content of osmoregulatory substances like soluble proteins and proline was raised. Moreover, the increase in reactive oxygen species (ROS) content indicated that the oxidative stress deepened. However, spraying Hemin increased the activities of antioxidants such as superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) and accelerated AsA-GSH cycling to remove excess ROS. In summary, Hemin improved salt stress's effects on rice leaves' physiological characteristics due to improved antioxidant defense mechanisms that impeded lipid peroxidation. Thus, Hemin could somewhat decelerate the damages caused by salt stress.

Keywords: AsA-GSH cycle, Enzymatic defense system, Hemin, Rice

Introduction

In the background of global warming, soil salinization has accelerated due to various factors such as seawater back-up, over-exploitation of groundwater and unreasonable development of arable land (Alkharabsheh et al., 2021). Saline land accounts for about one-fifth of the cultivated land and one-third of the irrigated farmland on the planet, and the area is increasing at an even faster rate (Mukhopadhyay et al., 2021). Salinity stress is one of the most widespread and severe abiotic stresses in the world. It has destructive effects on plant growth and physiological and biochemical processes and causes a decrease in grain production. According to incomplete statistics, the yield loss caused by salt stress accounts for about 20% of global yield (Ding et al., 2021).

With salt stress increasing soil osmotic pressure, plant roots fail to absorb water and nutrients, which causes delayed growth and development or even death (Liu et al., 2022). In addition, salt stress induces excessive production of ROS in plant cells. ROS is weakly stable and easily causes oxidative stress to cells. The excessive ROS enhances cell membrane lipid peroxidation and disrupts membrane system stability, which results in the expansion of cell membrane permeability and extravasation of intracellular materials (Seleiman et al., 2020; Hasanuzzaman et al., 2020). The researches showed that ROS responded with proteins, DNA, and lipids, which caused protein breakdown, DNA structural damage, and lipid peroxidation (Chandrakar et al., 2017; Lin et al., 2020). This disturbed the normal growth and physiological metabolic activities of plants. To avoid ROS accumulation, plants use antioxidant enzymes and non-enzymatic antioxidants to scavenge excess ROS (Alisofi et al., 2020). Among them, antioxidant enzymes include SOD, POD, CAT, and ascorbate peroxidase (APX). Non-enzymatic antioxidants include ascorbic acid and glutathione, which act as co-factors for different enzymes and participate in various metabolic processes (Hasanuzzaman et al., 2020). In addition, plants accumulate osmoregulatory substances to maintain the balance of inside and outside cell osmosis. There are two categories of osmoregulatory substances: inorganic ions (Na^+); and organic substances, including proline and soluble proteins (Athar et al., 2022). Under salt stress, plants balance the osmotic pressure between the plant and the external environment by conducting selective uptake of ions and promoting the accumulation of phase-soluble solutes.

Rice, a gramineous crop, has a long history of cultivation and consumption in China. Our consumption demand for rice is the first in the world, and more than half of the population eats rice as a major food (Huang et al., 2022; Zuo et al., 2022). However, salt stress has become one of the major abiotic stresses which limits rice production. Many studies showed that the seedling stage was an essential stage of individual development and was closely related to the later development of tillers and spikes. However, this stage is susceptible to the impact of salt stress (Zeng et al., 2001). Therefore, how to improve the salt tolerance of rice seedlings has become an urgent problem at present.

Plant growth regulators are a group of synthetic compounds with phytohormonal activity that improve the tolerance to abiotic stresses by affecting the expression of endogenous hormones in crops. Hemin is a small molecule with a porphyrin structure, consisting of nitrogen atoms on four pyrrole rings in a porphyrin ligated to a ferrous ion. In recent years, more and more researchers

have used Hemin in different crops for its natural, non-polluting, low cost, and high safety features. Hemin acts as a substrate and promoter of heme oxygenase 1 (HO-1), an initiator and rate-limiting enzyme for Hemin degradation, and has a specific inhibitor, zinc protoporphyrin (ZnPP). Hemin triggered salt acclimation in wheat by increasing HO-1 expression, while ZnPP, an inhibitor, interrupted the salinity tolerance (Xie et al., 2011). Under salt stress, Hemin increased proline and soluble protein content, enhanced antioxidant enzyme activities such as SOD, CAT, and APX, and alleviated oxidative damage in *Cassia obtusifolia* L (Zhang et al., 2012). In addition, under Zn, Pb, and Cr metal stress, Hemin activated the activities of various antioxidant enzymes (SOD, GR, and APX) in rice seedlings, improved the content of AsA and GSH, and reduced heavy metal accumulation.

At present, there are fewer studies on the mitigation of salt stress by Hemin on rice seedlings, and based on the mitigation effect of Hemin on stresses such as heavy metal stress of alfalfa (Fu et al., 2011), low-temperature stress of *Conyza blini* (Zheng et al., 2021), and salt stress in *Brassica juncea* L (Verma et al., 2015). Furthermore, spraying plant growth regulators can improve the resistance of rice seedlings during the critical period before transplanting, which is essential for the subsequent transplanting of rice seedlings on saline land. Hence, in this study, we used two rice varieties, Huang Huazhan and Xiang Liangyou 900, to research the impacts of foliar spraying with hemin at the three-leaf-one-heart stage on the growth and ROS metabolism (antioxidant enzymes and non-enzymatic antioxidants) of rice seedlings under salt stress, which aimed to reveal the mechanism of hemin in enhancing the salt tolerance of rice, and to provide theoretical basis and technical guidance for the cultivation of saline rice.

Materials and methods

Plant materials

The experiment was carried out in 2022 in Binhai Agricultural College of Guangdong Ocean University. The conventional rice Huanghuazhan (HHZ) and hybrid rice Xiangliangyou900 (XLY900), were used in this study to make the results more general. Hemin was provided from Shanghai Changdeduo Agricultural Technology Co., Ltd.

Experiment design

Seeds were selected for uniformity of size and color, sterilized with 3% H₂O₂ for about 15 min, and then rinsed 3-5 times with distilled water. These seeds were soaked and germinated for each 24 hours under dark conditions at 30°C. Sixty-five seeds were sown into pots containing 3 kg of test soil with 1:3 sand to latosol each. The plastic pot sizes were 19 cm for the upper diameter, 14 cm for the lower diameter, and 17 cm for the height, without holes in the bottom. Regular water irrigation was performed until the three leaves and one heart stage (about 18 days after planting). Rice leaves were foliar sprayed with 5 μmol·L⁻¹ Hemin and 25 μmol·L⁻¹ ZnPP alone or in combination, and plants were exposed to 25 mmol·L⁻¹ NaCl stress twice every 24 h after spraying, which meant that the salt concentration in the soil reached 50 mmol·L⁻¹ at 48 h after spraying. In subsequent experiment, concentrations were maintained by measuring soil conductivity (EC=5.0±0.5 dS·m⁻¹). There were five treatments for each variety, which were (1)CK (control): normal water; (2)S: 50 mmol·L⁻¹ NaCl treatment, (3)SH: 50 mmol·L⁻¹ NaCl+5 μmol·L⁻¹ Hemin,

(4)SZ: 50 mmol·L⁻¹ NaCl+25 μmol·L⁻¹ ZnPP, (5)SZH: 50 mmol·L⁻¹ NaCl+5 μmol·L⁻¹ Hemin+25 μmol·L⁻¹ ZnPP, 25 pots per treatment. The plant samples were harvested at 3, 5, 7, and 9 d after 50 mmol·L⁻¹ NaCl stress for morphological parameters and physiological were collected and frozen at -80 °C for biochemical analysis.

Morphological measurements

The plant height was measured with a ruler, stem diameter was measured with vernier, shoot fresh and dry weight were measured by a caliper electronic analytical balance. The shoots were dried for 30 min at 105 °C and 72 h at 85 °C.

Measurement of EL, MDA and H₂O₂ content

Electrolyte leakage (EL) was determined as described by Yu et al. (2021). The measurement of MDA content was carried out according to the method proposed by Ahmad et al. (2016). The frozen leaf sample (0.5g) was extracted in 10 ml phosphate buffer (0.05 mmol·L⁻¹ PBS, pH 7.8) and centrifuged at 6000 rpm for 20 min. 1 mL of the supernatant was added to 2 mL of 0.6% TBA, then boiled at 100 °C for 15 min. The mixture was cooled quickly with cold water and centrifuged at 4,000 rpm for 20 min. The absorption value was determined at 450 nm, 532 nm and 600 nm. The H₂O₂ content was determined by Rasheed et al. (2022). 0.5g of the frozen sample was ground into homogenate in 5 ml of TCA and centrifuged at 19,000 rpm for 20 min. 0.5 ml of supernatant was added to 0.5 ml PBS (10 mmol·L⁻¹ Ph 7.0) and 1 ml KI (1 mol·L⁻¹), then the reaction mixture was incubated at 28 °C for 1 h in the dark. The absorbance values were recorded at 410 nm.

Histochemical Detection of Hydrogen Peroxide and Superoxide Anion

The histochemical staining of hydrogen peroxide (H₂O₂) and superoxide radicle (O₂^{·-}) was measured by Zhang et al. (2009) and Sudhakar et al. (2015). On the 3rd day of stress, the second leave of CK, S, SH, SZ and SZH treatments of both varieties were taken for histochemical staining. The leaves were vacuumed and then kept at room temperature and dark conditions for 24 h until brown and blue spots appeared, respectively, then adding 95% ethanol and decolorized in a water bath at 80°C until the leaves were free of chlorophyll for observation and photography.

Measurement of the activities of SOD, POD, and CAT

The frozen leaf sample was extracted in 10 ml PBS (50 mmol·L⁻¹ pH 7.8) at 4°C, centrifuged at 12,000 rpm and 4 °C for 20 min. The supernatant was used as an enzyme solution to determine SOD (EC 1.15.1.1), POD (EC 1.11.1.7), and CAT (EC 1.11.1.6) (Habib et al., 2021). SOD activity was carried out according to the method by Lu et al. (2022). The supernatant was mixed with 14.5 mmol·L⁻¹ methionine solution, 3 mmol·L⁻¹ EDTA-Na₂ solution, 60 μmol·L⁻¹ riboflavin solution, and 2.25 mmol·L⁻¹ NBT solution. One unit of SOD activity was defined as the amount of enzyme that would inhibiting 50% of NBT photoreduction. POD was determined following the method given by Kenawy et al. (2022). The supernatant was mixed with PBS (pH 6.0), guaiacol, and 30% H₂O₂. The absorbance was measured at 470 nm. CAT was determined by the decreased absorbance rate of H₂O₂ at 240 nm, outlined by Basilio-Apolinar et al. (2021).

Measurement of AsA-GSH cycle products and substrate content

The procedure illustrated by Costa et al. (2002) and Yan et al. (2021) was followed to measure the contents of AsA and total AsA. The frozen leaf sample was extracted in 5% TCA and centrifuged at 12,000 rpm and 4°C for 15 min. The supernatant fluid was then moved to a calibration tube. The

supernatant was used to determine the content of AsA and total AsA. For AsA, the supernatant was mixed with a reaction solution containing 5% TCA, ethanol, H_3PO_4 -ethanol, BP-ethanol, and FeCl_3 -ethanol. The reaction was carried out at 30°C for 90 min. The absorbance was assayed at 534 nm. For total AsA, it was similar to the AsA assay. Still, it was first reacted with dithiothreitol (DTT)-ethanol solution and Na_2HPO_4 -NaOH solution for 10 min. Then, 20% TCA was added and mixed with the above reaction solution. The absorbance was assayed at 540 nm. Dehydroascorbate (DHA) content was calculated based on the difference between total AsA and reduced AsA. The GSH and GSSG content was determined according to the method described by Kaya et al. (2023). 0.5 g frozen sample was ground into homogenate in 5 mL of 5% metaphosphoric acid and centrifuged at 20000g for 20 min. The supernatant was used to determine the content of GSH+GSSG and GSSG. The supernatant was mixed homogeneously with the reaction solution, which contained $50 \text{ mmol}\cdot\text{L}^{-1}$ phosphate buffer, $10 \text{ mmol}\cdot\text{L}^{-1}$ NADPH, $12.5 \text{ mmol}\cdot\text{L}^{-1}$ DTNB, and 50 U GR, and the absorbance value of (GSH+GSSG) was measured at 412 nm. Besides adding the reaction solution, 2-vinylpyridine (2-VP) was added to eliminate GSH, and the GSSG absorbance value was measured at 412 nm. The GSH content was then calculated from the GSH+GSSG content reduced the GSSG content.

Measurement of the critical enzyme indexes of the AsA-GSH cycle

0.5 g of the frozen leaf sample was placed in a mortar, ground into a powder with $50 \text{ mmol}\cdot\text{L}^{-1}$ sodium phosphate buffer solution (pH7.8), and loaded into a centrifuge tube. The centrifuge tube was centrifuged at 12000g for 20 min. The resulting solution was used to measure the levels of APX (EC 1.11.1.11), monodehydroascorbate reductase (MDHAR, EC 1.6.5.4), dehydroascorbate reductase (DHAR, EC 1.8.5.1) and glutathione reductase (GR, EC 1.6.4.2).

The APX activity was determined according to the method described by Sharifi et al. (2021). The assay mixture contained 0.1 ml of enzyme extract, 2.6 ml EDTA- Na_2 ($0.1 \text{ mmol}\cdot\text{L}^{-1}$), 0.15 ml AsA ($5 \text{ mmol}\cdot\text{L}^{-1}$) and $20 \text{ mmol}\cdot\text{L}^{-1}$ H_2O_2 . The absorbance was assayed at 290 nm. ($E=2.8 \text{ mM}^{-1} \text{ cm}^{-1}$). MDHAR activity was measured using the method described by Hasanuzzaman et al. (2011). The reaction mixture consisted of $25 \text{ mmol}\cdot\text{L}^{-1}$ sodium phosphate buffer solution (pH7.8), $7.5 \text{ mmol}\cdot\text{L}^{-1}$ AsA, $2 \text{ mmol}\cdot\text{L}^{-1}$ NADPH, 50 U AsA oxidase (EC 1.10.3.3), and enzyme extract. The absorbance was assayed at 340 nm. ($E=6.2 \text{ mM}^{-1} \text{ cm}^{-1}$). DHAR activity was carried out by the method proposed by Shan and Liu (2017). DHAR was assayed in a mixed solution containing $25 \text{ mmol}\cdot\text{L}^{-1}$ sodium phosphate buffer solution (pH7.8), $20 \text{ mmol}\cdot\text{L}^{-1}$ GSH, $10 \text{ mmol}\cdot\text{L}^{-1}$ DHA, and enzyme extract. The absorbance was assayed at 340 nm. ($E=14 \text{ mM}^{-1} \text{ cm}^{-1}$). GR activity was done according to Keles and Oncel (2002). GR (EC 1.6.4.2) was assayed in a mixed solution containing $25 \text{ mmol}\cdot\text{L}^{-1}$ sodium phosphate buffer solution (pH 7.8), $2 \text{ mmol}\cdot\text{L}^{-1}$ EDTA, $10 \text{ mmol}\cdot\text{L}^{-1}$ GSSG, $24 \text{ mmol}\cdot\text{L}^{-1}$ NADPH, and enzyme extract. The absorbance was assayed at 340 nm. ($E=6.2 \text{ mM}^{-1} \text{ cm}^{-1}$).

Measurement of soluble protein and proline content

Soluble protein content was determined according to the method described by Tian et al. (2022). The absorbance value was measured at 595 nm using Coomassie brilliant blue. Proline content was carried out according to the method by Liu et al. (2020). The frozen sample (0.5g) was ground in 5 mL of 3% sulfosalicylic acid and then centrifuged at 3000g for 10 min. 2 ml of the supernatant

was added to 2 mL acetic acid and 2 mL acidic ninhydrin, boiling water bath at 100 °C for 30 min. After cooling, 4 mL of toluene was added and measured absorbance at 520nm.

Statistical analysis

The data was analyzed by Microsoft Excel 2019 and SPSS 25.0. The figures were drawn in Origin 2021. Duncan test ($p < 0.05$) was used to evaluate the difference within treatments, and the significant differences among different materials were determined.

Result

The morphological parameters of rice seedlings

There was significant inhibition of rice growth under NaCl stress, which showed a remarkable decrease in plant height, stem base width, shoot fresh weight, and shoot dry weight (Table 1 and Table 2). From days 3 through 9, in comparison to CK, the plant height, stem diameter, shoot fresh weight and shoot dry weight of HHZ under NaCl stress significantly decreased by 13.48%-16.58%, 23.08%-28.95%, 29.67%-32.41% and 21.14%-23.34% respectively; in XLY900, the above indicators decreased by 10.67%-13.98%, 17.43%-23.08%, 27.24%-30.71% and 18.22%-22.15%. Exogenous Hemin alleviated the inhibition of rice seedling growth by NaCl stress (Fig. 1). From days 3 through 9, in comparison to the S treatment, the plant height, stem diameter, shoot fresh weight and shoot dry weight of HHZ with SH treatment were significantly higher by 9.62%-12.38%, 20.00%-32.10%, 18.63%-27.43%, and 11.96%-15.84%, respectively; in XLY900, the above indicators were increased by 5.33%-8.01%, 15.56%-24.14%, 15.85%-26.58%, and 12.78%-14.26%, respectively. The aforementioned findings suggested that the hemin application can effectively mitigate the growth-inhibiting effects of NaCl stress on rice seedlings, and its regulatory influence on HHZ appeared to be more pronounced compared to XLY900. In contrast to the S treatment, the ZnPP did not lead to an increase in plant height, stem base width, shoot fresh weight, or shoot dry weight for both rice varieties. The addition of Hemin reversed the inhibition caused by ZnPP and enhanced the growth of rice seedlings. From days 3 through 9, the plant height of both HHZ and XLY900 was notably elevated in the SZH treatment, exhibiting a rise of 5.15% to 7.16% and 2.54% to 4.92%, respectively, as compared to the SZ treatment; the stem base width was enhanced by 7.59%-12.20% and 7.78%-15.91%, respectively; shoot fresh weight was increased by 9.12%-19.43% and 9.86%-16.19%, respectively and shoot dry weight was increased by 8.56%-10.66%, 6.70%-10.32%, respectively.

The membrane damage and ROS accumulation in rice seedlings

Compared to the CK, electrolyte leakage (EL), malondialdehyde (MDA), and H_2O_2 contents in two rice varieties gradually increased with the extension of NaCl stress time (Fig. 2). Compared to CK, the EL of HHZ and XLY900 under NaCl stress significantly increased by 16.26%-126.50% and 35.25%-71.98% from days 3 through 9, respectively. After NaCl treatment, there was a significant rise in the MDA and H_2O_2 content of HHZ in the S treatment. This increase ranged from 31.79% to 51.73% for MDA and 13.92% to 30.29% for H_2O_2 during the period from day 3 to 9, as compared to CK. In the S treatment of XLY900, the content of MDA and H_2O_2 was significantly enhanced by 22.25%-40.52% and 20.26%-25.09%, compared with CK, from days 3

through 9, respectively. The H_2O_2 and MDA contents of HHZ were higher than that of XLY900 on 9 d after NaCl stress, showing that NaCl stress was more harmful to HHZ, which was more sensitive to NaCl stress than XLY900. Compared with S treatment, spraying Hemin effectively reduced EL and rice varieties' MDA and H_2O_2 content.

In contrast to the S treatment, the EL of both HHZ and XLY900 exhibited a noticeable decrease in the SH treatment, showcasing reductions of 9.64% to 28.20% and 8.78% to 18.41%, respectively. The MDA and H_2O_2 content in the SH treatment of HHZ than in the S treatment decreased by 15.20%-20.28% and 11.59%-18.14%, while in the SH treatment of XLY900, they were 8.30%-16.52% and 5.97%-15.72% less than those in the S treatment from day 3 through 9, respectively. EL, MDA, and H_2O_2 remained high in both varieties under ZnPP treatment. Throughout the stress period, the SZH treatment led to a reduction in EL, MDA content, and H_2O_2 content of both HHZ and XLY900, when compared to the SZ treatment. On days 3 and 9, compared to SZ treatment, EL of HHZ exhibited noticeable decreases of 9.21% and 10.43%, respectively, in SZH treatment. From days 3 through 9, compared to SZ treatment, the EL of XLY900 with SZH treatment dramatically declined by 6.09%-9.01%. From days 3 through 9, compared with SZ treatment, the MDA and H_2O_2 content were decreased by 6.51%-7.15% and 3.51%-10.99% in HHZ with SZH treatment, were reduced by 1.44%-7.71% and 1.22%-9.71% in XLY900 with SZH treatment, respectively.

The histochemical localization of H_2O_2 and $O_2^{\cdot-}$ in rice leaves

The distribution of H_2O_2 and superoxide anion ($O_2^{\cdot-}$) were localized and expressed visually by histochemical localization of HHZ and XLY900 rice leaves. It was shown that H_2O_2 and $O_2^{\cdot-}$ in the leaves were dark brown and blue spots (Fig. 3). Compared to CK, dark brown and dark blue spots were significantly increased in rice leaves of both varieties under NaCl stress. Compared to the S treatment, dark brown and dark blue spots on leaves were significantly decreased in HHZ and XLY900 with SH treatments, which indicated that foliar spraying of Hemin could potentially reduce the accumulation and distribution of H_2O_2 and $O_2^{\cdot-}$. ZnPP treatment failed to lower the accumulation of ROS in the leaves, and dark brown spots and dark blue spots remained remarkably increased. There was a reduced accumulation of ROS with the combination of ZnPP and Hemin. Compared to the ZnPP treatment, the number of dark brown spots and dark blue spots decreased in HHZ and XLY900 with SZH treatments.

The SOD, POD, and CAT activity in rice seedlings

With the extension of exposure time, the SOD and POD activities in the S treatment of HHZ showed an upward and downward trend, and CAT activity showed an increased trend compared to CK (Fig. 4). Compared to CK, the SOD, POD, and CAT activities in the S treatment of XLY900 showed an upward trend with the prolonged time of NaCl stress. The SOD and POD activities in S treatment of HHZ reached the maximum at 3 d of NaCl stress, which were significantly increased, by 13.82% and 13.64%, respectively, and CAT activities increased by 11.45%-21.71% from 3 to 9 d of NaCl stress compared to CK. In comparison to CK, the SOD, POD, and CAT activities of XLY900 under NaCl stress increased by 7.30%-26.63%, 6.64%-14.26%, and 15.97%-24.76% from day 3 through 9, respectively. The application of exogenous Hemin boosted the SOD, POD, and CAT activities of SH treatment in two rice varieties. Compared to S treatment, the SOD,

281 POD, and CAT activities of HHZ with SH treatment were increased by 4.41%-17.66%, 6.48%-
 282 12.67%, and 6.43%-17.33% from day 3 through 9, respectively. In comparison to S treatment, the
 283 SOD and CAT activities of XLY900 with SH treatment were increased, by 5.53%-27.47% and
 284 10.54%-18.12%, from day 3 through 9, respectively, while POD activity increased by 4.53%-
 285 9.20% except for the day 5. Compared with the S treatment, the ZnPP treatment did not enhance
 286 the enzyme activity under the stress environment but lowered the enzyme activity. For example,
 287 compared to S treatments, on day 3, the CAT activity in SZ treatment of HHZ was significantly
 288 decreased by 6.54%; on day 5, the SOD activity in SZ treatment of XLY900 was significantly
 289 reduced by 11.12%. The combination with Hemin relieved the adverse effects of ZnPP and
 290 improved the above enzyme activities. Compared with the SZ treatment, the SZH treatment of
 291 HHZ showed SOD activity increased by 3.10%-13.12% from day 3 through 9; POD activity
 292 significantly enhanced by 8.05% on day 9; CAT activity significantly raised by 11.52% on day 3.
 293 Compared with the SZ treatment, the SZH treatment of XLY900 showed SOD activity markedly
 294 increased by 15.79% and 22.93% on days 3 and 5, respectively; POD activity significantly
 295 enhanced by 7.47%-8.07% from day 5 through 9; CAT activity significantly raised by 13.67% and
 296 12.48% on day 5 and 7, respectively.

297 The AsA-GSH cycle none-enzymatic antioxidants in rice leaves

298 With the time of NaCl stress extension, the AsA content decreased and the DHA and AsA+DHA
 299 content increased in the leaves of HHZ and XLY900 (Fig. 5). From day 5 through 9, compared to
 300 CK, the AsA content in the S treatment of HHZ and XLY900 decreased by 2.16%-15.52% and
 301 4.62%-14.26%, significantly different between treatments and control. In comparison to CK, the
 302 DHA and AsA+DHA content of HHZ under NaCl stress increased by 21.16%-60.17% and 4.47%-
 303 34.18%; for XLY900, the abovementioned indicators increased by 57.73%-67.58% and 10.39%-
 304 32.46%, from day 3 through 9, respectively. The application of exogenous Hemin further boosted
 305 the AsA content and diminished the accumulation of DHA and AsA+DHA. Compared to the S
 306 treatment, the AsA content in SH treatment of HHZ and XLY900 increased, by 4.63%-15.54%
 307 and 5.46%-10.44% from day 5 through 9, respectively, significantly different between S
 308 treatments and control. In comparison to S treatment, the DHA and AsA+DHA content in SH
 309 treatment of HHZ decreased by 15.53%-30.23% and 5.06%-19.87% from day 3 through 9,
 310 respectively; for XLY900, the abovementioned indicators decreased by 19.87%-29.67% and
 311 5.43%-12.57%. Under NaCl stress, ZnPP treatment mainly raised DHA and AsA+DHA contents
 312 in the leaves of two rice varieties. In comparison to S treatments, on day 7, the DHA and
 313 AsA+DHA content in the SZ treatment of HHZ were significantly increased by 15.00% and
 314 8.49%; on day 9, the DHA content in the SZ treatment of XLY900 was significantly increased by
 315 8.00%. In the combination of ZnPP and Hemin, the AsA content was higher, and the DHA and
 316 AsA+DHA contents were lower in both rice varieties compared to the ZnPP treatment. In
 317 comparison to SZ treatments, on days 5 and 9, the AsA content in the SZH treatment of HHZ was
 318 significantly increased by 11.53% and 3.22%, respectively; on days 5 and 7, the AsA content in
 319 the SZ treatment of XLY900 was significantly increased by 7.15% and 9.09%, respectively. In
 320 comparison to SZ treatment, the DHA and AsA+DHA content in SZH treatment of HHZ decreased
 321 by 12.39%-26.77% and 2.81%-14.35% from day 3 through 9, respectively; for XLY900, the

abovementioned indicators decreased by 8.08%-16.27% and 1.72%-7.34%.

It can be seen from Fig. 6 that, with the extension of the stress time, the contents of GSH and GSH+GSSG in S treatment leaves of both HHZ and XLY900 decreased; GSSG in S treatment leaves of both HHZ and XLY900 increased. On day 3, day 5, and day 9, compared to CK, the GSH content in the S treatment of HHZ significantly decreased 5.83%, 8.27% and 2.28%, respectively; in XLY900, the GSH content significantly decreased 3.49%, 7.17% and 8.68%, respectively. From days 3 through 9, compared to CK, the GSSG content in the S treatment of HHZ and XLY900 increased by 7.25%-22.36% and 8.20%-16.87%, significantly different between treatments and control. On day 3, and day 5, compared to CK, the GSH+GSSG content in the S treatment of HHZ significantly decreased 4.63% and 6.96%, respectively; On day 5 and day 9 in XLY900, the GSH content significantly decreased 5.22% and 6.93%, respectively. The application of exogenous Hemin further boosted the GSH and GSH+GSSG contents diminished the accumulation of DHA. Compared to the S treatment, the GSH content in SH treatment of HHZ and XLY900 increased, by 1.96%-14.31% and 3.60%-8.69% from day 5 through 9, respectively. In comparison to S treatment, the GSSG content in SH treatment of HHZ and decreased by 8.57%-22.36% and 5.74%-6.35% from day 3 through 9, respectively. Under NaCl stress, ZnPP treatment mainly raised GSSG content in the leaves. In comparison to S treatments, on day 3 and day 7, the GSSG content in the SZ treatment of XLY900 were significantly increased by 4.71% and 8.34%. In the combination of ZnPP and Hemin, the GSH and GSH+GSSG content were higher, and the GSSG content was lower in both rice varieties compared to the ZnPP treatment. Compared to the SZ treatment, the GSH content in SZH treatment of HHZ and XLY900 increased, by 1.77%-10.55% and 1.80%-8.16% from day 3 through 9, respectively. Compared to the SZ treatment, the GSSG content in SZH treatment of HHZ and XLY900 decreased, by 4.90%-5.82% and 6.71%-8.33% from day 3 through 9, respectively. In comparison to SZ treatments, on days 3, 5 and 7, the GSSG content in the SZH treatment of HHZ were significantly increased by 4.90%, 9.11% and 3.22%, respectively; on days 3 and 9, the GSSG content in the SZH treatment of XLY900 was significantly increased by 6.41% and 4.34%, respectively.

The AsA-GSH cycle enzymatic activities in rice leaves

As shown in Figure 7, APX, MDHAR, DHAR, and GR activities were increased with the prolongation of NaCl stress time. Compared with CK, during the stress period, the activities of the above four enzymes in the S treatment of HHZ were markedly enhanced by 11.00%-18.88%, 14.95%-54.23%, 23.19%-56.82% and 12.22%-27.96% respectively, and of XLY900 were significantly increased 18.82%-21.21%, 29.84%-51.15%, 19.62%-46.87% and 10.48%-13.56%, respectively. The use of Hemin further improved the activity of APX, MDHAR, DHAR, and DHAR. Compared with S treatment, from day 3 through 9, the activities of the above four enzymes in the SH treatment of HHZ were enhanced by 15.18%-25.33%, 19.95%-58.63%, 7.10%-33.25% and 8.65%-14.11%, while in SH treatment of XLY900 were increased 17.76%-26.90%, 11.84%-50.44%, 15.92%-24.11% and 7.47%-12.26%, respectively. However, with the use of ZnPP the activity of APX, MDHAR, DHAR, and GR was diminished. On day 3, in comparison to S treatment, the APX activity of HHZ in SZ treatment was significantly decreased by 17.03%. On day 9, in comparison to CK, the GR activity of HHZ and XLY900 in SZ treatment was

significantly decreased, by 7.14% and 6.46%, respectively. The combination of ZnPP with Hemin increased the above enzyme activities. In HHZ with SZH treatment, compared with SZ treatment, the APX activity was significantly increased by 11.41% and 21.15% on day 3 and day 7, respectively; the MDHAR activity was markedly increased by 21.78%-38.70%, from day 3 through 9; the DHAR activity was dramatically increased by 9.98%-29.65%, from day 3 through 7; the GR activity was remarkably increased by 13.47%, and 8.81%, on day 7, and day 9, respectively. In XLY900 with SZH treatment, compared with SZ treatment, the APX activity significantly increased by 21.60% and 29.99% on day 5 and day 9, respectively; the MDHAR activity was markedly increased by 32.81% and 20.13% on day 5, and day 7, respectively; the DHAR activity was dramatically increased by 14.37%-16.89%, from day 5 through 9; the GR activity was remarkably increased by 7.18%-9.02%, from day 5 through 9.

The content of osmoregulatory substances in rice leaves

NaCl stress caused a significant increase in proline content in the leaves of HHZ and XLY900 (Fig. 8 a and b). Compared to CK, the proline content of HHZ under NaCl stress was significantly increased by 34.95%-65.34%, from days 3 through 9. From 3 to 9 d, compared to CK, the proline content of XLY900 with S treatment dramatically increased by 18.95%-54.16%. Under NaCl stress, the increase in proline was more in HHZ than in XLY900. Hemin treatment further enhanced the proline content in the leaves of two rice varieties. Compared to S treatment, the proline content of HHZ and XLY900 with SH treatment significantly improved by 8.38%-27.10%, and 15.02%-24.35%, respectively, from days 3 through 9. Proline content of rice leaves was not improved by ZnPP treatment. For example, on day 3, compared to S treatment, the proline content of XLY900 with SZ treatment decreased by 8.64%. In combination with ZnPP and Hemin, the proline content was enhanced. Compared to SZ treatment, the proline content of HHZ with SZH treatment had a maximum improvement of 26.87% on day 9, and XLY900 with SZH treatment had a maximum growth of 26.51% on day 7. The soluble protein content of HHZ markedly increased in the early stage (3 d) and then decreased in the later stage (5-9 d) compared with CK under NaCl stress (Fig.8 c and d). The soluble protein content in XLY900 increased during the stress period with the difference reaching significant levels at all four-time points. Applying of Hemin enhanced soluble protein content in the leaves of two rice varieties. Compared with S treatment, soluble protein content noticeably increased by 2.75%, in SH treatment of HHZ, on day 9, while significantly improved by 3.93%, and 1.17%, respectively in XLY900. Spraying ZnPP did not increase soluble protein content. For example, compared with S treatment, soluble protein content significantly decreased by 3.20%, in SH treatment of HHZ, on day 3. When ZnPP was combined with Hemin, soluble protein content was enhanced. On day 3, soluble protein content was increased by 3.02% and 3.21%, respectively, of HHZ and XLY900 in SZH treatment.

Discussion

Salt stress is the most prevalent abiotic stress that limits crop growth and development. Research has shown that salt stress impedes the growth of several crops, such as wheat (Ashraf et al., 2023), sorghum (Liu et al., 2023), and soybean (Feng et al., 2021). Excessive salt interferes with normal biological and physiological processes and influences normal plant growth (Talubaghi et al.,

2022), such as reduced plant height, narrowed stem base width and diminished biomass. The results of this test were similar to the above studies. Under salt stress, the seedling growth of both HHZ and XLY900 was significantly inhibited, and all the morphological indexes were decreased (Table 1 and Table 2). Foliar spraying of Hemin positively regulated various morphological indicators and promoted aboveground growth and biomass accumulation in rice seedlings. Liu et al. (2021) showed that Hemin improved the growth of maize seedlings and increased biomass accumulation under drought stress. Furthermore, Hemin degraded in plants to produce CO, which alleviated the inhibition of wheat growth by NaCl stress (Ling et al., 2009). Exogenous ZnPP was unable to promote rice growth under salt stress in this study, which was consistent with the research of Cao et al. (2011).

ROS can be used as low concentrations of secondary messengers and signaling molecules (Antoniou et al., 2016). Plants generate and remove ROS in dynamic balances under normal growth conditions. Under abiotic stress conditions, ROS are surged, which in large quantities are destructive, leading to changes in the structure of DNA, proteins and enzymes, ultimately resulting in programmed cell death (Gill and Tuteja 2010; Singh et al., 2019). MDA is one of the membrane lipid peroxidation products whose content can reflect the level of ROS and the degree of membrane lipid peroxidation. EL can evaluate cell membrane permeability; larger represents stronger osmotic cell membranes (Ben Youssef et al., 2021). In this experiment, the findings showed that salt stress caused higher leaf EL, increased MDA and H₂O₂ contents in two rice varieties and that the results were positively correlated with stress duration (Fig. 2). Compared with 900, HHZ had a much stronger increase in the above three indexes, indicating that HHZ was more intensely exposed to the stress. This was similar to the findings of previous works, which indicated that HHZ was more deeply stressed (Chen et al., 2022). The localization of H₂O₂ and O₂^{·-} in leaves was measured by histochemical methods. Salt stress induced the accumulation of H₂O₂ and O₂^{·-} in the leaves of HHZ and XLY900 compared with CK (Fig. 3). This is in conformity with the findings of Jabeen et al. (2020) who worked on cultivated rice under salt stress. Hemin has the ability to mitigate the damage caused by adversity stress to plants, reducing ROS accumulation, MDA content and cell membrane permeability (Chen et al., 2009; Cui et al., 2012). The results of this experiment were in agreement with this conclusion. Foliar spraying of Hemin effectively diminished EL levels, H₂O₂ and MDA contents (Fig. 2), reduced H₂O₂ and O₂^{·-} accumulation (Fig. 3), and alleviated the damage of salt stress on cell membranes. Exogenous ZnPP could not scavenge excess reactive oxygen species and maintain cell membrane stability. When ZnPP was combined with Hemin, it scavenged part of the ROS and alleviated oxidative damage, this finding is in agreement with Zhang et al. (2012).

Facing stress, plants activate antioxidant defense systems to minimize damage caused by oxidative stress. Among them, antioxidant enzymes mainly include SOD, CAT, and POD. SOD represents the first barrier for plants to resist ROS damage caused by abiotic stresses and catalyzes the transformation of O₂^{·-} to O₂ (Karuppanapandian and Kim 2013). CAT eliminates H₂O₂ with minimal energy consumption and very high conversion rates for large-scale scavenging of ROS (Zamocky et al., 2012). POD has a strong affinity for H₂O₂ and is used for the delicate regulation of H₂O₂ (Abogadallah 2010). In this study, the effect of salt stress on SOD and POD activities of

HHZ showed an upward followed by a downward trend, and CAT activity was upward compared with CK (Fig. 4 a, c and e); SOD, POD and CAT activities of XLY900 were on an upward trend (Fig. 4 b, d and f). This indicates that in the short term of salt stress, rice eliminates ROS by increasing the activity of antioxidant enzymes; in the long term of salt stress, rice accumulates more ROS, which can't able to be scavenged in time by decreasing the activity of antioxidant enzymes. This is consistent with the results of previous findings (Vaidyanathan et al., 2003; Seckin et al., 2009 and Kumari et al., 2023). Foliar spraying of Hemin enhanced SOD, POD and CAT activities in leaves of two rice varieties under salt stress (Fig. 4). This demonstrates that exogenous Hemin stimulates the antioxidant enzyme system in rice and facilitates the increase of enzyme activity, which avoids oxidative damage and ensures normal plant growth. The inhibitor ZnPP was unable and even reduced some of the antioxidant enzyme activities. ZnPP combined with Hemin mitigated the inhibitory effect caused by ZnPP, with improved SOD, POD and CAT activities (Fig. 4). Based on a previous study (Zhang et al., 2012), it is hypothesized that Hemin enhances antioxidant enzyme activity in rice leaves by promoting HO expression and thereby increasing the antioxidant enzyme activity, while ZnPP acts as an inhibitor of HO hindering its expression which restricts the increase in antioxidant enzyme activity.

The AsA-GSH cycle is an essential ROS scavenging mechanism, mainly consists of the antioxidant enzymes APX, MDHAR, DHAR, and GR and the nonenzymatic antioxidants AsA and GSH, which can alleviate the oxidative damage caused by salt stress (Wang et al., 2022). The cycling process: AsA is catalyzed by APX, which converts H_2O_2 to H_2O and is oxidized to MDHA. MDHA is converted to AsA by reduction-oxidation reaction with MDHAR or to DHA by non-enzymatic disproportionation reaction. DHAR catalyzes DHA and GSH to produce AsA and GSSG, while GSSG can be restored to GSH by GR (Nahar et al., 2015; Tan et al., 2022). AsA and GSH act as nonenzymatic antioxidants and assist other antioxidant enzymes in scavenging ROS. In this study, salt stress decreased AsA content and increased DHA and AsA+DHA content in rice leaves (Fig. 5), indicating that APX activity enhancement decreased in AsA content. Foliar spraying of Hemin significantly improved AsA content and diminished DHA and AsA+DHA content. The findings suggest that the increase in MDHAR and DHAR activities causes an increase in AsA content and a decrease in DHA content. Under salt stress, GSH and GSH+GSSG content decreased, and GSSG content increased, while exogenous Hemin treatment increased GSH and GSH+GSSG content and decreased GSSG content in rice leaves (Fig. 6). The experiment showed that the enhanced GR activity facilitated the conversion of GSSG to GSH and maintained a high level of reduction state GSH, which was in agreement with the research of Piao et al. (2022). These indicated that Hemin sustained cellular reduction force at a high level, which resisted oxidative damage. In addition, in this experiment, salt stress increased APX, MDHAR, DHAR, and GR activities in two rice varieties compared with the control (Fig. 7). It indicated that salinity stress increased the H_2O_2 content of rice leaves, which prompted APX to accelerate the scavenging of H_2O_2 ; while the increased activities of MDHAR, DHAR and GR were beneficial to leaves' resistance to oxidative damage, which was a stress response to excess H_2O_2 . Foliar spraying of Hemin further induced the activities of APX, MDHAR, DHAR, and GR (Fig. 7). Previous studies have suggested that this might be possible by upregulating the transcription of genes for enzymes

related to the metabolism of the degradation products CO and GSH, which could increase the enzyme activity to help plants mitigate the oxidative damage caused by the stresses (Zhang et al., 2016). ZnPP cannot be degraded to CO₂ in plants and reduces endogenous CO₂ production by blocking HO expression; thus, could not enhance the activities of MDHAR, DHAR, and GR under salt stress. Moreover, Hemin induced HO gene expression and enhanced gene expression of critical enzymes in the AsA-GSH cycle, while ZnPP prevented HO expression and even strengthened the inhibitory effect of NaCl stress on the AsA-GSH cycle in rice seedlings (Cui et al., 2012). These results reflect that Hemin improved the efficiency of ROS scavenging in rice leaves, which maintained cell membrane stability and enhanced the resistance of rice.

Although saline soils contain water, plants cannot absorb it, mainly because the soil has a high level of ions that increase the osmotic pressure of the external environment, which prevents plant cells from absorbing water or even losing it. Therefore, plants ensure water absorption by increasing osmoregulatory substances and decreasing the difference in osmotic potential between the inside and outside cells. The important osmoregulatory substances, soluble proteins, and proline have different physiological functions in maintaining osmotic balance in plants. Soluble proteins can help the bound water in plant cells to combine with the liner and maintain the stability of the cell structure (Hao et al., 2021). Proline is a potential non-enzymatic antioxidant that functions as a scavenger of single-linear oxygen molecules and hydroxyl radicals; thus, proline prevents lipid peroxidation of cell membranes and avoids exposure of plants to ROS-induced oxidative damage (Szabados and Savoure 2010). In this study, we found that with the increase of NaCl stress time, soluble protein content initially increased and then decreased in HHZ, while it continuously increased in XLY900 (Fig. 7). Previous study have shown that salt stress disrupts the protein synthesis pathway at later stages, accelerating its catabolism, generating large amounts of amino acids, and ultimately reducing protein content (Alisofi et al., 2020). This could be the reason for the decrease in soluble protein content in HHZ leaves. The soluble protein content in XLY900 leaves was enriched to relieve the difference in osmotic potential. Two rice varieties under salt stress had significantly increased proline content. Compared with XLY900, salt stress caused HHZ to produce much more proline (Fig. 8 c and d). This was similar to the results of Gao et al. (2016), in which salt-sensitive varieties had high proline content in stressful environments. Foliar spraying of Hemin promoted the accumulation of osmoregulatory substances in rice leaves, which significantly increased soluble protein and proline contents. However, in the ZnPP treatment, the content of osmoregulatory substances was reduced instead of increased. Similar results were confirmed in the research of Zhao et al. (2022). It indicates that Hemin induces a large accumulation of proline and soluble proteins, which is beneficial for the absorption of water and the maintenance of cellular osmotic pressure in rice leaves under salt stress.

Conclusions

During the seedling stage, the activity of antioxidant enzymes and the contents of non-enzymatic antioxidants initially rose in response to salt stress. This response effectively countered the accumulation of reactive oxygen species (ROS) induced by the stress. However, with prolonged

exposure to stress, the enzyme activity continued to increase while the content of the antioxidants decreased, failing to adequately alleviate the stress in a timely manner. The accumulated ROS and membrane lipid peroxides exacerbated the damage caused by salt stress on rice, eventually leading to a decrease in growth. The application of Hemin through foliar treatment additionally enhanced the antioxidant enzymes activity and elevated the non-enzymatic antioxidants contents. This augmentation contributed to an overall improvement in the rice's antioxidant capacity, resulting in a reduction of membrane lipid peroxidation. The consistent functionality of the AsA-GSH cycle was ensured, consequently enhancing rice's resistance to salt stress.

Funding

Special Projects in Key Areas of Ordinary Colleges of the Educational Commission of Guangdong Province (2021ZDZX4027), Innovation Team Project of ordinary colleges of the Educational Commission of Guangdong Province (2021KCXTD011), Zhanjiang Science and Technology Bureau (2022A01016), Research start-up project of Guangdong Ocean University (R20046), Research start-up project of Guangdong Ocean University (060302052012).

Competing interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

Fengyan Meng conceived and designed the study performed the experiments, analyzed the data, authored or reviewed drafts of the article, interpreted the results and improve manuscript, and approved the final draft.

Naijie Feng conceived and designed the experiment performed the experiments, authored or reviewed drafts of the article, and approved the final draft.

Dianfeng Zheng analyzed the data, authored or reviewed drafts of the article, and approved the final draft.

Meiling Liu carried out experiments, prepared figures and/or tables, and approved the final draft.

Hang Zhou reviewed drafts of the article, and approved the final draft.

Rongjun Zhang performed the experiments, prepared figures and/or tables, and approved the final draft.

Xixin Huang analyzed the data, prepared figures and/or tables, and approved the final draft.

Anqi Huang analyzed the data, prepared figures and/or tables, and approved the final draft.

References

Abogadallah G M. 2010. Antioxidative defense under salt stress. *Plant signaling behavior* 5: 369-374 DOI: 10.4161/psb.5.4.10873.

Ahmad P, Abdel Latef A A, Hashem A, Abd Allah E F, Gucel S, Tran L S P. 2016. Nitric oxide mitigates salt stress by regulating levels of osmolytes and antioxidant enzymes in chickpea. *Frontiers in Plant science* 7: 347 DOI: 10.3389/fpls.2016.00347.

Alisofi S, Einalia A, Sangtarash M H. 2020. Jasmonic acid-induced metabolic responses in bitter melon (*Momordica charantia*) seedlings under salt stress. *Journal of horticultural science biotechnology* 95: 247-259 DOI: 10.1080/14620316.2019.1663135.

Alkharabsheh H M, Seleiman M F, Hewedy O A, Battaglia M L, Jalal R S, Alhammad B A, Schillaci C, Ali N, Al-Doss, A. 2021. Field crop responses and management strategies to mitigate soil salinity in modern agriculture: a review. *Agronomy-basel* 11: 2299 DOI: 10.3390/agronomy11112299.

Antoniou C, Savvides A, Christou A, Fotopoulos V. 2016. Unravelling chemical priming machinery in plants: the role of reactive oxygen-nitrogen-sulfur species in abiotic stress tolerance enhancement. *Current opinion in plant biology* 33: 101-107 DOI: 10.1016/j.pbi.2016.06.020.

Ashraf M A, Hafeez A, Rasheed R, Hussain I, Farooq U, Rizwan M, Ali S. 2023. Evaluation of physiological and biochemical responses for salt tolerance in wheat (*Triticum aestivum* L.) cultivars. *Journal of plant growth regulation* 43: 4402-4422 DOI: 10.1007/s00344-023-10905-4.

Athar H U, Zulfiar F, Moosa A, Ashraf M, Zafar Z U, Zhang L X, Ahmed N, Kalaji H M, Nafees M, Hossain M A, Islam M S, Elsabagh A, Siddique K H M. 2022. Salt stress proteins in plants: an overview. *Frontiers in plant science* 13: 999058 DOI: 10.3389/fpls.2022.999058.

Basilio-apolinar A, Eugenio Gonzalez-de La Vara L, Gabriel Ramirez-pimentel J, Aguirre-Mantilla Cesar L, Iturriaga Gabriel, Covarrubias-Prieto Jorge, Carlos Raya-Perez Juan. 2021. Silicon induces changes in the antioxidant system of millet cultivated in drought and salinity. *Chilean journal of agricultural research* 81: 655-663 DOI: 10.4067/S0718-58392021000400655.

Ben Youssef R, Jelali N, Boukari N, Albacete A, Martinez C, Alfocea F P, Abdely C. 2021. The Efficiency of different priming agents for improving germination and early seedling growth of local tunisian barley under salinity stress. *Plants basel* 10: 2264 DOI: 10.3390/plants10112264.

Cao Z, Geng B, Xu S, Xuan, W, Nie, L, Shen, W B, Liang, Y C, Guan, R Z. 2011. BnHO1, a haem oxygenase-1 gene from *Brassica napus*, is required for salinity and osmotic stress-induced lateral root formation. *Journal of experimental botany* 62: 4675-4689 DOI: 10.1093/jxb/err190.

Chandrakar V, Yadu B, Meena R K, Dubey A, Keshavkant S. 2017. Arsenic-induced genotoxic responses and their amelioration by diphenylene iodonium, 24-epibrassinolide and proline in *Glycine max* L. *Plant physiology biochemistry* 112: 74-86 DOI: 10.1016/j.plaphy.2016.12.023.

Chen G, Zheng D, Feng N, Zhou H, Mu D W, Zhao L M, Shen X F, Rao G S, Meng F Y, Huang A Q. 2022. Physiological mechanisms of ABA-induced salinity tolerance in leaves and roots of rice. *Scientific reports* 12: 8228 DOI: 10.1038/s41598-022-11408-0.

Chen X Y, Ding X, Xu S, Wang R, Xuan W, Cao Z Y, Chen J, Wu H H, Ye M B, Shen W B. 2009. Endogenous hydrogen peroxide plays a positive role in the upregulation of heme oxygenase and acclimation to oxidative stress in wheat seedling leaves. *Journal of integrative plant biology* 51: 951-960 DOI: 10.1111/j.1744-7909.2009.00869.x.

Costa H, Gallego S M, Tomaro M L. 2002. Effect of UV-B radiation on antioxidant defense system in sunflower

603 cotyledons. Plant science 162: 939-945 DOI: 10.1016/S0168-9452(02)00051-1.

604 Cui W T, Li L, Gao Z Z, Wu H H, Xie Y J, Shen W B. 2012. Haem oxygenase-1 is involved in salicylic acid-
605 induced alleviation of oxidative stress due to cadmium stress in *Medicago sativa*. Journal of experimental botany
606 63: 5521-5534 DOI: 10.1093/jxb/ers201.

607 Ding Z, Kheir A M S, Ali O A M, Hafez E M, ElShamey E A, Zhou ZnX, Wang B Z, Lin X E, Ge, Y, Fahmy
608 A E, Seleiman M F. 2021. A vermicompost and deep tillage system to improve saline-sodic soil quality and
609 wheat productivity. Journal environmental management 277: 111388 DOI: 10.1016/j.jenvman.2020.111388.

610 Feng N, Yu M, Li Y, Jin D, Zheng D F. 2021. Prohexadione-calcium alleviates saline-alkali stress in soybean
611 seedlings by improving the photosynthesis and up-regulating antioxidant defense. Ecotoxicology environmental
612 safety 220: 112369 DOI: 10.1016/j.ecoenv.2021.112369.

613 Fu G, Zhang L, Cui W, Wang Y Q, Shen W B, Ren Y; Zheng T Q. 2011. Induction of heme oxygenase-1 with
614 β -CD-hemin complex mitigates cadmium-induced oxidative damage in the roots of *Medicago sativa*. Plant and
615 soil 345: 271-285 DOI: 10.1007/s11104-011-0779-x.

616 Gao Y, Lu Y, Wu M Q, Liang E X, Li Y, Zhang D P, Yin Z T, Ren X Y, Dai Y, Deng D X. 2016. Ability to
617 remove Na^+ and retain K^+ correlates with salt tolerance in two maize inbred lines seedlings. Frontiers in plant
618 science 7: 1716 DOI: 10.3389/fpls.2016.01716.

619 Gill S S, Tuteja N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop
620 plants. Plant physiology biochemistry 48: 909-930 DOI: 10.1016/j.plaphy.2010.08.016.

621 Habib N, Ali Q, Ali S, Haider M Z, Javed M T, Khalid M, Perveen R, Alsahli A A, Alyemeni M N. 2021. Seed
622 priming with sodium nitroprusside and H_2O_2 confers better yield in wheat under salinity: water relations,
623 antioxidative defense mechanism and ion homeostasis. Journal of plant growth regulation 40: 2433-2453 DOI:
624 10.1007/s00344-021-10378-3.

625 Hao S H, Wang Y R, Yan Y X, Liu Y H, Wang J Y, Chen S. 2021. A review on plant responses to salt stress
626 and their mechanisms of salt resistance. Horticulturae 7: 132 DOI: 10.3390/horticulturae7060132.

627 Hasnuzzaman M, Bhuyan M, Zulfiqar F, Raza A, Mohsin S M, Al Mahmud J, Fujita M, Fotopoulos V. 2020.
628 Reactive oxygen species and antioxidant defense in plants under abiotic stress: revisiting the crucial role of a
629 universal defense regulator. Antioxidants 9: 681 DOI: 10.3390/antiox9080681.

630 Hasanuzzaman M, Hossain M A, Fujita M. 2011. Nitric oxide modulates antioxidant defense and the
631 methylglyoxal detoxification system and reduces salinity-induced damage of wheat seedlings. Plant
632 biotechnology reports 5: 353-365 DOI: 10.1007/s11816-011-0189-9.

633 Huang M, Zeng L, Liu C, Li X Y, Wang H L. 2022. Research on the eco-efficiency of rice production and its
634 improvement path: a case study from China. International journal of environmental research and public health
635 19: 8645 DOI: 10.3390/ijerph19148645.

636 Jabeen Z, Hussain N, Irshad F, Zeng J B, Tahir A, Zhang G P. 2020. Physiological and antioxidant responses of
637 cultivated and wild barley under salt stress. Plant soil and environment 66: 334-344. DOI: 10.17221/169/2020-
638 PSE.

639 Karuppanapandian T, Kim W. 2013. Cobalt-induced oxidative stress causes growth inhibition associated with
640 enhanced lipid peroxidation and activates antioxidant responses in Indian mustard (*Brassica juncea* L.) leaves.
641 Acta physiologiae plantarum 35: 2429-2443 DOI: 10.1007/s11738-013-1277-y.

642 Kaya C, Ugurlar F, Ashraf M, Alam P, Ahmad P. 2023. Nitric oxide and hydrogen sulfide work together to
643 improve tolerance to salinity stress in wheat plants by upraising the AsA-GSH cycle. Plant physiology and

644 biochemistry. 194: 651-663 DOI: 10.1016/j.plaphy.2022.11.041.

645 Keles Y, Oncel I. 2002. Response of antioxidative defence system to temperature and water stress combinations
646 in wheat seedlings. Plant science 163: 783-790 DOI: 10.1016/S0168-9452(02)00213-3.

647 Kenawy E-R, Rashad M, Hosny A, Shendy S, Gad D, Saad-Allah K M. 2022. Enhancement of growth and
648 physiological traits under drought stress in Faba bean (*Vicia faba* L.) using nanocomposite. Journal of plant
649 interactions 17: 404-41 DOI: 10.1080/17429145.2022.2038293.

650 Kumari S, Nazir F, Jain K, Khan M I R. 2023. GABA and potassium modulates defence systems, assimilation
651 of nitrogen and carbon, and yield traits under salt stress in wheat. Journal of plant growth regulation DOI:
652 10.1007/s00344-023-10992-3.

653 Lin Y J, Yu X Z, Li Y H, Yang L. 2020. Inhibition of the mitochondrial respiratory components (Complex I and
654 Complex III) as stimuli to induce oxidative damage in *Oryza sativa* L. under thiocyanate exposure. Chemosphere
655 243: 125472 DOI: 10.1016/j.chemosphere.2019.125472.

656 Ling T, Zhang B, Cui W, Wu M Z, Lin J S, Zhou W T, Huang J J, Shen W B. 2009. Carbon monoxide mitigates
657 salt-induced inhibition of root growth and suppresses programmed cell death in wheat primary roots by inhibiting
658 superoxide anion overproduction. Plant science 177: 331-340 DOI: 10.1016/j.plantsci.2009.06.004.

659 Liu C T, Mao B G, Yuan D Y, Chu C C, Duan M J. 2022. Salt tolerance in rice: physiological responses and
660 molecular mechanisms. Crop journal 10: 13-25 DOI: 10.1016/j.cj.2021.02.010.

661 Liu J, Wi Y Q, Dong G C, Zhu G L, Zhou G S. 2023. Progress of research on the physiology and molecular
662 regulation of sorghum growth under salt stress by gibberellin. International journal of molecular sciences
663 24:6777 DOI: 10.3390/ijms24076777.

664 Liu L J, Huang L, Lin X Y, Sun C L. 2020. Hydrogen peroxide alleviates salinity-induced damage through
665 enhancing proline accumulation in wheat seedlings. Plant cell reports. 39: 567-575. DOI: 10.1007/s00299-020-
666 02513-3.

667 Liu X, Meng Y, Wei S, Gu W R. 2021. Exogenous Hemin confers cadmium tolerance by decreasing cadmium
668 accumulation and modulating water status and matter accumulation in maize seedlings. Agronomy-basel 11
669 DOI: 10.3390/agronomy11040739.

670 Lu X P, Min W F, Shi Y F, Tian L, Li P F, Ma T L, Zhang Y X, Luo C K. 2022. Front plant science 13:849553
671 DOI: 10.3389/fpls.2022.849553.

672 Mukhopadhyay R, Sarkar B, Jat H S, Sharma P C, Bolan N S. 2021. Soil salinity under climate change:
673 Challenges for sustainable agriculture and food security. Journal of environmental management 280: 111736
674 DOI: 10.1016/j.jenvman.2020.111736.

675 Nahar K, Hasanuzzaman M, Alam M M, Fujita M. 2015. Exogenous spermidine alleviates low temperature
676 injury in mung bean (*Vigna radiata* L.) seedlings by modulating ascorbate-glutathione and glyoxalase pathway.
677 International Journal of molecular sciences 16: 30117-30132 DOI: 10.3390/ijms161226220.

678 Piao L, Wang Y, Liu X M, Sun G Y, Zhang S Y, Yan J Y, Chen Y, Meng Y, Li M, Gu W R. 2022. Exogenous
679 Hemin alleviated cadmium stress in maize (*Zea mays* L.) by enhancing leaf photosynthesis, AsA-GSH cycle and
680 polyamine metabolism. Front in plant science 13: 993675 DOI: 10.3389/fpls.2022.993675.

681 Rasheed R, Ashraf M A, Ahmad S J N, Parveen N, Hussain I, Bashir R. 2022. Taurine regulates ROS
682 metabolism, osmotic adjustment, and nutrient uptake to lessen the effects of alkaline stress on *Trifolium*
683 *alexandrinum* L. plants. South african journal of botany 148: 482-498 DOI: 10.1016/j.sajb.2022.05.023.

684 Seekin B, Sekmen A H, Turkan I. 2009. An Enhancing effect of exogenous mannitol on the antioxidant enzyme

activities in roots of wheat under salt stress. Journal plant growth regulation 28: 12-20 DOI: 10.1007/s00344-008-9068-1.

Seleiman M F, Semida W M, Rady M M, Mohamed G F, Hemida K A, Alhammad B A, Hassan M M, Shami A. 2020. Sequential application of antioxidants rectifies ion imbalance and strengthens antioxidant systems in salt-stressed cucumber. Plants 9: 1783 DOI: 10.3390/plants9121783.

Shan C J, Liu R Q. 2017. Exogenous hydrogen peroxide up-regulates the contents of ascorbate and glutathione in the leaves of *Vigna radiata* (Linn.) Wilczek. exposed to salt stress. Brazilian journal botany 40: 583-589 DOI: 10.1007/s40415-016-0354-z.

Sharifi P, Amirnia R, Torkian M, Bidabadi S S. 2021. Protective role of exogenous selenium on salinity-stressed stachys byzantine plants. Journal of soil science and plant nutrition 21: 2660-2672 DOI: 10.1007/s42729-021-00554-5.

Singh A, Kumar A, Yadav S, Singh I K. 2019. Reactive oxygen species-mediated signaling during abiotic stress. Plant gene 18: 100173 DOI: 10.1016/j.plgene.2019.100173.

Sudhakar C, Veeranagamallaiah G, Nareshkumar A, Sudhakarbabu O, Sivakumar M, Pandurangaiah M, Kiranmai K, Lokesh U. 2015. Polyamine metabolism influences antioxidant defense mechanism in foxtail millet (*Setaria italica* L.) cultivars with different salinity tolerance. Plant cell reports 34: 141-156 DOI: 10.1007/s00299-014-1695-3.

Szabados L, Savoure A. 2010. Proline: a multifunctional amino acid. Trends in plant science 15: 89-97 DOI: 10.1016/j.tplants.2009.11.009.

Talubaghi M J, Daliri M S, Mazloun P, Rameeh V, Mousavi A. 2022. Effect of salt stress on growth, physiological and biochemical parameters and activities of antioxidative enzymes of rice cultivars. Cereal research communications 51:403-411 DOI: 10.1007/s42976-022-00314-w.

Tan Z M, Xuan Z Y, Wu C Y, Cheng Y X, Xu C Z, Ma X C, Wang D S. 2022. Effects of selenium on the AsA-GSH system and photosynthesis of pakchoi (*Brassica chinensis* L.) under lead stress. Journal of soil science and plant nutrition 22: 5111-5122 DOI: 10.1007/s42729-022-00987-6.

Tian T, Wang J, Wang H, Cui J, Shi X Y, Song J H, Li W D, Zhong M T, Qiu Y, Xu T. 2022. Nitrogen application alleviates salt stress by enhancing osmotic balance, ROS scavenging, and photosynthesis of rapeseed seedlings (*Brassica napus*). Plant signaling behavior 17: e2081419 DOI: 10.1080/15592324.2022.2081419.

Vaidyanathan H, Sivakumar P, Chakrabarty R, Thomas G. 2003. Scavenging of reactive oxygen species in NaCl stressed rice (*Oryza sativa* L.) differential response in salt-tolerant and sensitive varieties. Plant science 165: 1411-1418 DOI: 10.1016/j.plantsci.2003.08.005.

Verma K, Dixit S, Shekhawat G S, Alam A. 2015. Antioxidant activity of heme oxygenase 1 in *Brassica juncea* (L.) Czern.(Indian mustard) under salt stress. Turkish journal of biology 39: 540-549 DOI: 10.3906/biy-1501-28.

Wang S, Zhou H, Feng N, Xiang H T, Liu Y, Wang F, Li W, Feng S J, Liu M L, Zheng D F. 2022. Physiological response of soybean leaves to uniconazole under waterlogging stress at R1 stage. Journal of plant physiology 268: 153579 DOI: 10.1016/j.jplph.2021.153579.

Xie Y, Cui W, Yuan X, Shen W B, Yang Q. 2011. Heme oxygenase-1 is associated with wheat salinity acclimation by modulating reactive oxygen species homeostasis. Journal integrative plant biology 53: 653-670 DOI: 10.1111/j.1744-7909.2011.01052.x.

Yan F, Wei H, Ding Y, Li W W, Liu Z H, Chen L, Tang S, Ding C Q, Jiang Y, Li G H. 2021. Melatonin regulates

antioxidant strategy in response to continuous salt stress in rice seedlings. *Plant physiology and biochemistry* 165: 239-250 DOI: 10.1016/j.plaphy.2021.05.003.

Yu M L, Wu Q, Zheng D F, Feng N J, Liang X L, Liu M L, Li Y, Mou B M. 2021. Plant growth regulators enhance saline–alkali tolerance by upregulating the levels of antioxidants and osmolytes in soybean seedlings. *Journal of plant growth regulation* 41: 3218-3232 DOI: 10.1007/s00344-021-10507-y.

Zamocky M, Gasselhuber B, Furtmuller P G, Obinger C. 2012. Molecular evolution of hydrogen peroxide degrading enzymes. *Archives of biochemistry biophysics* 525: 131-144 DOI: 10.1016/j.abb.2012.01.017.

Zeng L, Shannon M C, Lesch S M. 2001. Timing of salinity stress affects rice growth and yield components. *Agricultural water management* 48(3): 191-206 DOI: 10.1016/S0378-3774(00)00146-3.

Zhang C P, Li Y C, Yuan F G, Hu S J, He P. 2012. Effects of hematin and carbon monoxide on the salinity stress responses of *Cassia obtusifolia* L. seeds and seedlings. *Plant and soil* 359: 85-105 DOI: 10.1007/s11104-012-1194-7.

Zhang J, Yang X, Ren Y, Yang B, Liu Z W, You B W, Zhang H X, Shen W B, Chen X P. 2016. β -Cyclodextrin–hemin enhances tolerance against salinity in tobacco seedlings by reestablishment of ion and redox homeostasis. *Plant growth regulation* 81: 533-542 DOI: 10.1007/s10725-016-0230-7.

Zhang W, Jiang B, Li W, Song H, Yu Y S, Chen J F. 2009. Polyamines enhance chilling tolerance of cucumber (*Cucumis sativus* L.) through modulating antioxidative system. *Scientia horticulturae* 122: 200-208 DOI: 10.1016/j.scienta.2009.10.001.

Zhao M, Meng Y, Wang Y, Sun G Y, Liu X M, Li J, Wei S, Gu W R. 2022. Exogenous Hemin alleviates cadmium stress in maize by enhancing sucrose and nitrogen metabolism and regulating endogenous hormones. *International journal of phytoremediation* 25:368-380 DOI: 10.1080/15226514.2022.2086212.

Zheng T R, Zhan J Y, Yang M, Wang M J, Sun W J, Shan Z, Chen H. 2021. Hemin-induced increase in saponin content contributes to the alleviation of osmotic and cold stress damage to *Conyza blinii* in a heme oxygenase 1-dependent manner. *Journal of Zhejiang university-science B* 22: 682-694 DOI: 10.1631/jzus.B2000697.

Zuo X, Dai J, Wu W, Jin J H, Ge W, Wang Y P, Ren L, Lin Y J, Pei Y Y, Xie H. 2022. Microfossil evidence of rice cultivation on the southeast China coast 7500 years ago. *Science china earth sciences* 65: 2115-2126 DOI: 10.3389/fpls.2016.00347.

Table 1(on next page)

Effects of exogenous Hemin on the morphological indexes of rice seedlings under NaCl stress

Notes: Data in this table is mean±standard error of at least three replicates. According to Duncan's multiple range tests, different letters indicate significant difference at the five percent significant level Within each column.

Table 1
Effects of exogenous Hemin on the morphological indexes of rice seedlings under NaCl stress

morphological indexes	Varieties	Treatments	NaCl stress time (d)			
			3	5	7	9
Plant height (cm)	HHZ	CK	31.67±0.20a	32.63±0.19a	33.57±0.07a	33.63±0.09a
		S	27.13±0.71c	27.93±0.13d	28.00±0.00d	29.10±0.21d
		SH	29.83±0.19b	30.97±0.17b	31.47±0.03b	31.90±0.15b
		SZ	27.00±1.00c	27.63±0.57d	27.90±0.57d	28.50±0.32d
		SZH	28.93±0.03b	29.57±0.03c	29.80±0.30c	29.97±0.09c
	XLY900	CK	32.90±0.31a	33.43±0.18a	33.80±0.46a	34.67±0.03a
		S	28.30±0.06d	29.40±0.12d	29.70±0.06d	30.97±0.27d
		SH	30.57±0.03b	30.97±0.23b	31.33±0.18b	33.17±0.33b
		SZ	28.33±0.03d	29.50±0.06d	29.50±0.06d	30.50±0.06d
		SZH	29.10±0.10c	30.25±0.14c	30.53±0.03c	32.00±0.00c
Stem diameter (mm)	HHZ	CK	3.27±0.06a	3.47±0.06a	3.63±0.06a	3.80±0.10a
		S	2.43±0.00d	2.67±0.12d	2.63±0.06d	2.70±0.12d
		SH	3.07±0.06b	3.20±0.00b	3.40±0.00b	3.57±0.06b
		SZ	2.63±0.15d	2.67±0.12d	2.73±0.06d	2.87±0.21d
		SZH	2.83±0.06c	2.93±0.06c	3.07±0.06c	3.20±0.00c
	XLY900	CK	3.53±0.03a	3.63±0.03a	3.77±0.03a	3.90±0.00a
		S	2.87±0.03d	3.00±0.00d	2.90±0.06d	3.00±0.00d
		SH	3.37±0.03b	3.47±0.09b	3.60±0.00b	3.63±0.03b
		SZ	2.93±0.07d	3.00±0.06d	2.93±0.03d	3.10±0.06d
		SZH	3.20±0.00c	3.23±0.03c	3.40±0.00c	3.47±0.03c

Notes: Data in this table is mean ± standard error of at least three replicates. According to Duncan's multiple range tests, different letters indicate significant difference at the five percent significant level Within each column.

15
16
17
18

Table 2 (on next page)

Effects of exogenous Hemin on the biomass of rice seedlings under NaCl stress

Notes: Data in this table is mean $\pm$ standard error of at least three replicates. According to Duncan's multiple range tests, different letters indicate significant difference at the five percent significant level Within each column.

1 **Table 2**

2 **Effects of exogenous Hemin on the biomass of rice seedlings under NaCl stress**

morphological indexes	Varieties	Treatments	NaCl stress time (d)			
			3	5	7	9
Shoot fresh weight (g)	HHZ	CK	0.4771±0.0060a	0.4742±0.0109a	0.4829±0.0009a	0.5147±0.0041a
		S	0.3225±0.0092d	0.3335±0.0025d	0.3467±0.0043d	0.3617±0.0007d
		SH	0.4109±0.0094b	0.3956±0.0029b	0.4270±0.0043b	0.4430±0.0029b
		SZ	0.3169±0.0032d	0.3399±0.0056d	0.3398±0.0024d	0.3610±0.0091d
		SZH	0.3557±0.0109c	0.3709±0.0040c	0.4058±0.0018c	0.4180±0.0068c
	XLY900	CK	0.5147±0.0021a	0.5110±0.0262a	0.5225±0.0053a	0.5477±0.0098a
		S	0.3566±0.0101d	0.3577±0.0068c	0.3718±0.0051c	0.3985±0.0014d
		SH	0.4385±0.0076b	0.4528±0.0195b	0.4546±0.0318b	0.4617±0.0160b
		SZ	0.3623±0.0033d	0.3602±0.0220c	0.3732±0.0132c	0.3923±0.0094d
		SZH	0.4034±0.0057c	0.4185±0.0088b	0.4260±0.0016b	0.4310±0.0025c
Shoot dry weight (g)	HHZ	CK	0.0938±0.0014a	0.0966±0.0022a	0.1004±0.0007a	0.1016±0.0015a
		S	0.0740±0.0023c	0.0761±0.0025c	0.0770±0.0007c	0.0783±0.0013d
		SH	0.0840±0.0012b	0.0852±0.0013b	0.0870±0.0015b	0.0907±0.0013b
		SZ	0.0725±0.0024c	0.0751±0.0004c	0.0767±0.0014c	0.0779±0.0011d
		SZH	0.0824±0.0006b	0.0831±0.0015b	0.0841±0.0012b	0.0846±0.0015c
	XLY900	CK	0.0933±0.0023a	0.1016±0.0017a	0.1044±0.0005a	0.1096±0.0045a
		S	0.0817±0.0017d	0.0825±0.0015d	0.0818±0.0008cd	0.0853±0.0000cd
		SH	0.0933±0.0006b	0.0943±0.0003b	0.0934±0.0029b	0.0962±0.0009b
		SZ	0.0804±0.0023d	0.0816±0.0009d	0.0802±0.0043d	0.0811±0.0010d
		SZH	0.0876±0.0006c	0.0871±0.0015c	0.0880±0.0000bc	0.0895±0.0006bc

3 Notes: Data in this table is mean ± standard error of at least three replicates. According to Duncan's
 4 multiple range tests, different letters indicate significant difference at the five percent significant
 5 level Within each column.

6
7

Figure 1

Figure 1. Effect of Hemin on growth of rice seedlings under NaCl (9 d) in HHZ (a) and XLY900 (b).




Figure 2

Figure 2. Effect of Hemin on membrane damage and ROS accumulation of rice seedlings under NaCl.

EL in HHZ (a) and XLY900 (b); MDA in HHZ (c) and XLY900 (d) and H₂O₂ in HHZ (e) and XLY900 (f). Values are the means±SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.




Figure 3

Figure 3. Effect of Hemin on histochemical localization of H_2O_2 and $O_2^{\cdot-}$ on rice leaves under NaCl stress (3 d).

H_2O_2 in HHZ (a) and XLY900 (b) and $O_2^{\cdot-}$ in HHZ (c) and XLY900 (d).




Figure 4

Figure 4. Effect of Hemin on SOD, POD, and CAT activity of rice seedlings under NaCl stress.

SOD in HHZ (a) and XLY900 (b); POD in HHZ (c) and XLY900 (d) and CAT in HHZ (e) and XLY900 (f). Values are the means $\pm$ SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.




Figure 5

Figure 5. Effect of Hemin on ascorbic acid content of rice seedlings under NaCl stress.

AsA in HHZ (a) and XLY900 (b); DHA in HHZ (c) and XLY900 (d) and AsA+DHA in HHZ (e) and XLY900 (f). Values are the means $\pm$ SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.




Figure 6

Figure 6. Effect of Hemin on glutathione content of rice seedlings under NaCl stress.

GSH in HHZ (a) and XLY900 (b); GSSG in HHZ (c) and XLY900 (d) and GSH+GSSG in HHZ (e) and XLY900 (f). Values are the means $\pm$ SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.




Figure 7

Figure 7. Effect of Hemin on key enzyme activities in the AsA-GSH defense system of rice seedlings under NaCl stress.

APX in HHZ (a) and XLY900 (b); MDHAR in HHZ (c) and XLY900 (d) DHAR in HHZ (e) and XLY900 (f) and GR in HHZ (g) and XLY900 (h). Values are the means $\pm$ SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.




Figure 8

Figure 8. Effect of Hemin on osmoregulatory substances of rice seedlings under NaCl stress.

Proline content in HHZ (a) and XLY900 (b); soluble protein content in HHZ (c) and XLY900 (d). Values are the means $\pm$ SD of three replicate samples. Different letters in the data column indicate significant differences ($p<0.05$) according to Duncan's test.

