

Marine introgressions and Andean uplift drives diversification in neotropical Monkey tree frogs (Anura, Phyllomedusinae) (#91653)

1

First submission

Guidance from your Editor

Please submit by **9 Nov 2023** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

4 Figure file(s)
2 Table file(s)
1 Raw data file(s)
2 Other file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Marine introgressions and Andean uplift drives diversification in neotropical Monkey tree frogs (Anura, Phyllomedusinae)

Diego Almeida-Silva^{1,2}, Leonardo Matheus Servino^{Corresp., 1, 3}, Matheus Pontes-Nogueira¹, Ricardo J Sawaya¹

¹ Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, São Bernardo do Campo, São Paulo, Brazil

² Miguel Lillo, Unidad Ejecutora Lillo, San Miguel de Tucumán, Tucumán, Argentina

³ Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil

Corresponding Author: Leonardo Matheus Servino

Email address: leonardo.servino@ib.usp.br

The species richness in Neotropics has been linked to environmental heterogeneity and a complex geological history. We evaluated which biogeographical processes were more associated with the diversification of Monkey tree frogs, an endemic clade from the Neotropics. We tested the hypothesis that the diversification of Phyllomedusinae occurred in a south-north direction in the Neotropics, and that marine introgressions and Andean uplift had a crucial role promoting their diversification. We used 13 molecular markers on a bayesian analysis to infer phylogenetic relationships among 57 species of Phyllomedusinae and to estimate their divergence times. We hypothesized the ancestral range based in 12 biogeographical units defined considering the distribution of the phyllomedusine species and potential biogeographical barriers. Ancestral range estimations were made by models implemented in BioGeoBEARS. We found that the Phyllomedusinae hypothetical ancestor range was probably widespread through the Neotropics, from Central America to Southern Atlantic Forest, at 38.6 Mya. Phyllomedusines' ancestors diverged mostly through vicariance during early stages of speciation, generally followed by jump-dispersals and speciation in sympatry. Dispersal among areas mostly occurred from Western Amazonia towards Northern Andes and the diagonal of dry landscapes, rejecting our south-north diversification hypothesis. Our results revealed a complex diversification of Monkey tree frogs, occurring simultaneously with the orogeny of Northern Andes and the South American marine introgressions in the last 30 million years.

Marine introgressions and Andean uplift drives diversification in neotropical Monkey tree frogs (Anura, Phyllomedusinae)

Diego Almeida-Silva^{1,2}, Leonardo Matheus Servino^{2,3}, Matheus Pontes-Nogueira², Ricardo J. Sawaya²

¹Miguel Lillo, Unidad Ejecutora Lillo, San Miguel de Tucumán, Tucumán, Argentina

²Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, São Bernardo do Campo, São Paulo, Brazil

³Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil

Corresponding Author:

Leonardo Matheus Servino

Universidade de São Paulo (USP), Rua do Matão, Travessa 14, n° 101 Cidade Universitária, São Paulo, CEP: 05508-090, São Paulo, Brasil.

Email address: leonardo.servino@ib.usp.br

ABSTRACT

The species richness in Neotropics has been linked to environmental heterogeneity and a complex geological history. We evaluated which biogeographical processes were more associated with the diversification of Monkey tree frogs, an endemic clade from the Neotropics. We tested the hypothesis that the diversification of Phyllomedusinae occurred in a south-north direction in the Neotropics, and that marine introgressions and Andean uplift had a crucial role promoting their diversification. We used 13 molecular markers on a bayesian analysis to infer phylogenetic relationships among 57 species of Phyllomedusinae and to estimate their divergence times. We hypothesized the ancestral range based in 12 biogeographical units defined considering the distribution of the phyllomedusine species and potential biogeographical barriers. Ancestral range estimations were made by models implemented in BioGeoBEARS. We

found that the Phyllomedusinae hypothetical ancestor range was probably widespread through the Neotropics, from Central America to Southern Atlantic Forest, at 38.6 Mya. Phyllomedusines' ancestors diverged mostly through vicariance during early stages of speciation, generally followed by jump-dispersals and speciation in sympatry. Dispersal among areas mostly occurred from Western Amazonia towards Northern Andes and the diagonal of dry landscapes, rejecting our south-north diversification hypothesis. Our results revealed a complex diversification of Monkey tree frogs, occurring simultaneously with the orogeny of Northern Andes and the South American marine introgressions in the last 30 million years.

Keywords. Amphibia - Northern Andes - Biogeography - Neotropics - Paranaense sea - Pebas system

INTRODUCTION

Extending from the central portion of Mexico through the entire Central and South America (Morrone, 2014), the Neotropical region hosts the greatest biodiversity on Earth (Myers *et al.*, 2000; Antonelli & Sanmartín, 2011). The environmental heterogeneity in Neotropics, in association with its complex geological history from the early-Cenozoic, have driven patterns of species diversification, contributing to high levels of species richness and endemism for different taxa (Antonelli, 2016). Regarding the northern portion of South America, well-documented marine introgressions occurred from mid to the late-Cenozoic (~25–5 million years ago – Mya), the so-called Pebas and Acre systems (e.g., Hoorn *et al.*, 2010; Salas-Gismondi *et al.*, 2015). Probably related to global sea-level fluctuations (Hoorn, 1993), both flooding processes turned

the Western Amazonia into a lacustrine environment during the Miocene (23–7 mya; Hoorn *et al.*, 2010; Salas-Gismondi *et al.*, 2015), affecting Magdalena River delta, paleo-Orinoco, and proto-Amazonas River basins. Henceforth, Western Amazonia suffered drastic changes until the emergence of current fluvial systems, including flowing changes of its main rivers (Díaz de Gamero, 1996; Albert *et al.*, 2018). Moreover, some orogenic processes also promoted important changes into Neotropics, as the accelerated uplift of Eastern Cordillera of Andes during the Miocene (Hoorn, 1993; ~10–4 mya; Gregory-Wodzicki, 2000), that led to changes in the climatic and sedimentary sources for Western Amazonia (Insel *et al.*, 2010; Poulsen *et al.*, 2010; Hoorn *et al.*, 2017).

The entire uplift of Andes has been playing similar role for the southwestern part of Neotropics, since it started in the early-mid Cenozoic (Giambiagi *et al.*, 2016; Rodríguez Tribaldos *et al.*, 2017; Sundell *et al.*, 2019). The Andean orogeny, in addition to climatic factors throughout the entire Neogene and Quaternary (Garzzone *et al.*, 2006; Hoorn *et al.*, 2020), has been responsible for the emergence of the South American diagonal of open/dry landscapes (DODL; Zanella, 2011; Azevedo *et al.*, 2020), especially during the Oligocene (~25 mya) and late Neogene (~5–3 mya). As DODL expanded, a single large forest block must have been splitted into Amazonian and Atlantic forests (Costa, 2003; Sobral-Souza *et al.*, 2015; Peres *et al.*, 2017), the last being southern confined by marine introgressions (Hernández *et al.*, 2005; Abello *et al.*, 2010).

Most studies examining the processes leading to biota diversification in the Neotropics focus on ecologically or geographically restricted groups (Smith *et al.*, 2014; Werneck *et al.*, 2015; e.g., Guarnizo *et al.*, 2016), in a local-scale approach. Studies focusing on widespread clades, on the other hand, could elucidate the role of multiple processes over space and deep

time, contributing to the macroevolutionary framework (e.g., Vicente *et al.*, 2017; Hamdan *et al.*, 2019; Prieto-Torres *et al.*, 2019; Pontes-Nogueira *et al.*, 2021), given the complex dynamic of biogeographical processes in the Neotropical region. Among anurans, this scenario fits well for Phyllomedusinae, a subfamily virtually occurring throughout the entire Neotropics. This charismatic group, commonly known as Monkey tree frogs, comprises 67 species (Frost, 2023), distributed from Argentina to Mexico (Duellman *et al.*, 2016; Frost, 2023).

Systematics of the subfamily seems to be well defined, regarding closely related frog groups. Except for phylogenetic analyses exclusively based on morphology (Haas, 2003; Wiens *et al.*, 2005), phyllomedusines are consistently recovered as monophyletic, and a sister taxon to Pelodyadinae from the Australo-Papuan region, both constituting subfamilies of Hylidae treefrogs (Wiens *et al.*, 2005; Frost *et al.*, 2006; Faivovich *et al.*, 2010; Pyron & Wiens, 2011; Duellman *et al.*, 2016; Jetz & Pyron, 2018; Dubois *et al.*, 2021). Phylogenetic relationships for some clades of Phyllomedusinae are also consistent in the most comprehensive phylogenetic approaches (Faivovich *et al.*, 2010; Pyron & Wiens, 2011). Some discussion occurs regarding the early branching events in the group, since molecular approaches on phyllomedusine's phylogeny show a low sampling for *Phrynomedusa* Miranda-Ribeiro, 1923, a rare genus only known from a few localities of the Serra do Mar and Serra da Mantiqueira on the South American Atlantic Forest (Baêta *et al.*, 2016). On the other hand, Pelodyadinae is widespread in Australia and Papua New Guinea (Frost, 2023). More speciose than Phyllomedusinae, the pelodyadine frog subfamily is a clade composed by 223 species, whose systematics is object of a long-term debate regarding its internal clades relationships (Dubois *et al.*, 2021; Frost, 2023).

Molecular estimates indicated that the split between phyllomedusines and pelodyadines occurred during the late Paleocene (~55 Ma), a time when both the Neotropics and Australo-

Papuan regions were connected to Antarctica via a land bridge (Duellman *et al.*, 2016; Van Den Ende *et al.*, 2017). Such estimates also suggest the Most Recent Common Ancestor (MRCA) of Phyllomedusinae emerging during the late Eocene, following the appearance of the MRCA of Pelodryadinae. Diversification within Phyllomedusinae started in the Oligocene (Duellman *et al.*, 2016; Jetz & Pyron, 2018) as South America began separating from Antarctica. Considering that some lineages diversified in South America after their ancestral lineages arrived via Antarctica (e.g., orchids, Givnish *et al.*, 2016; sea spiders, Dietz *et al.*, 2019; ungulate mammals, Reguero *et al.*, 2014), it is plausible to hypothesize that the MRCA of all phyllomedusines also diversified in a south-north direction. Furthermore, certain speciation processes within phyllomedusines have tentatively been associated with the uplift of the Eastern Cordillera of the Andes (e.g., Duellman *et al.*, 2016; Ron *et al.*, 2013). However, the historical biogeography of this clade has not yet been explored in a statistical framework.

Herein, we combined sequences of multiple molecular markers of 53 species of Phyllomedusinae to produce a time-calibrated phylogeny. Then we reconstructed the subfamily diversification throughout Neotropical region, aiming to answer the following questions: (1) How were the hypothetical ancestors of Phyllomedusinae clade distributed along Neotropics? And (2) which biogeographical processes drove the current subfamily distribution? We tested the hypothesis that speciation in Phyllomedusinae occurred in a south-north direction in South America, and that diversification was associated to marine introgressions and Andean orogeny.

MATERIALS AND METHODS

Sequence data and phylogenetic analyses

We performed a phylogenetic inference using sequences retrieved from GenBank (Supplementary data S1). The molecular data available for the group warrants careful consideration. In the last decade, taxonomic reviews, phylogenies, and the description of new species based on molecular data have led to frequent species synonymy and the recognition of populations as new species within phylomedusines (e.g. Faivovich *et al.*, 2011; Baêta *et al.*, 2016; Castroviejo-Fisher *et al.*, 2017, Pereira *et al.*, 2018; Andrade *et al.*, 2020). However, GenBank curation has not been able to keep up to date with the taxonomy of the group. To avoid problems with misidentification and taxonomic inconsistencies, we took special care when selecting the sequences for Phyllomedusinae, considering the collection site of each accession number and cross-validating the species distribution following Frost (2023).

Our analysis included 57 species of Phyllomedusinae in the ingroup, in addition to 20 Pelodryadinae and 18 Hylinae, both subfamilies composing the outgroup. Molecular sampling covered 85% of all known species of Phyllomedusinae, including all genera of the subfamily (Frost, 2023). Considering the data availability in GenBank, we included as much data as possible in our analysis, excluding only the sequences that showed no significant similarity due to the absence of query coverage (BLAST utility; Zhang *et al.*, 2000). Our analysis comprised all the species of *Callimedusa* Duellman, Marion, and Hedges, 2016 (6 spp.), *Cruziohyla* Faivovich, Haddad, Garcia, Frost, Campbell, and Wheeler, 2005 (3 spp.), *Hylomantis* (2 spp.), and *Pithecopus* Cope, 1866 (12 spp.), as well as a representative selection of *Agalychnis* (9 species sampled from 14 spp. described), *Phasmahyla* Cruz, 1991 (7 species sampled from 8 spp. described), *Phyllomedusa* Wagler, 1830 (15 species sampled from 16 spp. described), and *Phrynomedusa* (3 species sampled from 6 spp. described). We searched for 13 molecular

markers, both nuclear (CXCR4, POMC, RAG1, RHOD, SIAH, and Tyr) and mitochondrial genes (12S, tRNA-Val, 16S, tRNA-Leu, ND1, tRNA-Ile, and CytB) genes.

The amount of missing information ($\bar{x} = 33\%$, ranging from 6% to 92%; not accounting for gaps) **does not seem alarming**, considering that the two best-represented genes in our analyses (12S/16S) provided a strong backbone for placing most species, as stated by Pyron & Wiens (2011). In fact, 81% of the species had complete data for the 12S gene partition ($\bar{x} = 12\%$ missing data), while the 16S partition was fully represented for 79% of the species ($\bar{x} = 9\%$ missing data), and all species were represented in at least one of them. Previous studies in the literature have supported this sample design for conducting model-based phylogenetic analyses, both theoretically and empirically (e.g., Wiens, 2003; Driskell *et al.*, 2004; Thomson & Shaffer, 2010; Wiens & Morrill, 2011), yielding taxonomically highly congruent and well-supported results (for a detailed discussion, see Pyron & Wiens, 2011). We used *MAFFT* by EMBL-EBI web toolkit (Li *et al.*, 2015) for aligning our sequences. For nuclear markers, we employed the G-INS-i strategy, assuming global homology and aligning the entire region. For mitochondrial fragments, we used the E-INS-i algorithm, which is recommended for sequences with conserved domains and a high number of gaps. We concatenated all genes into a single matrix (7,290 bp, 95 terminals) using *SequenceMatrix* v.1.7 (Vaidya *et al.*, 2011; Supplementary data S2 and S3). To determine the appropriate nucleotide substitution models, we conducted a search using *PartitionFinder* (Lanfear *et al.*, 2017; 38 partitions; Supplementary data S1 and S2), under greedy algorithm.

For the phylogenetic analysis, we conducted a Bayesian inference concurrent with node dating using the software *BEAST* v2.7.4 (Bouckaert *et al.*, 2019). We performed two independent Markov chain Monte Carlo (MCMC) simulations with a chain length of 200,000,000 generations

and a pre-burn-in of 20% using the *CIPRES Science Gateway* (Miller *et al.*, 2010). We linked the partitions into one phylogenetic species tree and incorporated clock models specific to each gene. Due to the limited fossil record in Hylidae, we followed the approach of Hime *et al.* (2021) and calibrated our node dating using a fossil-calibrated phylogenomic tree. Specifically, we set a prior for the split between Phyllomedusinae and Pelodryadinae to the late Paleocene to early Eocene period, approximately 44.5 million years ago (95% HPD: 39.3–49.5 Mya). Unfortunately, both phyllomedusines and pelodryadines are underrepresented in Hime's study ($n = 2$ for both subfamilies). Given the low taxon-sampling represent a problem to the divergence time estimation (Soares & Schrago, 2012), we also incorporated molecular clock models based on substitution rates reported in the literature for a nuclear (POMC: 0.0043/site/Myr; de Magalhães *et al.*, 2017) and a mitochondrial gene (CytB: 0.0161/site/Myr; Stöck *et al.*, 2012). Thus, we performed our search using optimised relaxed clock models (Douglas *et al.*, 2021), with a substitution rate of 0.0161/site/Myr for CytB and 0.0043/site/Myr for POMC (Stöck *et al.*, 2012; Magalhães *et al.*, 2017). We estimated the mean clock rates for all the other partitions under weak priors ($1/x$ distribution). We inferred the species tree using a Yule model prior, while keeping all other priors at their default values. We assessed the convergence of the MCMC chains by examining the estimated sample size ($ESS > 200$) and checking for model parameter stationarity using *TRACER* 1.7 (Rambaut *et al.*, 2018). We discarded the initial 25% of each chain as burn-in and summarized the output as a maximum clade credibility (MCC) tree using mean node heights in *TreeAnnotator* v.2.6.2 (Bouckaert *et al.*, 2014). We pruned the MCC tree using the *ape* R package (Paradis *et al.*, 2004), retaining only the species of Phyllomedusinae and Pelodryadinae for subsequent ancestral geographical range estimation.

Geographical distribution data

Our geographical dataset consists of georeferenced points obtained from the *Global Biodiversity Information Facility* (GBIF, 2023; <https://www.gbif.org/>), which is the largest online source of distributional records (Zizka *et al.*, 2020). The geographical points were obtained using the R software (R Core Team, 2023) package *rgbif* (Chamberlain & Boettiger, 2017), and the records were later treated in the software to eliminate duplicates and remove uncertain records.

Study area and regionalization

Multiple regionalization schemes have been proposed in the literature (e.g., Olson *et al.*, 2001; Morrone, 2006, 2014; Dinerstein *et al.*, 2017; Escalante, 2017). Our regionalization scheme primarily relies on the terrestrial ecoregions of the world, which were defined based on floristic maps and vegetational types (Olson *et al.*, 2001; Dinerstein *et al.*, 2017). Studies focusing on neotropical species often involve a considerable number of biogeographical units due to the landscape heterogeneity of the Neotropical region (see Carneiro *et al.*, 2018; Réjaud *et al.*, 2020; Pontes-Nogueira *et al.*, 2021). Thereafter, we defined 12 biogeographical units (Fig. 1A) based on relevant landscape modifications that could have influenced Phyllomedusinae diversification (Fig. 1B), such as the uplift of mountain ranges (e.g., Cordillera of Andes), riverine barriers (e.g., Amazonas and Madeira rivers) and phytophysognomic differences (e.g., DODL). Given that the Neotropics were once connected to Oceania through an Antarctic land bridge, we included the Australo-Papuan Pelodyadinae subfamily (the sister clade to the

Phyllomedusinae subfamily) sampled in our phylogeny for estimating the ancestral geographical range.

In order to refine the biogeographical results regarding the MRCA of Phyllomedusinae + Pelodryadinae, we also included Oceania in our regionalization scheme. Therefore, our regionalization scheme encompasses the following regions (Fig. 1A): Central America (A, being southern limited by Chocó Department and Pacific Coast of Colombia; its connectivity southwards having been enhanced over time due to the formation of Isthmus of Panama), Northern Andes (B, encompassing Western, Central and Eastern Cordilleras of Northern Andes; it becomes a geographical barrier through the Miocene, due to the acceleration on the uplift of Eastern Cordillera), Western (C), Eastern (D), and Southern Amazonia (E; limited by Amazonas and Madeira rivers; the three areas were differently affected by marine introgressions occurred along the Miocene), Caatinga (F; reducing connectivity between forested areas as the DODL expanded), Cerrado (G; reducing connectivity between forested areas as the DODL expanded), Northern (H), Central (I), and Southern Atlantic Forest (J; divided by Serra do Mar and Mantiqueira Mountain Ranges; the three areas being differently affected by the late uplift of both mountain ranges), Chaco/Pampas (K, encompassing Chaco, Pantanal, and the Uruguayan savanna, northern limited by Araucaria moist forests; reducing connectivity between forested areas as the DODL expanded), and Oceania (L, comprising the whole New Guinea island, the Wallacea region and Australia; continent with a complex history of connectivity with South America through Antarctica over geological time).

We used the R package *SpeciesGeoCoder* (Töpel *et al.*, 2017) to code the presence of all species in each biogeographical unit using a threshold of 5%. This means that for a species to be considered present in a unit, its distribution in that unit must be higher than 5% of all the

distributional records for that species. The result was later revised using Frost's (2023) description of the species distribution.

Ancestral geographical range estimation

Ancestral range estimation is performed based on the current distribution of sampled species and their phylogenetic relationships (Sanmartín, 2016). Several models for ancestral range estimation have been proposed in the literature, with the Dispersal-Vicariance Analysis (DIVA; Ronquist, 1997), the Dispersal-Extinction-Cladogenesis (DEC; Ree *et al.*, 2005; Ree & Smith, 2008), and the BayArea model (Landis *et al.*, 2013) being the most widely employed. These models have been incorporated into the *BioGeoBEARS* R package (Matzke, 2013, 2014) which provides a unified Maximum Likelihood (ML) environment for biogeographical analysis. This allows for the use of parameters that control biogeographical processes and model testing, eliminating the need for arbitrary model selection. In our study, we implemented eight models in *BioGeoBEARS*, all of them being variations of DEC and DIVALIKE (the ML version of the original DIVA included in *BioGeoBEARS*). We chose not to use BAYAREALIKE (the ML version of the original BayArea included in *BioGeoBEARS*) because it does not consider vicariance in its estimation (Matzke, 2013). Some models considered time-stratified dispersal matrices (TS), which are multipliers based on the landscape evolution of the study region (see Fig. 1C, and Supplementary data S4 and Table S4.1 for details). The values in the TS matrices restrict the probabilities of dispersal between geographical units, ranging from 0 (when a geographical barrier completely prevents dispersal) to 1 (when there are no dispersal limitations between units). To weigh the relative significance of the TS matrices, we also included the free parameter w , which acts as an exponent on the matrices (see Dupin *et al.*, 2017). Additionally, to

account for the colonization of novel biogeographical areas at the time of cladogenesis (Matzke, 2014, 2022; Klaus & Matzke, 2020), we included the parameter j . We set the maximum range size to 5, which corresponds to the number of areas occupied by the most widespread species in our clade. We compared all the models using AIC (Akaike Information Criterion) and calculated Akaike weights (Akaike, 1974; Burnham & Anderson, 2004; Wagenmakers & Farrell, 2004).

RESULTS

Phylogeny and divergence time estimation

Bayesian inference recovered all the genera in our sample with high posterior probabilities (PP = 1.00 for all genera; Fig. 2), both in Phyllomedusinae and Pelodryadinae. We recovered *Cruziohyla* as the sister clade to the other Phyllomedusinae genera (PP = 1.00; Fig. 2), with the MRCA of the genus dating from 16.2 Mya (HPD 95%: 4.4–30.6 Mya). *Phrynomedusa*, the next diverging lineage (PP = 0.59; Fig. 2), exhibited an MRCA that diversified from 17.8 Mya (HPD 95%: 8.0–28.7 Mya). We found *Agalychnis* Cope, 1864 as the sister to *Hylomantis* Peters, 1873 (PP = 1.00; Fig. 2), a clade age estimated to be 29.8 Mya (HPD 95%: 24.3–35.7 Mya). Regarding the core of Phyllomedusinae (i.e., MRCA of *Callimedusa*, *Phasmahyla*, *Phyllomedusa*, and *Pithecopus*; PP = 1.00; Fig. 2), a clade mainly diversified in forested areas, our estimation suggests an age of 32.4 Mya (HPD 95%: 27.1–38.0 Mya). We recovered *Phasmahyla* as sister group to the other three genera, being the MRCA of the clade *Phyllomedusa* (*Callimedusa* + *Pithecopus*) (PP = 1.00; Fig. 2) estimated to be 27.1 Mya (HPD 95%: 22.5–32.0 Mya).

Ancestral geographical range estimation

The best-fitted model in our analysis was DECTS+*j* (Table 1; AIC = 406.9; AICw ~ 1.00), incorporating landscape evolution in the Neotropics and jump dispersal processes (Fig. 3; see Supplementary data S5 for details). We found the divergence of Phyllomedusinae + Pelodryadinae (52.4 Mya; HPD 95%: 46.6–58.6 Mya) occurred through vicariance, with the MRCA of Pelodryadinae subsequently dispersing through Oceania (unit L; Figs. 3, 4a; Supplementary data S5). Meanwhile, the MRCA of Phyllomedusinae exhibited a wide distribution across the Neotropics, encompassing units ABHI (Figs. 3, 4a).

The earliest diversification event within the clade occurred when sympatric populations of the MRCA gave rise to *Cruziohyla* in Central America (unit A, 38.6 Mya; HPD 95%: 32.7–45.0 Mya). Subsequently, a vicariant event isolated populations in the Central Atlantic Forest at 36.9 Mya (HPD 95%: 31.3–43.0 Mya), leading to the origin of the *Phrynomedusa* genus (unit I; Figs. 3, 4b; Supplementary data S5). Within the ABH range, ancestral populations underwent another vicariant process at 29.8 Mya (HPD 95%: 24.3–35.7 Mya; Figs. 4c), resulting in the emergence of *Hylomantis* in the Northern Atlantic Forest (unit H) with species diversifying in sympatry, and *Agalychnis* in Central America and the Northern Andes (units AB). The MRCA of *Agalychnis* underwent an early vicariant process (27.4 Mya; HPD 95%: 21.9–33.1 Mya; Fig. 4d), resulting in the divergence of the Andean species *A. hulli* Duellman and Mendelson, 1995 (unit B) from other species (unit A). Subsequently, the Central American species of *Agalychnis* began to diversify primarily in sympatry in the last 24 million years.

Table 1. AIC comparisons of the Ancestral range estimation models.

Models	LnL*	n*	d*	e*	j*	w*	AIC	AIC weights
--------	------	----	----	----	----	----	-----	-------------

DECTS+j**	-200.5	3	0.0072	<0.001	0.051	1	406.9	~1
DECTS	-208	2	0.0086	<0.001	0	1	420	0.0015
DIVALIKETS+j	-210.7	3	0.0079	<0.001	0.052	1	427.3	<0.001
DECTS+j+w	-215.7	4	0.004	<0.001	0.017	0.0067	439.4	<0.001
DIVALIKETS	-218.4	2	0.011	<0.001	0	1	440.7	<0.001
DECTS+w	-220.6	3	0.0048	<0.001	0	0.0031	447.2	<0.001
DIVALIKETS+j+w	-225.6	4	0.0048	<0.001	0.018	0.0049	459.3	<0.001
DIVALIKETS+w	-230	3	0.0068	<0.001	0	0.083	466.1	<0.001
DEC+j	-236.9	3	0.0025	<0.001	0.0087	1	479.9	<0.001
DEC	-240.3	2	0.0029	<0.001	0	1	484.7	<0.001
DIVALIKE+j	-244.4	3	0.0028	<0.001	0.012	1	494.8	<0.001
DIVALIKE	-250.3	2	0.0037	<0.001	0	1	504.7	<0.001

*LnL = log-likelihood of the model. n = number of free parameters in the model (that being d , e , j and w); d = rate of range

expansion (i.e. anagenetic dispersal); e = rate of range contraction (i.e. extinction); j = jump dispersal process; and w = dispersal

multiplier parameter (for TS models).

**DECTS+j is shown in bold and represents the best model under AIC and AIC weights.

On the other hand, the MRCA of *Phasmahyla* + (*Phyllomedusa* (*Callimedusa* + *Pithecopus*)) colonised the Central Atlantic Forest (unit I; 32.4 Mya; HPD 95%: 27.1–38.0; Fig. 4b) through a jump dispersal event from the ancestral range ABH. Sympatric populations in the Central Atlantic Forest gave rise to the genus *Phasmahyla* (unit I). This genus displayed two distinct clades: one diversified through jump dispersal to the Northern Atlantic Forest and Cerrado (units G and H) in the past 7 Mya, while the other clade began diversifying through dispersal within the Central Atlantic Forest (unit I) and subsequent jump dispersal to the Northern Atlantic Forest approximately 5 Mya. Additionally, ancestral populations from the Central Atlantic Forest (unit I) underwent jump dispersal to Western Amazonia (unit C) at 27.1

311 Mya (HPD 95%: 22.5–32.0 Mya; Fig. 4b), leading to the MRCA of (*Phyllomedusa* (*Callimedusa*
312 + *Pithecopus*)) in sympatry.

313 We identified at least three major diversification patterns within the *Phyllomedusa* genus
314 (Fig. 3). Firstly, the clade comprising *Ph. vaillanti* Boulenger, 1882 and *Ph. bicolor* Boddaert,
315 1772 originated at 13.2 Mya (HPD 95%: 8.4–18.1 Mya) through sympatric speciation within
316 Western Amazonia, followed by subsequent dispersal throughout Amazonia and the Northern
317 Andes. Secondly, the *Ph. tarsius* Cope, 1868 group experienced jump dispersal events, initially
318 colonizing the Northern Andes at 6.1 Mya (HPD 95%: 2.7–9.9 Mya) and later moving to Eastern
319 Amazonia from 4.5 Mya (HPD 95%: 1.8–7.7 Mya). Species within this group then dispersed to
320 Southern Amazonia (*Ph. camba* De la Riva, 1999; units CE) and the Northern Andes (*Ph.*
321 *tarsius*; units BCD). Thirdly, the *Ph. burmeisteri* Boulenger, 1882 group underwent jump
322 dispersal events, colonizing Chaco (unit K), followed by subsequent dispersal to the Northern
323 Atlantic Forest (unit H) and Southern Atlantic Forest (unit J) between 16.2 (HPD 95%: 12.0–
324 20.7 Mya; Fig. 4e) to 7.6 (HPD 95%: 5.3–10.1 Mya; Fig. 4e) million years ago. Species from
325 this group further dispersed to Chaco (*Ph. iheringii* Boulenger, 1885; units JK), Caatinga (*Ph.*
326 *bahiana* Lutz, 1925; units FH), Central Atlantic Forest (*Ph. tetraploidea* Pombal and Haddad,
327 1992 and *Ph. distincta* Lutz, 1950; units IJ), Central America (*Ph. venusta* Duellman and Trueb,
328 1967; units AB), and Cerrado (*Ph. burmeisteri*; units GHII) within the last 2 million years.

329 The MRCA of *Pithecopus* and *Callimedusa* remained in Western Amazonia (unit C) at
330 23.6 Mya (HPD 95%: 19.1–28.2 Mya). Both genera exhibited distinct patterns of diversification.
331 *Pithecopus* displayed a colonization pattern in the Cerrado region (unit G) through two separate
332 jump dispersal events. The first jump dispersal to the Cerrado occurred at 15.7 Mya (HPD 95%:
333 12.2–19.3 Mya; Fig. 4f), leading to a clade that diversified through sympatric speciation in

Cerrado and additional jump dispersals to the Atlantic Forest. The second jump dispersal to the Cerrado took place around 14.2 Mya (HPD 95%: 10.9–17.8 Mya; Fig. 4f), resulting in a clade primarily diversifying through sympatric speciation in Cerrado and range expansions (i.e., anagenetic dispersals) to the Atlantic Forest and Chaco. Additionally, *Pi. hypochondrialis* Daudin, 1800 reached the Northern Andes and Eastern/Southern Amazonia. However, *Pi. gonzagai* Andrade, Haga, Ferreira, Recco-Pimentel, Toledo, and Bruschi, 2020, a species within this clade, originated from a jump dispersal event from the Cerrado to the Northern Atlantic Forest and Caatinga at 5.3 Mya (HPD 95%: 1.6–9.1 Mya). On the other hand, the *Callimedusa* genus had an early sympatric speciation that gave rise to *C. tomopterna* Cope, 1868. This species expanded its range through the entire Amazonia region (units CDE) at 18.6 Mya (HPD 95%: 13.4–24.0 Mya) and colonised the Northern Andes (unit B). Subsequently, another sympatric speciation occurred within Western Amazonia around 14.2 Mya (HPD 95%: 8.2–20.3 Mya; Fig. 4f). This cladogenesis resulted in the origin of *C. atelopoides* Duellman, Cadle, and Cannatella, 1988 and the MRCA of the other *Callimedusa* species, who reached the Northern Andes by jump dispersal, afterward diversifying in sympatry since 10.4 Mya (HPD 95%: 5.1–15.8 Mya; Fig. 4f).

DISCUSSION

In the present study, we provided a detailed analysis of the diversification and colonisation history of Monkey tree frogs across the Neotropics. By sampling 85% of the formally described phyllomedusine species, using mitochondrial and nuclear markers, our results represent a robust framework to discuss the processes concerning the biogeographical history of the group. Regarding the phylogenetic relationships among phyllomedusine genera, our topology is mostly congruent with previous studies in literature based on molecular data (e.g., Faivovich *et*

al., 2005; 2010; Pyron & Wiens, 2011; Rivera-Correa *et al.*, 2013; Duellman *et al.*, 2016; Jetz & Pyron, 2018). Furthermore, our results on node dating and ancestral geographical range estimation indicates the diversification of phyllomedusines was markedly influenced by environmental changes resulting from the Miocene marine introgressions and Andean orogeny. **In detail**, most lineages in Phyllomedusinae diversified in a dynamic scenario that occurred in Western Amazonia, which was limited by the Pebas System and the Paranaense sea (north/westward and southward, respectively) during the Miocene.

The most evident divergence recovered in phylogenetic inference was the relative position of *Cruziophyla* and *Phrynomedusa* genera. Previous studies usually recover *Phrynomedusa* as the first branching lineage in Phyllomedusinae (Faivovich *et al.*, 2005; 2010; Pyron & Wiens, 2011) or composing a clade *Cruziophyla* + *Phrynomedusa* (Duellman *et al.*, 2016; Jetz & Pyron, 2018). Since the split between the lineage that will give rise to *Cruziophyla* and the other phyllomedusines occurred at 38.6 Mya (HPD 95%: 32.7–45.0 Mya), our results imply a first conquest of the northern Neotropics between the late Eocene and middle Oligocene. At the genus level, the topology we found for *Phasmahyla* deviates from the previous studies, but none of them coincide (Faivovich *et al.*, 2005; 2010; Pyron & Wiens, 2011; Duellman *et al.*, 2016; Jetz & Pyron, 2018; Pereira *et al.*, 2018), and therefore the phylogeny of the genus remains under debate. Also, some inconsistencies regarding *Phasmahyla* argue for a careful taxonomic review. *Phasmahyla cruzi* Carvalho-e-Silva, Silva, and Carvalho-e-Silva, 2009, for instance, is assumed to be known only from its type locality (Rio das Pedras Reserve, Municipality of Mangaratiba, state of Rio de Janeiro, Brazil; Frost, 2023). However, there are no molecular data on Genbank from this population, and the species was represented by a specimen assigned to other locality (Picinguaba, municipality of Ubatuba, state of São Paulo, Brazil; Pyron & Wiens,

2011; Faivovich *et al.*, 2005; 2010; Duellman *et al.*, 2016; Jetz & Pyron, 2018; Pereira *et al.*, 2018; present study, Supplementary data S1). We found other occasional divergences regarding the relative position of some species in *Pithecopus* and *Phyllomedusa*, compared to other studies (Faivovich *et al.*, 2010; Pyron & Wiens, 2011; Duellman *et al.*, 2016; Jetz & Pyron, 2018; Pereira *et al.*, 2018).

Much of the current information provided on the node dating of phyllomedusine' frogs come from studies focusing on large-scale approaches, but with a low representativeness for the clade. Phylogenomic studies using similar sets of fossil calibrations point to a split between Phyllomedusinae and Pelodryadinae occurring at 47.4 Mya (Feng *et al.*, 2017) or 44.5 Mya (Hime *et al.*, 2020; Portik *et al.*, 2023), implying in an early diversification of phyllomedusines at 24.5 Mya (CI 95%: 20.0–29.7 Mya; Feng *et al.*, 2017), 23.2 Mya (HPD 95%: 18.4–28.1 Mya; Hime *et al.*, 2020) or 27.5 Mya (HPD 95%: 24.0–35.8 Mya; Portik *et al.*, 2023). The divergence time estimation of the clade may be taken carefully, given the low representativeness of phyllomedusines in some of these previous works (Feng *et al.*, 2017; Hime *et al.*, 2020) and the recognised impact of taxon-sampling on time estimations under bayesian frameworks (Soares & Schrago, 2012; Matschiner, 2019; Luo *et al.*, 2023). Our results set both the split phyllomedusine-pelodryadine and the early onset of diversification in Phyllomedusidae to the past, occurring at 52.4 Mya (HPD 95%: 46.6–58.6 Mya) and 38.6 Mya (HPD 95%: 32.7–45.0 Mya), respectively. In other words, our findings assign the phyllomedusines the status of a clade that dates from the late-Eocene/early-Oligocene rather than the late-Oligocene. Similar results were obtained previously by a maximum likelihood approach performed using an extensive sampling of the subfamily (Duellman *et al.*, 2016), suggesting the split phyllomedusine-

pelodryadine to be 61.8 Mya (HPD 95%: 57.5–66.1 Mya) and the Phyllomedusinae dated to 33.3 Mya (HPD 95%: 29.0–37.6 Mya).

The split we found for the MRCA of Phyllomedusinae + Pelodryadinae occurred by vicariance (Figures 3 and 4A; Supplementary data S5). At the early Eocene, our results suggest that the MRCA of Phyllomedusinae + Pelodryadinae was widely distributed throughout South America, Oceania, and, therefore, Antarctica (Fig. 4A). The diversification between the two subfamilies during the passage from late-Eocene to early-Oligocene occurred concurrently to the convoluted process of separation between the three continents, as proposed by Duellman et al. (2016). Hence, the vicariance associated was probably caused by landmasses movement, promoting the initial divergence of pelodryadines and phyllomedusines. Moreover, our results emphasize much of the early diversification of the Phyllomedusinae was influenced by vicariances from this widespread Neotropical ancestor (e.g., MRCA of the clade *Hylomantis* + *Agalychnis*, diversification of *Phrynomedusa* and *Agalychnis* genera; Figure 3 and 4B-D). Hence, our results reject the “south-north” diversification hypothesis of the Phyllomedusinae subfamily, as the diversification within the group occurred from Western Amazonia towards Northern Andes and the diagonal of dry landscapes. Our findings show a different scenario from Duellman et al. (2016) proposal, where the authors argued the split between the two subfamilies may took place within the Neotropics, with “protopelodryadines” dispersing to Australia afterward. Since our sampling focused in Phyllomedusinae, it is difficult to extend the discussion to the context of the whole Hylidae family. We encourage future biogeographical studies to examine the question in more detail.

Our results on the DECTS+*j* model suggest that Western Amazonia (unit C) acted as a species pump for *Phyllomedusa*, *Pithecopus* and *Callimedusa* genera. This area is frequently

recovered for ancestral ranges in several animal groups (e.g., lizards, Prates *et al.*, 2017; snakes, Dal Vechio *et al.*, 2018; Pontes-Nogueira *et al.*, 2021). Western Amazonia was recovered as the ancestral area for the clade *Phyllomedusa* (*Callimedusa* + *Pithecopus*) after a jump dispersal in the split between *Phasmahyla* and the clade *Phyllomedusa* (*Callimedusa* + *Pithecopus*). This scenario took place from the early to the middle Miocene, concomitantly to the occurrence of lacustrine conditions due to the Pebas system (Hoorn *et al.*, 2010, 2017; Jaramillo *et al.*, 2010; Figs. 4e, 4f), and virtually affecting the entire Western Amazonia. Previous studies also emphasise the influence of Pebas system over the biogeography of neotropical anuran fauna, driving their evolutionary history in different ways (Carvalho, 1954; Zimmermann and Zimmermann, 1988; Fouquet *et al.*, 2012; 2021a; 2022; Réjaud *et al.*, 2020). The absence of an overall pattern in Anura seems to be related to the diversity of natural history traits. Hence, Pebas system turned Western Amazonia an unsuitable environment for ground-dweller frogs, negatively affecting the diversification of terrestrial (Phyzelaphryninae and *Allobates* Zimmermann and Zimmermann, 1988; Fouquet *et al.*, 2012; Réjaud *et al.*, 2020) and burrowing (*Synapturanus* Carvalho, 1954; Fouquet *et al.*, 2021a) clades. On the other hand, the marine incursion was crucial for the origin and diversification of aquatic clades (*Pipa* Linnaeus, 1758; Fouquet *et al.*, 2022). Here, we found evidence of Pebas system promoting the diversification of arboreal anurans, given that the most speciose clade in Phyllomedusinae had origin and diversification in this area. This is an important finding, since other studies regarding arboreal frogs have shown diversification after the regression of the Pebas system (e.g., *Boana albopunctata* Spix, 1824 species group; Fouquet *et al.*, 2021b). As far as we know, our present study is the most comprehensive framework on this issue.

From Western Amazonia, we found that phyllomedusines conquered the central and southern regions of the Neotropics in the last 16 million years (Figs. 4E and 4F), encompassing the genera *Pithecopus* and *Phyllomedusa*. *Pithecopus* lineages jump-dispersed from Western Amazonia twice during the middle Miocene, occupying areas that would later become the diagonal of open/dry landscapes (DODL). We observed a similar pattern in the most speciose clade in *Phyllomedusa*, encompassing the *Ph. burmeisteri* group (sensu Faivovich *et al.*, 2010): *Ph. boliviana* Boulenger, 1902, *Ph. sauvagii* Boulenger, 1882, and *Ph. venusta* Duellman and Trueb, 1967. Both cases exemplify a “north-southern” biogeographical colonisation of the Neotropics, far mentioned on literature for several groups (snakes, Wüster *et al.*, 2002; Hamdan *et al.*, 2019; termites, Carrijo *et al.*, 2020), including frogs (e.g., Fouquet *et al.*, 2012, 2014). This further contradicts the hypothesis of a “south-north” diversification pattern for this subfamily.

By occurring during the middle Miocene, these biogeographical processes are concurrent to the transition from Pebas into the Acre system, taking place in an Amazonian wetland. Firstly, all those lineages left Western Amazonia coincidently with a period of recurring marine incursions of the Paranaense sea (Hernández *et al.*, 2005), that may have limited the Amazonia region southward (Figs. 4E and F). These findings suggest a progressive isolation of the core Phyllomedusinae in northern South America, agreeing with previous studies regarding the influence of Paranaense sea in biogeographical history of herpetofauna (e.g., Magalhães, 2012; Seger *et al.*, 2021). The subsequent regression of the Paranaense sea could have facilitated the colonisation of southern areas, helping to explain the diversification pattern we found. Secondly, the subsequent colonisation of South American Atlantic Forest by some of these Miocene lineages of Phyllomedusinae is congruent with the very early opening of DODL, and the late orogeny of the Serra do Mar Mountain Range. This pattern represents a jump dispersal response

to the retraction of forested areas in due to the environmental changes drove by the opening of DODL, commonly found in other anuran groups (e.g., Pirani *et al.*, 2020; Carvalho *et al.*, 2021). The whole biogeographical process also shed light on our findings about the diversification of *Phasmahyla* genus, that remained isolated in South American Atlantic Forest in the last 10 million years.

Also, we found that several phyllomedusine' lineages diversified along the Miocene from Western Amazonia to northward (Figs. 3 and 4F), through the Andes and Central America. The biogeographical history of these species suggests a colonisation of Amazonia and Northern Andes, occupying the forested lowlands from the Miocene to the Pleistocene. During a period of drastic changes in local drainage patterns due to the transition from Pebas into the Acre system (Hoorn *et al.*, 2010), we found the *Phyllomedusa* genus has been particularly successful colonising the entire Amazonian region. Together, these results seem to reinforce the idea that phyllomedusine frogs were able to survive in the lacustrine environment resulting from changes in the Amazonian drainage pattern when compared to other frog groups. Previous studies have already suggested the isolation and geographic expansion in other arboreal frogs may have been affected by the Miocene marine introgressions, depending on their capacity to exploit wetland environments for reproduction (e.g., Ortiz *et al.*, 2023).

Following this northward diversification (Fig. 4F), we found some aspects of the historical biogeography of phyllomedusines as closely related to the uplift of the Northern Andes. Populations of the MRCA of *Callimedusa* in Western Amazonia experienced an early divergence within this region, giving rise to *Callimedusa tomopterna* (Fig. 3). Subsequently, populations from the MRCA of other *Callimedusa* species jump-dispersed to the Northern Andes, where mountain uplifts potentially facilitated sympatric speciation during the mid-

Miocene (Fig. 4F). The diversification of *Pithecopus* in the Cerrado can also be directly attributed to the Andes uplift. Although the MRCA of *Pithecopus* primarily were in Western Amazonia, ancestral populations reached the Cerrado by two separate jump dispersal events (Fig. 3). Divergences coincided with the final opening of the DODL, where the late Andean uplift contributed to the uplift and dryness of the Brazilian Plateau and the subsidence of the Chaco region (Figs. 4E and 4F; Zanella, 2011; see also Silva, 1995; Pontes-Nogueira et al., 2022). Once ancestral lineages of *Pithecopus* colonised the Cerrado, they diversified in sympatry alongside the changing landscape of this region (Fig. 3). Thus, the uplift of the Andes probably played a main role in cladogenetic processes in phyllomedusine lineages during mountain uplift events, as well as in areas that suffered landscape changes influenced by the elevation of the mountain range.

The diversification of the *Agalychnis* and *Cruziohyla* genera during their conquest of Central America is intriguing. In *Agalychnis*, the colonisation of the Central America by the ancestors of the Phyllomedusinae frogs before Miocene precedes the formation of the Isthmus of Panama, proposed to occur in the Plio-Pleistocene (~3 mya; Fig. 3; Haug & Tiedemann, 1998; O’Dea et al., 2016). Bacon et al. (2015) demonstrated two significant waves of dispersal between South and North America at around 20 and 5 Mya, also preceding the recent formation hypothesis times. At a first glance, our results for *Cruziohyla* are in accordance with the first wave of dispersal, as the MRCA of *Cruziohyla* is synchronous to the very early formation of Isthmus of Panama and early uplift of the Eastern Cordillera of Northern Andes (Gregory-Wodzicki, 2000). Recent biological (Bacon et al., 2013, 2015; Bloch et al., 2016) and geological (Farris et al., 2011; Montes et al., 2012, 2015; Jaramillo et al., 2017) findings suggest an older formation for the Isthmus of Panama (early-mid Miocene), despite the discussion in literature

(e.g., O’Dea *et al.*, 2016). However, any statement in this sense is imprecise, as the biogeographical process associated with the early conquest of Central America during the split between phyllomedusines and pelodyadines is not known. It is thought that range expansion (i.e., anagenetic dispersal) is more plausible to occur by land (for land animals) and that jump dispersals are predominantly associated with geographical barriers (see Matzke, 2014 for more information). Therefore, further studies considering the whole Hylidae family may address this issue more precisely. The biogeography of the Neotropics is surely intriguing and intricate, and the history of the monkey tree frogs described here comes to add even more a level of certainty to this statement.

CONCLUSIONS

We found that the biogeographical history of Phyllomedusinae started with a vicariance splitting the Neotropical region, Oceania, and Antarctica. Indeed, vicariance was a common biogeographical process at the early diversification of phyllomedusines, while jump dispersals must have been responsible for the majority of colonizations in the group since the Miocene. Western Amazonia seems to have figured as a species pump for most of the monkey tree frogs, with species colonising the area and diversifying sympatrically even during the highly unstable environment of the Miocene. Also, the orogeny of Northern Andes could have played an important role in species diversification, promoting sympatric speciation both through the uplift of mountains and in areas with drastic landscape changes provoked by the elevation of the Andean Mountain range. Our results also reject the hypothesis of a “south-north” diversification for the group. In brief, we successfully encompassed most of the historical biogeography of this

speciose group, enabling a highly detailed description of the diversification of a charismatic frog subfamily.

ACKNOWLEDGMENTS

We are grateful to Pedro Paulo Goulart Taucce, Tiago Fernandes Carrijo, and Marcelo José Sturaro for their outstanding contributions on the systematics and node dating issues. MPN also thanks Felipe Grazziotin for his help with the R script that generated the Figure 3 of this work. Also, we thank Caio Vinícius de Mira-Mendes, Daniel Branch, Félix Salazar, Henry Miller Alexandre, Mauro Teixeira Jr, Miguel Trefaut Urbano Rodrigues, Ross Alford, Russel Barrett, and Sébastien Sant for providing the photos of anurans presented on Figure 2. Photos of *C. tomopterna* (© S. Sant/ Parc Amazonien de Guyane), *C. sylviae* (© Félix Salazar) and *P. guttata* (© Henry Miller Alexandre) were uploaded under Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) license. Photo of *P. rohdei* (© Caio Vinícius de Mira-Mendes) was uploaded under Attribution 4.0 International (CC BY 4.0) license. LMS dedicates this study in honour of Marco Antonio Servino. No ethics or permit approvals were required for this research.

REFERENCES

- Abello MA, Posadas PE, Ortiz Jaureguizar E. 2010.** Biogeografía histórica de los Caenolestidae (Marsupialia, Paucituberculata) del Cenozoico de América del Sur. *X Congreso Argentino de Paleontología y Bioestratigrafía y VII Congreso Latinoamericano de Paleontología (La Plata, 2010).*
- Akaike H. 1974.** A New Look at the Statistical Model Identification. *IEEE Transactions on*

560 *Automatic Control* 19: 716–723.

561 **Albert JS, Val P, Hoorn C. 2018.** The changing course of the amazon river in the neogene:
 562 Center stage for neotropical diversification. *Neotropical Ichthyology* 16: 1–24.

563 **Andrade FS, Haga IA, Ferreira JS, Recco-Pimentel SM, Toledo LF, Bruschi DP. 2020.** A
 564 new cryptic species of *Pithecopus* (Anura, Phyllomedusidae) in north-eastern Brazil. *European*
 565 *Journal of Taxonomy* 723: 108–134.

566 **Antonelli A. 2016.** Biogeografia Neotropical: Possíveis Metas e Desafios para os Próximos Dez
 567 Anos. In: Carvalho CJB., In: Almeida EAB, eds. *Biogeografia da América do Sul*. Rio de
 568 Janeiro: ROCA, 279–287.

569 **Antonelli A, Sanmartín I. 2011.** Why are there so many plant species in the Neotropics? *Taxon*
 570 60: 403–414.

571 **Azevedo JAR, Collevatti RG, Jaramillo CA, Strömberg CAE, Guedes TB, Matos-Maraví**
 572 **P, Bacon CD, Carillo JD, Faurby S, Antonelli A. 2020.** On the young savannas in the land of
 573 ancient forests. *Neotropical diversification: Patterns and processes*. Springer, 271–298.

574 **Bacon CD, Mora A, Wagner WL, Jaramillo CA. 2013.** Testing geological models of
 575 evolution of the Isthmus of Panama in a phylogenetic framework. *Botanical Journal of the*
 576 *Linnean Society* 171: 287–300.

577 **Bacon CD, Silvestro D, Jaramillo CA, Smith BT, Chakrabarty P, Antonelli A. 2015.**
 578 Biological evidence supports an early and complex emergence of the Isthmus of Panama.
 579 *Proceedings of the National Academy of Sciences* 112: 6110–6115.

580 **Baêta D, Giasson LOM, Pombal Jr JP, Haddad CFB. 2016.** Review of the rare genus

Phrynomedusa Miranda-Ribeiro, 1923 (Anura: Phyllomedusidae) with description of a new species. *Herpetological Monographs* 30: 49–78.

Bloch JI, Woodruff ED, Wood AR, Rincon AF, Harrington AR, Morgan GS, Foster DA, Montes C, Jaramillo CA, Jud NA, Jones DS, MacFadden BJ. 2016. First North American fossil monkey and early Miocene tropical biotic interchange. *Nature* 533: 243.

Boddaert P. 1772. Brief ... aan ... Johannes Oosterdyk Schacht naar het Leven Vervaaardige Afbeelding, van den Twee-koleurigen Kikvorsch, Uit des Vezameling van ... Johannes Albertus Schlosser /Epistola ad ... Johannem Oosterdyk Schacht ... de Rana bicolore, Descripta Atque Accuratissima Icone Illustrata Ex museo ... Johannis Alberti Schlosseri. M. Magérus.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, Suchard MA, Rambaut A, Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS computational biology* 10: e1003537.

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert D, De Maio N, Matschiner M, Mendes FK, Müller NF, Ogilvie HA, du Plessis L, Poppinga A, Rambaut A, Rasmussen D, Siveroni I, Suchard MA, Wu CH, Xie D, Zhang C, Stadler T, Drummond AJ. 2019. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLOS Computational Biology* 15: e1006650.

Boulenger GA. 1882. Catalogue of the snakes in the British Museum (Natural History). order of the Trustees.

Boulenger GA. 1885. IX.—Second list of reptiles and batrachians from the province Rio Grande do Sul, Brazil, sent to the Natural-History Museum by Dr. H. von Ihering. *Journal of Natural History* 16: 85–88.

- 603 **Boulenger GA. 1902.** LIV.—Descriptions of new batrachians and reptiles from the Andes of
604 Peru and Bolivia. *Journal of Natural History* 10: 394–402.
- 605 **Burnham KP, Anderson DR. 2004.** Multimodel inference: Understanding AIC and BIC in
606 model selection. *Sociological Methods and Research* 33: 261–304.
- 607 **Carneiro L, Bravo GA, Aristizabal N, Cuervo AM, Aleixo A. 2018.** Molecular systematics
608 and biogeography of lowland antpittas (Aves, Grallariidae): The role of vicariance and dispersal
609 in the diversification of a widespread Neotropical lineage. *Molecular phylogenetics and*
610 *evolution* 120: 375–389.
- 611 **Carrijo TF, Pontes-Nogueira M, Santos RG, Morales AC, Cancellato EM, Scheffrahn RH.**
612 **2020.** New World Heterotermes (Isoptera, Rhinotermitidae): molecular phylogeny, biogeography
613 and description of a new species. *Systematic Entomology* 45: 527–539.
- 614 **Carvalho AL. 1954.** A preliminary synopsis of the genera of American mycrohylid frogs.
615 *Occasional papers of the Museum of Zoology, University of Michigan* 555: 1–19.
- 616 **Carvalho-e-Silva ANAMPT de, Silva GR da, Carvalho-e-Silva SP. 2009.** A new species of
617 Phasmahyla Cruz, 1990 from the Atlantic Forest in the state of Rio de Janeiro, Brazil (Amphibia,
618 Hylidae, Phyllomedusinae). *Zootaxa* 2120: 15–26.
- 619 **Carvalho TRD, Moraes LJCL, Lima AP, Fouquet A, Peloso PL V, Pavan D, Drummond**
620 **LO, Rodrigues MT, Giaretta AA, Gordo M, Neckel-Oliveira S, Haddad CFB. 2021.**
621 Systematics and historical biogeography of Neotropical foam-nesting frogs of the Adenomera
622 heyeri clade (Leptodactylidae), with the description of six new Amazonian species. *Zoological*
623 *Journal of the Linnean Society* 191: 395–433.

- 624 **Castroviejo-Fisher S, Koehler J, De La Riva I, Padial JM. 2017.** A new morphologically
- 625 cryptic species of Phyllomedusa (Anura: Phyllomedusidae) from Amazonian forests of northern
- 626 Peru revealed by DNA sequences. *Zootaxa* 4269: 245–264.
- 627 **Chamberlain SA, Boettiger C. 2017.** *R Python, and Ruby clients for GBIF species occurrence*
- 628 *data*. PeerJ Preprints.
- 629 **Cope ED. 1864.** Contributions to the herpetology of tropical America. *Proceedings of the*
- 630 *Academy of natural Sciences of Philadelphia*: 166–176.
- 631 **Cope ED. 1866.** Fourth contribution to the herpetology of tropical America. *Proceedings of the*
- 632 *Academy of natural Sciences of Philadelphia*: 123–132.
- 633 **Cope ED. 1868.** An examination of the Reptilia and Batrachia obtained by the Orton Expedition
- 634 to Equador and the Upper Amazon, with notes on other species. *Proceedings of the Academy of*
- 635 *Natural Sciences of Philadelphia*: 96–140.
- 636 **Costa LP. 2003.** The historical bridge between the Amazon and the forest of brazil a study of
- 637 molecular phylogeography with small mammals. *Journal of Biogeography* 30: 71–86.
- 638 **Cruz CAG da. 1991.** Sobre as relaoes intergenericas de Phyllomedusinae da floresta atlntica
- 639 (Amphibia, Anura, Hylidae). *Revista Brasileira de Biologia* 50: 709–726.
- 640 **Dal Vechio F, Prates I, Grazziotin FG, Zaher H, Rodrigues MT. 2018.** Phylogeography and
- 641 historical demography of the arboreal pit viper *Bothrops bilineatus* (Serpentes, Crotalinae) reveal
- 642 multiple connections between Amazonian and Atlantic rain forests. *Journal of Biogeography*: 1–
- 643 12.
- 644 **Daudin FM. 1800.** Histoire Naturelle des Quadrupèdes Ovipares. Livraison 2. Marchant et Cie,

645 Paris.

646 **Díaz de Gamero ML. 1996.** The changing course of the Orinoco River during the Neogene: a
647 review. *Palaeogeography, Palaeoclimatology, Palaeoecology* 123: 385–402.

648 **Dietz L, Dömel JS, Leese F, Mahon AR, Mayer C. 2019.** Phylogenomics of the longitarsal
649 Colossendeidae: The evolutionary history of an Antarctic sea spider radiation. *Molecular*
650 *Phylogenetics and Evolution* 136: 206–214.

651 **Dinerstein E, Olson D, Joshi A, Vynne C, Burgess ND, Wikramanayake E, Hahn N,**
652 **Palminteri S, Hedao P, Noss R, Hansen M, Locke H, Ellis EC, Jones B, Barber CV, Hayes**
653 **R, Kormos C, Martin V, Crist E, Sechrest W, Price L, Baillie JEM, Weeden D, Suckling K,**
654 **Davis C, Sizer N, Moore R, Thau D, Birch T, Potapov P, Turubanova S, Tyukavina A, De**
655 **Souza N, Pintea L, Brito JC, Llewellyn OA, Miller AG, Patzelt A, Ghazanfar SA,**
656 **Timberlake J, Klöser H, Shennan-Farpón Y, Kindt R, Lillesø JPB, Van Breugel P, Graudal**
657 **L, Voge M, Al-Shammari KF, Saleem M. 2017.** An Ecoregion-Based Approach to Protecting
658 Half the Terrestrial Realm. *BioScience* 67: 534–545.

659 **De La Riva I. 1999.** A new Phyllomedusa from southwestern Amazonia (Amphibia: Anura:
660 Hylidae). *Revista Española de Herpetología* 13: 123–131.

661 **Douglas J, Zhang R, Bouckaert R. 2021.** Adaptive dating and fast proposals: Revisiting the
662 phylogenetic relaxed clock model. *PLOS Computational Biology* 17: e1008322.

663 **Driskell AC, Ané C, Burleigh JG, McMahon MM, O’meara BC, Sanderson MJ. 2004.**
664 Prospects for building the tree of life from large sequence databases. *Science* 306: 1172–1174.

665 **Dubois A, Ohler A, Pyron RA. 2021.** *New concepts and methods for phylogenetic taxonomy*

666 *and nomenclature in zoology, exemplified by a new ranked cladonomy of recent amphibians*
 667 *(Lissamphibia): corrigenda and addenda.*

668 **Duellman WE, Trueb L. 1967.** Two new species of tree frogs (genus *Phyllomedusa*) from
 669 Panamá. *Copeia*: 125–131.

670 **Duellman WE, Cadle JE, Cannatella DC. 1988.** A new species of terrestrial *Phyllomedusa*
 671 (*Anura*: *Hylidae*) from southern Peru. *Herpetologica*: 91–95.

672 **Duellman WE, Mendelson JR. 1995.** Amphibians and reptiles from northern Departamento
 673 Loreto, Peru: taxonomy and biogeography. *The University of Kansas Science Bulletin* 55: 329–
 674 376.

675 **Duellman WE, Marion AB, Hedges SB. 2016.** Phylogenetics, classification, and biogeography
 676 of the treefrogs (*Amphibia*: *Anura*: *Arboranae*). *Zootaxa* 4104: 1–109.

677 **Dupin J, Matzke NJ, Särkinen T, Knapp S, Olmstead RG, Bohs L, Smith SD. 2017.**
 678 Bayesian estimation of the global biogeographical history of the *Solanaceae*. *Journal of*
 679 *Biogeography* 44: 887–899.

680 **Escalante T. 2017.** A natural regionalization of the world based on primary biogeographic
 681 homology of terrestrial mammals. *Biological Journal of the Linnean Society* 120: 349–362.

682 **Faivovich J, Haddad CFBB, Garcia PC a. A, Frost DR, Campbell J a., Wheeler WC. 2005.**
 683 Systematic review of the frog family *Hylidae*, with special reference to *Hylinae*: phylogenetic
 684 analysis and taxonomic revision. *Bulletin of the American Museum of natural History* 2005: 1–
 685 240.

686 **Faivovich J, Haddad CFB, Baêta D, Jungfer K, Álvares GFR, Brandão RA, Sheil C,**

- Barrientos LS, Barrio-Amorós CL, Cruz CAG. 2010. The phylogenetic relationships of the charismatic poster frogs, Phyllomedusinae (Anura, Hylidae). *Cladistics* 26: 227–261.
- Farris DW, Jaramillo CA, Bayona G, Restrepo-Moreno SA, Montes C, Cardona A, Mora A, Speakman RJ, Glascock MD, Valencia V. 2011. Fracturing of the Panamanian Isthmus during initial collision with: South America. *Geology* 39: 1007–1010.
- Feng YJ, Blackburn DC, Liang D, Hillis DM, Wake DB, Cannatella DC, Zhang P. 2017. Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proceedings of the national Academy of Sciences* 114: E5864–E5870.
- Fouquet A, Loebmann D, Castroviejo-Fisher S, Padial JM, Orrico VGD, Lyra ML, Roberto IJ, Kok PJR, Haddad CFB, Rodrigues MT. 2012. From Amazonia to the Atlantic forest: Molecular phylogeny of Phyzelaphryninae frogs reveals unexpected diversity and a striking biogeographic pattern emphasizing conservation challenges. *Molecular Phylogenetics and Evolution* 65: 547–561.
- Fouquet A, Santana Cassini C, Fernando Baptista Haddad C, Pech N, Trefaut Rodrigues M. 2014. Species delimitation, patterns of diversification and historical biogeography of the Neotropical frog genus Adenomera (Anura, Leptodactylidae). *Journal of Biogeography* 41: 855–870.
- Fouquet A, Leblanc K, Framit M, Réjaud A, Rodrigues MT, Castroviejo-Fisher S, Peloso PL V, Prates I, Manzi S, Suescun U. 2021a. Species diversity and biogeography of an ancient frog clade from the Guiana Shield (Anura: Microhylidae: Adelastes, Otophryne, Synapturanus) exhibiting spectacular phenotypic diversification. *Biological Journal of the Linnean Society* 132:

709 233–256.

710 **Fouquet A, Marinho P, Réjaud A, Carvalho TR, Caminer MA, Jansen M, Rainha RN,**
 711 **Rodrigues MT, Werneck FP, Lima AP. 2021b.** Systematics and biogeography of the *Boana*
 712 *albopunctata* species group (Anura, Hylidae), with the description of two new species from
 713 Amazonia. *Systematics and Biodiversity* 19: 375–399.

714 **Fouquet A, Cornuault J, Rodrigues MT, Werneck FP, Hrbek T, Acosta-Galvis AR,**
 715 **Massemin D, Kok PJR, Ernst R. 2022.** Diversity, biogeography, and reproductive evolution in
 716 the genus *Pipa* (Amphibia: Anura: Pipidae). *Molecular Phylogenetics and Evolution* 170:
 717 107442.

718 **Frost DR. 2023.** Amphibian Species of the World: an Online Reference. Version 6.1.

719 **Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CF, De Sa RO, Channing A,**
 720 **Wilkinson M, Donnellan SC. 2006.** The amphibian tree of life. *Bulletin of the American*
 721 *Museum of natural History* 2006: 1–291.

722 **Garzione CN, Molnar P, Libarkin JC, MacFadden BJ. 2006.** Rapid late Miocene rise of the
 723 Bolivian Altiplano: Evidence for removal of mantle lithosphere. *Earth and Planetary Science*
 724 *Letters* 241: 543–556.

725 **GBIF. 2023.** Global Biodiversity Information Facility.

726 **Giambiagi L, Alvarez P, Spagnotto S. 2016.** Temporal variation of the stress field during the
 727 construction of the central Andes: Constrains from the volcanic arc region (22–26 S), Western
 728 Cordillera, Chile, during the last 20 Ma. *Tectonics* 35: 2014–2033.

729 **Givnish TJ, Spalink D, Ames M, Lyon SP, Hunter SJ, Zuluaga A, Doucette A, Caro GG,**

730 **McDaniel J, Clements MA, Arroyo MTK, Endara L, Kriebel R, Williams NH, Cameron**
 731 **KM. 2016.** Orchid historical biogeography, diversification, Antarctica and the paradox of orchid
 732 dispersal. *Journal of Biogeography* 43: 1905–1916.

733 **Gregory-Wodzicki KM. 2000.** Uplift history of the Central and Northern Andes: A review.
 734 *Geological Society of America Bulletin* 112: 1091–1105.

735 **Guarnizo CE, Werneck FP, Giugliano LG, Santos MG, Fenker J, Sousa L, D’Angiolella**
 736 **AB, Dos Santos AR, Strüssmann C, Rodrigues MT. 2016.** Cryptic lineages and diversification
 737 of an endemic anole lizard (Squamata, Dactyloidae) of the Cerrado hotspot. *Molecular*
 738 *Phylogenetics and Evolution* 94: 279–289.

739 **Haas A. 2003.** Phylogeny of frogs as inferred from primarily larval characters (Amphibia:
 740 Anura) . *Cladistics* 19: 23–89.

741 **Hamdan B, Guedes TB, Carrasco PA, Melville J. 2019.** A complex biogeographic history of
 742 diversification in Neotropical lancehead pitvipers (Serpentes, Viperidae). *Zoologica Scripta* 00:
 743 1–14.

744 **Haug GH, Tiedemann R. 1998.** Effect of the formation of the Isthmus of Panama on Atlantic
 745 Ocean thermohaline circulation. *Nature* 393: 673–676.

746 **Hernández RM, Jordan TE, Dalenz Farjat A, Echavarría L, Idleman BD, Reynolds JH,**
 747 **Farjat AD, Echavarría L, Idleman BD, Reynolds JH. 2005.** Age, distribution, tectonics, and
 748 eustatic controls of the Paranense and Caribbean marine transgressions in southern Bolivia and
 749 Argentina. *Journal of South American Earth Sciences* 19: 495–512.

750 **Hime PM, Lemmon AR, Lemmon ECM, Prendini E, Brown JM, Thomson RC, Kratovil**

751 **JD, Noonan BP, Pyron RA, Peloso PL V. 2021.** Phylogenomics reveals ancient gene tree
752 discordance in the amphibian tree of life. *Systematic biology* 70: 49–66.

753 **Hoorn C. 1993.** Marine incursions and the influence of Andean tectonics on the Miocene
754 depositional history of northwestern Amazonia: results of a palynostratigraphic study.
755 *Palaeogeography, Palaeoclimatology, Palaeoecology* 105: 267–309.

756 **Hoorn C, Wesselingh FP, ter Steege H, Bermudez MA, Mora A, Sevink J, Sanmartín I,**
757 **Sanchez-Meseguer A, Anderson CL, Figueiredo JP, Jaramillo CA, Riff D, Negri FR,**
758 **Hooghiemstra H, Lundberg JG, Stadler T, Sarkinen T, Antonelli A. 2010.** Amazonia
759 Through Time: Andean Uplift, Climate Change, Landscape Evolution, and Biodiversity. *Science*
760 330: 927–931.

761 **Hoorn C, Bogotá-A GR, Romero-Baez M, Lammertsma EI, Flantua SGA, Dantas EL, Dino**
762 **R, do Carmo DA, Chemale F. 2017.** The Amazon at sea: Onset and stages of the Amazon River
763 from a marine record, with special reference to Neogene plant turnover in the drainage basin.
764 *Global and Planetary Change* 153: 51–65.

765 **Hoorn C, Kirschner J, Beer M, Wei C, Kukla T, Jardine P. 2020.** Grass development in the
766 Amazon drainage basin, evidence from the fossil and phytochemical record. *EGU General*
767 *Assembly Conference Abstracts*.13822.

768 **Insel N, Poulsen CJ, Ehlers TA. 2010.** Influence of the Andes Mountains on South American
769 moisture transport, convection, and precipitation. *Climate Dynamics* 35: 1477–1492.

770 **Jaramillo CA, Hoorn C, Silva SAF, Leite F, Herrera F, Quiroz L, Dino R, Antonioli L.**
771 **2010.** The Origin of the Modern Amazon Rainforest: Implications of the Palynological and
772 Palaeobotanical Record. *Amazonia, Landscape and Species Evolution: A Look into the Past:*

773 317–334.

774 **Jaramillo CA, Montes C, Cardona A, Silvestro D, Antonelli A, Bacon CD. 2017.** Comment
 775 (1) on “Formation of the Isthmus of Panama” by O’Dea et al. *Science Advances* 3: 1–8.

776 **Jetz W, Pyron RA. 2018.** The interplay of past diversification and evolutionary isolation with
 777 present imperilment across the amphibian tree of life. *Nature Ecology and Evolution* 2: 850–858.

778 **Klaus K V., Matzke NJ. 2020.** Statistical comparison of trait-dependent biogeographical
 779 models indicates that Podocarpaceae dispersal is influenced by both seed cone traits and
 780 geographical distance. *Systematic Biology* 69: 61–75.

781 **Landis MJ, Matzke NJ, Moore BR, Huelsenbeck JP. 2013.** Bayesian analysis of
 782 biogeography when the number of areas is large. *Systematic Biology* 62: 789–804.

783 **Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. 2017.** Partitionfinder 2: New
 784 methods for selecting partitioned models of evolution for molecular and morphological
 785 phylogenetic analyses. *Molecular Biology and Evolution* 34: 772–773.

786 **Li W, Cowley A, Uludag M, Gur T, McWilliam H, Squizzato S, Park YM, Buso N, Lopez**
 787 **R. 2015.** The EMBL-EBI bioinformatics web and programmatic tools framework. *Nucleic Acids*
 788 *Research* 43: W580–W584.

789 **Linnaeus C. 1758.** Systema Naturae per regna tria naturae, secundum classes, ordines, genera,
 790 species; cum characteribus, differentiis, synonymis, locis.

791 **Luo, A, Zhang, C, Zhou, QS, Ho, SY, & Zhu, CD. 2023.** Impacts of Taxon-Sampling Schemes
 792 on Bayesian Tip Dating Under the Fossilized Birth-Death Process. *Systematic Biology*, 72: 781–
 793 801.

794 **Lutz A. 1925.** Batraciens du Brésil. *Comptes Rendus et Mémoires Hebdomadaires des Séances*
795 *de la Société de Biologie et des ses Filiales* 93: 137–139.

796 **Lutz B. 1950.** Anfíbios Anuros da coleção Adolpho Lutz: V. locomoção e estrutura das
797 extremidades Va. *Phyllomedusa* (P.) *burmeisteri* distincta A. lutz: Vb. *Aplastodiscus perviridis*
798 A. Lutz. *Memórias do Instituto Oswaldo Cruz* 48: 599–637.

799 **Magalhães RF de. 2012.** Filogeografia em áreas abertas da América do Sul: origem e
800 diversificação de linhagens de *Scinax squalirostris* (Anura, Hylidae). Unpublished thesis,
801 Universidade Federal de Goiás.

802 **Magalhães RF de, Lemes P, Camargo A, Oliveira U, Brandão RA, Thomassen H, Garcia**
803 **PC de A, Leite FSF, Santos FR. 2017.** Evolutionarily significant units of the critically
804 endangered leaf frog *Pithecopus ayeaye* (Anura, Phyllomedusidae) are not effectively preserved
805 by the Brazilian protected areas network. *Ecology and Evolution* 7: 8812–8828.

806 **Matschiner, M. 2019.** Selective sampling of species and fossils influences age estimates under
807 the fossilized birth–death model. *Frontiers in Genetics*, 10: 1064.

808 **Matzke NJ. 2013.** Probabilistic historical biogeography: new models for founder-event
809 speciation, imperfect detection, and fossils allow improved accuracy and model-testing.
810 *Frontiers of Biogeography* 5: 242–248.

811 **Matzke NJ. 2014.** Model selection in historical biogeography reveals that founder-event
812 speciation is a crucial process in island clades. *Systematic Biology* 63: 951–970.

813 **Matzke NJ. 2022.** Statistical comparison of DEC and DEC+J is identical to comparison of two
814 ClaSSE submodels, and is therefore valid. *Journal of Biogeography* 49: 1805–1824.

- 815 **Miller MA, Pfeiffer W, Schwartz T. 2010.** Creating the CIPRES Science Gateway for
816 inference of large phylogenetic trees. *2010 Gateway Computing Environments Workshop*. New
817 Orleans, LA: IEEE, 1–8.
- 818 **Miranda-Ribeiro A de. 1923.** As Phyllomedusas do Museu Paulista. *Boletim do Museu*
819 *Nacional do Rio de Janeiro* 1: 3–6.
- 820 **Montes C, Cardona A, McFadden R, Morón SE, Silva CA, Restrepo-moreno SA, Ramírez**
821 **DA, Hoyos N, Wilson J, Farris DW, Bayona GA, Jaramillo CA, Valencia V, Bryan J,**
822 **Flores JA. 2012.** Evidence for middle Eocene and younger land emergence in central Panama:
823 Implications for Isthmus closure. *Bulletin of the Geological Society of America* 124: 780–799.
- 824 **Montes C, Cardona A, Jaramillo CA, Pardo A, Silva JC, Valencia V, Ayala C, Pérez-Angel**
825 **LC, Rodriguez-Parra LA, Ramirez V, Niño H. 2015.** Middle Miocene closure of the Central
826 American Seaway. *Science* 348: 226–229.
- 827 **Morrone JJ. 2006.** Biogeographic Areas and Transition Zones of Latin America and the
828 Caribbean Islands Based on Panbiogeographic and Cladistic Analyses of the Entomofauna.
829 *Annual Review of Entomology* 51: 467–494.
- 830 **Morrone JJ. 2014.** *Biogeographical regionalisation of the neotropical region*.
- 831 **Myers N, Mittermeyer RA, Mittermeyer CG, Da Fonseca GAB, Kent J. 2000.** Biodiversity
832 hotspots for conservation priorities. *Nature* 403: 853–858.
- 833 **O’Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-moreno SA, Cione AL, Collins LS,**
834 **Queiroz A De, Farris DW, Norris RD, Stallard RF, Woodburne MO, Aguilera O, Aubry M**
835 **pierre, Berggren WA, Budd AF, Cozzuol MA, Coppard SE, Duque-caro H, Finnegan S,**

836 **Gasparini GM, Grossman EL, Johnson KG, Keigwin LD, Knowlton N, Leigh EG,**
 837 **Leonard-pingel JS, Marko PB, Pyenson ND, Rachello-dolmen PG, Soibelzon E, Soibelzon**
 838 **L, Todd JA, Vermeij GJ, Jackson JBC. 2016.** Formation of the Isthmus of Panama. *Science* 2:
 839 1–12.

840 **Olson D, Dinerstein E, Wikramanayake E, Burgess N, V. N. Powell G, C. Underwood E, A.**
 841 **D’amico J, Itoua I, E. Strand H, Morrison J, J. Loucks C, F. Allnutt T, Ricketts T, Kura Y,**
 842 **Lamoreux J, W. Wettengel W, Hedao P, Kassem K. 2001.** Terrestrial Ecoregions of the
 843 World: A New Map of Life on Earth. *BioScience* 51: 933–938.

844 **Ortiz DA, Hoskin CJ, Werneck FP, Réjaud A, Manzi S, Ron SR, Fouquet A. 2023.**
 845 Historical biogeography highlights the role of Miocene landscape changes on the diversification
 846 of a clade of Amazonian tree frogs. *Organisms Diversity & Evolution* 23: 395–414.

847 **Paradis E, Claude J, Strimmer K. 2004.** APE: Analyses of phylogenetics and evolution in R
 848 language. *Bioinformatics* 20: 289–290.

849 **Pereira EA, Rocha LCL, Folly H, da Silva HR, Santana DJ. 2018.** A new species of spotted
 850 leaf frog, genus *Phasmahyla* (Amphibia, Phyllomedusidae) from Southeast Brazil. *PeerJ* 6:
 851 e4900.

852 **Peres EA, Silva MJ, Solferini VN. 2017.** Phylogeography of the spider *Araneus venatrix*
 853 (*Araneidae*) suggests past connections between Amazon and Atlantic rainforests. *Biological*
 854 *Journal of the Linnean Society* 121: 771–785.

855 **Peters WCH. 1873.** Über eine, zwei neue Gattungen enthaltende, Sammlung von Batrachiern
 856 des Hrn. Dr. O. Wucherer aus Bahia, so wie Über einige neue oder weniger bekannte Saurier. 1.
 857 Über Hrn. Dr. O. Wucherer’s Batrachiersammlung aus Bahia. *Monatsberichte der königlich*

- 858 *Akademie der Wissenschaften zu Berlin* 1972: 768–774.
- 859 **Pirani RM, Peloso PL V, Prado JR, Polo ÉM, Knowles LL, Ron SR, Rodrigues MT,**
860 **Sturaro MJ, Werneck FP. 2020.** Diversification history of clown tree frogs in Neotropical
861 rainforests (Anura, Hylidae, *Dendropsophus leucophyllatus* group). *Molecular Phylogenetics*
862 *and Evolution* 150: 106877.
- 863 **Pombal Jr Jr JP, Haddad CFB. 1992.** Espécies de Phyllomedusa do grupo burmeisteri do
864 Brasil oriental, com descrição de uma espécie nova (Amphibia, Hylidae). *Revista Brasileira de*
865 *Biologia* 52: 217–229.
- 866 **Pontes-Nogueira M, Martins M, Alencar LR V., Sawaya RJ. 2021.** The role of vicariance and
867 dispersal on the temporal range dynamics of forest vipers in the Neotropical region. *PLOS ONE*
868 16: e0257519.
- 869 **Portik, DM, Streicher, JW, & Wiens, JJ. 2023.** Frog phylogeny: A time-calibrated, species-
870 level tree based on hundreds of loci and 5,242 species. *Molecular Phylogenetics and Evolution*
871 188, 107907.
- 872 **Poulsen CJ, Ehlers TA, Insel N. 2010.** Onset of Convective Rainfall During Gradual Late
873 Miocene Rise of the Central Andes. *Science* 328: 490–493.
- 874 **Prates I, Melo-Sampaio PR, Drummond L de O, Teixeira M, Rodrigues MT, Carnaval AC.**
875 **2017.** Biogeographic links between southern Atlantic Forest and western South America:
876 Rediscovery, re-description, and phylogenetic relationships of two rare montane anole lizards
877 from Brazil. *Molecular Phylogenetics and Evolution* 113: 49–58.
- 878 **Prieto-Torres DA, Rojas-Soto OR, Bonaccorso E, Santiago-Alarcon D, Navarro-Sigüenza**

879 **AG. 2019.** Distributional patterns of Neotropical seasonally dry forest birds: a biogeographical
880 regionalization. *Cladistics* 35: 446–460.

881 **Pyron RA, Wiens JJ. 2011.** A large-scale phylogeny of Amphibia including over 2800 species,
882 and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics*
883 *and Evolution* 61: 543–583.

884 **R Core Team. 2023.** R: A language and environment for statistical computing. *R Foundation for*
885 *Statistical Computing*.

886 **Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018.** Posterior summarization in
887 Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67: 901–904.

888 **Ree RH, Moore BR, Webb CO, Donoghue MJ. 2005.** A likelihood framework for inferring the
889 evolution of geographic range on phylogenetic trees. *Evolution* 59: 2299–2311.

890 **Ree RH, Smith SA. 2008.** Maximum Likelihood Inference of Geographic Range Evolution by
891 Dispersal, Local Extinction, and Cladogenesis. *Systematic Biology* 57: 4–14.

892 **Reguero MA, Gelfo JN, López GM, Bond M, Abello A, Santillana SN, Marensi SA. 2014.**
893 Final Gondwana breakup: The Paleogene South American native ungulates and the demise of the
894 South America–Antarctica land connection. *Global and Planetary Change* 123: 400–413.

895 **Réjaud A, Rodrigues MT, Crawford AJ, Castroviejo-Fisher S, Jaramillo AF, Chaparro JC,**
896 **Glaw F, Gagliardi-Urrutia G, Moravec J, Ignacio J. 2020.** Historical biogeography identifies
897 a possible role of Miocene wetlands in the diversification of the Amazonian rocket frogs
898 (Aromobatidae: Allobates). *Journal of Biogeography* 47: 2472–2482.

899 **Rivera-Correa M, Duarte-Cubides F, Rueda-Almonacid JV, Daza JM. 2013.** A new red-

- 900 eyed treefrog of *Agalychnis* (Anura: Hylidae: Phyllomedusinae) from middle Magdalena River
- 901 valley of Colombia with comments on its phylogenetic position. *Zootaxa* 3636: 85–100.
- 902 **Rodríguez Tribaldos V, White NJ, Roberts GG, Hoggard MJ. 2017.** Spatial and temporal
- 903 uplift history of South America from calibrated drainage analysis. *Geochemistry, Geophysics,*
- 904 *Geosystems* 18: 2321–2353.
- 905 **Ron SR, Almendariz A, Cannatella DC. 2013.** The Phyllomedusa perinesos group (Anura:
- 906 Hylidae) is derived from a Miocene Amazonian lineage. *Zootaxa* 3741: 289–294.
- 907 **Ronquist F. 1997.** Dispersal-Vicariance Analysis: A New Approach to the Quantification of
- 908 Historical Biogeography. *Syst Biol* 46: 195–203.
- 909 **Salas-Gismondi R, Flynn JJ, Baby P, Wesselingh FP, Antoine PO, Salas-gismondi R. 2015.**
- 910 A Miocene hyperdiverse crocodylian community reveals peculiar trophic dynamics in proto-
- 911 Amazonian mega-wetlands. *Proceedings of the Royal Society B: Biological Sciences* 282: 4–8.
- 912 **Sanmartín I. 2016.** Modelos Probabilísticos em Biogeografia. In: Carvalho CJB,, In: Almeida
- 913 EAB, eds. *Biogeografia da América do Sul*. São Paulo: ROCA, 119–128.
- 914 **Seger KR, da Veiga Teixeira BF, Annibale FS, Rossa-Feres D de C, Lima AP, Andrade**
- 915 **GV, Giaretta AA, Lourenço LB. 2021.** Five Independent Lineages Revealed by Integrative
- 916 Taxonomy in the *Dendropsophus nanus*–*Dendropsophus walfordi* Species Complex. *Diversity*
- 917 13: 522.
- 918 **Silva JMC. 1995.** Biogeographic analysis of the South American Cerrado avifauna. *Steenstrupia*
- 919 21: 49–67.
- 920 **Smith BT, McCormack JE, Cuervo AM, Hickerson M, Aleixo A, Cadena CD, Perez-Eman**

- 921 **J, Burney CW, Xie X, Harvey MG. 2014.** The drivers of tropical speciation. *Nature* 515: 406–
922 409.
- 923 **Soares, AE, & Schrago, CG. 2012.** The influence of taxon sampling and tree shape on
924 molecular dating: an empirical example from Mammalian mitochondrial genomes.
925 *Bioinformatics and Biology Insights*, 6, BBI-S9677.
- 926 **Sobral-Souza T, Lima-Ribeiro MS, Solferini VN. 2015.** Biogeography of Neotropical
927 Rainforests: past connections between Amazon and Atlantic Forest detected by ecological niche
928 modeling. *Evolutionary Ecology* 29: 643–655.
- 929 **Spix JB von. 1824.** Animalia nova sive Species novae testudinum et ranarum, quas in itinere per
930 Brasiliam annis MDCCCXVII-MDCCCXX... FS Hübschmann.
- 931 **Stöck M, Dufresnes C, Litvinchuk SN, Lymberakis P, Biollay S, Berroneau M, Borzée A,**
932 **Ghali K, Ogielska M, Perrin N. 2012.** Cryptic diversity among Western Palearctic tree frogs:
933 postglacial range expansion, range limits, and secondary contacts of three European tree frog
934 lineages (*Hyla arborea* group). *Molecular phylogenetics and evolution* 65: 1–9.
- 935 **Sundell KE, Saylor JE, Lapen TJ, Horton BK. 2019.** Implications of variable late Cenozoic
936 surface uplift across the Peruvian central Andes. *Scientific reports* 9: 1–12.
- 937 **Thomson RC, Shaffer HB. 2010.** Sparse supermatrices for phylogenetic inference: taxonomy,
938 alignment, rogue taxa, and the phylogeny of living turtles. *Systematic biology* 59: 42–58.
- 939 **Töpel M, Zizka A, Calió MF, Scharn R, Silvestro D, Antonelli A. 2017.** SpeciesGeoCoder:
940 Fast categorization of species occurrences for analyses of biodiversity, biogeography, ecology,
941 and evolution. *Systematic Biology* 66: 145–151.

942 **Vaidya G, Lohman DJ, Meier R. 2011.** SequenceMatrix: concatenation software for the fast
 943 assembly of multi-gene datasets with character set and codon information. *Cladistics* 27: 171–
 944 180.

945 **Van Den Ende C, White LT, van Welzen PC. 2017.** The existence and break-up of the
 946 Antarctic land bridge as indicated by both amphi-Pacific distributions and tectonics. *Gondwana*
 947 *Research* 44: 219–227.

948 **Vicente N, Kergoat GJ, Dong J, Yotoko K, Legendre F, Nattier R, Robillard T. 2017.** In and
 949 out of the Neotropics: historical biogeography of Eneopterinae crickets. *Journal of Biogeography*
 950 44: 2199–2210.

951 **Wagenmakers EJ, Farrell S. 2004.** AIC model selection using Akaike weights. *Psychonomic*
 952 *Bulletin & Review* 11: 192–196.

953 **Wagler J. 1830.** Natürliches system der Amphibien mit vorangehender classification der
 954 Säugethiere und Vögel: Ein Beitrag zur vergleichenden zoologie. JG Cotta.

955 **Werneck FP, Leite RN, Geurgas SR, Rodrigues MT. 2015.** Biogeographic history and cryptic
 956 diversity of saxicolous Tropicuridae lizards endemic to the semiarid Caatinga. *BMC*
 957 *Evolutionary Biology* 15: 1–24.

958 **Wiens JJ. 2003.** Missing data, incomplete taxa, and phylogenetic accuracy. *Systematic biology*
 959 52: 528–538.

960 **Wiens JJ, Fetzner Jr JW, Parkinson CL, Reeder TW. 2005.** Hylid frog phylogeny and
 961 sampling strategies for speciose clades. *Systematic biology* 54: 778–807.

962 **Wiens JJ, Morrill MC. 2011.** Missing data in phylogenetic analysis: reconciling results from

simulations and empirical data. *Systematic biology* 60: 719–731.

Wüster W, Salomão MDG, Quijada-Mascareñas JA, Thorpe RS, BBBSP. 2002. Origin and evolution of the South American pitviper fauna: evidence from mitochondrial DNA sequence analysis. *Biology of the Vipers*. Eagle Mountain Publishing Eagle Mountain, Utah, 111–128.

Zanella FC V. 2011. Evolução da biota da diagonal de formações abertas secas da América do Sul. In: Carvalho CJB,, In: Almeida EAB, eds. *Biogeografia da América do Sul: padrões e processos*. São Paulo: Roca, 198–220.

Zhang Z, Schwartz S, Wagner L, Miller W. 2000. A Greedy Algorithm for Aligning DNA Sequences. *Journal of Computational Biology* 7: 203–214.

Zimmermann H, Zimmermann E. 1988. Etho-Taxonomie und zoogeographische artengruppenbildung bei pfeilgiftfröschen (Anura: Dendrobatidae). *Salamandra (Frankfurt am Main)* 24: 125–160.

Zizka A, Antunes Carvalho F, Calvente A, Rocio Baez-Lizarazo M, Cabral A, Coelho JFR, Colli-Silva M, Fantinati MR, Fernandes MF, Ferreira-Araújo T, Gondim Lambert Moreira F, Santos NMC, Santos TAB, dos Santos-Costa RC, Serrano FC, Alves da Silva AP, de Souza Soares A, Cavalcante de Souza PG, Calisto Tomaz E, Vale VF, Vieira TL, Antonelli A. 2020. No one-size-fits-all solution to clean GBIF (M Costello, Ed.). *PeerJ* 8: e9916.

.

DATA AVAILABILITY STATEMENTS

984 Supplementary data is available at Figshare, under the following DOI:
 985 <https://doi.org/10.6084/m9.figshare.24282592.v1>. The third-party data used in this study is
 986 available at GBIF under the following URL: <https://www.gbif.org/species/4817115>.

987

Figure 1

A, Map of biogeographical areas in Neotropical region;

B, Map of biogeographical areas in Oceania. The areas are: Central America (A), Northern Andes (B), Western Amazonia (C), Eastern Amazonia (D), Southern Amazonia (E), Caatinga (F), Cerrado (G), Northern Atlantic Forest (H), Central Atlantic Forest (I), Southern Atlantic Forest (J), Chaco/Pampas (K), and Oceania (L); C, The summary of events acting as potential geographical barriers. We have used double arrows to specify instances where dispersal probabilities between the areas denoted were the only ones being affected; in contrast, we used commas to specify the cases where any dispersal through those areas had their probability decreased. Time stratification was applied to address landscape dynamics to our analysis, being indicated by the gray-white transition in geological time scale. In Neotropics, the complex Amazonas/Madeira was denoted as a geographical barrier to dispersal between Amazonian areas since late-Miocene; previously, Pebas and Acre systems were actuating on the same region along the entire Miocene. Moreover, Paranaense sea was another marine introgression occurring in Neotropics along the Miocene. The increase in connectivity between North and South America since the mid-Miocene, due to the formation of Panamá Isthmus bring another example of a geographical barrier “softened” through time. On the other hand, the uplift rates of Northern Andes and Serra do Mar and Mantiqueira Mountain Ranges had increased since mid-Miocene, becoming a harsher barrier. This is also the case of DODL, that reduced connectivity between Amazonian and Atlantic forested areas by the expansion of aridity since the very late-Miocene. In Oceania, the connectivity between Australia and both New Guinea and Zealand has also varied over the time. Furthermore, deserts had firstly isolated Western Mesic Biome, subsequently expanding to the East since Oligocene and affecting dispersals to the Eastern Mesic Biome. See Supplementary data S4 and Tables S4.1 and S4.2 for details.




Figure 2

Bayesian dated phylogenetic tree of Phyllomedusidae, based on 13 mitochondrial and nuclear concatenated loci (7,290 bp, 95 terminals).

Horizontal green bars represent the 95% HPD (height posterior density) intervals for the divergence date estimates. Black asterisks indicate clades where the value of posterior probability (pp) is lower than 0.9. See Supplementary data S1 for details about partitioning and the models of nucleotide substitution.




Figure 3

Ancestral range estimations from DECTS + j model implemented in BioGeoBEARS.

Colored boxes with letters represent the most probable range estimated by the model (boxes for nodes in the last 20 million years are not shown for better visualization of these nodes).

Ancestral area estimations at nodes represent areas before an inferred instantaneous speciation event; colored rectangles at the tips with letters indicate the current distribution of extant species. Pie charts on nodes represent probabilities of ranges. Only the four most probable states at each node are shown for better visualization, and blank spaces represents all other probabilities (see Supplementary data S5 for more information).




Figure 4

Summary of the recovered biogeographical processes using arrows, following our time stratification (from the oldest to the newest).

A, a vicariant event at the split between Pelodryadinae + Phyllomedusinae, isolating both subfamilies from a wide-distributed MRCA; B, a vicariant event originating the *Phrynomedusa* genus by the isolation of populations in the Central Atlantic Forest, the same biogeographical unit from where jump-dispersed the populations who conquest the Western Amazonia and lead to the MRCA of (*Phyllomedusa* (*Callimedusa* + *Pithecopus*)) at 32.4 Mya; C, a vicariant event, resulting in the emergence of *Hylomantis* and *Agalychnis* genera from a wide-distributed MRCA; D, another vicariant event at the early diversification of *Agalychnis* genus, at the divergence of the Andean species *A. hulli*; E, north-south pattern of diversification of *P. burmeisteri* group, whose MRCA jump dispersed from Western Amazonia giving rise to a diversification along Chaco and, subsequently, Atlantic Forest' units during the early opening of DODL; F, the divergent patterns of diversification in *Pithecopus* (north-south, colonizing the Cerrado by jump dispersal in two waves) and *Callimedusa* (south-north, colonizing the Northern Andes by range expansion) genera, both taking place in Miocene concurrently to the Pebas system. See Fig. 3 for info about the units' colors and letters.




Table 1 (on next page)

AIC comparisons of the Ancestral range estimation models.

LnL = log-likelihood of the model. n = number of free parameters in the model (that being d, e, j and w); d = rate of range expansion (i.e. anagenetic dispersal); e = rate of range contraction (i.e. extinction); j = jump dispersal process; and w = dispersal multiplier parameter (for TS models). DECTS+j is shown in bold and represents the best model under AIC and AIC weights.

Table 1. AIC comparisons of the Ancestral range estimation models. LnL = log-likelihood of the model. n = number of free parameters in the model (that being d, e, j and w); d = rate of range expansion (i.e. anagenetic dispersal); e = rate of range contraction (i.e. extinction); j = jump dispersal process; and w = dispersal multiplier parameter (for TS models). DECTS+j is shown in bold and represents the best model under AIC and AIC weights.

Models	LnL	n	<i>d</i>	<i>e</i>	<i>j</i>	<i>w</i>	AIC	AIC weights
DECTS+j	-200.5	3	0.0072	<0.001	0.051	1	406.9	~1
DECTS	-208	2	0.0086	<0.001	0	1	420	0.0015
DIVALIKETS+j	-210.7	3	0.0079	<0.001	0.052	1	427.3	<0.001
DECTS+j+w	-215.7	4	0.004	<0.001	0.017	0.0067	439.4	<0.001
DIVALIKETS	-218.4	2	0.011	<0.001	0	1	440.7	<0.001
DECTS+w	-220.6	3	0.0048	<0.001	0	0.0031	447.2	<0.001
DIVALIKETS+j+w	-225.6	4	0.0048	<0.001	0.018	0.0049	459.3	<0.001
DIVALIKETS+w	-230	3	0.0068	<0.001	0	0.083	466.1	<0.001
DEC+j	-236.9	3	0.0025	<0.001	0.0087	1	479.9	<0.001
DEC	-240.3	2	0.0029	<0.001	0	1	484.7	<0.001
DIVALIKE+j	-244.4	3	0.0028	<0.001	0.012	1	494.8	<0.001
DIVALIKE	-250.3	2	0.0037	<0.001	0	1	504.7	<0.001