

# Assessing the impact of two conventional wastewater treatment plants on small streams with effect-based methods

Catalina Trejos Delgado <sup>Corresp., 1</sup>, Andrea Dombrowski <sup>1</sup>, Jörg Oehlmann <sup>1, 2</sup>

<sup>1</sup> Department Aquatic Ecotoxicology, Johann Wolfgang Goethe Universität Frankfurt am Main, Frankfurt, Germany

<sup>2</sup> Kompetenzzentrum Wasser Hessen, Frankfurt, Germany

Corresponding Author: Catalina Trejos Delgado  
Email address: mctrejosd@stud.uni-frankfurt.de

60% of surface water bodies in Europe do not meet the requirements for good ecological and chemical status and in Germany the situation is even worse with over 90% of surface waters. In addition to hydromorphological degradation, intensive land use and invasive species, chemical pollution is primarily considered to be responsible for the inadequate ecological status of the water bodies. As a quantitatively important source of micropollutants in particular, wastewater treatment plants (WWTPs) represent an important entry path for chemical stressors. It is therefore important to analyze the effectiveness of the WWTPs in eliminating micropollutants and other chemical stressors to mitigate the negative impacts of the treated wastewater (WW) in the aquatic ecosystems. Accordingly, in this study, we evaluated the impacts of two conventional, medium-sized WWTPs on their small receiving water systems in the southwestern region of Hessen in Germany during two sampling campaigns (spring and fall) using effect-based methods (EBM). The results showed that the untreated WW had a very high baseline toxicity and also high endocrine and mutagenic activities. Conventional WW treatment, consisting of mechanical and biological treatment with nitrification, denitrification and phosphate precipitation, reduced baseline toxicity by more than 90% and endocrine activities by more than 80% in both WWTPs. Despite these high elimination rates, the remaining baseline toxicity, the endocrine, dioxin-like and mutagenic activities of the conventionally treated WW was so high that negative effects on the two receiving waters were to be expected. This was confirmed in active monitoring with the amphipod *Gammarus fossarum* and the mud snail *Potamopyrgus antipodarum*, as mortality of both species increased downstream of the WWTPs and reproduction in *P. antipodarum* was also affected. These results indicate that advanced WW treatment is needed to more effectively eliminate chemical stressors to prevent negative impacts of treated WW particularly in small receiving waters.

# Assessing the impact of two conventional wastewater treatment plants on small streams with effect-based methods

Catalina Trejos-Delgado<sup>1</sup>, Andrea Dombrowski<sup>1</sup>, Jörg Oehlmann<sup>1,2</sup>,

<sup>1</sup> Department Aquatic Ecotoxicology, Goethe University Frankfurt am Main, Frankfurt am Main, Germany

<sup>2</sup> Kompetenzzentrum Wasser Hessen, Frankfurt am Main, Germany

Corresponding Author:

Catalina Trejos-Delgado<sup>1</sup>

Max-von-Laue-Str. 13, Frankfurt am Main, 60438, Germany

Email address: mctrejosd@stud.uni-frankfurt.de

## Abstract

60% of surface water bodies in Europe do not meet the requirements for good ecological and chemical status and in Germany the situation is even worse with over 90% of surface waters. In addition to hydromorphological degradation, intensive land use and invasive species, chemical pollution is primarily considered to be responsible for the inadequate ecological status of the water bodies. As a quantitatively important source of micropollutants in particular, wastewater treatment plants (WWTPs) represent an important entry path for chemical stressors. It is therefore important to analyze the effectiveness of the WWTPs in eliminating micropollutants and other chemical stressors to mitigate the negative impacts of the treated wastewater (WW) in the aquatic ecosystems. Accordingly, in this study, we evaluated the impacts of two conventional, medium-sized WWTPs on their small receiving water systems in the southwestern region of Hessen in Germany during two sampling campaigns (spring and fall) using effect-based methods (EBM). The results showed that the untreated WW had a very high baseline toxicity and also high endocrine and mutagenic activities. Conventional WW treatment, consisting of mechanical and biological treatment with nitrification, denitrification and phosphate precipitation, reduced baseline toxicity by more than 90% and endocrine activities by more than 80% in both WWTPs. Despite these high elimination rates, the remaining baseline toxicity, the endocrine, dioxin-like and mutagenic activities of the conventionally treated WW was so high that negative effects on the two receiving waters were to be expected. This was confirmed in active monitoring with the amphipod *Gammarus fossarum* and the mud snail *Potamopyrgus antipodarum*, as mortality of both species increased downstream of the WWTPs and reproduction in *P. antipodarum* was also affected. These results indicate that advanced WW treatment is needed to more effectively

eliminate chemical stressors to prevent negative impacts of treated WW particularly in small receiving waters.

# Keywords

Endocrine disruption, baseline toxicity, dioxin-like activity, mutagenicity, active monitoring, *Potamopyrgus antipodarum*, *Gammarus fossarum*

# Introduction

In many areas worldwide, the ecological status of small streams is impacted by morphological degradation (e.g., channelization and straightening), agricultural land use in the catchment, and a high load of treated wastewater (WW) as the major source of aquatic pollution (Reemtsma et al., 2006; Stalter et al., 2013; Ternes et al., 2017). Next to other anthropogenic chemicals entering the water cycle via municipal and industrial wastewater treatment plants (WWTPs), micropollutants, which include pharmaceuticals, personal care products, biocides, pesticides and endocrine disrupting chemicals (EDCs), among others, have been identified as a main driver that, even at low concentrations, can affect ecosystems and human health (Khan et al., 2022; Yang et al., 2022). These stressors have led to significant declines in aquatic biodiversity, which in turn has had a profound impact on the ecological integrity of many aquatic ecosystems (Enns et al., 2023; Jähnig et al., 2021). Currently, 60% of the surface water bodies in Europe do not meet the requirements of a good ecological and chemical status according to the European Water Framework Directive (European Commission, 2019). In Germany the situation is even worse because more than 90% of surface waters have not yet reached a good ecological status (German Environment Agency, 2017). Municipal WWTPs have been implemented to reduce high nitrogen, phosphorus, and organic matter levels. However, many studies have already shown that the elimination rate for micropollutants often remains poor (Giebner et al., 2018; Gosset et al., 2021; Long & Bonefeld-Jørgensen, 2012). These pollutants are thus continuously released in trace amounts (typically from ng/L to µg/L) into receiving watercourses (Wolf et al., 2022). Particularly, estrone, 17β-estradiol, 17α-ethinylestradiol and bisphenol A have been identified by various studies both in the influent and in the effluent of WWTPs (Körner et al., 2000; Kusk et al., 2011). Besides other compounds, such as polycyclic aromatic hydrocarbons (PAHs) which could be aryl hydrocarbon receptor (AhR) agonists and have a high mutagenic potential, have been measured in industrial and domestic discharges entering rivers directly or indirectly via sewage effluents (Denison & Nagy, 2003; Reifferscheid et al., 2011). Additionally, biologically highly active chemicals such as pharmaceuticals and pesticides have been detected in the effluents from WWTPs on a regular basis (Lopez et al., 2022; Ternes et al., 2015, 2017). Due to the prominent role of WWTPs as a source of micropollutants in surface waters, there are currently significant efforts to improve the elimination of micropollutants in WW treatment. Oxidative processes such as ozonation and filtration processes with activated carbon have proven

to be particularly promising (Prasse et al., 2015). Also, in our study area, the Hessian Ried in the south of Frankfurt am Main, Germany, WW-borne micropollutants have been detected not only in high concentration in the receiving surface waters but also in groundwater, due to the specific geological and hydrological situation with a high infiltration rate. With the “Hessian Ried Trace Substance Strategy”, the Hessian state government is currently implementing various measures to reduce micropollutant contamination in surface waters and groundwater and thereby minimize the ecological consequences and risks to drinking water resources. A central measure is the upgrading of selected municipal WWTPs with an advanced WW treatment step (so-called 4<sup>th</sup> treatment step) to achieve improved micropollutant elimination (HMUKLV, 2018a).

As part of a scientific accompanying program, we will examine the effectiveness of this measure, which will be implemented in the next years, with eBM-based methods (EBM). The aim of our work is to describe the situation before and after implementation of the 4<sup>th</sup> treatment step with a combination of ozonation, followed by activated carbon filtration in two WWTPs. In a first step, we characterized both the impact profile of the treated WW and the effects of WW discharge on the receiving surface waters and thus established a baseline for subsequent investigations after the implementation of the WWTP upgrade. We hypothesize that due to the insufficient elimination of micropollutants (1) a broad spectrum of baseline toxic and specific toxic activities can be detected in conventionally treated WW using EBMs and (2) the corresponding effects can also be detected in the receiving surface water downstream the WWTP effluents. As EBMs we use a battery of *in vivo* and *in vitro* bioassays: The baseline or unspecific toxicity is assessed with the Microtox assay with *Aliivibrio fischeri*, mutagenicity with the Ames fluctuation assay (*Salmonella typhimurium*), endocrine and dioxin-like activities with recombinant yeast reporter gene assays (*Saccharomyces cerevisiae*), and reproduction and survival in an active biological effect monitoring with the stream amphipod *Gammarus fossarum* and the mudsnail *Potamopyrgus antipodarum* in two sampling campaigns (spring and fall) to take into account possible seasonal effects.

## Materials & Methods

### Characterization of the wastewater treatment plants Mörfelden-Walldorf and Bickenbach

The medium-sized WWTPs Mörfelden Walldorf (MW) and Bickenbach (B) are located in the southwestern region of Hessen in Germany and have approximately 48,000 and 32,000 population equivalents (PE), respectively. The WW is currently processed with conventional treatment (mechanical, biological treatment, nitrification, denitrification, phosphate precipitation) in both WWTPs. The receiving watercourses for the treated wastewater are the Geräthsbach (waterbody code 239818) for the WWTP MW, and the Landbach (waterbody code 239628) for the WWTP B. They are classified as small streams in riverine floodplains (type 19) and exhibit a poor chemical status, a strongly or completely altered morphological structure, as well as a moderate and poor ecological status respectively (HLNUG, 2022). Additionally, the mean flow (MQ) in these streams

is 125 L/s (Geräthsbach) and 238 L/s (Landbach) (HLNUG, 2022). Also, it is important to mention that 10 km upstream WWTP MW, there is another WWTP in Langen (Figure 1).

## Collection and storage of the wastewater samples

For the study, four sampling sites were chosen for each of the two WWTPs (Table 1).

In two sampling campaigns in April/May 2022 (spring) and October/November 2022 (fall), 2 L water samples were taken at the four sampling sites at each WWTP at the beginning and at the end of the four weeks active biomonitoring period so that two water samples were obtained and analyzed per season. During the sampling campaign in fall 2022, the WWTP in Langen, which is located 10 km upstream of WWTP MW at the Geräthsbach, was also taken into account (sample code BT-L), because the first collection campaign had shown that the Geräthsbach already showed high activities in the EBM upstream the WWTP MW. At each sampling site, physico-chemical parameters such as temperature, pH, dissolved oxygen, and conductivity were measured in the field with a portable multi-meter (HQ40d, Hach, Germany). The water samples were collected in glass vessels precleaned with deionized water, acetone, and ethanol and heated to 200°C before use. The samples were transported and stored at 7°C in darkness. Phosphate, ammonium, nitrite, and nitrate concentrations were determined using spectroquant test kits (Merck, Darmstadt, Germany) immediately after returning the samples to the laboratory.

## Sample preparation

Water samples were filtered through glass microfiber filters (VWR International GmbH, No. 692, European Cat. No. 516-0885, 90 mm, particle retention: 1.5 µm, Darmstadt, Germany). Then, 2000 mL of each water sample was solid-phase extracted (SPE) with OASIS HLB cartridges (6cc, 200 mg, Waters, Milford, MA, USA) according to Giebner et al. (2018) and transferred to a final volume of 400 µL DMSO so that the final extracts were 5,000-fold concentrated compared to the aqueous samples. Extracts were stored in glass vials at -25°C until further processing. To check a potential contamination of the samples during the sample preparation, 2000 mL of ultrapure water was extracted in parallel (SPE blank) (Giebner et al., 2018).

## Test systems

### *In vitro* assays

#### Baseline toxicity test

The Microtox assay or bioluminescence inhibition test with the bacterium *Aliivibrio fischeri* was conducted to assess the baseline toxicity in the extracts. Based on the International Organization for Standardization (ISO) guideline 11348-3 (ISO, 2007) the assay was performed in 96-well

microplates according to Völker et al. (2017). Samples were analyzed in dilution series in a saline buffer with seven consecutive steps (spacing factor 2 between concentrations), resulting in a 25-fold to 0.2-fold final concentration (Völker et al., 2017). Negative and solvent controls, SPE blank, reference compound (3,5-dichlorophenol), and SPE extracts were serially diluted (1:2) in a saline buffer. 100 µL sample was added to 50 µL of *A. fischeri* solution (not exceeding 1% DMSO in the final medium volume). To detect inhibition, luminescence was measured prior to sample addition and after 30 min incubation using a microplate reader (Infinite 200 Pro, Tecan, Crailsheim, Germany) as described by Völker et al. (2017). The inhibition of luminescence is expressed as the arithmetic mean value of the 50% effect concentration (EC<sub>50</sub>) of three independent experiments, referring to the relative enrichment factor (REF) of the water sample. Nontoxic water samples exhibit an EC<sub>50</sub> threshold value of 300 (REF).

### Recombinant yeast reporter gene assays

The Yeast Estrogen Screen (YES), Yeast Androgen Screen (YAS) Yeast Anti-Estrogen Screen (YAES), Yeast Anti-Androgen Screen (YAAS), and Yeast Dioxin Screen (YDS) were used to assess the receptor-mediated agonistic and antagonistic endocrine activity of the extracted water samples as well as the activation of the aryl hydrocarbon receptor assay according to Brettschneider et al. (2019a, b), Giebner et al. (2018) and Schneider et al. (2020). These assays use genetically modified strains of *Saccharomyces cerevisiae*, which contain the human estrogen receptor  $\alpha$ , androgen and aryl hydrocarbon receptor, respectively. The YAES and YAAS were performed with native samples because SPE enrichment of water samples from surface waters and WWTPs leads to a strong loss of anti-estrogenic and anti-androgenic activities (Brettschneider et al., 2019a; Giebner et al., 2018; Schneider et al., 2020). The YES, YAS and YDS were performed in three independent experiments with every sample, the YAES and YAAS in two independent experiments, with eight technical replicates per sample. The measured activities are expressed as equivalent concentrations (EQ) for the positive substances 17 $\beta$ -estradiol (YES), testosterone (YAS), 4-hydroxytamoxifen (YAES), flutamide (YAAS) and  $\beta$ -naphthoflavone (YDS) and were corrected regarding dilution and enrichment so that equivalent concentrations relate back to the native water sample.

### Ames fluctuation test

The Ames fluctuation assay (ISO, 2012) was performed to assess the mutagenic potential of the SPE samples, using *Salmonella typhimurium* strains YG1041 and YG1042 (Hagiwara et al., 1993) with and without metabolic activation by rodent liver enzymes (S9-mix; Envigo CRS, Roßdorf, Germany). The test is based on the inability of genetically modified *S. typhimurium* strains to synthesize the essential amino acid histidine and, therefore, not being able to survive in a histidine-free medium (Maron & Ames, 1983). The bacteria are exposed to the SPE extracts in a 16.7-fold final sample

concentration (0.2% v/v solvent) and incubated for 72 hours at 37°C in a histidine-free medium. After this time, a color change of the medium indicates the survival of the bacteria by regaining the ability to synthesize histidine because of a back mutation. YG1041 and YG1042 strains without the S9 mix were tested with 2-nitrofluorene as a positive control. Conversely, these strains with S9-mix were tested with 2-amino anthracene as a positive control. The mutagenic potential of the samples is assessed via the percentage of reverted wells, subtracted by the mutants of negative historical controls with samples exceeding the threshold of 20.8% revertant wells are considered as mutagenic. Every sample was tested with both strains, with and without the S9-mix. If an effect was observed, the test was consequently conducted two times.

### ***In vivo* assays**

The *in vivo* assays were applied as an active in-situ monitoring at the upstream and downstream samplings sites of both WWTPs as described by Brettschneider et al. (2019a) and Harth et al. (2018).

### **Active monitoring with *Gammarus fossarum***

The gammarids for the active monitoring were collected at the upper reach of the river Urselbach (Hessen, Germany, 50°13'30.1"N 8°31'06.1"E) one week before the start of the spring and fall campaigns. At the upstream and downstream samplings sites of both WWTPs, two stainless steel cages each containing three enclosures (12.5 cm x 6 cm) were fixed on the riverbed in the direction of flow. The enclosures were capped with a net (mesh size 1.0 mm) at the ends to ensure a water flow. Each enclosure, representing a replicate, contained ten gammarids, resulting in a total of 60 gammarids per site which were exposed *in situ* for four weeks. Black alder leaves (*Alnus glutinosa*) were provided as food in the enclosures *ad libitum* and two small nets (polytetrafluorethylene, 8.2 cm x 3.3 cm) served as hiding places to prevent cannibalism.

After four weeks of exposure, the cages were taken to the laboratory, and the gammarids were fixed in 70% ethanol and stored separately in Eppendorf tubes. As biological endpoints the mortality and fecundity index (number of eggs divided by the body length) were assessed.

For all measurements, a stereomicroscope (Olympus SZ61 R, Olympus Corporation, Tokio, Japan) with a digital camera (JVC Digital Camera KY-F75U, Victor Company of Japan Ltd., Yokohama, Japan) and the Software Diskus (Version 4.50.1458, Carl H. Hilgers, Königswinter, Germany) was used.

### **Active monitoring with *Potamopyrgus antipodarum***

The mudsnails were collected at the rivers Lumda and Horloff (Hessen, Germany) in spring and at the rivers Gambach and Waschbach (Hessen, Germany) in fall. Snails with a shell length

between 3.5 and 4.5 mm were used, following the protocol by Brettschneider et al. (2019a). At the upstream and downstream samplings sites of both WWTPs, two stainless steel cages containing three smaller steel cages (4.5 cm × 3.5 cm, mesh size 0.7 mm) were fixed on the riverbed in the direction of flow. Ten snails were placed in each small cage, representing the six replicates and resulting in a total of 60 snails which were exposed *in situ* for four weeks. Organic carrot cubes served as food supply and were refilled after two weeks.

As biological endpoints the mortality of the snails, the shell length and the number of embryos of the surviving snails were determined using a stereomicroscope with an ocular micrometer (SMZ-168, Motic, Xiamen, China). The active monitoring with *P. antipodarum* followed the principle of the OECD guideline 242 (OECD, 2016) to assess the impact of reproductive toxicants, including EDCs, on this species.

## Statistical analyses

Statistical analyses were performed using GraphPad Prism version 5.03 for Windows (GraphPad Software, San Diego, California, USA). Normal distribution was tested using the D'Agostino and Pearson test. Continuous data were analyzed with t-tests for normally distributed data and Welch's correction for unequal variances. All data were tested as unpaired data and with 95% confidence interval. Quantal data (mortality) were analyzed with Fisher's exact test. The level of significance was defined as  $\alpha = 0.005$  and indicated at the graphs with asterisks as follows:  $\downarrow = p < 0.05$ ;  $\downarrow\downarrow = p < 0.01$  and  $\downarrow\downarrow\downarrow = p < 0.001$

## Results

### Physicochemical parameters

The conventional treatment process in both WWTPs removed more than 90% of the concentrations of ammonium and phosphate and around 33% of the conductivity, independent of the season. Moreover, the oxygen saturation increased significantly during the fall campaign at the WWTP Mörfelden-Walldorf (MW) by 79% and in spring at the WWTP Bickenbach (B) by 89%. The nitrification in the biological treatment step not only significantly reduced the ammonium concentrations, but the nitrate concentrations increased significantly by factor 9.6 (spring) to 19.6 (fall) in MW and by factor 2.5 (spring) to 2.9 times (fall) in B. Considering the sum concentrations of ammonium-N, nitrite-N and nitrate-N, inorganic N was removed by 90% and 94 % in MW and 97 and 94% in B in spring and fall, respectively.

Regardless of the effectiveness of both WWTPs in reducing the concentrations of phosphate and inorganic nitrogen, they still had a strong impact on the physicochemical parameters in the receiving streams, which carried between 78% and 92% treated WW downstream of the discharger. The oxygen saturation decreased in the receiving streams during both seasons. The conductivity and the nitrate concentrations exceeded the reference value of 800  $\mu\text{S}/\text{cm}$  and 0.05



mg/L, respectively for river type 19 downstream of both WWTPs during the spring and fall campaign (Halle & Müller, 2014). Despite the efficient ammonium removal in both WWTPs, the ammonium concentrations in the receiving waters exceeded the reference value of 0.4 mg/L by up to factor 2.8 (MW in spring). Conversely, the downstream concentrations for nitrate and phosphate are below the reference value during spring and fall in both WWTPs (10 and 0.05 mg/L, respectively) (German Environment Agency, 2017; Halle & Müller, 2014). Finally, the mean values of the pH (not shown in the table 2) meet the requirements for good chemical status.

## ***In vitro* assays**

### **Baseline toxicity test**

The results of the Microtox assay are shown in Figure 2. The baseline toxicity increased in the receiving streams downstream of both WWTPs compared with the upstream sites significantly, except at WWTP MW in fall. Furthermore, all sample sites, except upstream of WWTP B in both seasons showed a significant baseline toxicity. The baseline toxicity was removed by 90% in the WWTPs MW and B in spring and by 96% and 97% in fall. After including the effluent from the WWTP Langen (BT-L in Fig. 2A; located upstream of the WWTP MW) in the analyzes in autumn, it became apparent that the basic toxicity, which was already significantly increased at the upstream site of Mörfelden-Walldorf, can be attributed to the discharges from the WWTP Langen.

### **Estrogenic and anti-estrogenic activity**

The estrogenic activity as determined with the YES is shown in Figure 3. All samples exhibited a significant estrogenic activity above the limit of quantification (LOD = 0.148 ng E-EQ/L). Except the fall campaign at the WWTP MW with a decrease by 6%, the estrogenic activities at the downstream sample sites increased significantly at both WWTPs when compared to the upstream sites in the receiving streams. In the case of the WWTP MW, the increment was around 22% in spring, while at the WWTP B the estrogen activity increased by 60% in the receiving stream in both seasons, despite the fact that the WWTPs removed more than 90% of the estrogenic activity during the biological treatment.

The anti-estrogenic activities in the YAES are presented in Figure 4. The WWTPs did not increase the anti-estrogenic activity in the receiving streams significantly. Both WWTPs had a high removal efficiency of anti-estrogenic activities of over 80% in MW and 90% in Bickenbach in both seasons.

### **Androgenic and anti-androgenic activity**

Androgenic and anti-androgenic activities were analyzed in 5000-fold concentrated SPE extracts and native water samples, respectively. While no cytotoxic effects occurred in the native water samples, the observed cytotoxicity in the SPE extracts necessitated the analysis of diluted extracts.

With the exception of the MT samples, no other sample from the two WWTPs and in the receiving streams showed an androgenic and anti-androgenic activity with a LOD of 25.6 ng testosterone-EQ/L and 771 µg flutamide-EQ/L, respectively during the spring and fall campaigns.

### **Dioxin-like activity**

Figure 5 shows the agonistic activity at the AhR. The MT samples were also cytotoxic for the yeast cells in the YDS, so that the dioxin-like activity had to be assessed in serial dilutions. However, in the case of Bickenbach, even with a dilution series, it was not possible to calculate the dioxin-like activity of the MT samples (Fig. 5B).

All sample sites showed dioxin-like activity above the limit of detection (LOD = 33.8 ng β-naphthoflavone-EQ/L). Dioxin-like activity increased significantly in the receiving streams due to the effluents from the two WWTPs at the downstream sites compared to the upstream sites, with the exception of the fall campaign at the WWTP MW (Fig. 5A). The increases were 21% and 11% in MW and 85% and 72% in B during the spring and fall campaigns, respectively. The calculation of the removal efficiency of dioxin-like activity was only possible for the WWTP MW because of the cytotoxicity even with highly diluted MT samples of the WWTP B. In MW, elimination rates of 60% and 53% were observed.

### **Ames fluctuation test**

The results of the Ames fluctuation test using the strains YG1041 and YG1042 with S9-mix are documented in Table 3. The tests with both strains without the S9-mix resulted in negative results for all samples (less than 20.8% revertant colonies; data not shown). Due to their high cytotoxicity, the MT samples from both WWTPs could not be analyzed with the Ames fluctuation test. All analyzed wastewater samples from the WWTPs MW, B and Langen proved to be mutagenic in the tests with the strains YG1041 and YG1042 with S9 mix, as did the samples taken from the receiving water below the sewage treatment plants. The upstream (US) samples in the receiving water of the WWTP MW were also mutagenic with both strains, while the US sample at WWTP B was not mutagenic in the spring campaign, but mutagenic with the YG1042 strain with S9-mix in the fall campaign.

### **Active monitoring with *Gammarus fossarum***

The mortality in *Gammarus fossarum* during the 4 weeks active monitoring period in the receiving rivers of both WWTPs in spring and fall is shown in figure 6. In both rivers the mortality in the upstream site was higher in fall than in spring. There was also a significantly higher mortality in the downstream sites ( $p < 0.05$ ) compared to the upstream sites (except WWTP B in fall) with an increase of 66% and 77% at WWTP MW and B, respectively.

The fecundity index was assessed for the spring campaign but did not show any significant difference between the downstream and corresponding upstream site. For the fall campaign, the fecundity index could not be calculated because the gammarid populations were outside their reproductive phase.

### Active monitoring with *Potamopyrgus antipodarum*

The mortality and number of embryos in *Potamopyrgus antipodarum* during both campaigns are shown in Figures 7 and 8. The mortality increased significantly at the downstream sites at both WWTPs in fall, while there was no effect on mortality during the spring campaign.

The embryo numbers as the parameter for reproduction in *P. antipodarum* increased at the downstream sites of both WWTPs in the spring campaign, although this increase was not statistically significant. In contrast, the embryo numbers at the downstream site of the WWTP MW were on the same level as upstream in the fall campaign but increased significantly ( $p < 0.01$ ) at the WWTP B in the same period.

## Discussion

### Baseline toxicity

The bioluminescence of *Aliivibrio fischeri* is directly proportional to their metabolic activity, so any disruption by toxic substances results in a decreased luminescence. Considering this, a marked inhibition of the bioluminescence by the raw WW (MT) in both WWTPs was observed ( $EC_{50} < 6.45 \pm 4.16$  REF). This observation is in accordance with previous reports (Macova et al., 2011; Völker et al., 2017). Both conventional WWTPs in B and MW eliminated the baseline toxicity of the raw WW during biological treatment by more than 90%. This elimination performance is also comparable to the findings of other studies, which report elimination rates between 86% and 94% (Völker et al., 2017; Yu et al., 2014).

Despite the considerable elimination of baseline toxicity during the WW treatment process in both WWTPs, the effluents had still a major impact on the receiving waterbodies, so that the downstream sites during the spring and fall campaign were under the non-toxic limit in the Microtox assay. At MW the baseline toxicity increased by 75% in spring and 12% in fall at DS compared to US, while the corresponding values in B were 68% in spring and 44% in fall. These results suggest that the level of baseline toxicity was higher during spring than in the fall campaign in both WWTPs, which is in line with previous studies that concluded that the toxicity of effluents was usually higher in spring and summer (Vasquez & Fatta-Kassinos, 2013). This can be explained

by the fact that during the spring, there is an increase in agricultural activity. Also, due to the increase in runoff, some pesticides or fertilizers may be washed into the water bodies and may contribute to the higher activity in the Microtox assay as a bioassay detecting the unspecific toxicity of a wide range of substances (Tang et al., 2013).

### *Endocrine activity*

The endocrine profile of the influent samples indicates that WW contains compounds with estrogenic, androgenic, and anti-estrogenic activities. However, the conventional treatment process in both WWTP efficiently reduced the estrogenic and androgenic activities by more than 90%. This finding is in line with the results of other studies which report an elimination of estrogenic and androgenic activities between 88% and 95% (Archer et al., 2020; Arlos et al., 2018; Giebner et al., 2018). In the case of anti-estrogenic activity, the efficiency at the WWTPs MW (above 80%) and B (above 90%) was even higher compared to the results of Giebner et al. (2018) reporting a reduction of anti-estrogenic activity during conventional treatment of only 15%.

Although the estrogenic activity of the WW was reduced by more than 90% in both WWTPs, the residual activity was sufficient to adversely affect the receiving waters. This is particularly noticeable at the Landbach, the receiving water of WWTP B, where the estrogenic activity increased most significantly downstream of the WWTP ( $p < 0.001$ ). At the Geräthsbach, the estrogenic activity increased significantly only in spring downstream of the WWTP MW ( $p < 0.01$ ) and in autumn it even decreased significantly ( $p < 0.05$ ). These results are in line with results in a survey of 36 Danish WWTPs and their effluents (Danish Ministry of the Environment, 2005). In almost 70% of the water courses receiving effluents from WWTPs, the level of estrogenic activity downstream the discharge point was higher than the activity upstream of the WWTP effluent. Also the study conducted by Aerni et al. (2004) who compared the estrogenic effect upstream and downstream of five Swiss and French WWTPs support our findings. In all the samples sites, the river upstream of the point of discharge of each WWTP had not estrogen activity ( $< \text{LOQ}$ ) while in the effluent the activity was on average between 0.4-5.5 ng E-EQ/L.

The reason for the lower impact of the WWTP MW on the estrogenic activity in the Geräthsbach compared to the WWTP B and the Landbach is that the Geräthsbach is already influenced by the WWTP Langen upstream of WWTP MW. Accordingly, the WWTP MW is not the only and first source of contamination. This pattern was observed during the spring sampling campaign for almost all analyzed activities (baseline toxicity, estrogenic and dioxin-like activity, mutagenicity). The WWTP Langen is located 10 km upstream of MW and is with 54,600 population equivalents even larger than MW (HMUKLV, 2018b). For this reason, we decided to take an additional sample at the effluent of WWTP Langen during the fall campaign. The result for the estrogenic activity supports the assumption that the effluent from the WWTP Langen is responsible for the higher activity in the US site of the Geräthsbach compared to the Landbach.

We observed that the estrogenic activity in both WWTPs effluents was significantly higher during the fall campaign than in spring ( $p < 0.001$ ). Also this finding is in line with the Danish survey at 36 WWTPs which reports the highest levels of estrogenic activity in summer and fall and the lowest during winter and spring (Danish Ministry of the Environment, 2005). A possible explanation is the higher precipitation in winter and spring so that the concentrations of estrogenically active compounds in the streams are lower than in the dry summer and autumn months. Also, the fall season provides additional phyto-estrogenic material from the leaves (Janeczko, 2021), increasing the estrogenic load in the water bodies.

In terms of agonistic and antagonistic androgen activity, values under the LOD (25.6 ng T-EQ /L and 771.06  $\mu$ g Flu-EQ /L) were detected in both effluents. The lack of a measurable androgenic and anti-androgenic activity is supported by previous studies (Brzezinska et al., 2023; Li et al., 2010). The same was observed for the anti-estrogenic activity, where no significant difference was found between upstream and downstream sample points (values close to the LOD=1,84 mg OHT-EQ/L). This finding contrasts the report by Brzezinska et al. (2023) where an antagonistic activity up to 100 mg OHT/L was found in the effluent of the WWTP Lodz in Poland.

### **Dioxin-like activity**

The raw WW at the WWTP MW contained compounds with an agonistic activity at the aryl-hydrocarbon receptor, resulting in a high dioxin-like activity. Although the conventional biological treatment decreased these activities by more 60% in spring and 53% in fall, this decrease is lower than in the study of Magdeburg et al., (2014) reporting an elimination of dioxin-like activity of more than 81%. For WWTP B, the elimination of dioxin-like activity through conventional WW treatment could not be determined because the raw WW samples were cytotoxic after mechanical treatment or the enzymes responsible for luminescence were directly inhibited by the samples (Reifferscheid et al., 2011). Völker et al. (2016) found a similar result for the raw WW of a WWTP in their study.

Despite the significant reduction of dioxin-like activity in both WWTPs, the remaining activity was sufficient to significantly increase the activity in the Geräthsbach and Landbach, except for the autumn sample from the Geräthsbach, where the activity only increased slightly but not statistically significantly by 12.6%. The measured activities in both rivers are comparable to the rivers Horloff (up to 1150 ng  $\beta$ -NF-EQ/L) and Nidda (up to 1180 ng  $\beta$ -NF-EQ/L) in the north of Frankfurt am Main, which also receive effluents from WWTPs but higher compared to rivers not impacted by WWTP effluents with activities less than 200 ng  $\beta$ -NF-EQ/L (Brettschneider et al., 2019b, 2023).

The seasonal comparison shows significantly higher dioxin-like activities ( $p < 0.001$ ) at all sampling points in MW in spring compared to autumn, while for the WWTP B there is only a corresponding trend without significant differences. Although both WWTPs contribute to the high dioxin-like activity in the rivers, other sources are relevant. The dioxin-like activity in the water phase and in the sediments of rivers is primarily caused by polycyclic aromatic hydrocarbons (PAHs), which enter the waters also via atmospheric deposition and surface runoff from traffic areas (Boxall & Maltby, 1997; Maltby et al., 1995).

### **Mutagenicity (Ames fluctuation test)**

The investigations at both WWTPs showed high mutagenic activities for all BT effluent samples and for all DS samples taken downstream of the WWTPs in the two rivers, using the strains YG1041 and YG1042 with S9 mix, while the same samples were negative in the tests without S9 mix. Due to the high cytotoxicity, the MT samples could not be evaluated. Only the spring sample from the Landbach upstream of WWTP B was negative in all Ames tests.

These findings show that the conventionally treated WW contains substances that are only converted into mutagenic compounds through metabolic activation, as typically occurs by enzymes in the liver of vertebrates or the midgut gland of invertebrates. Furthermore, it shows that conventional WW treatment is not sufficient to eliminate these compounds. The high mutagenic activity at the sampling points in both rivers influenced by the discharge of treated WW demonstrates the urgent need for action to reduce this undesirable effect through advanced WW treatment. Giebner et al. (2018) were able to show that activated carbon filtration in particular is suitable for removing mutagenic substances from WW, while oxidative processes such as ozonation can lead to the formation of alkylating agents, nitrosamines and other mutagenic substances in the treated WW (Mestankova et al., 2014; Schmidt & Brauch, 2008; Tsutsumi et al., 2010).

### **Active monitoring with *Gammarus fossarum* and *Potamopyrgus antipodarum***

To investigate potential pollution effects on the invertebrate community, we conducted an active monitoring study with the crustacean *Gammarus fossarum* and the gastropod *Potamopyrgus antipodarum*. Both species have widely been used for ecotoxicological studies, including the assessment of WWTP effluents and their impacts on the biocenosis in receiving waters (Brettschneider et al., 2019a, 2023; Harth et al., 2018; Schneider et al., 2020). *G. fossarum* is an indicator species of a low-mountain range biocenosis (LANUV, 2015), is particularly sensitive to anthropogenic pressures on water bodies and plays an important role in the food web as a shredder and food for macroinvertebrates, fish, or birds (Besse et al., 2013). *P. antipodarum* is an euryoecious snail inhabiting small creeks, streams, lakes, and estuaries which is particularly sensitive to endocrine active chemicals (Duft et al., 2007).

Given the prominent ecological function of *G. fossarum* in the aquatic community of low-mountain range rivers, the significant increased mortality downstream of the WWTP MW to 59% in spring and 81% in fall is worrying. Although the same tendency was also observed downstream of the WWTP B in both seasons, the mortality levels were lower and increased only significantly during the spring campaign. These findings are in line with results of other active monitoring studies such as Brettschneider et al. (2019a), who found increased mortality rates of 42% and 68% in *G. fossarum* downstream of WWTPs with 43,800 PE and 78,000 PE, respectively.

Also, the mortality of *P. antipodarum* increased significantly downstream of both WWTPs in fall, while there was no significant change in mortality in spring. This finding corresponds to previous studies on other WW-influenced rivers, in which a significant increase in the mortality of *P. antipodarum* occurred downstream of WWTPs, especially in autumn, while the effects were significantly smaller in spring (Brettschneider et al. 2019a). In addition, in previous studies in which both species were used together for active monitoring, mortality was generally lower for *P. antipodarum* than for *G. fossarum*, possibly because the snail can avoid exposure peaks by temporarily closing its shell with the operculum (Brettschneider et al., 2023).

While no influence on reproduction could be determined for *G. fossarum*, the number of embryos for *P. antipodarum* increased downstream of the two WWTPs, except for the autumn campaign in the Landbach. Since it is known that *P. antipodarum* responds to exposure to substances with estrogenic activity by increasing embryo production (Duft et al., 2003, 2007; Jobling et al., 2004) and the estrogenic activity determined by the YES was significantly increased at the study sites located downstream the WWTPs, this indicates that the estrogenic exposure is the responsible factor for the higher embryo numbers. The very high toxicity below the WWTP MW in autumn, which is also reflected in the highest mortalities in *G. fossarum* and *P. antipodarum*, possibly masks the estrogenic effect on the number of embryos in the snail. Also in other studies, the number of embryos in *P. antipodarum* increased below WWTP effluents or when exposed to conventionally treated WW (Brettschneider et al., 2019a; Schneider et al., 2020; Stalter et al., 2010).

Together, these findings demonstrate that the effluents from conventional WWTPs negatively affect key life cycle parameters of representative macroinvertebrate species in the water bodies.

## Conclusions

The findings of this study showed that the EBM are an efficient tool to analyze the multiple effects that complex mixtures, such as WW, has in the aquatic environment. These methods revealed that although both WWTPs had high toxicity removal rates, remaining baseline toxicity, endocrine activity (especially of estrogenic activity) and dioxin-like activity, had a negative impact on both

streams. In the specific case of mutagenic effects, high activities were found in both WWTPs, where the effluents had high mutagenic activity compared with the sample taken upstream. The negative impact of the conventionally treated WW on the receiving streams was not only demonstrated by the *in vitro* test battery but also in the active *in vivo* biomonitoring. In summary, both hypotheses are supported by our findings: conventionally treated WW showed a wide range of activities in assays addressing specific modes of toxicological action but also a high baseline toxicity. These effects were also measured in the receiving surface waters downstream the WWTPs. Therefore, our study demonstrated that conventional WW treatment is insufficient to prevent negative impacts on the receiving aquatic ecosystem. This implies that further advanced WW treatment is urgently needed to guarantee safe effluents in these streams.

# References

- Aerni, H.-R., Kobler, B., Rutishauser, B. V., Wettstein, F. E., Fischer, R., Giger, W., Hungerböhler, A., Marazuela, M. D., Peter, A., Schönenberger, R., Vögeli, A. C., Suter, M. J.-F., & Eggen, R. I. L. (2004). Combined biological and chemical assessment of estrogenic activities in wastewater treatment plant effluents. *Analytical and Bioanalytical Chemistry*, 378(3), 688-696. <https://doi.org/10.1007/s00216-003-2276-4>
- Archer, E., Wolfaardt, G. M., Van Wyk, J. H., & Van Blerk, N. (2020). Investigating (anti)estrogenic activities within South African wastewater and receiving surface waters: Implication for reliable monitoring. *Environmental Pollution*, 263, 114424. <https://doi.org/10.1016/j.envpol.2020.114424>
- Arlos, M. J., Parker, W. J., Bicudo, J. R., Law, P., Marjan, P., Andrews, S. A., & Servos, M. R. (2018). Multi-year prediction of estrogenicity in municipal wastewater effluents. *Science of The Total Environment*, 610–611, 1103-1112. <https://doi.org/10.1016/j.scitotenv.2017.08.171>
- Besse, J.-P., Coquery, M., Lopes, C., Chaumot, A., Budzinski, H., Labadie, P., & Geffard, O. (2013). Caged *Gammarus fossarum* (Crustacea) as a robust tool for the characterization of bioavailable contamination levels in continental waters: Towards the determination of threshold values. *Water Research*, 47(2), 650-660. <https://doi.org/10.1016/j.watres.2012.10.024>
- Boxall, A. B. A., & Maltby, L. (1997). The effects of motorway runoff on freshwater ecosystems: 3. Toxicant confirmation. *Archives of Environmental Contamination and Toxicology*, 33(1), 9-16. <https://doi.org/10.1007/s002449900216>
- Brettschneider, D. J., Misovic, A., Schulte-Oehlmann, U., Oetken, M., & Oehlmann, J. (2019a). Detection of chemically induced ecotoxicological effects in rivers of the Nidda catchment (Hessen, Germany) and development of an ecotoxicological, Water Framework Directive–compliant assessment system. *Environmental Sciences Europe*, 31(1), 7. <https://doi.org/10.1186/s12302-019-0190-4>
- Brettschneider, D. J., Misovic, A., Schulte-Oehlmann, U., Oetken, M., & Oehlmann, J. (2019b). Poison in paradise: Increase of toxic effects in restored sections of two rivers jeopardizes the success of hydromorphological restoration measures. *Environmental Sciences Europe*, 31(1), 36. <https://doi.org/10.1186/s12302-019-0218-9>
- Brettschneider, D. J., Spring, T., Blumer, M., Welge, L., Dombrowski, A., Schulte-Oehlmann, U., Sundermann, A., Oetken, M., & Oehlmann, J. (2023). Much effort, little success: Causes for the



- low ecological efficacy of restoration measures in German surface waters. *Environmental Sciences Europe*, 35(1), 31. <https://doi.org/10.1186/s12302-023-00736-1>
- Brzezinska, A., Sakson, G., & Olejnik, D. (2023). Analysis of (anti-)oestrogenic and (anti-)androgenic activities in wastewater from the Lodz sewer system. *Water*, 15(13), 2454. <https://doi.org/10.3390/w15132454>
- Danish Ministry of the Environment. (2005). *Survey of Estrogenic Activity in the Danish Aquatic Environment*. Report of Environmental Project No. 977 2005. Copenhagen. <https://www2.mst.dk/udgiv/publications/2005/87-7614-505-0/pdf/87-7614-506-9.pdf>
- Denison, M. S., & Nagy, S. R. (2003). Activation of the aryl hydrocarbon receptor by structurally diverse exogenous and endogenous chemicals. *Annual Review of Pharmacology and Toxicology*, 43(1), 309-334. <https://doi.org/10.1146/annurev.pharmtox.43.100901.135828>
- Duft, M., Schulte-Oehlmann, U., Weltje, L., Tillmann, M., & Oehlmann, J. (2003). Stimulated embryo production as a parameter of estrogenic exposure via sediments in the freshwater mudsnail *Potamopyrgus antipodarum*. *Aquatic Toxicology*, 64(4), 437-449. [https://doi.org/10.1016/S0166-445X\(03\)00102-4](https://doi.org/10.1016/S0166-445X(03)00102-4)
- Duft, M., Schmitt, C., Bachmann, J., Brandelik, C., Schulte-Oehlmann, U., & Oehlmann, J. (2007). Prosobranch snails as test organisms for the assessment of endocrine active chemicals - An overview and a guideline proposal for a reproduction test with the freshwater mudsnail *Potamopyrgus antipodarum*. *Ecotoxicology*, 16(1), 169-182. <https://doi.org/10.1007/s10646-006-0106-0>
- Enns, D., Cunze, S., Baker, N. J., Oehlmann, J., & Jourdan, J. (2023). Flushing away the future: The effects of wastewater treatment plants on aquatic invertebrates. *Water Research*, 243, 120388. <https://doi.org/10.1016/j.watres.2023.120388>
- European Commission. (2019). *Report from the Commission to the European Parliament and the Council on the Implementation of the Water Framework Directive (2000/ 60/EC) and the Floods Directive (2007/60/EC). Second River Basin Management Plans. First Flood Risk Management Plans*. Brussels.
- German Environment Agency. (2017). *Waters in Germany Status and assessment*. Dessau-Roßlau. [https://www.gewaesser-bewertung.de/files/171018\\_uba\\_gewasserdlt\\_engl\\_bf.pdf](https://www.gewaesser-bewertung.de/files/171018_uba_gewasserdlt_engl_bf.pdf)
- Giebner, S., Ostermann, S., Straskraba, S., Oetken, M., Oehlmann, J., & Wagner, M. (2018). Effectivity of advanced wastewater treatment: Reduction of *in vitro* endocrine activity and mutagenicity but not of *in vivo* reproductive toxicity. *Environmental Science and Pollution Research*, 25(5), 3965-3976. <https://doi.org/10.1007/s11356-016-7540-1>
- Gosset, A., Wiest, L., Fildier, A., Libert, C., Giroud, B., Hammada, M., Hervé, M., Sibeud, E., Vulliet, E., Polomé, P., & Perrodin, Y. (2021). Ecotoxicological risk assessment of contaminants of emerging concern identified by “suspect screening” from urban wastewater treatment plant effluents at a territorial scale. *Science of the Total Environment*, 778, 146275. <https://doi.org/10.1016/j.scitotenv.2021.146275>
- Hagiwara, Y., Watanabe, M., Yoshimitsu, O., Toshio, S., & Takehiko, N. (1993). Specificity and sensitivity of *Salmonella typhimurium* YG1041 and YG1042 strains possessing elevated levels of both nitroreductase and acetyltransferase activity. *Mutation Research*, 291, 171-180. [https://doi.org/10.1016/0165-1161\(93\)90157-u](https://doi.org/10.1016/0165-1161(93)90157-u)
- Halle, M., & Müller, A. (2014). *Korrelationen zwischen biologischen Qualitätskomponenten und allgemeinen chemischen und physikalisch-chemischen Parametern in Fließgewässern*. Final

report of LAWA project O 3.12. 190 pages.  
<http://www.laenderfinanzierungsprogramm.de/static/LFP/Dateien/LAWA/AO/o-3-12-endbericht.pdf>

Harth, F. U. R., Arras, C., Brettschneider, D. J., Misovic, A., Oehlmann, J., Schulte-Oehlmann, U., & Oetken, M. (2018). Small but with big impact? Ecotoxicological effects of a municipal wastewater effluent on a small creek. *Journal of Environmental Science and Health, Part A*, 53(13), 1149-1160. <https://doi.org/10.1080/10934529.2018.1530328>

HLNUG. (2022). *Hessisches Karteninformationssystem: WRR-Viewer*. Hessian Agency for Nature Conservation, Environment and Geology, Wiesbaden. <https://wrrl.hessen.de/>

HMUKLV. (2018a). *Spurenstoffstrategie Hessische Ried*. Hessian Ministry for the Environment, Climate Protection, Agriculture and Consumer Protection, Wiesbaden. [https://umwelt.hessen.de/sites/umwelt.hessen.de/files/2021-07/spurenstoffstrategie\\_hessisches\\_ried.pdf](https://umwelt.hessen.de/sites/umwelt.hessen.de/files/2021-07/spurenstoffstrategie_hessisches_ried.pdf)

HMUKLV. (2018b). *Spurenstoffstrategie Hessisches Ried. Anhang*. Hessian Ministry for the Environment, Climate Protection, Agriculture and Consumer Protection, Wiesbaden. [https://umwelt.hessen.de/sites/umwelt.hessen.de/files/2021-07/anhang\\_zur\\_spurenstoffstrategie\\_hessisches\\_ried.pdf](https://umwelt.hessen.de/sites/umwelt.hessen.de/files/2021-07/anhang_zur_spurenstoffstrategie_hessisches_ried.pdf)

ISO. (2007). *ISO 11348-3. Water quality: Determination of the inhibitory effect of water samples on the light emission of Vibrio fischeri (Luminescent bacteria test) - Part 3: Method using freeze-dried bacteria*. Geneva.

ISO. (2012). *ISO 11350. Water quality: Determination of the genotoxicity of water and waste water - Salmonella/microsome fluctuation test (Ames fluctuation test)*. Geneva.

Jähnig, S. C., Baranov, V., Altermatt, F., Cranston, P., Friedrichs-Manthey, M., Geist, J., He, F., Heino, J., Hering, D., Hölker, F., Jourdan, J., Kalinkat, G., Kiesel, J., Leese, F., Maasri, A., Monaghan, M. T., Schäfer, R. B., Tockner, K., Tonkin, J. D., & Domisch, S. (2021). Revisiting global trends in freshwater insect biodiversity. *WIREs Water*, 8(2), e1506. <https://doi.org/10.1002/wat2.1506>

Janeczko, A. (2021). Estrogens and androgens in plants: The last 20 years of studies. *Plants*, 10(12), 2783. <https://doi.org/10.3390/plants10122783>

Jobling, S., Casey, D., Rodgers-Gray, T., Oehlmann, J., Schulte-Oehlmann, U., Pawlowski, S., Baunbeck, T., Turner, A. P., & Tyler, C. R. (2004). Comparative responses of molluscs and fish to environmental estrogens and an estrogenic effluent. *Aquatic Toxicology*, 66(2), 207-222. <https://doi.org/10.1016/j.aquatox.2004.01.002>

Khan, S., Naushad, Mu., Govarthan, M., Iqbal, J., & Alfadul, S. M. (2022). Emerging contaminants of high concern for the environment: Current trends and future research. *Environmental Research*, 207, 112609. <https://doi.org/10.1016/j.envres.2021.112609>

Körner, W., Bolz, U., Süßmuth, W., Hiller, G., Schuller, W., Hanf, V., & Hagenmaier, H. (2000). Input/output balance of estrogenic active compounds in a major municipal sewage plant in Germany. *Chemosphere*, 40(9-11), 1131-1142. [https://doi.org/10.1016/S0045-6535\(99\)00362-8](https://doi.org/10.1016/S0045-6535(99)00362-8)

Kusk, K. O., Krüger, T., Long, M., Taxvig, C., Lykkesfeldt, A. E., Frederiksen, H., Andersson, A.-M., Andersen, H. R., Hansen, K. M. S., Nellemann, C., & Bonefeld-Jørgensen, E. C. (2011). Endocrine potency of wastewater: Contents of endocrine disrupting chemicals and effects measured by *in vivo* and *in vitro* assays. *Environmental Toxicology and Chemistry*, 30(2), 413-426. <https://doi.org/10.1002/etc.385>

- LANUV. (2015). *Fliessgewässertypenkarten Nordrhein-Westfalens: LANUV-Arbeitsblatt 25*. Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, Recklinghausen. 105 pages. [https://www.lanuv.nrw.de/fileadmin/lanuvpubl/4\\_arbeitsblaetter/40025.pdf](https://www.lanuv.nrw.de/fileadmin/lanuvpubl/4_arbeitsblaetter/40025.pdf)
- Li, J., Wang, Z., Ma, M., & Peng, X. (2010). Analysis of environmental endocrine disrupting activities using recombinant yeast assay in wastewater treatment plant effluents. *Bulletin of Environmental Contamination and Toxicology*, 84(5), 529-535. <https://doi.org/10.1007/s00128-010-0004-2>
- Long, M., & Bonefeld-Jørgensen, E. C. (2012). Dioxin-like activity in environmental and human samples from Greenland and Denmark. *Chemosphere*, 89(8), 919-928. <https://doi.org/10.1016/j.chemosphere.2012.06.055>
- Lopez, F. J., Pitarch, E., Botero-Coy, A. M., Fabregat-Safont, D., Ibáñez, M., Marin, J. M., Peruga, A., Ontañón, N., Martínez-Morcillo, S., Olalla, A., Valcárcel, Y., Varó, I., & Hernández, F. (2022). Removal efficiency for emerging contaminants in a WWTP from Madrid (Spain) after secondary and tertiary treatment and environmental impact on the Manzanares River. *Science of The Total Environment*, 812, 152567. <https://doi.org/10.1016/j.scitotenv.2021.152567>
- Macova, M., Toze, S., Hodggers, L., Mueller, J. F., Bartkow, M., & Escher, B. I. (2011). Bioanalytical tools for the evaluation of organic micropollutants during sewage treatment, water recycling and drinking water generation. *Water Research*, 45(14), 4238-4247. <https://doi.org/10.1016/j.watres.2011.05.032>
- Magdeburg, A., Stalter, D., Schlüsener, M., Ternes, T., & Oehlmann, J. (2014). Evaluating the efficiency of advanced wastewater treatment: Target analysis of organic contaminants and (geno-)toxicity assessment tell a different story. *Water Research*, 50, 35-47. <https://doi.org/10.1016/j.watres.2013.11.041>
- Maltby, L., Forrow, D. M., Boxall, A. B. A., Calow, P., & Betton, C. I. (1995). The effects of motorway runoff on freshwater ecosystems: 1. Field study. *Environmental Toxicology and Chemistry*, 14(6), 1079-1092. <https://doi.org/10.1002/etc.5620140620>
- Maron, D. M., & Ames, B. N. (1983). Revised methods for the *Salmonella* mutagenicity test. *Mutation Research/Environmental Mutagenesis and Related Subjects*, 113(3-4), 173-215. [https://doi.org/10.1016/0165-1161\(83\)90010-9](https://doi.org/10.1016/0165-1161(83)90010-9)
- Mestankova, H., Schirmer, K., Canonica, S., & Von Gunten, U. (2014). Development of mutagenicity during degradation of N-nitrosamines by advanced oxidation processes. *Water Research*, 66, 399-410. <https://doi.org/10.1016/j.watres.2014.08.012>
- OECD. (2016). *Test No. 242: Potamopyrgus antipodarum Reproduction Test*. OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264264311-en>.
- Prasse, C., Stalter, D., Schulte-Oehlmann, U., Oehlmann, J., & Ternes, T. A. (2015). A critical review on the chemical and biological assessment of current wastewater treatment technologies. *Water Research*, 87, 237-270. <https://doi.org/10.1016/j.watres.2015.09.023>
- Reemtsma, T., Weiss, S., Mueller, J., Petrovic, M., González, S., Barcelo, D., Ventura, F., & Knepper, T. P. (2006). Polar pollutants entry into the water cycle by municipal wastewater: A European perspective. *Environmental Science & Technology*, 40(17), 5451-5458. <https://doi.org/10.1021/es060908a>
- Reifferscheid, G., Buchinger, S., Cao, Z., & Claus, E. (2011). Identification of mutagens in freshwater sediments by the Ames-fluctuation assay using nitroreductase and acetyltransferase

- overproducing test strains. *Environmental and Molecular Mutagenesis*, 52(5), 397-408.  
<https://doi.org/10.1002/em.20638>
- Schmidt, C. K., & Brauch, H.-J. (2008). *N,N*-Dimethylsulfamide as precursor for *N*-nitrosodimethylamine (NDMA) formation upon ozonation and its fate during drinking water treatment. *Environmental Science & Technology*, 42(17), 6340-6346.  
<https://doi.org/10.1021/es7030467>
- Schneider, I., Abbas, A., Bollmann, A., Dombrowski, A., Knopp, G., Schulte-Oehlmann, U., Seitz, W., Wagner, M., & Oehlmann, J. (2020). Post-treatment of ozonated wastewater with activated carbon and biofiltration compared to membrane bioreactors: Toxicity removal *in vitro* and in *Potamopyrgus antipodarum*. *Water Research*, 185, 116104.  
<https://doi.org/10.1016/j.watres.2020.116104>
- Stalter, D., Magdeburg, A., & Oehlmann, J. (2010). Comparative toxicity assessment of ozone and activated carbon treated sewage effluents using an *in vivo* test battery. *Water Research*, 44(8), 2610-2620. <https://doi.org/10.1016/j.watres.2010.01.023>
- Stalter, D., Magdeburg, A., Quednow, K., Botzat, A., & Oehlmann, J. (2013). Do contaminants originating from state-of-the-art treated wastewater impact the ecological quality of surface waters? *PLoS ONE*, 8(4), e60616. <https://doi.org/10.1371/journal.pone.0060616>
- Tang, J. Y. M., McCarty, S., Glenn, E., Neale, P. A., Warne, M. S. J., & Escher, B. I. (2013). Mixture effects of organic micropollutants present in water: Towards the development of effect-based water quality trigger values for baseline toxicity. *Water Research*, 47(10), 3300-3314.  
<https://doi.org/10.1016/j.watres.2013.03.011>
- Ternes, T., Joss, A., & Oehlmann, J. (2015). Occurrence, fate, removal and assessment of emerging contaminants in water in the water cycle (from wastewater to drinking water). *Water Research*, 72, 1-2. <https://doi.org/10.1016/j.watres.2015.02.055>
- Ternes, T. A., Prasse, C., Eversloh, C. L., Knopp, G., Cornel, P., Schulte-Oehlmann, U., Schwartz, T., Alexander, J., Seitz, W., Coors, A., & Oehlmann, J. (2017). Integrated evaluation concept to assess the efficacy of advanced wastewater treatment processes for the elimination of micropollutants and pathogens. *Environmental Science & Technology*, 51(1), 308-319.  
<https://doi.org/10.1021/acs.est.6b04855>
- Tsutsumi, N., Inami, K., & Mochizuki, M. (2010). Activation mechanism for N-nitroso-N-methylbutylamine mutagenicity by radical species. *Bioorganic & Medicinal Chemistry*, 18(23), 8284-8288. <https://doi.org/10.1016/j.bmc.2010.10.002>
- Umwelt Bundesamt. (2017). *Waters in Germany: Status and assessment*. German Environment Agency, Dessau-Roßlau. 132 pages.  
[https://www.umweltbundesamt.de/sites/default/files/medien/377/publikationen/171018\\_uba\\_gewasserdtl\\_engl\\_bf.pdf](https://www.umweltbundesamt.de/sites/default/files/medien/377/publikationen/171018_uba_gewasserdtl_engl_bf.pdf)
- Vasquez, M. I., & Fatta-Kassinos, D. (2013). Is the evaluation of “traditional” physicochemical parameters sufficient to explain the potential toxicity of the treated wastewater at sewage treatment plants? *Environmental Science and Pollution Research*, 20(6), 3516-3528.  
<https://doi.org/10.1007/s11356-013-1637-6>
- Völker, J., Castronovo, S., Wick, A., Ternes, T. A., Joss, A., Oehlmann, J., & Wagner, M. (2016). Advancing biological wastewater treatment: Extended anaerobic conditions enhance the removal of endocrine and dioxin-like activities. *Environmental Science & Technology*, 50(19), 10606-10615. <https://doi.org/10.1021/acs.est.5b05732>

- Völker, J., Vogt, T., Castronovo, S., Wick, A., Ternes, T. A., Joss, A., Oehlmann, J., & Wagner, M. (2017). Extended anaerobic conditions in the biological wastewater treatment: Higher reduction of toxicity compared to target organic micropollutants. *Water Research*, 116, 220-230. <https://doi.org/10.1016/j.watres.2017.03.030>
- Wolf, Y., Oster, S., Shuliakovich, A., Brückner, I., Dolny, R., Linnemann, V., Pinnekamp, J., Hollert, H., & Schiwy, S. (2022). Improvement of wastewater and water quality via a full-scale ozonation plant? – A comprehensive analysis of the endocrine potential using effect-based methods. *Science of the Total Environment*, 803, 149756. <https://doi.org/10.1016/j.scitotenv.2021.149756>
- Yang, Y., Zhang, X., Jiang, J., Han, J., Li, W., Li, X., Yee Leung, K. M., Snyder, S. A., & Alvarez, P. J. J. (2022). Which micropollutants in water environments deserve more attention globally? *Environmental Science & Technology*, 56(1), 13-29. <https://doi.org/10.1021/acs.est.1c04250>
- Yu, X., Zuo, J., Li, R., Gan, L., Li, Z., & Zhang, F. (2014). A combined evaluation of the characteristics and acute toxicity of antibiotic wastewater. *Ecotoxicology and Environmental Safety*, 106, 40-45. <https://doi.org/10.1016/j.ecoenv.2014.04.035>

# **Table 1**(on next page)

Description of sampling sites

**Table 1** . Sampling sites at the WWTPs Mörfelden-Walldorf and Bickenbach and in the receiving rivers.

1

Sampling site [code]	Characteristic
Upstream [US-MW]	Geräthsbach 20 m upstream of the effluent of WWTP Mörfelden-Walldorf (49°58'14.6"N 8°32'59.3"E)
Mechanical treatment [MT-MW]	24 hours composite sample after mechanical treatment at WWTP Mörfelden-Walldorf
Biological treatment [BT-MW]	24 hours composite sample after biological treatment and phosphate precipitation at WWTP Mörfelden-Walldorf
Downstream [DS-MW]	Geräthsbach 20 m downstream of the effluent of WWTP Mörfelden-Walldorf (49°58'14.7"N 8°32'56.6"E)
Upstream [US-B]	Landbach 20 m upstream of the effluent of WWTP Bickenbach (49°45'33.9"N 8°35'52.0"E)
Mechanical treatment [MT-B]	72 hours composite sample after mechanical treatment at WWTP Bickenbach
Biological treatment [BT-B]	72 hours composite sample after biological treatment and phosphate precipitation at WWTP Bickenbach
Downstream [DS-B]	Landbach 20 m downstream of the effluent of WWTP Bickenbach (49°45'33.8"N 8°35'49.4"E)

2

## Table 2 (on next page)

Physicochemical parameters at the sampling sites

**Table 2. Mean values of physicochemical parameters during the spring and fall campaign.** All data were measured on two occasions (4 weeks difference) during each campaign. BT-L data during the spring campaign was not measured.



1

Parameter	Season	Sample sites								
		BT-L	US-MW	MT-MW	BT-MW	DS-MW	US-B	MT-B	BT-B	DS-B
Oxygen saturation [%]	Spring	--	99.1	65.0	68.5	78.7	92.6	9.00	82.5	65.0
	Fall	104	83.2	20.0	93.6	77.7	85.8	58.3	61.2	68.1
Conductivity [μS/cm]	Spring	--	781	1190	772	817	493	1560	952	967
	Fall	1080	976	1270	934	1000	532	1500	1040	932
Phosphate P [PO <sub>4</sub> <sup>3-</sup> -P mg/L]	Spring		0.04	1.44	0.04	0.03	0.05	0.68	0.02	0.02
	Fall	0.03	0.03	0.87	0.04	0.02	0.02	0.67	0.02	0.02
Ammonium-N [NH <sub>4</sub> <sup>+</sup> -N mg/L]	Spring		0.03	34.5	1.81	1.11	0.02	52.5	0.26	0.15
	Fall	0.61	0.19	44.7	0.05	0.18	0.11	23.2	0.29	0.32
Nitrite-N [NO <sub>2</sub> <sup>-</sup> -N mg/L]	Spring		0.02	0.07	0.06	0.07	0.02	0.03	0.10	0.03
	Fall	0.05	0.05	0.04	0.01	0.03	0.02	0.04	0.04	0.04
Nitrate-N [NO <sub>3</sub> <sup>-</sup> -N mg/L]	Spring		1.35	0.16	1.53	1.08	0.45	0.46	1.13	1.03
	Fall	2.21	1.72	0.13	2.55	1.78	0.51	0.32	0.93	0.67

2

# Table 3 (on next page)

Overview of the results of the mutagenicity test

Table 3 : Mutagenicity at the WWTPs Mörfelden-Walldorf and Bickenbach in the Ames fluctuation test with Salmonella typhimurium strains YG1041 and YG1042 with S9 mix. Mutagenic samples (> 20.8% revertant colonies) are labelled in red, non-mutagenic samples in green. n.a.: not analyzed due to high cytotoxicity.

YG1041+S9 mix				
Season	WWTP	Sample site	Test response	Revertant percentage mean (%)
Spring	Mörfelden Walldorf	US	+	39.3
		MT	n.a.	n.a.
		BT	+	49.7
		DS	+	56.5
	Bickenbach	US	-	4.8
		MT	n.a.	n.a.
		BT	+	49.2
		DS	+	52.3
Fall	Mörfelden Walldorf	BT-L	+	52.8
		US	-	22.1
		MT	n.a.	n.a.
		BT	+	57.5
		DS	+	54.4
	Bickenbach	US	-	15.3
		MT	n.a.	n.a.
		BT	+	47.1
		DS	+	53.3

YG1042+S9 mix				
Season	WWTP	Sample site	Test response	Revertant percentage mean (%)
Spring	Mörfelden Walldorf	US	+	41.9
		MT	n.a.	n.a.
		BT	+	47.1
		DS	+	41.4
	Bickenbach	US	-	20.5
		MT	n.a.	n.a.
		BT	+	45.0
		DS	+	42.4
Fall	Mörfelden Walldorf	BT-L	+	54.9
		US	+	32.5
		MT	n.a.	n.a.
		BT	+	52.8
		DS	+	51.3
	Bickenbach	US	+	36.2
		MT	n.a.	n.a.
		BT	+	51.8
		DS	+	56.0


1

2

# Figure 1

Map of the sampling locations


Figure 1. **Map of the sampling location of the two WWTPs in Germany.**



# Figure 2

## Baseline toxicity of water samples at the WWTPs

Baseline toxicity of water samples at the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B) and in the receiving streams. Toxicity was determined by the Microtox assay and is expressed as EC50 values, referring to the relative enrichment factor (REF) of the water. Low values indicate higher toxicity. Dashed line: non toxic limit (NTL). Significant differences refer to downstream values (DS) compared to upstream values (US) (unpaired t-test with Welch's correction for unequal variances; ★  $p < 0.05$ , ★★  $p < 0.01$ , ★★★  $p < 0.001$ ). Numbers in blue indicate the mean decrease in % of measured baseline toxicity during the biological treatment.





# Figure 3

Estrogenic activity

*Figure 3 . **Estrogenic activity at the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B) and the receiving streams, determined with the Yeast Estrogen Screen (YES).** Activities are expressed as 17 $\beta$ -estrogen equivalent concentrations (E-EQ). Dashed line: limit of quantification (LOD: 0.148 ng E-EQ/L). Significant differences refer to downstream values (DS) compared to upstream values (US) (unpaired t-test with Welch's correction for unequal variances; ★  $p < 0.05$ , ★★  $p < 0.01$ , ★★★  $p < 0.001$ ). Numbers in blue indicate the mean decrease in % of measured estrogenic activity during the biological treatment*







# Figure 4

Anti-estrogenic activity



*Figure 4. **Anti-estrogenic activity at the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B) and the receiving rivers, determined with the Yeast Anti Estrogen Screen (YEAS).** Activities are expressed as hydroxy-tamoxifen equivalent concentrations (OHT-EQ). Dashed line: limit of quantification (LOD: 1.84 mg OHT-EQ/L). None of the downstream values (DS) exhibits statistically significant differences to upstream values (US) (unpaired t-test with Welch's correction for unequal variances;  $p > 0.05$ ). Numbers in blue indicate the mean decrease in % of measured estrogenic activity during the biological treatment*



# Figure 5

AhR agonistic activity



**Figure 5. AhR agonistic activity at the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B) and in the receiving rivers, determined with the Yeast Dioxin Screen (YDS).** Activities are expressed as  $\beta$ -naphthoflavone equivalent concentrations ( $\beta$ -NF-EQ). Dashed line: limit of quantification (LOD: 33.8 ng  $\beta$ -NF-EQ/L). Significant differences refer to downstream values (DS) compared to upstream values (US) (unpaired t-test with Welch's correction for unequal variances; ★  $p < 0.05$ , ★★★  $p < 0.001$ ). Numbers in blue indicate the mean decrease in % of measured estrogenic activity during the biological treatment; skull and crossbones symbol indicate cytotoxic samples



# Figure 6

Mortality of *Gammarus fossarum*

**Figure 6. Mortality of *Gammarus fossarum* in the active monitoring campaign over one month in the upstream (US) and downstream (DS) site in the receiving rivers of the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B).** Significant differences refer to DS compared to US (Fisher's exact test; ★  $p < 0.05$ , ★★★  $p < 0.001$ ).





# Figure 7

Mortality of *Potamopyrgus antipodarum*

**Figure 7. Mortality of *Potamopyrgus antipodarum* in the active monitoring campaign over one month in the upstream (US) and downstream (DS) site in the receiving rivers of the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B).**

Significant differences refer to DS compared to US (Fisher's exact test; ★★★  $p < 0.001$ ).





# Figure 8

Number of embryos in *Potamopyrgus antipodarum*

**Figure 8. Number of embryos in *Potamopyrgus antipodarum* in the active monitoring campaign over one month in the upstream (US) and downstream (DS) site in the receiving rivers of the WWTPs Mörfelden-Walldorf (A) and Bickenbach (B).** Significant differences refer to DS compared to US (unpaired t-test with Welch's correction for unequal variances; ★★  $p < 0.01$ ).

