

Benthic and fish community composition on mesophotic reefs in Grand Cayman (#98134)

1

First submission

Guidance from your Editor

Please submit by **26 May 2024** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

8 Figure file(s)

1 Table file(s)

4 Raw data file(s)



Custom checks

Field study



Have you checked the authors [field study permits](#)?



Are the field study permits appropriate?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Benthic and fish community composition on mesophotic reefs in Grand Cayman

Lucas Le Gall¹, Jack V. Johnson¹, Alex D Chequer¹, Matt Doherty¹, Gretchen Goodbody-Gringley^{Corresp. 1}

¹ Reef Ecology And Evolution Lab, Central Caribbean Marine Institute, Little Cayman, Cayman Islands

Corresponding Author: Gretchen Goodbody-Gringley
Email address: ggoodbody@reefresearch.org

Mesophotic Coral Ecosystems (MCEs) represent unique ecological habitats that range from 30m to 150m deep, harbouring phylogenetically distinct species and offering refuge for many taxa during times of environmental stress. Yet owing to inaccessibility of ecosystems at these depths, most MCEs remain unexplored, with quantifications of ecological communities in these habitats lacking across many regions. Here, using open- and closed-circuit technical diving, we quantified benthic and fish community composition at four mesophotic reef sites (45m depth) in Grand Cayman. We show significant differences in benthic community composition over a small spatial scale driven by disparate coverage of sponges, crustose coralline algae, and sand/rubble, yet consistent patterns of macroalgal dominance representing >50% coverage at each site and low hard coral cover at an average of 2.4%. Reef fish species richness, biomass, and density was consistent across sites, however the relative contribution of individual species to community composition differed significantly. Macrocarnivores were found to be the dominant contributors to biomass, with invertivores the most speciose, and omnivores and planktivores at the highest densities, consistent with previous descriptions of mesophotic fish assemblages in other regions. Similarly, the low hard coral cover and high macroalgae and sponge cover of the benthic communities also appear ecologically similar to several described mesophotic reefs yet is not uniform across the Caribbean. The ecological organisation of Grand Cayman's MCEs may result from a variety of factors such as isolation from other major land masses, geology, local geography, and anthropogenic activity at both the local and global scale and highlight the importance of continued exploration and documentation of MCE communities.

**Benthic and fish community composition on mesophotic reefs in
Grand Cayman**

Lucas Le Gall¹, Jack V Johnson¹, Alex Chequer¹, Matt Doherty¹, Gretchen Goodbody-Gringley^{1*}

¹Reef Ecology and Evolution Lab, Central Caribbean Marine Institute, Little Cayman, Cayman Islands

*Corresponding Author:

Gretchen Goodbody-Gringley

CCMI, Box 37, Little Cayman, Cayman Islands KY3-2501

Email address: ggoodbody@reefresearch.org


Abstract

Mesophotic Coral Ecosystems (MCEs) represent unique ecological habitats that range from 30m to 150m deep, harbouring phylogenetically distinct species and offering refuge for many taxa during times of environmental stress. Yet owing to inaccessibility of ecosystems at these depths, most MCEs remain unexplored, with quantifications of ecological communities in these habitats lacking across many regions. Here, using open- and closed-circuit technical diving, we quantified benthic and fish community composition at four mesophotic reef sites (45m depth) in Grand Cayman. We show significant differences in benthic community composition over a small spatial scale driven by disparate coverage of sponges, crustose coralline algae, and sand/rubble, yet consistent patterns of macroalgal dominance representing >50% coverage at each site and low hard coral cover at an average of 2.4%. Reef fish species richness, biomass, and density was consistent across sites, however the relative contribution of individual species to community composition differed significantly. Macrocarnivores were found to be the dominant contributors to biomass, with invertivores the most speciose, and omnivores and planktivores at the highest densities, consistent with previous descriptions of mesophotic fish assemblages in other regions. Similarly, the low hard coral cover and high macroalgae and sponge cover of the benthic communities also appear ecologically similar to several described mesophotic reefs yet is not uniform across the Caribbean. The ecological organisation of Grand Cayman's MCEs may result from a variety of factors such as isolation from other major land masses, geology, local geography, and anthropogenic activity at both the local and global scale and highlight the importance of continued exploration and documentation of MCE communities.

Introduction

Mesophotic Coral Ecosystems (MCEs) represent globally distributed, deeper water ecosystems, typically found between depths of 30 and 150 meters (Pyle and Copus 2019). Comprised of light-dependent scleractinian corals, these ecosystems also contain a variety of taxa including sponges, macroalgae, encrusting algae, and soft coral colonies (Hinderstein et al. 2010). Despite the ecological importance of MCEs for harbouring marine biodiversity, which contributes to coastal ecosystem services (Holstein et al. 2019), MCEs have historically received less attention than their shallow water counterparts due primarily to limited access using traditional SCUBA. However, as diving and submersible technology has advanced, so has access to the mesophotic, resulting in a dramatic increase in scientific investigation and biodiversity assessments (Pyle and Copus, 2019). This recent surge has primarily focused on Caribbean MCEs, which are typically located in regions with high anthropogenic activity, comprising roughly 55% of MCE focused publications as of 2018 (Laverick et al. 2018). These studies have been crucial in documenting the biodiversity of MCEs, highlighting the ecologically and biologically distinct nature of these systems (Kahng et al. 2010) with prominent differences in both benthic (Stefanoudis, et al. 2019, Lesser et al, 2018) and fish assemblages (Garcia-Sais et al. 2008, Stefanoudis, et al. 2019, Lesser et al, 2018, Pinheiro et al, 2016) compared to shallow water ecosystems. Within MCEs, observed breaks in biodiversity are documented at approximately 60m, distinguishing the upper and lower mesophotic zones (Lesser et al. 2018). While the defined depth range is influenced by topography, light intensity is also a major driving factor affecting community composition and defining species boundaries (Perez Castro et al. 2022; Laverick et al. 2017). Such physical factors of the environment differ among and within locations, however, resulting in strong differences in communities and distributions among various MCEs, warranting continued assessments of community composition.

The benthic composition of MCEs of the Cayman Islands has been the focus of previous work by Slattery & Lesser (2019), who focused primarily on sponge communities over limited spatial distribution and described a “sponge belt” (Slattery & Lesser, 2019). Mesophotic sponges in Grand Cayman were also the focus of Macartney et al. (2021), whose work was concentrated on a single species. While Carpenter et al. (2022) assessed the benthic composition of corals and key functional groups on MCEs across Little Cayman Island, their study did not include Grand Cayman. Importantly, none of the previous mesophotic studies in the Cayman Islands included assessments of fish communities, nor the interaction of benthic habitat and fish assemblages, highlighting a critical knowledge gap in our understanding of these ecosystems in Grand Cayman.

Variations in benthic habitat often corresponds to changes in fish community composition. For example, in the Seychelles shallow reefs with high macroalgal cover had higher densities of generalist fish species and lacked specialists, whereas reefs with high coral cover had higher abundances of specialised corallivores species (Chong-Seng et al, 2012). In the Philippines, a positive association was found between hard coral cover and biomass of planktivores, piscivores, and corallivores, while a negative association was found between hard coral cover and detritivores and sand feeders (Russ et al, 2021). In the Caribbean, studies have found a positive correlation between macroalgae density and herbivorous fish assemblages on shallow reefs but fail to show an impact of coral cover on fish populations (Sandin et al, 2008,  mesophotic studies

in the Gulf of Mexico have also shown a difference of fish assemblages in relation with benthic composition, with high fish biomass associated with coralline algal reefs (Voss et al, 2014). As these patterns vary across locations and depths, it is important to include assessments of both benthic habitat and fish communities to accurately characterize MCEs, which was previously lacking for MCEs in Grand Cayman.

Here, we characterize the biodiversity and community composition of mesophotic reefs across several sites on Grand Cayman Island. Using benthic photogrammetry and *in situ* visual surveys we provide a holistic description of benthic community composition, including scleractinian corals and key functional groups, as well as of fish community assemblages. Given the hypothesized role of MCEs as areas of refuge for preserving biodiversity under future climate change scenarios (e.g. DRRH; Bongaerts et al. 2010), baseline characterizations of these habitats are crucial for increasing our understanding of mesophotic ecosystem function and informing marine spatial planning and management.

Materials and Methods:

The present study was carried out off the coast of Grand Cayman (19.329858 -81.252361), the biggest and most populated of the 3 islands that compose the Cayman Islands (Cayman Islands' 2021 Census Report, ESO). Four sites situated around the West Bay peninsula were surveyed: Ghost Mountain (19.402300 -81.385680), Roundabout (19.384333 -81.318300), La Mesa (19.321680 -81.393080), and Lighthouse Point (19.372954 -81.421855) (Fig. 1). The sites were situated off the wall, with a steep slope except for lighthouse point that is characterized by a more gradual slope (personal observation). They were composed of an alternance of hard-bottom and sandy substrate.

Fish surveys were conducted along 30m by 2m transects at a depth of 45m (n=3-5 site; total area 60m²). All fish encountered along the transects were identified to species, counted, and categorised into size classes based on total length: 0-5, 5-10, 10-20, 20-30, 30-40, and >40 cm. Following the surveys, fish data were classified into one of five food web classes: herbivore, invertivore, omnivore, macrocarnivore, and planktivore based on data available on www.fishbase.org (Houde and Zastrow 1993; Welcomme 1988), a data base by Froese and Pauly (2000). Density was calculated as the total number of individuals per transect (no. of fish/60m² transect). Biomass was calculated using the formula:

$$\frac{(a \times L^b)}{N}$$

Where L is the estimated length using the upper end of the size class category, and N is the number of individuals of that size class (Bohnsack and Harper 1988; Kulbicki et al. 1993). Values for a and b were obtained for each species on www.fishbase.org (Houde and Zastrow, 1993; Welcomme, 1988). When these values were not available for a particular species, data for a congeneric species were used. Species richness is the total number of species per transect and diversity is based on the Shannon-Weaver index of diversity (Shannon & Weaver, 1949):

$$- \sum P_i (\log P_i)$$

with P_i being the proportion of each species in the sample.

At each of the four sites, a 10m x 10m plot was laid on the benthos using transect tapes, with the lower edge at 45m depths. Time lapse images were taken of benthos within each plot at one second intervals by two divers that swam in a lawnmower pattern approximately 1m above the benthos, using two GoPro Hero 10 cameras mounted onto a PVC frame with 2 lights (Sola Video 2500 Flood). Eight targets were laid haphazardly on the substrate within the plot to assist with building a digital 3D model of the studied area using the software AGIsoft Metashape Pro (version 1.8.3). This was then converted into a 2D orthomosaics and exported as a PNG that was subsequently split into ~1m² replicate quadrats using Photoshop (version 24.3) and scaled on the pictured measuring tape. ImageJ (version 1.53t) software was then used to scale each image for further analysis. Within each quadrat all scleractinian corals were identified to the highest taxonomic level possible based on image quality. Species within the general *Agaricia*, *Siderastrea*, and *Orbicella*, were limited to genus. Surface area of each colony was then

measured using the freehand tool (in cm²). Total surface area of all corals relative to the size of the quadrat was then used to calculate percent coral cover.

Benthic composition of other key functional groups was then assessed visually for eight categories: Crustose Coralline Algae (CCA), Soft Coral (SC), Sand and Rubble (S/R), Turf Algae (TA), Recently Dead Coral (RDC), Macroalgae (MA), and Other (soft coral plumes, fish targets, blurry and indistinguishable area, etc.). Measurements and visual assessment of the functional groups were conducted by the same person throughout the data collection phase to limit bias and errors.

Statistical analyses were conducted on R version 4.1.0 (R Core Team, 2023).

Fish density, biomass, diversity, coral diversity index and richness were compared among sites and among food webs with non-parametric Kruskal-Wallis' tests (Kruskal and Wallis 1952), as they did not meet the assumption of normality (Shapiro and Wilk 1965), followed by post-hoc analysis using the Dunns test (Dunn 1964) in the FSA package (Ogle, 2021). The fish coral diversity index met the assumption of normality (Shapiro and Wilk 1965) and were compared among sites with a one-way analysis of variance (ANOVA), followed by a post hoc analysis using Tuckey's test (Tuckey 1949).

Benthic functional group composition among sites did not meet the assumptions of normality (Shapiro and Wilk 1965) and was therefore compared using a Kruskal Wallis with a Dunn's test as a post hoc. "Other", while being measured as a functional group, is not integrated into any of the mentioned statistical analyses. By being an imprecise category that is compensating for a various array of non-benthic category measures or photomosaic inconsistency, it was judged irrelevant to the aim of the study and the measured percentages have been scaled accordingly.

To assess dissimilarity in the benthic composition and fish community composition among sites, we used non-Metric Multi-Dimensional scaling (nMDS) models from the Vegan package (Oksanen et al. 2011). Models were based on a Bray-Curtis dissimilarity matrix of community composition for both benthic composition, fish species composition, and reef fish trophic community. A dissimilarity matrix rather than distance matrix (Jaccard 1901) is preferable for our dataset which reflects community composition, as Bray-Curtis dissimilarity has less sensitivity to the absence of species within the matrix compared to the distance (metric) based counterparts. To statistically compare the dissimilarity between sites for community composition, we used a PERMANOVA test with the dissimilarity matrix as the response, and site as the predicting factor.

Results

In total 48 fish species were recorded across all four sites ranging from 30 at Lighthouse Point to 24 at Roundabout, however, median species richness per transects did not differ significantly among sites ($P>0.05$; Kruskal Wallis, Fig. 2A). However, according to the species accumulation curve, the number of surveys conducted in [redacted] enough to account for the full species composition, and thus this is likely an underestimate of diversity (Supplemental Fig. S [redacted]). Fish biomass also did not differ significantly among sites ($P>0.05$; Kruskal Wallis), Lighthouse Point had the highest median biomass (5936.296 ± 1026.23 g/transect), while Ghost Mountain had the lowest (3211.59 ± 797.70 g/transect) (Fig. 2B). The median fish density was significantly different ($X^2=-2.74$, $P=0.03$) between Lighthouse Point (157 ± 30.39 fish per transect) and La Mesa (63 ± 10.08 fish per transect) (Fig. 2C). Shannon-Weiner diversity index was significantly different between Ghost Mountain and La Mesa ($X^2= 0.726$, $P=0.019$), being respectively the least diverse with a median index value of 1.65 ± 0.14 and the most diverse with a median value of 2.31 ± 0.15 (Fig. 2D).

For trophic guilds specifically, invertivores were the most species rich group of fish with a median of 10 ± 1.31 species recorded and planktivores were the least with a median species richness of 3 ± 0.4 (Fig. 3A). Macrocarivores had the highest median biomass (6693.41 ± 1434.09 g/transect; Fig. 3B) while planktivores had the lowest median biomass (789.6238 ± 394.81 g/transect). Omnivores had the highest median density (200 ± 4.42 of fish per transect; Fig. 3C). Significant differences were found in species richness ($X^2=12.169$, $P=0.016$), fish density ($X^2=17.031$, $P=0.001$) and biomass ($X^2=9.622$, $P=0.047$) among trophic guilds. Higher species richness was observed for invertivores compared to planktivores ($X^2=3.13$, $P=0.017$), while macrocarivores had significantly lower density compared to planktivores ($X^2=-2.816$, $P=0.043$) and omnivores ($X^2=-3.697$, $P=0.002$). Shannon diversity index values did not differ significantly ($P>0.05$, ANOVA) among trophic groups.

Across sites, 20 families of fish were recorded (Fig. 4). The basslet family (*Grammatidae*), had the highest density per square meter with a mean value of 19.048 ± 7.554 % (Fig. 4A), accounting for 0.59 ± 0.317 % of the total fish biomass (Fig. 4B). Members of the snapper family (*Lutjanidae*) had the highest average biomass (10.432 ± 3.974 %) but only accounted for 0.071 ± 0.031 % of the recorded density (Fig 4A and 4B).

Across the surveyed sites, *Gramma melacara*, had the highest frequency by density at three sites, with 53.78% at Ghost Mountain, 35.91% at Roundabout and 24.75% at Lighthouse Point, while *Chromis cyanea* had the highest frequency at La Mesa of 20.61% (Fig. 4C). *Scarus taeniopterus* had the highest frequency by biomass at Ghost Mountain at 28.02%, while *Lutjanus analis* had the highest frequency by biomass at La Mesa at 41.85%, and Roundabout at 25.46% (Fig. 4D). At Lighthouse Point, *Lutjanus mahogoni* had the highest frequency by biomass at 20.91% (Fig. 4D). Out of the 48 observed species, only 13 were found across all sites and 22 species were endemic to one site.


Macroalgae percent cover was significantly higher than all other functional groups across all four sites (Fig 5; Table 1) with a median surface coverage of 63.378 ± 1.125 %, while hard coral cover was significantly lower than all other functional groups, with a median of 0.662 ± 0.285 % (Fig.


5; Table 1). Sponge cover did not differ significantly from either sand/rubble or CCA cover, however, sand/rubble cover was significantly higher than CCA (Fig. 5; Table 1).


Agaricia was the dominant coral genus at all four sites (Fig. 6) ranging from 77% at Ghost Mountain to 54% at Lighthouse Point. *Siderastrea* and *Orbicella* were present in moderate frequency at all sites, *Montastraea* and *Porites* were consistently present but in low coverage, and *Mycetophyllia* was only recorded in La Mesa. Shannon-Weiner diversity of coral species showed no significant differences between sites ($P > 0.05$, Kruskal-Wallis)

Multivariate analyses revealed the benthic composition of sites were significantly dissimilar (PERMNOVA value = 11.69, $P = 0.0001$). Lighthouse Point appears to be the most unique site based on benthic composition (Fig 6A), related to high sand/rubble coverage (Fig 6B) the trophic composition of fish communities (Fig 6C) was also significantly different among sites (PERMANOVA, F value = 2.76, $P = 0.0058$) as was the species composition (Fig. 6E; PERMANOVA, F value = 2.55, $P = 0.003$)

Discussion

Here, we provide the first characterization of mesophotic reefs in the Cayman Islands that combines benthic habitat and fish assemblages. Our results show that fish populations are primarily composed of basslet (*Grammidae*) and damselfish (*Pomacentridae>Stegastinae*) species, with species of wrasse (*Labridae*) and chromis (*Pomacentridae>Chrominae*) also present in high relative abundance. Although present in lower relative densities, species of snapper (*Lutjanidae*) and parrotfish (*Scaridae*) represent the highest contribution to community biomass on the mesophotic reefs surveyed. The benthos at 45m is dominated by macroalgae and sand/rubble, which can be generally described as algae-covered hard substrate dispersed among patches of sand and rubble. Sponges, seafans, and crustose coralline algae are also found in high frequency  however, hard coral cover represents one of the lowest percent covers. Among the corals present, the dominant contributing species belong to the genera *Agaricia*, *Siderastrea*, and *Orbicella*.

Among all the mesophotic sites surveyed, *Gramma melacara* (blackcap basslet) and *Chromis insolata* (sunshinefish) are the most abundant in terms of number of fish. These species, however, are absent on shallow reefs in the same region of Grand Cayman (Johnson et al. 2023). Previous surveys of Caribbean fish communities also note high densities of *C. insolata* on mesophotic reefs with their presence generally lacking on shallow reefs, suggesting this species is a mesophotic specialist (Baldwin et al. 2018; Garcia-Hernandez et al. 2018; Gress et al. 2018; Pinheiro et al. 2016). Likewise, *G. melacara* is also a mesophotic specialist and is considered common on MCEs in the Western/Central Caribbean (Chasqui et al. 2020; Dustan and Lang 2019), however it is not present on MCE species lists from Cozumel (Gress et al. 2018), Curacao (Baldwin et al. 2018; Pinheiro et al. 2016), or Bermuda (Pinheiro et al. 2016; Goodbody-Gringley et al. 2019), and thus its geographic distribution may be limited to the Central Caribbean. Although species of snapper (*Lutjanidae*) are the most abundant by biomass across all sites, this is not consistent among the 4 sites surveyed, with these species being generally absent from Ghost Mountain. Rather, parrotfish are the dominant group at this site, where the most abundant species by biomass is *Scarus taenopterus*. Within the other three sites, variation also exists among the dominant species of snapper with *Lutjanus analis* being the highest contributor to biomass at La Mesa and Roundabout, while *Lutjanus mahogoni*, is the highest contributor at Lighthouse Point. As macrocarnivores, such as snappers, are often transient it is likely that variations in species contributions of these groups and the absence of snappers from surveys at Ghost Mountain and may be an artifact of the visual survey method, which is limited to a short period of time and may miss transient species 

The overall benthic cover of MCEs in Grand Cayman is similar to mesophotic reefs at other Central Caribbean locations (Dustan and Lang 2019; Reed et al. 2019; Carpenter et al. 2022). For example, sponges are the highest contributing benthic invertebrate to percent cover on MCEs in Grand Cayman, constituting an average of $10.587 \pm 0.560\%$ of the benthos, which conforms to previous reports from Lesser et al. (2018) who described an increase in sponge abundance and diversity with increasing depth. However, while macroalgae cover in Grand Cayman is similar to that on MCEs at Pulley Ridge (Florida, United States of America) and in Puerto Rico (Reed et al. 2019; Apeldoorn et al. 2019)  is higher than that reported for several other Caribbean MCEs (Gress et al. 2019; Dustan and Lang 2019), including Little Cayman (Carpenter et al. 2022).

Differences in macroalgae cover between these neighbouring islands may be related to differences in levels of marine protection, as roughly 57% of the nearshore environment in Little Cayman is classified as no-take marine protected areas compared to less than 20% in Grand Cayman. While comparable fish data for mesophotic reefs in Little Cayman is not available, protection of ecologically important reef fish populations, such as herbivores, is expected to lead to reduced algal cover (Jackson et al. 2014).

While coral cover at 45m depth in Grand Cayman is low, at an average of 2.6%, similar percent coral cover is noted on MCEs in Honduras, Mexico, and at Pulley Ridge, yet is lower than the 8% cover documented in Bonaire and substantially lower than in Puerto Rico and the US Virgin Islands, where percent cover reaches nearly 30% on upper mesophotic sites (Reed et al. 2019; Gress et al. 2019; Appeldoorn et al. 2019; Smith et al. 2019; Frade et al. 2019). Despite regional differences in percent cover of scleractinian corals, species presence on MCEs appears relatively homogenous across the Caribbean, with species in the genus *Agaricia* being predominant (Reed et al. 2019; Gress et al. 2019; Appeldoorn et al. 2019; Smith et al. 2019; Frade et al. 2019). In Grand Cayman, coral species assemblages closely resemble those of nearby Little Cayman, where species in the genera *Agaricia*, *Siderastraea* and *Orbicella* have the highest contribution to percent cover at 45m depth (Carpenter et al. 2022). However, *Madracis* spp. occur in relatively high frequency in Little Cayman yet are absent from the surveys in Grand Cayman. Conversely, the genus *Mycetophyllia* is a top contributor to MCEs in Grand Cayman but not in Little Cayman. Variations in percent coral cover and species contributions among locations are likely driven by the availability of incident light as a function of reef topography. For example, on the *Orbicella* reefs in the US Virgin Islands, the reef slope is gradual with extensive flat regions in the upper mesophotic zone (Smith et al. 2019). However, the Cayman Islands are surrounded by deep trenches, with near vertical walls beyond the shallow fore reef leading to limited light availability beyond 30m (Lesser et al. 2018; Slattery and Lesser 2019).

Alternatively, differences in benthic composition, such as the dominance of *Agaricia* spp on Grand Cayman MCEs, may be related to disturbance history. The proximity of these MCEs to the Island of Grand Cayman exposes these sites to heavy levels of tourism and overfishing, impacting both coral and fish communities (Stallings 2009; Johnson et al. 2023; Rocha et al., 2023). In fact, benthic composition of MCEs is documented to affect the fish community, with high coral cover being linked to high fish density and richness (Garcia-Sais 2010). Thus, MCEs likely do not provide a refuge from disturbance for coral and fish communities in regions with high disturbance frequency or severity, and will likely be composed of weedy, fast-growing coral species that can tolerate high disturbance conditions, associated with less dense and diverse fish populations.

Conclusion

In conclusion, this study provides an overview of the fish and benthic assemblage at mesophotic depth off the coast of Grand Cayman. Our study showed that the benthic composition is highly variable across a small geographical scale with the exception of macroalgae, being the dominant benthic component across all sites. Fish assemblages were also distinct, with differences for biomass and density between trophic guilds across sites. Our study highlights the unique benthic

327 and fish communities of Grand Cayman's mesophotic reefs at 45m, which qualitatively appear
 328 different to other regions within the Caribbean (Scott 2019; Garcia-Hernandez 2018; Smith
 329 2010). Whether the unique composition of benthic and fish communities is a result of geological
 330 and biological processes, or the influence of local scale anthropogenic activity remains to be
 331 seen. Future research on remote mesophotic reefs within the same ecoregion will provide further
 332 insight into why Grand Caymans mesophotic reefs appear as distinct communities and will allow
 333 assessments of the relative influence from local anthropogenic activity.

334 **Supplementary**

335

336 Supplementary I

References

- Appeldoorn R, Alfaro M, Ballantine D, Bejarano I, Ruiz H, Schizas N, Schmidt W, Sherman C, Weil E. 2019. Puerto Rico. In: 111–129. DOI: [10.1007/978-3-319-92735-0_7](https://doi.org/10.1007/978-3-319-92735-0_7).
- Baldwin CC, Tornabene L, Robertson DR. 2018. Below the Mesophotic. *Scientific Reports* 8:4920. DOI: [10.1038/s41598-018-23067-1](https://doi.org/10.1038/s41598-018-23067-1).
- Bohnsack JA, Harper DE. 1988.Length-weight relationships of selected marine reef fishes from the southeastern United States and the Caribbean. *Available at* <https://repository.library.noaa.gov/view/noaa/3027> (accessed October 9, 2023).
- Bongaerts P, Ridgway T, Sampayo EM, Hoegh-Guldberg O. 2010. Assessing the ‘deep reef refugia’ hypothesis: focus on Caribbean reefs. *Coral Reefs* 29:309–327. DOI: [10.1007/s00338-009-0581-x](https://doi.org/10.1007/s00338-009-0581-x).
- Carpenter GE, Chequer AD, Weber S, Mass T, Goodbody-Gringley G. 2022. Light and photoacclimatization drive distinct differences between shallow and mesophotic coral communities. *Ecosphere* 13:e4200. DOI: [10.1002/ecs2.4200](https://doi.org/10.1002/ecs2.4200).
- Chasqui L, Rincón-Díaz N, Vanegas MJ, Chasqui L, Rincón-Díaz N, Vanegas MJ. 2020. Abundance of the invasive lionfish *Pterois volitans* in the coastal coral reefs from Colombian Caribbean. *Boletín de Investigaciones Marinas y Costeras - INVEMAR* 49:157–170. DOI: [10.25268/bimc.invemar.2020.49.1.779](https://doi.org/10.25268/bimc.invemar.2020.49.1.779).
- Chong-Seng KM, Mannering TD, Pratchett MS, Bellwood DR, Graham NAJ. 2012. The Influence of Coral Reef Benthic Condition on Associated Fish Assemblages. *PLOS ONE* 7:e42167. DOI: [10.1371/journal.pone.0042167](https://doi.org/10.1371/journal.pone.0042167).
- Dunn OJ. 1964. Multiple Comparisons Using Rank Sums. *TECHNOMETRICS* 6.
- Dustan P, Lang JC. 2019. Discovery Bay, Jamaica. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 85–109. DOI: [10.1007/978-3-319-92735-0_6](https://doi.org/10.1007/978-3-319-92735-0_6).
- Frade PR, Bongaerts P, Baldwin CC, Trembanis AC, Bak RPM, Vermeij MJA. 2019. Bonaire and Curaçao. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 149–162. DOI: [10.1007/978-3-319-92735-0_9](https://doi.org/10.1007/978-3-319-92735-0_9).
- Froese R, Pauly D. 2000. FishBase 2000: Concepts Designs and Data Sources. WorldFish.
- García-Hernández JE, Sanchez PJ, Hammerman NM, Schizas NV. 2018. Fish, Coral, and Sponge Assemblages Associated With Altiphotic and Mesophotic Reefs Along the Guánica Biosphere Reserve Continental Shelf Edge, Southwest Puerto Rico. *Frontiers in Marine Science* 5.
- Garcia-Sais JR. 2010. Reef habitats and associated sessile-benthic and fish assemblages across a euphotic–mesophotic depth gradient in Isla Desecheo, Puerto Rico. *Coral Reefs* 29:277–288. DOI: [10.1007/s00338-009-0582-9](https://doi.org/10.1007/s00338-009-0582-9).
- García-Sais J, Appeldoorn R, Battista T, Bauer L, Bruckner A, Caldow C, Carrubba L, Corredor J, Diaz E, Lilyestrom C, García-Moliner G, Hernández-Delgado E, Menza C, Morell J, Pait A, Sabater J, Weil E, Williams E, Williams S. 2008. The State of Coral Reef Ecosystems of Puerto Rico.
- Goodbody-Gringley G, Eddy C, Pitt JM, Chequer AD, Smith SR. 2019. Ecological Drivers of Invasive Lionfish (*Pterois volitans* and *Pterois miles*) Distribution Across Mesophotic Reefs in Bermuda. *Frontiers in Marine Science* 6. DOI: [10.3389/fmars.2019.00258](https://doi.org/10.3389/fmars.2019.00258).

Gress E, Arroyo-Gerez MJ, Wright G, Andradi-Brown DA. 2018. Assessing mesophotic coral ecosystems inside and outside a Caribbean marine protected area. *Royal Society Open Science* 5:180835. DOI: [10.1098/rsos.180835](https://doi.org/10.1098/rsos.180835).

Gress E, Voss JD, Eckert RJ, Rowlands G, Andradi-Brown DA. 2019. The Mesoamerican Reef. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 71–84. DOI: [10.1007/978-3-319-92735-0_5](https://doi.org/10.1007/978-3-319-92735-0_5).

Hinderstein LM, Marr JCA, Martinez FA, Dowgiallo MJ, Puglise KA, Pyle RL, Zawada DG, Appeldoorn R. 2010. Theme section on “Mesophotic Coral Ecosystems: Characterization, Ecology, and Management.” *Coral Reefs* 29:247–251. DOI: [10.1007/s00338-010-0614-5](https://doi.org/10.1007/s00338-010-0614-5).

Holstein DM, Fletcher P, Groves SH, Smith TB. 2019. Ecosystem Services of Mesophotic Coral Ecosystems and a Call for Better Accounting. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Coral Reefs of the World. Cham: Springer International Publishing, 943–956. DOI: [10.1007/978-3-319-92735-0_49](https://doi.org/10.1007/978-3-319-92735-0_49).

Houde ED, Zastrow CE. 1993. Ecosystem- and Taxon-specific Dynamic and Energetics Properties of Larval Fish Assemblages. *Bulletin of Marine Science* 53:290–335.

Jaccard, P. 1901. Étude comparative de la distribution florale dans une portion des Alpes et des Jura. *Bulletin de la Société Vaudoise des Sciences Naturelles*.

Jackson J (ed), Donovan M (ed), Cramer K (ed), Lam V (ed). 2014. *Status and trends of Caribbean coral reefs : 1970-2012*. Gland, Switzerland: Global Coral Reef Monitoring Network; International Union for the Conservation of Nature (IUCN).

Johnson JV, Chequer AD, Goodbody-Gringley G. 2023. Insights from the 2-year-long human confinement experiment in Grand Cayman reveal the resilience of coral reef fish communities. *Scientific Reports* 13:1–10. DOI: [10.1038/s41598-023-49221-y](https://doi.org/10.1038/s41598-023-49221-y).

Kahng SE, Garcia-Sais JR, Spalding HL, Brokovich E, Wagner D, Weil E, Hinderstein L, Toonen RJ. 2010. Community ecology of mesophotic coral reef ecosystems. *Coral Reefs* 29:255–275. DOI: [10.1007/s00338-010-0593-6](https://doi.org/10.1007/s00338-010-0593-6).

Kruskal WH, Wallis WA. 1952. Use of Ranks in One-Criterion Variance Analysis. *Journal of the American Statistical Association* 47.

Kulbicki M, Moutham G, Thollot P, Wantiez L. 1993. Length-weight relationships of fish from the lagoon of New Caledonia. *Naga Iclarm Q*.

Laverick JH, Andradi-Brown DA, Rogers AD. 2017. Using light-dependent scleractinia to define the upper boundary of mesophotic coral ecosystems on the reefs of Utila, Honduras. *PLOS ONE* 12:e0183075. DOI: [10.1371/journal.pone.0183075](https://doi.org/10.1371/journal.pone.0183075).

Laverick JH, Piango S, Andradi-Brown DA, Exton DA, Bongaerts P, Bridge TCL, Lesser MP, Pyle RL, Slattery M, Wagner D, Rogers AD. 2018. To what extent do mesophotic coral ecosystems and shallow reefs share species of conservation interest? A systematic review. *Environmental Evidence* 7:15. DOI: [10.1186/s13750-018-0127-1](https://doi.org/10.1186/s13750-018-0127-1).

Lesser MP, Slattery M, Leichter JJ. 2018. Ecology of mesophotic coral reefs. *Journal of Experimental Marine Biology and Ecology*. DOI: [10.1016/j.jembe.2009.05.009](https://doi.org/10.1016/j.jembe.2009.05.009).

Lesser MP, Slattery M, Mobley CD. 2018. Biodiversity and Functional Ecology of Mesophotic Coral Reefs. *Annual Review of Ecology, Evolution, and Systematics* 49:49–71. DOI: [10.1146/annurev-ecolsys-110617-062423](https://doi.org/10.1146/annurev-ecolsys-110617-062423).

Macartney KJ, Clayshulte Abraham A, Slattery M, Lesser MP. 2021. Growth and feeding in the sponge *Agelas tubulata* from shallow to mesophotic depths on Grand Cayman Island. *Ecosphere* 12:e03764. DOI: [10.1002/ecs2.3764](https://doi.org/10.1002/ecs2.3764).

Pérez-Castro MÁ, Schubert N, Ang-Montes de Oca G, Leyte-Morales GE, Eyal G, Hinojosa-Arango G. 2022. Mesophotic Coral Ecosystems in the Eastern Tropical Pacific: The current state of knowledge and the spatial variability of their depth boundaries. *Science of The Total Environment* 806:150576. DOI: [10.1016/j.scitotenv.2021.150576](https://doi.org/10.1016/j.scitotenv.2021.150576).

Pinheiro HT, Goodbody-Gringley G, Jessup ME, Shepherd B, Chequer AD, Rocha LA. 2016. Upper and lower mesophotic coral reef fish communities evaluated by underwater visual censuses in two Caribbean locations. *Coral Reefs* 35:139–151. DOI: [10.1007/s00338-015-1381-0](https://doi.org/10.1007/s00338-015-1381-0).

Pyle RL, Copus JM. 2019. Mesophotic Coral Ecosystems: Introduction and Overview. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 3–27. DOI: [10.1007/978-3-319-92735-0_1](https://doi.org/10.1007/978-3-319-92735-0_1).

Reed JK, Farrington S, David A, Harter S, Pomponi SA, Cristina Diaz M, Voss JD, Spring KD, Hine AC, Kourafalou VH, Smith RH, Vaz AC, Paris CB, Dennis Hanisak M. 2019. Pulley Ridge, Gulf of Mexico, USA. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 57–69. DOI: [10.1007/978-3-319-92735-0_4](https://doi.org/10.1007/978-3-319-92735-0_4).

Russ GR, Rizzari JR, Abesamis RA, Alcalá AC. 2021. Coral cover a stronger driver of reef fish trophic biomass than fishing. *Ecological Applications* 31:e02224. DOI: [10.1002/eap.2224](https://doi.org/10.1002/eap.2224).

Sandin SA, Sampayo EM, Vermeij MJA. 2008. Coral reef fish and benthic community structure of Bonaire and Curaçao, Netherlands Antilles. *Caribbean Journal of Science* 44:137–144. DOI: [10.18475/cjos.v44i2.a2](https://doi.org/10.18475/cjos.v44i2.a2).

Scott AR, Battista TA, Blum JE, Noren LN, Pawlik JR. 2019. Patterns of benthic cover with depth on Caribbean mesophotic reefs. *Coral Reefs* 38:961–972. DOI: [10.1007/s00338-019-01824-6](https://doi.org/10.1007/s00338-019-01824-6).

Shannon C, Weaver W. 1949. The Mathematical Theory of Communication. *Urban. Univ. Illinois Press. Illinois. p. 125*.

Shapiro SS, Wilk MB. 1965. An Analysis of Variance Test for Normality (Complete Samples). *Biometrika* 52:591. DOI: [10.2307/2333709](https://doi.org/10.2307/2333709).

Slattery M, Lesser MP. 2019. The Bahamas and Cayman Islands. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Coral Reefs of the World. Cham: Springer International Publishing, 47–56. DOI: [10.1007/978-3-319-92735-0_3](https://doi.org/10.1007/978-3-319-92735-0_3).

Smith TB, Blondeau J, Nemeth RS, Pittman SJ, Calnan JM, Kadison E, Gass J. 2010. Benthic structure and cryptic mortality in a Caribbean mesophotic coral reef bank system, the Hind Bank Marine Conservation District, U.S. Virgin Islands. *Coral Reefs* 29:289–308. DOI: [10.1007/s00338-009-0575-8](https://doi.org/10.1007/s00338-009-0575-8).

Smith TB, Holstein DM, Ennis RS. 2019. Disturbance in Mesophotic Coral Ecosystems and Linkages to Conservation and Management. In: Loya Y, Puglise KA, Bridge TCL eds. *Mesophotic Coral Ecosystems*. Cham: Springer International Publishing, 911–929. DOI: [10.1007/978-3-319-92735-0_47](https://doi.org/10.1007/978-3-319-92735-0_47).

Stallings CD. 2009. Fishery-Independent Data Reveal Negative Effect of Human Population Density on Caribbean Predatory Fish Communities. *PLOS ONE* 4:e5333. DOI: [10.1371/journal.pone.0005333](https://doi.org/10.1371/journal.pone.0005333).

Stefanoudis PV, Rivers M, Smith SR, Schneider CW, Wagner D, Ford H, Rogers AD, Woodall LC. 2019. Low connectivity between shallow, mesophotic and rariphotic zone benthos. *Royal Society Open Science* 6:190958. DOI: [10.1098/rsos.190958](https://doi.org/10.1098/rsos.190958).

471 Tukey JW. 1949. Comparing Individual Means in the Analysis of Variance. *Biometrics* 5:99.
 472 DOI: [10.2307/3001913](https://doi.org/10.2307/3001913).
 473 Voss J, Williams M, Reed J, Clark R. 2014. Chapter 5 Benthic and Fish Communities in the Mid
 474 and Lower Mesophotic Zone of the Sanctuary. In: *Fish and Benthic Communities of the Flower*
 475 *Garden Banks National Marine Sanctuary: Science to Support Sanctuary Management*. 201–
 476 260.
 477 Welcomme RL. 1988. International Introductions of Inland Aquatic Species. Food & Agriculture
 478 Org.

Figure 1

Context of the study and sampling techniques.

Maps of the study sites, Lighthouse Point, Ghost Mountain, La Mesa and Roundabout and the position of Grand Cayman amidst the Caribbean Sea (A). Picture of the dive team laying transect and taking pictures of the quadrat during a 15 meters practice dive (B).




Figure 2

Comparison of Species richness (number of unique species per transect) (A), Biomass (estimated weight of fish in g per transect) (B), Density (number of fish per transect) (C) and Diversity (Shannon diversity index) (D) between Sites.

Boxplots' horizontal bars represent the sites median values, the upper and lower sections (box outlines) represent the interquartile range, the whiskers extend to 90% of the data, with dots being outliers.




Figure 3

Comparison of Species richness (number of unique species per transect) (A), Biomass (estimated weight of fish in mg per transect) (B), Density (number of fish per transect) (C), and Diversity (Shannon diversity index) (D) between Trophic Guilds.

Boxplots' horizontal bars represent the sites median values, the upper and lower sections (box outlines) represent the interquartile range, the whiskers extend to 90% of the data, with dots being outliers.




Figure 4

Biomass (A) and Density (B) of the common fish families across all sites. Biomass (C) and Density (D) of each species across all sites.

Solid lines in plots A and B are the average per common family, and the whiskers are the range of the data. Each rectangle in C and D shows at least one occurrence of a species on one of the transects. The absence of a rectangle represents the absence of a species on a site.




Figure 5

Comparison of the percentage cover of the different functional groups (macroalgae, sand/rubbles, sponge, CCA and hard coral).

The horizontal middle bars represent the median values, whilst the upper and lower sections (box outlines) represent the interquartile range. The whiskers extend to 90% of the data, with dots being outliers.




Figure 6

Comparison of the percent contribution of each Scleractinian coral genus observed per site.

With *Agaricia* being the most common genus recorded across all sites.




Figure 7

Ordination plots of the dissimilarity between the sites regarding benthic composition (A), fish trophic guild composition (C) and fish species (E).

Each dot represents a quadrat for the benthic ordination plot (A) and a transect for the fish trophic plots (C and E). Roundabout's ellipse is not present owing to a deficit of replicates which does not allow the projection (C and E). Each ellipse represents the tendency of a site to drift along the two axes following the vectors on the right of each ordination plot (B and D).




Table 1 (on next page)

Benthic component's Adjusted P-values throughout sites.

	Sand/Rubble	Sponge	CCA	Hard Coral
Macroalgae	<0.001	<0.001	<0.001	<0.001
Sand/Rubble		0.067	0.001	<0.001
Sponge			0.157	<0.001
CCA				<0.001

1