

1 **Small population size and possible extirpation of the**
2 **threatened Malagasy poison frog *Mantella cowanii***

3 Devin Edmonds^{1,2}, Raphali R. Andriantsimanarilafy³, Angelica Crottini^{4,5,6}, Michael J.
4 Dreslik^{1,2}, Jade Newton-Youens⁷, Andoniana Ramahefason⁸, Christian
5 Randrianantoandro⁸, Franco Andreone⁹

6 ¹Department of Natural Resources and Environmental Sciences, University of Illinois at
7 Urbana-Champaign, Champaign, Illinois, USA

8 ²Illinois Natural History Survey, Prairie Research Institute, University of Illinois at
9 Urbana-Champaign, Champaign, IL, USA

10 ³Madagasikara Voakajy, Antananarivo, Madagascar

11 ⁴CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Vairão,
12 Portugal

13 ⁵Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, Porto,
14 Portugal

15 ⁶BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de
16 Vairão, Vairão, Portugal

17 ⁷Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK

18 ⁸Mention Zoologie et Biodiversité Animale, Université d'Antananarivo, Antananarivo,
19 Madagascar

20 ⁹Museo Regionale di Scienze Naturali, Torino, Italy

21

22 Corresponding Author:

23 Devin Edmonds^{1,2}

24 1816 S Oak Street, Champaign, Illinois, 61820, USA

25

26 Email address: dae2@illinois.edu

Abstract

Amphibians are experiencing severe population declines, requiring targeted conservation action for the most threatened species and habitats. Unfortunately, we do not know the basic demographic traits of most species, which hinders population recovery efforts. We studied one of Madagascar's most threatened frog species, the harlequin mantella (*Mantella cowanii*), to confirm it is still present at historic localities and estimate annual survival and population sizes. We surveyed eleven of all thirteen known localities and were able to detect the species at eight. Using a naïve estimate of detection probability from sites with confirmed presence, we estimated 1.54 surveys (95% CI: 1.10–2.37) are needed to infer absence with 95% confidence, suggesting the three populations where we did not detect *M. cowanii* are now extirpated. However, we also discovered two new populations. Repeated annual surveys at three sites showed population sizes ranged from 13–137 adults over 3–8 years, with the most intensively surveyed site experiencing a >80% reduction in population size during 2015–2023. Annual adult survival was moderately high (0.529–0.618) and we recaptured five individuals in 2022 and one in 2023 first captured as adults in 2015, revealing the maximum lifespan of the species in nature can reach nine years and beyond. Our results show *M. cowanii* is characterized by a slower life history pace than other *Mantella* species, putting it at greater extinction risk. Illegal collection for the international pet trade and continued habitat degradation are the main threats to the species. We recommend conservation efforts continue monitoring populations and reassess the IUCN Red List status because it may be Critically Endangered rather than Endangered based on population size and trends.

Introduction

Amphibian species are facing extinction rates at least 22 times faster than the average rate during the 10 millennia before industrialization, resulting in their status as the most threatened vertebrate class (Ceballos et al., 2015; Luedtke et al., 2023). Many species are experiencing severe population declines, leading to widespread range contractions through extirpation (e.g., Beyer and Manica 2020; Granados-Martínez et al. 2021; Patla and Peterson 2022). Habitat loss is the largest threat to amphibians, but infectious diseases, invasive species, climate change, overexploitation, and pollution are all responsible for declines and interact in complex ways (Collins, 2010; Campbell Grant, Miller & Muths, 2020). Such threats and population trends highlight the immediate need for increased conservation, especially targeted toward the most threatened species and their habitats.

Commenté [a1]: Be consistent "and or &" see above
Line 56 & 57

63 The island of Madagascar supports extraordinary amphibian species richness
64 and endemism, with more than 415 described endemic frog species representing five
65 anuran clades of independent origin (Crottini et al., 2012; Antonelli et al., 2022;
66 AmphibiaWeb 2023). Alarmingly, 46.4% of assessed Malagasy frog species are
67 threatened, owing largely to deforestation (Ralimanana et al., 2022; IUCN, 2023).
68 Deforestation has eliminated as much as a quarter of the tree cover on the island over
69 the last 25 years and the rate has only increased since 2005 (Vieilledent et al., 2018;
70 Suzzi-Simmons, 2023). Consequently, many frog species in Madagascar have patchy
71 distributions restricted to isolated pockets of forest in an otherwise inhospitable
72 landscape (Lehtinen & Ramanamanjato, 2006). So far, there have been no documented
73 modern frog species extinctions in Madagascar (Andreone et al., 2008, 2021), but many
74 records of species presence are from biological inventories conducted decades ago in
75 areas with high rates of land use change. Verifying species presence and confirming the
76 extant distribution of threatened species is some of the most vital information for
77 informing conservation (Villero et al., 2017). Relatedly, we know little about frog
78 population trends in Madagascar, even for highly threatened species. The lack of
79 demographic information is not unique to Madagascar; we do not know survival or
80 fertility rates for 87.5% of amphibian species globally (Conde et al., 2019). Baseline
81 estimates of survival, recruitment, and other demographic traits are urgently needed to
82 improve conservation efforts and inform management decisions (Campbell Grant, Miller
83 & Muths, 2020).

Mis en forme : Surlignage

Commenté [a2]: Ecological requirement is another important aspect to consider.

84 Some of the most well-known amphibians in Madagascar are the Malagasy
85 poison frogs in the genus *Mantella*. One species (*M. laevis*) exhibits parental care
86 and all display aposematic coloration to warn predators of their poisonous skin alkaloids
87 sequestered from prey (Vences et al., 2022). As such, they are familiar examples of
88 convergent evolution with Neotropical dendrobatids (Daly, Highet & Myers, 1984; Chiari
89 et al., 2004; Fischer et al., 2019). While several species are widespread and have been
90 found in degraded habitat and agricultural plantations (e.g., *M. betsileo*, *M. ebenau*, and
91 *M. viridis*, Vences et al., 1999; Andreone et al., 2006; Crottini et al., 2012), most are
92 restricted to small areas, have highly localized populations, show a recent dramatic
93 demographic decline, and are threatened by ongoing habitat degradation (e.g., Crottini
94 et al. 2019). Compounding the threat of habitat loss is overexploitation; thousands of
95 wild poison frogs are exported annually from Madagascar for the international pet trade
96 (Rabemananjara et al., 2007b), though export quotas have been restricted recently to
97 smaller quantities of just six species (CITES, 2022).

Commenté [a3]: If so, may be important to add more information about demographic traits, survival, recruitment and ecological requirement for this species. So, do these information improve the management and conservation of this species specialist of Bamboo?

Commenté [a4]: These species are more generalists in terms of habitat and seem prefer forest edge and open habitats and high tolerance without strict requirement ecologically.

Commenté [a5]: It is not necessary habitat degradation only but it may be a habitat change too (related to climate for example)

Commenté [a6]: There was no 2007a before

98 The harlequin mantella frog (*M. cowanii*) is one of the most threatened *Mantella*
99 species, with a small and fragmented distribution in the central highlands. This region of
100 Madagascar was formerly a mosaic of grassland, woodland, and subhumid forest,
101 covering the mountainous area between the island's humid east and dryer west (Yoder

Commenté [a7]: Because it has a very specific ecological requirement.

et al., 2016). Today the central highlands consist mostly of secondary grasses and land converted for subsistence agriculture and cattle grazing, with little humid forest remaining (Andriambeloson et al., 2021; Ranarilalaitiana et al., 2022). Thirteen localities of *M. cowanii* are known from the region: six in a cluster around the town of Antoetra, five 80 km northwest on the Itremo Massif, and two isolated localities located >100 km north of all other known populations, one near Betafo and the other east of Antakasina (Rabibisoa, 2008; Rabibisoa et al., 2009). All populations occur along mountainous streams with large boulders and adjacent wet rockfaces covered wet by moss and bryophytes. While some sites still have intact gallery forests, others are almost entirely devoid of trees (Fig. 1). Additionally, due to its striking black and orange coloration (Fig. 2), *M. cowanii* was heavily exploited for the international pet trade during the 1990s and early 2000s (Andreone, Mercurio & Mattioli, 2006). From 1994–2003, several thousand frogs recorded as *M. cowanii* were exported from Madagascar for commercial purposes, after which legal trade was halted and the export quota set to zero (Rabemananjara et al., 2007b; CITES, 2022).

Commenté [a8]: It is not something new in Madagascar

Commenté [a9]: I think Antoetra is a village.

Commenté [a10]: It is a bit ambiguous, as if there is forest or gallery forest anywhere before.

Mis en forme : Surlignage

Despite heavy collection pressure and ongoing habitat loss, the demographic characteristics of the remaining *M. cowanii* populations have not been studied. Additionally, some populations are known from only one or two scientific expeditions carried out decades ago and could already be extirpated. In 2021 the *Mantella cowanii* Action Plan (McAP) was launched to draw attention to the species and develop a blueprint for its conservation (Andreone et al., 2020; Rakotoarison, Ndriantsoa & Rabemananjara, 2022). The McAP builds on the initial conservation strategy by Rabibisoa (2008). We aimed to fill critical research needs in the McAP by 1) confirming the presence of *M. cowanii* at localities across its range and 2) if present, estimating the key demographic traits of survival and population size.

Commenté [a11]: There was but rather scattered: Fades project in 2003-2004 with the Animal Biology Department, Faculty of Sciences, University of Antananarivo. Andrianasolo Tsiorisoa Ginah Master in 2016 on ecology and biology of *M. cowanii* in Fohisokina. The NGO Mavoa carried-out different researches in this area too.

Materials & Methods

Study Sites

We surveyed 11 of 13 known *M. cowanii* localities, nine in the Amoron'i Mania Region and two in the Vakinankaratra Region (Fig. 3). Sites ranged in elevation ~1380–2120 m asl. We repeatedly surveyed three sites (Ambatofotsy, Soamasaka, and Fohisokina) over 3–8 years for 2–20 days/year to estimate demographic traits (Table 1). The remaining eight sites (Ambinanitelo, Andraholoma, Antakasina, Antsirakambiaty, Bekaraka, Farihimazava, Tsimabeomby, and Vatolampy) were visited up to two times for 1–6 days to confirm species presence. Only two sites have some form of habitat protection: Fohisokina (also known as Vohisokina) is community-managed with an NGO (Nowakowski & Angulo, 2015), and Antsirakambiaty falls within the boundaries of Itremo Massif Protected Area (Alvarado, Silva & Archibald, 2018). The other sites are unmanaged and without legal protection.

Mis en forme : Surlignage

140 Data Collection

141 Fieldwork was conducted during the onset of the rainy season from late
142 November to mid-January when *M. cowanii* is active and detectable. Surveys were
143 carried out in the morning during 5–8h, but at Fohisokina in 2015, we also surveyed just
144 before dusk during 16–18h. We searched for frogs visually, walking together in teams of
145 2–7 people along the stream or adjacent wet rockface. If we heard a frog calling, we
146 used the call to help find its location. Stream segment length varied from 38 m at
147 Bekaraka to 690 m at Ambatofotsy. At Fohisokina, we surveyed along six 50
148 m-long transects rather than opportunistically throughout the entire site to accomplish
149 additional research objectives (see Newton-Youens 2017).

150 After capturing a frog, we held it in a plastic bag (or a petri dish in 2015 at
151 Fohisokina), marking the location with a GPS point, flagging tape, and a unique number.
152 We measured snout-to-vent length with digital calipers or a ruler. Individuals <22 mm
153 were recorded as subadults under one-year post-metamorphosis based on Guarino et
154 al. 2008. Weight was recorded with a digital scale to the nearest 0.01 g, and sex was
155 recorded based on whether an individual had been calling before capture and, if not,
156 body size (Tessa et al. 2009). We also took dorsal and ventral photographs, allowing us
157 to identify recaptured individuals because each frog has a unique ventral color pattern
158 (Fig. 4). Such photographic capture-mark-recapture techniques can be more accurate
159 and are less invasive than traditional toe-clipping or visual implant elastomers (Caorsi,
160 Santos, and Grant 2012; Davis, VanCompernelle, and Dickens 2020). Initially, we tried
161 using photographic matching software to automate identifying individuals (sensu
162 Edmonds et al. 2019), but photograph angle and quality varied between days, sites, and
163 years, so the process was prone to false negatives. Instead, one of us (D. Edmonds)
164 visually examined all ventral photographs side-by-side and recorded when there was a
165 match. Following photographs and measurements, frogs were released in the location
166 where they were captured.

167 We followed all applicable international, national, and institutional guidelines for
168 the care and use of animals, so the animal ethics are respected. The study methods
169 were approved by the Ministère de l'Environnement et du Développement Durable in
170 permits N°173/20/MEDD/SG/DGGGE/DAPRNE/SCBE.Re, N°439/21/
171 MEDD/SG/DGGGE/DAPRNE/SCBE.Re, and N°173/22/
172 MEDD/SG/DGGGE/DAPRNE/SCBE.Re and by the University of Illinois Urbana-
173 Champaign Institutional Animal Care and Use Committee in protocol #21180.

174 Analysis

175 We used a robust design capture-mark-recapture model to estimate population
176 size ($\hat{N}$) and apparent annual survival (ϕ) at Ambatofotsy, Fohisokina, and Soamasaka.
177 These sites were selected because we had sampled them for at least three years, the

Commenté [a12]: How are these people arranged during the observation? How far from the stream side is covered? How to proceed for the other types of habitats (wet outcrops...)?

Commenté [a13]: Precision??

Commenté [a14]: Somewhere &

Mis en forme : Surlignage

Mis en forme : Surlignage

Commenté [a15]: The same day of capture?

minimum required for estimating annual survival. Our analysis compared 17 models, all incorporating site as a group-level effect. We considered models with either constant or site-specific φ and the temporary emigration parameters γ'' and γ' either constrained to 0 assuming no movement or set equal assuming random emigration. To account for individual heterogeneity in capture probability (p), we included a random effect of individual on p . Additionally, we compared models with capture probability covariates of site, year, number of surveyors, and survey effort calculated as the number of surveyors multiplied by the survey duration in minutes. We could not include environmental covariates that might be associated with capture probability because they were not collected consistently across all sites and years. To compare candidate models, we ranked them using Akaike's Information Criterion adjusted for small sample sizes (AIC_c; Burnham and Anderson 2002). Models with $\Delta \text{AIC}_c < 2$ were considered to have support. We analyzed capture-mark-recapture data in program MARK through the *RMark* interface (White & Burnham, 1999; Laake, 2013) and assessed the goodness-of-fit with package *R2ucare* in R version 4.2.0 (Gimenez et al., 2018; R Core Team, 2022).

To infer absence if we did not detect *M. cowanii* at a historic locality, we estimated detection probability with a single-season occupancy model in package *unmarked* (Fiske and Chandler 2011). Data from nine localities with confirmed presence and at least three surveys were used to generate a naïve estimate of detection probability assuming constant detection and occupancy. We then followed Pellet and Schmidt (2005) to estimate the number of surveys needed to detect the species as:

$$N_{min} = \frac{\ln(0.05)}{\ln(1 - p)}$$

where N_{min} is the minimum number of surveys, p is the detection probability, and 0.05 is the confidence level needed to be 95% certain of absence, assuming independent and comparable surveys. Using the number of surveys, we then calculated the confidence level around an observed absence given N number of surveys as:

$$conf = e^{N \cdot \ln(p)}$$

Results

Verifying Species Presence

We confirmed *M. cowanii* presence at 8 of 11 surveyed localities and identified 2 previously unrecorded populations. However, we failed to detect the species at the historical localities of Andraholoma, Tsimabeomby, and Vatolampy. The naïve detection probability from sites with confirmed presence was 0.86 (95% CI: 0.72–0.93), showing it takes 1.54 surveys (95% CI: 1.10–2.37) to be 95% confident a population is extirpated and 2.37 surveys (95% CI: 1.69–3.65) to be 99% confident. With at least two days of

Commenté [a16]: It is necessary to underline in the materials and methods that these three sites are historical localities (why?) and to point out when the last time *M.c* was found there.

Commenté [a17]: As daily activity rhythm of the frog and ectothermic animals in general depend largely on daily meteorological conditions, it is important to precise that this is the case in optimal condition for biological activities.

213 surveys at Andraholoma and Vatolampy, we can be >97.9% (95% CI: 91.9–99.9%)
214 confident the populations are extirpated and 85.7% (95% CI: 71.7–93.4%) confident
215 there are no *M. cowanii* at Tsimabeomby.

216 *Capture Patterns Across Sites*

217 We made 764 captures of 280 individuals across all study sites and years,
218 excluding putative *M. baroni* x *M. cowanii* hybrids. All but 8 frogs were adults. Over half
219 of all captures were at Fohisokina, where we caught 149 individuals 481 times. Six
220 individuals at Fohisokina were recaptured 7 years after initially being caught in 2015 as
221 adults (Fig. 4), one of which was recaptured again in year 8 in 2023. At our second most
222 intensively surveyed site, Soamasaka, we made 137 captures of 40 individuals during
223 annual fieldwork in 2020–2023. Three frogs were recaptured 4 years after their initial
224 encounter in 2020, and 4 were recaptured 3 years apart. At Ambatofotsy, we conducted
225 surveys in 2021, 2022, and 2023 and made 76 captures of 32 individuals, all adults.

226 *Annual Survival and Population Sizes*

227 The most parsimonious capture-mark-recapture model had capture probability p
228 and annual adult survival φ varying by site (Table 2). Models with no movement
229 generally performed better than those with random emigration (Table 2). The top model
230 estimated population sizes ($\hat{N}$) ranging from 13–137 adult frogs per site across years
231 (Fig 5). The highest $\hat{N}$ estimate was from Fohisokina in 2015 (137, 95% CI: 120–170)
232 and the lowest from Soamasaka in 2023 (13, 95% CI: 11–22). Fohisokina showed a
233 decreasing population size during 2015–2023, whereas Soamasaka and Ambatofotsy
234 were relatively stable over a shorter period (Fig. 5). There was strong support for site-
235 varying survival (Table 2), with the estimated annual adult survival more precise for
236 Fohisokina than Soamasaka due to the longer study duration (Fig. 6). For Ambatofotsy,
237 the survival estimate was too imprecise to be informative because the data spanned
238 only three years and only a small number of frogs were captured each year. Overall, the
239 annual survival estimates at Fohisokina and Soamasaka were comparable, although the
240 estimate from Soamasaka was lower than Fohisokina (Fig. 6).

241 **Discussion**

242 Our results demonstrate *M. cowanii* was still present at no less than ten localities
243 in 2022–2023, but the population size is very small for at least three sites (<50 adults
244 per site). Extrapolating across all known localities, in the worst-case scenario, the total
245 adult population size for the species may number <500 individuals. However, frog
246 populations naturally fluctuate in abundance, and a snapshot over several years can
247 easily lead to erroneous conclusions that populations are declining when they are stable
248 (Pechmann et al., 1991; Blaustein, Wake & Sousa, 1994; Meyer, Schmidt &

Commenté [a18]: It is necessary to be prudent in the interpretation of such result with a cryptic species with biological activity largely influence by environmental conditions especially the climatic (P° & T°) conditions.

Commenté [a19]: Approximately how often an individual is recaptured? In the same area? And for how many successive days? That may help to understand the ecology and the behavior of this species.

Commenté [a20]: In the same area as before? A kind fidelity to?

Mis en forme : Surlignage

Mis en forme : Surlignage

Grossenbacher, 1998). Such fluctuations are typical of species with fast life histories, where fecundity is high and generation time short, thus, demographic rates tend to vary (Sæther et al., 2004, 2013). Additionally, amphibian populations fluctuate more for pond breeding species and less for terrestrial and stream breeding species (Green, 2003). Considering *M. cowanii* is a terrestrial stream-breeding frog with comparatively low reproductive output (20–57 eggs per egg mass; Tessa et al. 2009), relatively high annual adult survival, and a maximum lifespan in nature of at least nine years, we believe our results are not an artifact of stochastic fluctuations in population size.

We highlight three possible factors contributing to the >80% population decline at Fohisokina between 2015 and 2023. First, fires burnt much of Fohisokina in November 2020 at the start of their breeding season, just before our survey. Though *M. cowanii* presumably is protected from fire while sheltering in moist rock crevices and caves for much of the year, fire can cause mortality in terrestrial frogs when they are active above ground during the breeding season (e.g., Humphries and Sisson 2012; Potvin et al. 2017). Second, according to a European private breeder, in 2017 more than 100 *M. cowanii* were illicitly offered for sale in Germany, and an unknown number were offered again in 2021. As Fohisokina is the most easily accessed *M. cowanii* locality and was historically a collection site for the pet trade (Rabemananjara et al., 2007b), the frogs were possibly poached from Fohisokina. Lastly, although no records of chytridiomycosis have been confirmed in Madagascar and all frogs we sampled appeared healthy, the amphibian chytrid fungus *Batrachochytrium dendrobatidis* has been reported from the island, and in 2014 was detected on a single *M. cowanii* individual at Soamasaka ~5 km south of Fohisokina (Bletz et al., 2015). Therefore, we cannot rule out disease either.

At Vatolampy, we did not detect *M. cowanii* after six days of surveys and suspect the population is extirpated. Until our work, the site had not been surveyed since 2003–2004, when Andreone et al. (2007) and Rabemananjara et al. (2007a) collected tissue samples and voucher specimens from the population. Similarly, Andraholoma had not been surveyed since 2009 when the site was visited by one of us (C. Randrianantoandro) for one day and 5 individuals observed. Conversely, we question whether Tsimabeomby, the third site where we did not detect *M. cowanii*, ever supported a population. We believe the locality was possibly published in error by Rabibisoa (2008) because Tsimabeomby consists of a wet meadow, is without rocks or running water, is isolated from the next nearest population by >3 km, and the local people we worked with had never encountered *M. cowanii* there whereas they knew of the other populations. Nonetheless, *M. cowanii* could have been present but undetected during our surveys if the detection probability at Andraholoma, Tsimabeomby, and Vatolampy was lower than elsewhere. Additionally, the naïve estimate of detection probability we used to infer absence did not account for observer, environmental, or temporal factors influencing detection. Still, our team detected *M. cowanii* at other sites on the days we

Commenté [a21]: What about the effect of ash and sediment when it rains, it may affect the quality of the stream and change the ecological parameters disturbing eggs and tadpoles developments?

Commenté [a22]: So the zero quota defined by the CITES management authority in Madagascar was not respected?

Mis en forme : Surlignage

Commenté [a23]: But it seems that it is the only legally protected site with a NGO and local community as the managers.

Commenté [a24]: Is the ecological condition optimal for biological activities?

Commenté [a25]: Why 2007a after 2007b? Chronology

Commenté [a26]: This is the case when the conditions are favorable for activities

288 surveyed Andraholoma, Tsimabeomby, and Vatolampy, illustrating local environmental
289 conditions were favorable for detecting *Mantella*. Re-surveying the three sites in the
290 coming years will confirm if the populations are extirpated.

291 Some amphibian species are capable of dispersing long distances to recolonize
292 suitable habitat (Marsh and Trenham 2001; Fonte, Mayer, and Lötters 2019), but we do
293 not expect so for *M. cowanii* in Madagascar's degraded highland landscape. There are
294 no studies on the movement of Madagascar's poison frogs (but see Andreone et al.
295 2013), however research on their Neotropical dendrobatid counterparts shows adults
296 rarely move more than a few hundred meters from established territories (e.g., Ringler
297 et al. 2009, Pašukonis et al. 2013; Beck et al. 2017; Pašukonis et al. 2019). Moreover,
298 in a review of the dispersal ability of amphibians, Smith and Green (2005) found nearly
299 half of studied species moved <400m. Historically, adult frogs may have moved
300 between patches when there was more forest in the highlands. However, we suspect
301 *Mantella* usually passively disperse when tadpoles are flushed between habitat patches
302 during heavy rain. As such, natural recolonization of Vatolampy or Andraholoma is
303 unlikely, especially considering the physiological, movement, and site fidelity constraints
304 amphibians face (Blaustein, Wake & Sousa, 1994). At Vatolampy, the next closest
305 locality is Farihimazava, 1.5 km northeast in a different valley, which supports mostly *M.*
306 *baroni* and *M. baroni* x *M. cowanii* hybrids (Chiari et al., 2005; Andreone et al. 2007).
307 For Andraholoma, the next closest locality is Andaobatofofotsivava >7 km north, though
308 admittedly, the area of the Itremo Massif is poorly explored and there could be
309 additional unrecorded populations between the two. Indeed, we located two new
310 localities during fieldwork.

311 Previous skeletochronology research estimated the maximum lifespan of *M.*
312 *cowanii* at three years (Guarino et al., 2008; Andreone et al., 2011), but we identified
313 individuals at least eight years post-metamorphosis and one individual nine years.
314 Skeletochronology is known to underestimate the ages of older individuals because
315 skeletal growth rings progressively converge with age, and amphibian bone tissue is
316 prone to reabsorption (Eden et al., 2007; Sinsch, 2015). Our results show the
317 advantages of using capture-mark-recapture surveys for estimating demographic traits if
318 resources are available. The long lifespan of *M. cowanii* is notable when considered
319 together with their reproductive output and body size. The species is one of the largest
320 in the genus, has the largest egg diameter, and has the lowest number of eggs per
321 mass, exemplifying their slow life history compared to other *Mantella* species (Tessa et
322 al. 2009). All of this aligns with our discovery that *M. cowanii* has the longest lifespan for
323 the genus. Life history traits often follow altitudinal clines, with slower traits associated
324 with higher altitudes (Hille & Cooper, 2015; Laiolo & Obeso, 2017). Indeed, amphibians
325 tend to live longer and have larger body sizes at higher altitudes (Morrison & Hero,
326 2003; Andreone et al. 2004); *M. cowanii* is no exception (Tessa et al. 2009). Such

Commenté [a27]: It is necessary to extend the survey until the end of the rainy season. An article now in process report the finding of 39 individuals from March to April in Fohisokina if there is only 23 individuals during the rainy season November-December. The best way is to conduct repeated visits in different periods.

Commenté [a28]: But dispersion took place before degradation, that's why the distribution areas are now fragmented.

patterns also occur within species across altitudinal gradients (Zhang & Lu, 2012). Considering the *M. cowanii* from Betafo populations occur ~500m higher than where we recorded ~8–9-year-old individuals, frogs from Betafo may live even longer. Such slow life history traits also have conservation implications because they are associated with higher extinction risk (Webb, Brook & Shine, 2002). Given the relatively long lifespan and low reproductive output of *M. cowanii*, the success of recovery efforts for the species may not be as rapid as in other related amphibian species.

To ensure populations remain extant, we must better identify the causes of declines and the magnitude of threats. Screening populations for *Bd* and monitoring for illicit *M. cowanii* in the pet trade are obvious actions, but disentangling the threat of habitat loss is more complicated. The number of trees remaining at sites varies from intact closed-canopy forest to rocky landscapes almost entirely devoid of trees, so the degree to which deforestation is a threat may depend on additional habitat characteristics. Newton-Youens (2017) identified rock caves and refuges as essential habitat features, and speculated they might be used for breeding, though so far, no eggs, tadpoles, or newly metamorphosed individuals have been found in nature. Studies on the microhabitat preferences and activity levels of *M. cowanii*, like those carried out by Edwards et al. (2019, 2022) for *M. aurantiaca*, would further help identify the most critical habitat features to protect and the best time to survey sites. Likewise, better information about habitat requirements could be used to locate new sites with unprotected populations we do not know about. All known *M. cowanii* populations are centered around four isolated sites with likely past connectivity when the highlands were an intact forest-grassland mosaic (Bond, Silander & Ratsirarson, 2023). Fieldwork in the remote areas between the four population centers could uncover additional isolated *M. cowanii* populations, but time is running out.

Conclusions

We set out to verify the presence and estimate key demographic traits for one of Madagascar's most threatened frog species and found several historical localities are likely extirpated while others are extremely small. Unfortunately, our results are not unique to Madagascar but represent a global trend in amphibian populations (Stuart et al., 2004; Grant et al., 2016). Amphibians are at the forefront of the extinction crisis, and population monitoring is essential to measure responses to conservation actions and detect declines before recovery is impossible. We used capture-mark-recapture methods to estimate abundance at three localities (Ambatofotsy, Soamasaka, and Fohisokina; Fig. 3), but less costly approaches relying on presence-absence data are likely suitable for monitoring *M. cowanii* across its entire range (Joseph et al., 2006; Jones, 2011). When enacted with local people as part of a broader program, monitoring can galvanize conservation efforts by adding value to a threatened species and instilling pride in local communities (Andrianandrasana et al., 2005; Danielsen, Burgess &

Commenté [a29]: You missed three and found two right?

366 Balmford, 2005). As we have shown, the causes of declines are often unclear without
367 adequate information on a species' basic ecology and life history. For *M. cowanii*, we
368 recommend further research run concurrently with conservation efforts and focus on
369 determining the relative impact of disease, illegal trade, and habitat loss. We also
370 recommend reassessing the IUCN Red List status of *M. cowanii*. The species was last
371 assessed in 2014 as Endangered but it may qualify for the Critically Endangered status
372 based on our estimates of population sizes and trends.

373 **Acknowledgements**

374 We are grateful to Chloe Helsey who assisted with fieldwork in 2015, to L'Homme et
375 l'Environnement and V.O.I. FOMISAME for collaborative efforts at Antoetra, and to
376 Tiana Randriamboavonjy of the Kew Madagascar Conservation Centre for logistical
377 support at Itremo. Special thanks go to Association Mitsinjo members Frederic R.
378 Razafimahefa, Georges Ramarolahy, Samina S. Sam Edmonds, Edupsie
379 Tsimilaomanana, and Fanirihimbola J.E. Abraham for providing indispensable field
380 assistance. We thank R. Griffiths, E. Larson, and R. Schooley for offering feedback on
381 an early draft of the manuscript.

382

References

- Alvarado ST, Silva TSF, Archibald S. 2018. Management impacts on fire occurrence: A comparison of fire regimes of African and South American tropical savannas in different protected areas. *Journal of Environmental Management* 218:79–87. DOI: 10.1016/j.jenvman.2018.04.004.
- AmphibiaWeb. 2023. AmphibiaWeb: Information on amphibian biology and conservation. Available at <http://www.amphibiaweb.org> (accessed March 9, 2023).
- Andreone F, Andriantsimanarilafy RR, Crottini A, Edmonds D, Garcia G, Hansen-Hendrikx CM, Rakotoarison A, Razafimanahaka JH. 2020. *Mantella cowanii* Action Plan 2021–2025 / Plan d'Action *Mantella cowanii* 2021–2025. Museo Regionale di Scienze Naturali and the Amphibian Survival Alliance, Turin.
- Andreone F, Bergò PE, Mercurio V, Rosa GM. 2013. Spatial ecology of *Scaphiophryne gottlebei* in the canyons of the Isalo Massif, Madagascar. *Herpetologica* 69:11–21.
- Andreone F, Carpenter AI, Cox N, Du Preez L, Freeman K, Furrer S, Garcia G, Glaw F, et al. 2008. The challenge of conserving amphibian megadiversity in Madagascar. *PLoS Biology* 6:943–946. DOI: 10.1371/journal.pbio.0060118.
- Andreone F, Carpenter AI, Crottini A, D'Cruze N, Dubos N, Edmonds D, Garcia G, Luedtke J, et al. 2021. Amphibian Conservation in Madagascar: Old and Novel Threats to a Peculiar Fauna. In: Heatwole H, Rödel M-O eds. *Status and Threats of Afrotropical Amphibians*. Frankfurt am Main: Chimaira, 243.
- Andreone F, Giacoma C, Guarino FM, Mercurio V, Tessa G. 2011. Age profile in nine *Mantella* poison frogs from Madagascar, as revealed by skeletochronological analyses. *Alytes* 27:73–84.
- Andreone F, Mercurio V, Mattioli F. 2006. Between environmental degradation and international pet trade: conservation strategies for the threatened amphibians of Madagascar. *Natura - Soc. it. Sci. nat. Museo civ. Stor. nat. Milano* 95:81–96.
- Andreone F, Miaud C, Eusebio Bergò P, Doglio S, Stocco P, Riberon A, Gautier P. 2004. Living at high altitude: testing the effects of life history traits upon the conservation of *Salamandra lanzai* (Amphibia, Salamandridae). *The Italian Journal of Zoology* 71: 35–43.
- Andreone F, Vences M, Glaw F, Randrianirina JE. 2007. Remarkable records of amphibians and reptiles on Madagascar's central high plateau. *Tropical Zoology* 20:19–37.
- Andriambeloson JB, Blanco MB, Andriantsalohimisantatra A, Rivoharison T V., Walker N, Birkinshaw C, Yoder AD. 2021. Living in tiny fragments: a glimpse at the ecology

Mis en forme : Français (France)

418 of Goodman's mouse lemurs (*Microcebus lehilahytsara*) in the relic forest of
 419 Ankafoke, Central Highlands, Madagascar. *Primates* 62:887–896. DOI:
 420 10.1007/s10329-021-00947-1.

421 Andrianandrasana HT, Randriamahefasoa J, Durbin J, Lewis RE, Ratsimbazafy JH.
 422 2005. Participatory ecological monitoring of the Alaotra wetlands in Madagascar.
 423 *Biodiversity and Conservation* 14:2757–2774. DOI: 10.1007/s10531-005-8413-y.

424 Antonelli A, Smith RJ, Perrigo AL, Crottini A, Hackel J, Testo W, Farooq H, Torres
 425 Jiménez MF, et al. 2022. Madagascar's extraordinary biodiversity: Evolution,
 426 distribution, and use. *Science* 378:eaf0869.

427 Beck KB, Loretto MC, Ringler M, Hödl W, Pašukonis A. 2017. Relying on known or
 428 exploring for new? Movement patterns and reproductive resource use in a tadpole-
 429 transporting frog. *PeerJ*:e3745. DOI: 10.7717/peerj.3745

430 Beyer RM, Manica A. 2020. Historical and projected future range sizes of the world's
 431 mammals, birds, and amphibians. *Nature Communications* 11:1–8. DOI:
 432 10.1038/s41467-020-19455-9.

433 Blaustein AR, Wake DB, Sousa WP. 1994. Amphibian declines: judging stability,
 434 persistence, and susceptibility of populations to local and global extinctions.
 435 *Conservation Biology* 8:60–71. DOI: 10.1046/j.1523-1739.1994.08010060.x.

436 Bletz MC, Rosa GM, Andreone F, Courtois EA, Schmeller DS, Rabibisoa NHC,
 437 Rabemananjara FCE, Raharivololoniaina L, et al. 2015. Widespread presence of
 438 the pathogenic fungus *Batrachochytrium dendrobatidis* in wild amphibian
 439 communities in Madagascar. *Scientific Reports* 5:1–10. DOI: 10.1038/srep08633.

440 Bond WJ, Silander JA, Ratsirarson J. 2023. Madagascar's grassy biomes are ancient
 441 and there is much to learn about their ecology and evolution. *Journal of*
 442 *Biogeography* 50:614–621. DOI: 10.1111/jbi.14494.

443 Burnham KP, Anderson DR. 2002. *Model Selection and Multimodel Inference: A*
 444 *Practical Information-Theoretic Approach*. New York: Springer-Verlag.

445 Campbell Grant EH, Miller DAWW, Muths E. 2020. A Synthesis of evidence of drivers of
 446 amphibian declines. *Herpetologica* 76:101–107. DOI: 10.1655/0018-0831-76.2.101.

447 Caorsi VZ, Santos RR, Grant T. 2012. Clip or snap? An evaluation of toe-clipping and
 448 photo-identification methods for identifying individual Southern Red-Bellied Toads,
 449 *Melanophryniscus cambaraensis*. *South American Journal of Herpetology* 7(2):79–
 450 84. DOI: 10.2994/057.007.0210.

451 Ceballos G, Ehrlich PR, Barnosky AD, Garcia A, Pringle RM, Palmer TM. 2015.
 452 Accelerated modern human-induced species losses: Entering the sixth mass

extinction. *Science Advances* 1:e1400253–e1400253. DOI: 10.1126/sciadv.1400253.

Mis en forme : Français (France)

Chiari Y, Andreone F, Vences M, Meyer A. 2005. Genetic variation of an endangered Malagasy frog, *Mantella cowani*, and its phylogeographic relationship to the widespread *M. baroni*. *Conservation Genetics* 6:1041–1047. DOI: 10.1007/s10592-005-9085-6.

Mis en forme : Français (France)

Chiari Y, Vences M, Vieites DR, Rabemananjara F, Bora P, Ramilijaona Ravoahangimalala O, Meyer A. 2004. New evidence for parallel evolution of colour patterns in Malagasy poison frogs (*Mantella*). *Molecular Ecology* 13:3763–3774. DOI: 10.1111/j.1365-294X.2004.02367.x.

CITES. 2022. CITES Trade Database version 2022.1. Available at <https://trade.cites.org/>

Collins JP. 2010. Amphibian decline and extinction: What we know and what we need to learn. *Diseases of Aquatic Organisms* 92:93–99. DOI: 10.3354/dao02307.

Conde DA, Staerk J, Colchero F, da Silva R, Schöley J, Maria Baden H, Jouvet L, Fa JE, et al. 2019. Data gaps and opportunities for comparative and conservation biology. *Proceedings of the National Academy of Sciences of the United States of America* 116:9658–9664. DOI: 10.1073/pnas.1816367116.

Crottini A., Brown JL, Mercurio V, Glaw F, Vences M, Andreone F. 2012. Phylogeography of the poison frog *Mantella viridis* (Amphibia: Mantellidae) reveals chromatic and genetic differentiation across ecotones in northern Madagascar. *Journal of Zoological Systematics and Evolutionary Research* 50:305–314.

Crottini A, Madsen O, Poux C, Strauß A, Vieites DR, Vences M. 2012. Vertebrate time-tree elucidates the biogeographic pattern of a major biotic change around the K-T boundary in Madagascar. *Proceedings of the National Academy of Sciences of the United States of America* 109:5358–5363. DOI: 10.1073/pnas.1112487109.

Crottini A, Orozco-terWengel P, Rabemananjara FCE, Hauswaldt JS, Vences M. 2019. Mitochondrial introgression, color pattern variation and severe demographic bottlenecks in three species of Malagasy poison frogs, genus *Mantella*. *Genes* 10:317. DOI:10.3390/genes10040317.

Daly JW, Highet RJ, Myers CW. 1984. Occurrence of skin alkaloids in non-dendrobatid frogs from Brazil (Bufonidae), Australia (Myobatrachidae) and Madagascar (Mantellinae). *Toxicon* 22:905–919. DOI: 10.1016/0041-0101(84)90182-X.

Danielsen F, Burgess ND, Balmford A. 2005. Monitoring matters: Examining the potential of locally-based approaches. *Biodiversity and Conservation* 14:2507–

488 2542. DOI: 10.1007/s10531-005-8375-0.

489 Davis HP, VanCompernelle M, Dickens J. 2020. Effectiveness and reliability of
 490 photographic identification methods for identifying individuals of a cryptically
 491 patterned toad. *Herpetological Conservation and Biology* 15(1):204–211.

492 Eden CJ, Whiteman HH, Duobinis-Gray L, Wissinger SA. 2007. Accuracy assessment
 493 of skeletochronology in the Arizona Tiger Salamander (*Ambystoma tigrinum*
 494 *nebulosum*). *Copeia* 2007:471–477.

495 Edmonds D, Kessler E, Bolte L. 2019. How common is common? Rapidly assessing
 496 population size and structure of the frog *Mantidactylus betsileanus* at a site in east-
 497 central Madagascar. *Austral Ecology* 44:1196–1203. DOI: 10.1111/aec.12797.

498 Edwards WM, Bungard MJ, Griffiths RA. 2022. Daily activity profile of the golden
 499 mantella in the “Froggotron”—A replicated behavioral monitoring system for
 500 amphibians. *Zoo Biology* 41:3–9. DOI: 10.1002/zoo.21650.

501 Edwards WM, Griffiths RA, Bungard MJ, Rakotondrasoa EF, Razafimanahaka JH,
 502 Razafindraibe P, Andriantsimanarilafy RR, Randrianantoandro JC. 2019.
 503 Microhabitat preference of the critically endangered golden mantella frog in
 504 Madagascar. *Herpetological Journal* 29:207–213.

505 Fischer EK, Roland AB, Moskowitz NA, Vidoudez C, Ranaivorazo N, Tapia EE, Trauger
 506 SA, Vences M, et al. 2019. Mechanisms of convergent egg provisioning in poison
 507 frogs. *Current Biology* 29:4145–4151.e3. DOI: 10.1016/j.cub.2019.10.032.

508 Fonte LFM, Mayer M, Lötters S. 2019. Long-distance dispersal in amphibians. *Frontiers*
 509 *of Biogeography* 11(4):e44577. DOI: 10.21425/F5FBG44577.

510 Gimenez O, Lebreton JD, Choquet R, Pradel R. 2018. R2ucare: An R package to
 511 perform goodness-of-fit tests for capture–recapture models. *Methods in Ecology*
 512 *and Evolution* 9:1749–1754. DOI: 10.1111/2041-210X.13014.

513 Granados-Martínez S, Zumbado-Ulate H, Searle CL, Oliveira BF, García-Rodríguez A.
 514 2021. Niche contraction of an endangered frog driven by the amphibian chytrid
 515 fungus. *EcoHealth* 18:134–144. DOI: 10.1007/s10393-021-01525-z.

516 Grant EHC, Miller DAW, Schmidt BR, Adams MJ, Amburgey SM, Chambert T,
 517 Cruickshank SS, Fisher RN, et al. 2016. Quantitative evidence for the effects of
 518 multiple drivers on continental-scale amphibian declines. *Scientific Reports* 6:1–9.
 519 DOI: 10.1038/srep25625.

520 Green DM. 2003. The ecology of extinction: Population fluctuation and decline in
 521 amphibians. *Biological Conservation* 111:331–343. DOI: 10.1016/S0006-
 522 3207(02)00302-6.

Mis en forme : Français (France)

Mis en forme : Français (France)

523 Guarino FM, Sacco M, Tessa G, Andreone F. 2008. Short life span of two charismatic
 524 *Mantella* species: age-structure in the critically endangered *M. cowani* and in the
 525 syntopic *M. baroni*. In: Androne F ed. *A Conservation Strategy for the Amphibians*
 526 *of Madagascar*. Torino: Museo Regionale di Scienze Naturali, 265–276.

527 Hille SM, Cooper CB. 2015. Elevational trends in life histories: revising the pace-of-life
 528 framework. *Biological Reviews* 90:204–2013.

529 Humphries WJ, Sisson MA. 2012. Long distance migrations, landscape use, and
 530 vulnerability to prescribed fire of the Gopher Frog (*Lithobates capito*). *Journal of*
 531 *Herpetology* 46:665–670. DOI: 10.1670/11-124.

532 IUCN. 2023. The IUCN Red List of Threatened Species. Version 2022-2. Available at
 533 <http://www.iucnredlist.org> (accessed March 15, 2023).

534 Jones JPG. 2011. Monitoring species abundance and distribution at the landscape
 535 scale. *Journal of Applied Ecology* 48:9–13. DOI: 10.1111/j.1365-
 536 2664.2010.01917.x.

537 Joseph LN, Field SA, Wilcox C, Possingham HP. 2006. Presence-absence versus
 538 abundance data for monitoring threatened species. *Conservation Biology* 20:1679–
 539 1687. DOI: 10.1111/j.1523-1739.2006.00529.x.

540 Laake JL. 2013. *RMark: An R Interface for analysis of capture-recapture data with*
 541 *MARK*. AFSC Processed Rep. 2013-1. Alaska Fish. Sci. Cent., NOAA, Seattle,
 542 WA.

543 Laiolo P, Obeso JR. 2017. *Life-History Responses to the Altitudinal Gradient*. Cham,
 544 Switzerland: Springer Nature. DOI: 10.1007/978-3-319-55982-7_11.

545 Lehtinen R, Ramanamanjato J-B. 2006. Effects of rainforest fragmentation and
 546 correlates of local extinction in a herpetofauna from Madagascar. *Applied*
 547 *Herpetology* 3:95–110. DOI: 10.1163/157075406776984248.

548 Luedtke JA, Chanson J, Neam K, Hobin L, Maciel AO, Catenazzi A, Borzée A, Hamidy
 549 A, et al. 2023. Ongoing declines for the world's amphibians in the face of emerging
 550 threats. *Nature* 622:308–314. DOI: 10.1038/s41586-023-06578-4.

551 Marsh DM, Trenham PC. 2001. Metapopulation dynamics and amphibian conservation.
 552 *Conservation Biology* 15(1):40–49. DOI: 10.1111/j.1523-1739.2001.00129.x.

553 Meyer AH, Schmidt BR, Grossenbacher K. 1998. Analysis of three amphibian
 554 populations with quarter-century long time-series. *Proceedings of the Royal Society*
 555 *B: Biological Sciences* 265:523–528. DOI: 10.1098/rspb.1998.0326.

556 Morrison C, Hero J-M. 2003. Geographic variation in life-history characteristics of

Mis en forme : Allemand (Suisse)

557 amphibians: a review. *Journal of Animal Ecology* 72:270–279.

558 Newton-Youens J. 2017. The importance of microclimates in the ex situ management of
559 amphibians. MS Thesis. University of Manchester.

560 Nowakowski J, Angulo A. 2015. Targeted habitat protection and its effects on the
561 extinction risk of threatened amphibians. *FrogLog* 23:8–10.

562 Patla DA, Peterson CR. 2022. The slow decline of a Columbia Spotted Frog population
563 in Yellowstone National Park: A cautionary tale from a developed zone within a
564 large protected area. *Ecological Indicators* 136:108606. DOI:
565 10.1016/j.ecolind.2022.108606.

566 Pašukonis A, Ringler M, Brandl HB, Mangione R, Ringler E, Hödl W. 2013. The homing
567 frog: high homing performance in a territorial dendrobatid frog *Allobates femoralis*
568 (Dendrobatidae). *Ethology* 119(9):762–768. DOI: 10.1111/eth.12116.

569 Pašukonis A, Loretto MC, Rojas B. 2019. How far do tadpoles travel in the rainforest?
570 Parent-assisted dispersal in poison frogs. *Evolutionary Ecology* 33(4):613–623.
571 DOI: 10.1007/s10682-019-09994-z.

572 Pechmann JHK, Scott DE, Semlitsch RD, Caldwell JP, Vitt LJ, Gibbons JW. 1991.
573 Declining amphibian populations: The problem of separating human impacts from
574 natural fluctuations. *Science* 253:892–895. DOI: 10.1126/science.253.5022.892.

575 Pellet J, Schmidt BR. 2005. Monitoring distributions using call surveys: Estimating site
576 occupancy, detection probabilities and inferring absence. *Biological Conservation*
577 123:27–35. DOI: 10.1016/j.biocon.2004.10.005.

578 Potvin DA, Parris KM, Smith Date KL, Keely CC, Bray RD, Hale J, Hunjan S, Austin JJ,
579 et al. 2017. Genetic erosion and escalating extinction risk in frogs with increasing
580 wildfire frequency. *Journal of Applied Ecology* 54:945–954. DOI: 10.1111/1365-
581 2664.12809.

582 R Core Team. 2022. R: A Language and Environment for Statistical Computing. R
583 Foundation for Statistical Computer, Vienna, Austria. <https://www.R-project.org/>.

584 Rabemananjara FCE, Chiari Y, Ramilijaona OR, Vences M. 2007a. Evidence for recent
585 gene flow between north-eastern and south-eastern Madagascan poison frogs from
586 a phylogeography of the *Mantella cowani* group. *Frontiers in Zoology* 4:1. DOI:
587 10.1186/1742-9994-4-1.

588 Rabemananjara FCE, Raminosoa NR, Ravoahangimalala OR, Rakotondravony D,
589 Andreone F, Bora P, Carpenter AI, Glaw F, et al. 2007b. Malagasy poison frogs in
590 the pet trade: a survey of levels of exploitation of species in the genus *Mantella*.
591 *Amphibian and Reptile Conservation* 5:3–16. DOI: 10.1514/journal.arc.0050020.

Mis en forme : Français (France)

Mis en forme : Français (France)

592 Rabibisoa N. 2008. *Plan d'Action de Conservation de Mantella cowani (PACM)*.
 593 Conservation International and the Amphibian Specialist Group of Madagascar.

594 Rabibisoa N, Randrianasolo H, Anjeriniaina M, Mackinnon J, Andriamamonjisoa A,
 595 Randrianantoandro C, Andreone F. 2009. New findings of harlequin mantella
 596 improve the conservation status of Madagascar's most threatened frog. *Froglog*
 597 92:5–8.

598 Rakotoarison A, Ndriantsoa S, Rabemananjara F., Rabibisoa NHC, Rakotonanahary
 599 TF, Randriamahazo H, Andreone F. 2022. More than 15 years of amphibian
 600 conservation in Madagascar under the flag of IUCN SSC Amphibian Specialist
 601 Group. In: Biaggini M., Corti C., Giacobbe D., Lo Cascio P. eds. *Herpetologia*
 602 *Siciliae - XIII Congresso della Societas Herpetologica Italica (Lipari 22-26*
 603 *settembre 2021)*. *Naturalista Siciliano* 46:305–312.

604 Ralimanana H, Perrigo AL, Smith RJ, Borrell JS, Faurby S, Rajaonah MT,
 605 Randriamboavonjy T, Vorontsova MS, et al. (2022): Madagascar's extraordinary
 606 biodiversity: Threats and opportunities. *Science* 378:eadf1466.

607 Ranarilalaitiana T, Razafindraleva HA, Granath G, Bukontaite Malm R, Rakotonirina JC,
 608 Razafindranaivo V, Ravaomanarivo LHR, Johansson F, et al. 2022. Remaining
 609 forests on the Central Highlands of Madagascar—Endemic and endangered
 610 aquatic beetle fauna uncovered. *Ecology and Evolution* 12:1–26. DOI:
 611 10.1002/ece3.9580.

612 Ringler M, Ursprung E, Hödl W. 2009. Site fidelity and patterns of short- and long-term
 613 movement in the brilliant thighed poison frog *Allobates femoralis* (Aromobatidae).
 614 *Behavioral Ecology and Sociobiology* 63:1281–1293. DOI:
 615 10.1163/156853891X00329.

616 Sæther BE, Coulson T, Grøtan V, Engen S, Altwegg R, Armitage KB, Barbraud C,
 617 Becker PH, et al. 2013. How life history influences population dynamics in
 618 fluctuating environments. *American Naturalist* 182:743–759. DOI: 10.1086/673497.

619 Sæther BE, Engen S, Møller AP, Weimerskirch H, Visser ME, Fiedler W, Matthysen E,
 620 Lambrechts MM, et al. 2004. Life-history variation predicts the effects of
 621 demographic stochasticity on avian population dynamics. *American Naturalist*
 622 164:793–802. DOI: 10.1086/425371.

623 Sinsch U. 2015. Review: skeletochronological assessment of demographic life-history
 624 traits in amphibians. *Herpetological Journal* 25:5–13.

625 Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASLL, Fischman DL, Waller
 626 RW. 2004. Status and trends of amphibian declines and extinctions worldwide.
 627 *Science* 306:1783–1786. DOI: 10.1126/science.1103538.

628 Suzzi-Simmons A. 2023. Status of deforestation of Madagascar. *Global Ecology and*
629 *Conservation* 42:e02389. DOI: 10.1016/j.gecco.2023.e02389.

630 Tessa G, Mattioli F, Vincenzo M, Andreone F. 2009. Egg numbers and fecundity traits in
631 nine species of *Mantella* poison frogs from arid grasslands and rainforests of
632 Madagascar (Anura: Mantellidae). *Madagascar Conservation & Development*
633 4:113–119.

634 Vences M, Glaw F, Bohme W. 1999. A review of the genus *Mantella* (Anura, Ranidae,
635 Mantellinae): taxonomy, distribution and conservation of Malagasy poison frogs.
636 *Alytes* 17:3–72.

637 Vences M, Crottini A, Rabemananjara FCE, Randrianantoandro C, Tsiorisoa GA. 2022.
638 Mantellidae, Mantellinae: *Mantella* and *Wakea*. In: Goodman SM, ed. *The New*
639 *Natural History of Madagascar*. Princeton: Princeton University Press, 1360–1365.

640 Vieilledent G, Grinand C, Rakotomalala FA, Ranaivosoa R, Rakotoarijaona J-R, Allnutt
641 TF, Achard F. 2018. Combining global tree cover loss data with historical national
642 forest-cover maps to look at six decades of deforestation and forest fragmentation
643 in Madagascar. *Biological Conservation* 222:189–197. DOI: 10.1101/147827.

644 Villero D, Pla M, Camps D, Ruiz-Olmo J, Brotons L. 2017. Integrating species
645 distribution modelling into decision-making to inform conservation actions.
646 *Biodiversity and Conservation* 26:251–271. DOI: 10.1007/s10531-016-1243-2.

647 Webb JK, Brook BW, Shine R. 2002. What makes a species vulnerable to extinction?
648 Comparative life-history traits of two sympatric snakes. *Ecological Research* 17:59–
649 67. DOI: 10.1046/j.1440-1703.2002.00463.x.

650 White GC, Burnham KP. 1999. Program MARK: Survival estimation from populations of
651 marked animals. *Bird Study* 46:S120–S139. DOI: 10.1080/00063659909477239.

652 Yoder AD, Campbell CR, Blanco MB, Dos Reis M, Ganzhorn JU, Goodman SM,
653 Hunnicutt KE, Larsen PA, Kappeler PM, Rasoloarison RM, Ralison JM, Swofford
654 DL, Weisrock DW. 2016. Geogenetic patterns in mouse lemurs (genus *Microcebus*)
655 reveal the ghosts of Madagascar's forests past. *Proceedings of the National*
656 *Academy of Sciences of the United States of America* 113:8049–8056. DOI:
657 10.1073/pnas.1601081113.

658 Zhang L, Lu X. 2012. Amphibians live longer at higher altitudes but not at higher
659 latitudes. *Biological Journal of the Linnean Society* 106:623–632. DOI:
660 10.1111/j.1095-8312.2012.01876.x.

Mis en forme : Français (France)