

Decadal changes and delayed avian species losses due to deforestation in the northern Neotropics

How avifauna respond to the long-term loss and fragmentation of tropical forests is a critical issue in biodiversity management. We use data from over 30 years to gain insights into such changes in the northernmost Neotropical rainforest in the Sierra de Los Tuxtlas of southern Veracruz, Mexico. This region has been extensively deforested over the past half-century. The Estación de Biología Tropical Los Tuxtlas, of the Universidad Nacional Autónoma de México (UNAM), protects a 640 ha tract of lowland forest. It became relatively isolated from other forested tracts between 1975 and 1985, but it retains a corridor of forest to more extensive forests at higher elevations on Volcán San Martín. Most deforestation in this area occurred during the 1970s and early 1980s. Forest birds were sampled on the station and surrounding areas using mist nets during eight non-breeding seasons from 1973 to 2004 (though in some seasons netting extended into the local breeding season for some species). Our data suggested local extinctions or declines in 12 species of birds subject to capture in mist nets. Six of the eight species no longer present were captured in 1992-95, but not in 2003-2004. Presence/absence data from netting and observational data suggested that another four low-density species also disappeared since sampling began. This indicates a substantial time lag between the loss of habitat and the apparent extirpation of these species. Delayed species loss and the heterogeneous nature of the species affected will be important factors in tropical forest management and conservation.

1

2 David W. Shaw^{1,2}, Patricia Escalante³, John H. Rappole⁴, Mario A. Ramos⁵,
3 Richard J. Oehlenschläger⁶, Dwain W. Warner⁵, and Kevin Winker¹

4

5 ¹ University of Alaska Museum, 907 Yukon Drive, Fairbanks, Alaska 99775
6 USA.

7

8 ² Current Address: 1540 Jones Road, Fairbanks Alaska 99709 USA.

9

10

11 ³ Colección Nacional de Aves, Instituto de Biología, UNAM, Circuito Exterior
12 s/n, Ciudad Universitaria, Copilco, Coyoacán, A. P. 70-233 Mexico, Distrito
13 Federal. C. P. 14510 MEXICO.

14

15 ⁴ Smithsonian's National Zoological Park, Conservation and Research Center,
16 1500 Remount Road, Front Royal, VA 22630 USA.

17

18 ⁵ Deceased.

19

20 ⁶ Science Museum of Minnesota, 120 W. Kellogg Blvd. St. Paul, MN 55102 USA.

21

22 Corresponding Author:

23 Kevin Winker

24 University of Alaska Museum

25 907 Yukon Drive

26 Fairbanks, AK 99775

27 kevin.winker@alaska.edu

28 907-474-7027

29

INTRODUCTION

Deforestation is one of the main threats to biodiversity conservation. Forest loss and fragmentation have caused declines or local extinctions among animal species at many locations (Turner, 1996; Fahrig, 2003; Dirzo & Raven, 2003). Local population declines and extinctions may be the most important leading indicators of biodiversity loss (Ceballos & Ehrlich, 2002; O'Grady et al., 2004). Bird losses have been documented in many forest systems (e.g., Willis, 1974, 1979; Leck, 1979; Karr, 1982; Bierregaard & Lovejoy, 1989; Kattan, Halvarez-Lopez, & Giraldo, 1994; Robinson, 1999; Sodhi, Liow, & Bazzaz, 2004; Ferraz et al., 2007; Patten, Gómez de Silva, & Smith-Patten, 2010; Laurance et al., 2011). Perhaps nowhere has this phenomenon been more noticeable than among tropical forests, where species losses have been documented in numerous taxonomic groups (e.g., Zimmerman & Bierregaard, 1986; Powell & Powell, 1987; Malcolm, 1988; Pahl, Winter, & Heinsohn 1988; Becker, Moure, & Peralta, 1991; Daily & Ehrlich, 1995; Brook, Sodhi, & Ng, 2003, Dirzo & Raven, 2003, Stuart et al. 2004; Robinson & Sherry, 2012). Species losses can occur at the landscape or patch levels and depend on the intensity of the change in forest cover, the distance to and size of other forest fragments, shape and size of the fragment, and other factors (Robbins, 1980; Lovejoy et al., 1984, 1986, Rolstad, 1991; Andrén, 1994; Faaborg et al., 1995; Lees & Peres, 2006; Barlow et al., 2006; Robinson & Sherry 2012). Tropical forest species, which often occur in small, low-density populations, may be particularly vulnerable to local extinctions (Terborgh & Winter 1980; Pimm, Jones, & Diamond, 1988;

54 Stotz et al., 1996).

55 Relatively few studies have assessed changes through decades,
56 however (Ewers & Didham, 2006). And although deforestation and
57 fragmentation can occur over a short period, some time may pass before
58 species begin to disappear from an affected area (Leigh, 1975, 1981; Karr,
59 1982, Tilman et al., 1994; Brooks, Pimm, & Oyugi, 1999). Thus, to fully
60 document the impact of deforestation on a forest community, a site must be
61 studied for a substantial period of time after habitat alteration has occurred.
62 Detailing the process of local population decline and extinction over time
63 provides invaluable information about species' abilities to cope with habitat
64 fragmentation and how community composition itself stabilizes following this
65 disturbance.

66 Studies of species losses in birds have used a variety of methods,
67 including comparing species richness in different-sized fragments (Willis,
68 1979; Nemark, 1991; Blake, 1991), comparison of species composition at a
69 site pre- and post-fragmentation (Willis, 1974; Leck, 1979; Bierregaard &
70 Lovejoy, 1989; Kattan, 1994), and experimental fragmentation (Lovejoy et al.,
71 1986; Bierregaard & Lovejoy, 1988, 1989; Ferraz et al., 2003, 2007; Laurance
72 et al., 2011), and have often included scattered survey data prior to
73 fragmentation (Willis, 1974; Leck, 1979; Kattan, Halvarez-Lopez, & Giraldo,
74 1994; Robinson, 1999; Patten, Gómez de Silva, & Smith-Patten, 2010). Many
75 of these studies have relied on qualitative visual and audio survey
76 techniques, with multiple observers, though such techniques can allow
77 cryptic and low-density species to be overlooked (Whitman, Hagan, &

Brokaw, 1997). Additionally, observer skills and intensity of sampling may vary among surveys.

Mist netting offers the most consistent and quantitative method available to sample birds among years (Rappole, Winker, & Powell, 1998). However, mist nets have documented weaknesses; the most relevant is the limited stratum and size of birds they can effectively sample (Remsen & Good, 1996; Whitman, Hagan, & Brokaw, 1997; Rappole, Winker, & Powell, 1998). This is particularly noticeable in structurally diverse habitats such as tropical rainforests, where probability of detection using mist nets is unknown for most species. Mist net studies in the Neotropics are therefore biased toward understory, small- to mid-sized passerines. While mist nets, unlike other methods, are less prone to observer bias and variability, we augmented our analyses of netting data that suggested species losses with presence-absence observational data (daily checklists in later years); this becomes particularly important for low-density species and for those not readily captured.

The Sierra de Los Tuxtlas of southern Veracruz, Mexico provides a textbook case of deforestation. This small range of volcanic mountains is home to the northernmost Neotropical rainforest (Pennington & Sarukhan, 1968; Dirzo & Miranda, 1991). The region has lost more than 90% of its forests in the past century, with the majority of that loss occurring in the lowlands over the past fifty years (Dirzo & Garcia, 1992; Rappole, Powell, and Sader, 1994; Winker, 1997). Our study compares eight seasons of mist net sampling from Los Tuxtlas over the course of more than thirty years. This

allows us to at least partly answer the question of how species composition and relative abundance changed in and around a conserved core of local rainforest habitat on a decadal scale.

METHODS

The Sierra de Los Tuxtlas is located in southern Veracruz, Mexico, 90 km southeast of Veracruz city. This range of mountains lies in the northwestern portion of the Isthmus of Tehuantepec and is isolated from the Sierra Madre Oriental by extensive lowlands. Los Tuxtlas encompass approximately 4,200 km², and the range is dominated by Volcán San Martín and Volcán Santa Marta, each reaching more than 1,500 m elevation. The Gulf of Mexico lies a short distance from the mountains to the north and east. The northernmost Neotropical evergreen rainforest formerly dominated the habitat in the region (Andrle, 1966; Pennington & Sarukhan, 1968; Dirzo & Miranda, 1991), but due to deforestation it is now a mosaic with a high percentage of pasture, cropland, fencerows, and isolated trees (pers. obs.; Dirzo & Garcia 1992; Estrada, Coates-Estrada, & Merritt 1997). Andrle (1966) estimated that 50% of the region was forested in 1962. By 1975 Rappole & Warner (1980) estimated that a third of the forests still stood. Just 15% of forest remained in 1986 (Winker, Rappole, & Ramos, 1990; Dirzo & Garcia, 1992), and in 1994 only 7-10% of the region was forested (Winker, 1997). Remaining forest occurs primarily in the highlands, and below 500 m forest is scarce (Rappole, Powell, & Sader, 1994; Mendoza, Fay, & Dirzo, 2005; Figs. 1, 2).

The climate in Los Tuxtlas is warm and wet, with a mean annual

temperature of 25 C, and annual precipitation is 4,500-4,900 mm, with a short dry season from March-May (Soto & Gama 1997). Canopy heights in primary forest range from 30-35 m (Ibarra-Manriquez et al., 1997). Second growth areas generally have variable canopy heights from 3-20 m (pers. obs.).

In 1967 the Universidad Nacional Autónoma de México established the Estación de Biología Los Tuxtlas, protecting a 640-ha tract of lowland rainforest (18° 34'30"N, 95° 04'20"W; González-Soriano, Dirzo, & Vogt, 1997). Over the following decades this site became largely isolated from other tracts of forest, although a corridor of forest remains, connecting to the more extensive upland forests on Volcán San Martín (Dirzo & Garcia, 1992; Fig. 1). The first intensive sampling of birds in the region began in 1973, and data from that effort are included here (see Winker, 1997).

During the non-breeding seasons of 1973-74 and 1974-75 Oehlenschläger, Ramos, Rappole, and Warner conducted the first intensive mist-netting efforts in the area. Sites extended through what was then contiguous rainforest from the biological station eastward to the coast (Fig. 3). In 1986, Rappole, Ramos, and Winker operated mist nets at the biological station, and Winker and Escalante continued work there from 1992 to 1994. In 2003-04 as part of a study of migrant birds, Shaw operated mist nets at the same location as Winker and Escalante's work in the 1990s. This study was approved by the University of Alaska Fairbanks IACUC (approval numbers: #00-33 & #04-03). Fieldwork occurred primarily during the non-breeding season. Effort was made to equally sample the available forest

types throughout the study period, although, in order to do this, habitat changes precluded using the same sites across all years (see Winker 1995; Fig. 3). Field effort as gauged by net hours also varied among years (Table 1).

Our earliest sampling occurred over a wider area than later seasons (Fig. 3). During the earliest sampling, large tracts of contiguous forest consisting of various microhabitats dominated the region and were sampled accordingly (Fig. 3). This broader expanse of forest likely provided habitat to more species than the current distribution of forest. This increased detection probabilities for some species such as *S. turdina*, which was rare even during our earliest sampling. Two general types of forest were present after fragmentation: primary forest and acahual (second growth). Because our sampling was forest-oriented, our efforts tracked the distribution of these habitats. Primary rainforest and second growth habitats were sampled in all efforts. We were unable to separate capture data by site for the early sampling periods; our findings therefore include data from the somewhat larger area from the station east to the coast. This sampling heterogeneity leads us to be cautious and conservative in our analyses and interpretations. Importantly, however, the same site and net lanes were used in the 1992-2004 efforts (sample periods 4-8 in Table 1).

Only resident species were used in our analyses due to seasonal migration and the high levels of variance in abundance this causes among obligate migrants. Changes in relative abundance were detected by comparing capture rates (birds per 1000 net hours) from each year of sampling. Species not captured in later sampling efforts and those with

apparently declining or increasing rates of capture were selected for more detailed analyses (instead of applying statistical tests across all species). Presence/absence patterns and observational data (daily checklists in later years) were also considered to provide insight into changes in abundance in low-density species that did not have sufficient samples for statistical testing. Species were considered for examination for presence/absence if they had not been captured since at least 1986-87. Vagrants, defined as those rarely encountered species whose ranges do not normally include the Sierra de Los Tuxtlas, were excluded (Winker et al., 1992; Howell & Webb, 1995). Only first-time captures were used in statistical analyses. Simple linear regression was used to detect changes in abundance for selected species. We looked for newly appearing species using presence/absence netting, observational, and specimen data. Daily checklists were used to augment mist-net data as a check to determine whether absence from the mist-net data was indicative of reality.

Species showing statistically significant declines and those not captured or observed in later sampling periods were categorized by preferred habitat (edge, forest, or semi-open), food preference (fruit/nectar or insects), elevational range, and whether Los Tuxtlas was at the periphery or core of its geographic range (Howell & Webb, 1995). These characteristics were used to assess whether certain traits of the species increased their vulnerability to local extirpation.

RESULTS

During this study we accumulated 165,083 net hours, equivalent to

37.7 net years if netting with a single net occurred twelve hours per day (Table 1). A species accumulation curve for a representative year (1992) with below-average net hours (12,605; mean = 20,220) showed that the avifauna was effectively fully sampled during most field seasons (Fig. S1, though in documenting a species' absence it is the among-season, aggregate sampling that is important). In total, 122 nonmigratory species were captured (Appendix).

Seven species showed statistically significant declines during the sampling period: *Phaethornis striilularis*, *Xenops minutus*, *Glyphorhynchus spirurus*, *Onychorhynchus coronatus*, *Myiobius sulphureipygius*, *Henicorhina leucosticta*, and *Eucometis penicillata* (Table 2). Of these taxa, four were captured throughout the sampling period: *P. striilularis*, *X. minutus*, *E. penicillata*, and *H. leucosticta*. *G. spirurus* was last captured in 1975, *O. coronatus* in 1986, and *M. sulphureipygius* in 1994, the last season of autumn netting. Four other species were captured in substantial numbers during early sampling periods but were not captured in later years: *Lepidocolaptes souleyetii*, *Ornithion semiflavum*, *Leptopogon amaurocephalus*, and *Coereba flaveola* (the latter may be an intratropical migrant in this region; Ramos, 1983); however, these species failed to show statistically significant declines in linear regression analyses, perhaps due to nonlinear declines. *L. souleyetii* was last captured in 1993-94, and the others were last captured in 1994-95. One species, *Hylomanes momotula*, was captured from 1986-1995 but not in the 1970s or in 2003-04. Though there were no captures in the 1970s, one individual was collected on 17 May 1974 a few km northeast of the station. A

similar pattern occurred in *Anabacerthia variegaticeps*, with captures occurring only in the 1990s. Only two species (*Trogon collaris* and *Xiphorhynchus flavigaster*) showed significant increases during the study period.

Presence/absence mist-net capture data for low-density species not captured after 1986-87 could be interpreted as suggesting that an additional 23 taxa were extirpated during the study (Table 3). However, we know from observational data that not all of these species were absent. These taxa included rarely captured species that are too large for effective mist-net capture or that prefer the forest canopy (e.g., *Micrastur ruficollis*, *Cotinga amabilis*), mixed/open habitat specialists (e.g., *Thraupis abbas* and *T. episcopus*), a small-stream specialist (*Chloroceryle aenea*), and highland species (e.g., *Myadestes unicolor*) that are either not prone to capture in mist nets or at our site. Species such as *Tityra inquisitor*, both *Thraupis* tanagers, and others were known to be present on the site or nearby but were not captured in later sampling periods. Four species of hummingbirds are included in Table 3, but due to inconsistent capture probabilities of low-density hummingbird species and non-definitive observational data with respect to accurate identification, we provide no hypotheses regarding their possible extirpation or persistence at the site; further work focusing on these species is warranted. There were six other species not in Tables 2 or 3 in which mist net data alone might suggest declines or absences (Appendix) during the entire study but which were present throughout from observational data; netting is not an effective sampling tool for these taxa

because of body size or forest stratum occupied (e.g., *Glaucidium*
brasilianum, *Ciccaba virgata*, and *Celeus castaneus*) or because forest
understory is not preferred habitat (e.g., *Pitangus sulphuratus*, *Myiozetetes*
similis, and *Volatinia jacarina*; Appendix). The first three of these species
require more focused study to determine abundances and possible declines.

Four lower-density species have likely been extirpated: *Taraba major*,
Formicarius analis, *Grallaria guatemalensis*, and *Schiffornis turdina* (Table 3).
One low-density species that might seem to have been extirpated from our
data, *Elaenia flavogaster*, is likely an intratropical migrant here (pers. obs.;
Howell & Webb, 1995; Table 3). Several species were captured only in later
sampling periods (Appendix) but were observed or collected throughout,
suggesting that there were no additions to the biological station's resident
avifauna during the study.

Based on all available data during the study (netting and observational
data), a minimum of 11 species of birds appear to have been extirpated from
the biological station over the past three decades. This translates into an
average loss of 3.7 species per decade or a local loss of 2.0% of the entire
Los Tuxtlas avifauna (561 spp.; Schaldach & Escalante, 1997), 4.1% of the
resident avifauna (269 spp.; Schaldach & Escalante, 1997), or 9.0% of the
resident species captured in our study (122 spp.; Appendix). All 16 species
showing significant declines or no longer present on the site prefer some
degree of forest cover (Table 4). Three species are edge specialists: *O.*
semiflavum, *O. mexicanus*, and *C. flaveola*. Twelve prefer closed canopy
forest: *P. striilularis*, *H. momotula*, *X. minutus*, *G. spirurus*, *T. major*, *F. analis*,

G. guatemalensis, *L. amaurocephalus*, *M. sulphureipygius*, *S. turdina*, *H. leucosticta*, and *E. penicillata*. *L. souleyetii* prefers semi-open or partly cleared forest.

Eleven of 16, or 68.8%, of the species showing declines or extirpations in this study are insectivores, whereas among all species captured 41% are insectivores. This trend was not significant, however (*G*-test with Williams' correction, $P > 0.1$).

The Sierra de Los Tuxtlas is the northernmost limit of the ranges of 13 of the 16 species showing declines. *G. guatemalensis* and *H. leucosticta* are the only species with a distribution extending substantially to the north and west of the study site. The field site is well within the elevational limits for all 16 species (Table 4).

The two species that significantly increased in abundance over the sample period (Table 4) both occur here at the core of their ranges, elevational distributions, and in their preferred forest habitat. *T. collaris* is a frugivore, and *X. flavigaster* is an insectivore.

DISCUSSION

Although the absence of a species is not a clear indication of extirpation, our sampling effort, despite its heterogeneity, does suggest that at minimum a species' absence indicates a decline. It is possible that some of the species now apparently gone from the station may persist in other, unsampled fragments. If the data presented here and our interpretation of them are accurate, the extirpation of species from the Estación de Biología Los Tuxtlas has been ongoing since its isolation. Such an "extinction debt" is

a recognized component of deforestation, and models of empirical data show that in birds this occurs across decades, but the species affected and the mechanisms of species loss remain poorly understood (Tilman et al., 1994; Ewers & Didham, 2003, Robinson & Sherry, 2012). Since 1973, 16 species susceptible to capture in mist nets have either become locally extirpated or are showing significant declines in abundance. The total number of losses and declines is undoubtedly higher than presented, because species not regularly captured in mist nets, such as large-bodied and canopy species, were not adequately surveyed. Species known to have been extirpated from Los Tuxtlas include *Sarcoramphus papa*, *Harpia harpyja*, and *Ara macao*. Patten, Gómez de Silva, & Smith-Patten (2010) also documented the extirpation of the latter two in Chiapas, Mexico. Many additional species have also been categorized as endangered or threatened in the Sierra de Los Tuxtlas (see Winker, 1997).

Our estimate of the average rate of avian losses from the station of 3.7 species per decade may not be directly comparable to other studies due to differences in habitat and sampling, but it is similar to the rate of loss observed at Barro Colorado Island by Robinson (1999) of 3.3 species per decade. Our estimate, however, includes only those taxa captured in mist nets, whereas Robinson's work included all species detected through observation.

Of the eight species with data sufficient for statistical analysis that showed local extirpation, six were lost between 1992 and 2004 (on the same site), suggesting a continuing extirpation of species from the station.

Bierregaard & Lovejoy (1988, 1989) found that as surrounding habitat was lost, species richness in remaining fragments increased as individuals displaced from surrounding areas found their way to remaining forest patches. This increased richness was limited by the lifespan of the individual birds (Bierregaard & Lovejoy 1988, 1989). Unlike these studies, in which forest patches were suddenly and completely isolated, the forest of the Estación de Biología Los Tuxtlas was isolated gradually. Because extirpation seems to be continuing, we expect declines and extinctions to continue for some time at the station, even if no further deforestation occurs in the region (Willis, 1974; Brooks, Pimm, & Oyugi, 1999, Robinson, 1999; Ferraz et al., 2003).

Mechanisms for tropical bird species losses due to deforestation and fragmentation probably include factors such as greater specialization as compared to temperate birds, reduced dispersal abilities, lower population densities, and patchy distributions (Robinson et al., 2004; Stratford & Robinson, 2005; Moore et al., 2008; Rompré et al., 2007). Our assessment of possible causes for the loss of these species reveals no definite patterns, however, other than the predominant requirement of forested habitat. On Barro Colorado Island in Lake Gatún, Panama, maturation of habitat and loss of open areas was responsible for the decline in the island's avifauna (Willis, 1974; Karr, 1982). This is unlikely to be the case in Los Tuxtlas. Despite major degradation of surrounding forests, the station has remained primary forest with areas of second growth. A loss of sapling and seedling species has been described (Dirzo & Miranda, 1990), but the overall structure of the forest

appears to have remained fairly stable. Vetter et al. (2011) found in a meta-analysis of 30 studies that the effects of fragmentation are not subject to simple generalities, and that they are highly site specific.

Los Tuxtlas is at the northernmost extent of the ranges of 13 of the 16 species we found to be declining or extirpated (Tables 3 and 4). Evidence is mixed as to whether populations at the periphery of a species' range are more vulnerable to extinction (Terborgh & Winter, 1980; Kattan, Halvarez-Lopez, & Giraldo, 1994; Johnson, 1998). Los Tuxtlas is at the edge of all species' geographic ranges endemic to Neotropical rainforest, so it is not clear why this subset might be more subject to this phenomenon. The elevational distribution of each of these species encompasses sea level to 750 m or more (Howell & Webb, 1995), and we consider this factor unlikely to be responsible for the vulnerability of these particular taxa.

Although insectivores showed a trend toward being disproportionately affected in our study, it was not significant. Elsewhere insectivores have been shown to be particularly vulnerable to severe habitat change (e.g., Kattan, Halvarez-Lopez, & Giraldo, 1994; Canaday, 1996; Johnson & Winker, 2010; Vetter et al., 2011). Additionally, deforestation can negatively impact species found in multi-species foraging flocks (Van Houtan et al., 2006), which are important to many birds of tropical rainforest communities (Willis, 1966; Morton, 1973; Buskirk, 1976; Rappole et al., 1983). Rappole & Morton (1985) noted that *X. minutus*, one of the species showing a significant decline in our study (Table 2) was a regular member of mixed flocks in the Sierra de Los Tuxtlas.

We considered large-scale range shifts, perhaps from climate change, as a possible cause for species loss, but this seems unlikely. At least some of the species lost in our study appear to have persisted in the southern portion of Los Tuxtlas near Volcán Santa Marta at least into the mid-1990s (Winker, pers. obs.). If range shifts were the cause, species would likely have disappeared region-wide and we would not expect only forest-related species to be affected. Habitat loss and degradation seem to be the best explanations for the losses observed, but exactly how these changes affected each species remains unknown.

Another possible influence on mist-net captures, particularly in the most recent, late winter/spring sampling periods, would be intra-tropical and elevational movements in some of the study species (Ramos, 1983, 1988). There is evidence that *C. flaveola* and *E. flavogaster* move seasonally within the tropics, seemingly to breed in Los Tuxtlas then departing (Ramos & Rappole, pers. obs.). Vega Rivera (1982) found probable elevational movements in *M. sulphureipygius*. The local extinctions of seven of the 16 species are particularly notable. *C. flaveola* is a widely distributed species known to thrive in manipulated habitats such as gardens and forest edges and is a generalist frugivore and nectarivore (pers. obs.; Howell & Webb, 1995). This is not a species we would expect to decline due to forest fragmentation; both its habitat and food preferences are well suited to survival in a mosaic landscape, and it is known to persist in a fragmented landscape elsewhere in northern Middle America (Johnson & Winker, 2010). Intratropical migrations of *C. flaveola* may partially explain the changing

390 capture rates in this species (Ramos & Rappole, pers. obs.). *O. semiflavum*
 391 and *L. amaurocephalus* are both edge specialists; thus, limited
 392 fragmentation, creating an increase in edges, might *a priori* seem to benefit
 393 these species. Though the habitat protected by the station has remained
 394 relatively static, the intensity of lowland deforestation in Los Tuxtlas as a
 395 whole (Fig. 2) may be too extensive even for these edge specialists. *L.*
 396 *souleyetii* prefers open forest and partially cleared areas (Howell & Webb,
 397 1995). The habitat surrounding the station during the 1980s and 1990s was
 398 dominated by pasture scattered with isolated trees. In our later field seasons
 399 there was a noticeable decline in the number of isolated trees and fences
 400 constructed of living trees (Winker, pers. obs.). This loss may account for the
 401 extirpation of *L. souleyetii*. *G. spirurus* apparently disappeared from the
 402 station between the 1970s and 1986, the first of the documented
 403 extirpations. The majority of deforestation across the region took place
 404 during this period. This previously abundant species disappeared from our
 405 data in just over a decade. Interestingly, on the slopes of neighboring Volcán
 406 Santa Marta the species was present at least into the 1990s and probably still
 407 persists there (Winker, pers. obs.). Also, Estrada, Coates-Estrada, & Merritt
 408 (1997) had observational data of the species' presence in the station area in
 409 1990-1992, indicating at least a decline if not extirpation (Table 2). In Brazil,
 410 *G. spirurus* persisted in experimentally isolated fragments well after isolation
 411 (Stouffer & Bierregaard 1995), and the species persists in highly fragmented
 412 forest in southern Belize (Johnson & Winker, 2010). *H. momotula* was
 413 collected but not netted in 1974, was captured in substantial numbers during

1986 and 1992-94, but was absent in the last two seasons of sampling. This pattern is mysterious. This species has an elevational range extending to 1500 m and may persist in the forests of the upper slopes of Volcán San Martín. If so, we speculate that the station may serve as a sink for this species, where habitat is insufficient for a self-sustaining population but may occasionally be colonized by dispersing individuals (see also Winker et al., 1997). Continued sampling may provide more insight into its abundance patterns. It illustrates the need for improved understanding of species-specific dispersal behavior within and among forest fragments (e.g., Van Houtan et al., 2007; Moore et al., 2008; Ibarra-Macias, Robinson, & Gaines 2011), which may be an important driver for patterns such as those we observed.

Two other studies provide comparative value to our results. The four species we consider likely extirpated (*Taraba major*, *Formicarius analis*, *Grallaria guatemalensis*, and *Schiffornis turdina*) were not detected in the much broader census surveys of Estrada, Coates-Estrada, & Merritt (1997) in 1990-1992. Patten, Gómez de Silva, & Smith-Patten (2010) conducted the geographically closest long-term study to ours in their analysis of avian declines at Palenque, Chiapas, Mexico. Their results showed only three species that overlapped our results. They found *Eucometis penicillata* extirpated (to our decline) and two others that declined as our populations did (*Xenops minutus* and *Leptopogon amaurocephalus*). Indeed, the species-level heterogeneity between our studies is noteworthy.

Our analyses suggest that the Estación de Biología Tropical Los Tuxtlas

is too small to maintain its full, historic complement of bird species. If deforestation accelerated region-wide, eliminating other forest refugia, the station alone (640 ha) would be unable to maintain the historical avian diversity of the region or to provide source populations for restored forest habitats for many of its present bird species. Given the scale of deforestation in the region, it is surprising that there are not more species showing declines. Indeed, we may consider it good news that important forest seed dispersers such as *Habia* tanagers (Puebla & Winker, 2004) did not show significant declines. The overall size of the remaining forests, particularly in the highlands, may be ameliorating the effects of lowland deforestation. However, increasing or continued isolation of the station will probably limit recolonization from elsewhere, and species losses will likely continue.

In our study, although several species seemed to quickly succumb to local and regional deforestation, others showed delayed declines and extirpations, a phenomenon also known to happen at larger scales (Tilman et al., 1994, Pimm et al., 2006). Moreover, the effects of deforestation were remarkably heterogeneous among forest-related species, with no single clear pattern of why some species experienced declines or extirpation. Our long-term data suggest that predicting which species will be most affected by deforestation in the northern Neotropics, and thus effectively working to ameliorate the effects of forest loss, will be particularly challenging. Nevertheless, as similar long-term datasets accrue, subtle patterns may reveal how species-specific responses reflect underlying commonalities that can be exploited for effective management and conservation.

ACKNOWLEDGMENTS

We thank the many field assistants who have helped us over the years and those who have provided and helped obtain permits to conduct this research. R. Barry, A. Powell, S. Pimm, and three anonymous reviewers provided excellent advice and comments.

LITERATURE CITED

- Andrén H. 1994. Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos* 71:355-366.
- Andrle RF. 1966. North American migrants in the Sierra de Tuxtla of southern Veracruz, Mexico. *Condor* 68:177-184.
- Barlow J, Peres CA, Henriques LMP, Stouffer PC, Wunderle JM. 2006. The responses of understory birds to forest fragmentation, logging and wildfires: An Amazonian synthesis. *Biological Conservation* 128:182-192.
- Becker P, Moure JS, Peralta FJA. 1991. More about euglossine bees in Amazonian forest fragments. *Biotropica* 23:586-591.
- Bierregaard RO, Lovejoy TE. 1988. Birds in Amazonian forest fragments: Effects of insularization. Pages 1564-1579 in: *Acta XIX Congressus Internationalis Ornithologici* (H. Oellet, ed.). University of Ottawa Press, Ottawa, Canada.
- Bierregaard RO, Lovejoy TE. 1989. Effects of forest fragmentation on Amazonian understory bird communities. *Acta Amazonica* 19:215-241.
- Blake JG. 1991. Nested subsets and the distribution of birds in isolated

- 486 woodlots. *Conservation Biology* 5:58-86.
- 487 Brook BW, Sodhi NS, Ng PKL. 2003. Catastrophic extinctions follow
- 488 deforestation in Singapore. *Nature* 424:420-423.
- 489 Brooks TM, Pimm SL, Oyugi JO. 1999. Time lag between deforestation and
- 490 bird extinction in tropical forest fragments. *Conservation Biology*
- 491 13:1140-1150.
- 492 Buskirk WH. 1976. Social systems in a tropical forest avifauna. *American*
- 493 *Naturalist* 110:293-310.
- 494 Canaday C. 1996. Loss of insectivorous birds among a gradient of human
- 495 impact in Amazonia. *Biological Conservation* 77:63-77.
- 496 Ceballos G, Ehrlich PR. 2002. Mammal population losses and the extinction
- 497 crisis. *Science* 296:904-907.
- 498 Daily GC, Ehrlich PR. 1995. Preservation of biodiversity in small rainforest
- 499 patches: rapid evaluations using butterfly trapping. *Biodiversity and*
- 500 *Conservation* 4: 35-55.
- 501 Dirzo R, Garcia MC. 1992. Rates of deforestation in Los Tuxtlas, a Neotropical
- 502 area in southeast Mexico. *Conservation Biology* 6:84-90.
- 503 Dirzo R, Miranda A. 1990. Contemporary Neotropical defaunation and forest
- 504 structure, function, and diversity, a sequel to John Terborgh.
- 505 *Conservation Biology* 4:444-447.
- 506 Dirzo R, Miranda A. 1991. Altered patterns of herbivory and diversity in the
- 507 forest understory: a case study of the possible consequences of
- 508 contemporary defaunation. Pages 273-287 in: *Plant-Animal*
- 509 *Interactions: Evolutionary ecology in tropical and temperate regions*

- 510 (Price PW, Lewinsohn TM, Fernandes GW, Benson WW, eds.). Wiley and
511 Sons, Inc., New York.
- 512 Dirzo R, Raven PH. 2003. Global state of biodiversity and loss. *Annual Review*
513 *of Environment and Resources* 28:137-167.
- 514 Estrada A, Coates-Estrada R, Merritt Jr DA. 1997. Anthropogenic landscape
515 changes and avian diversity at Los Tuxtlas, Mexico. *Biodiversity and*
516 *Conservation* 6:19-43.
- 517 Ewers RM, Didham RK. 2006. Confounding factors in the detection of species
518 responses to fragmentation. *Biological Reviews* 81:117-142.
- 519 Faaborg J, Brittingham M, Donovan T, Blake J. 1995. Habitat fragmentation in
520 the temperate zone. Pages 357-380 in: *Ecology and management of*
521 *Neotropical migratory birds: a synthesis and review of the critical*
522 *issues* (Finch DM, Martin TE, eds.). Oxford University Press, Cambridge,
523 United Kingdom.
- 524 Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annual*
525 *Review of Ecology Evolution and Systematics* 34:487-515.
- 526 Ferraz G, Russell GJ, Stouffer PC, Bierregaard RO, Pimm SL, Lovejoy TE. 2003.
527 Rates of species loss from Amazonian forest fragments. *Proceedings of*
528 *the National Academy of Sciences USA* 100:14069-14073.
- 529 Ferraz G, Nichols JD, Hines JE, Stouffer PC, Bierregaard RO, Lovejoy TE. 2007.
530 A large-scale deforestation experiment: Effects of patch area and
531 isolation on Amazon birds. *Science* 315:238-241.
- 532 González-Soriano E, Dirzo R, Vogt RC, eds. 1997. *Historia Natural de Los*
533 *Tuxtlas*. Universidad Nacional Autónoma de México, México, D.F.

- 534 Howell S, Webb S. 1995. *A guide to the birds of Mexico and northern Central*
535 *America*. Oxford University Press, New York.
- 536 Ibarra-Macias A, Robinson WD, Gaines MS. 2011. Experimental evaluation of
537 bird movements in a fragmented Neotropical landscape. *Biological*
538 *Conservation* 144:703-712.
- 539 Ibarra-Manríquez G, Martinez-Ramos M, Dirzon R, Núñez-Farfán J. 1997 La
540 vegetación. Pages 61–74 in *Historia Natural de Los Tuxtlas*
541 (González-Soriano E, Dirzo R, Vogt RC, Eds). Universidad Nacional
542 Autónoma de México, México, D.F.
- 543 Johnson AB, Winker K. 2010. Short-term hurricane impacts on a Neotropical
544 community of marked birds. *PLoS ONE* 5:e15109.
- 545 Johnson CN. 1998. Species extinction and the relationship between
546 distribution and abundance. *Nature* 394:272-274.
- 547 Kattan GH, Halvarex-Lopez H, Giraldo M. 1994. Forest fragmentation and bird
548 extinctions: San Antonio eighty years later. *Conservation Biology*
549 8:138-146.
- 550 Karr JR. 1982. Population variability and extinction in a tropical land-bridge
551 island.
552 *Ecology* 63:1975-1978.
- 553 Laurance WF, et al. 2011. The fate of Amazonian forest fragments: A 32-year
554 investigation. *Biological Conservation* 144:56-67.
- 555 Leck CF. 1979. Avian extinctions in an isolated tropical wet-forest preserve,
556 Ecuador. *Auk* 96:343-352.
- 557 Lees AC, Peres CA. 2006. Rapid avifaunal collapse along the Amazonian

deforestation frontier. *Biological Conservation* 133:198-211.

Leigh EG, Jr. 1975. Population fluctuations, community stability, and environmental variability. Pages 52-73 in: *Ecology and evolution of communities* (Cody ML, Diamond JM, eds.). Belknap Press, Cambridge, Massachusetts.

Leigh EG, Jr. 1981. The average lifetime of a population in a varying environment. *Journal of Theoretical Biology* 90:213-239.

Lovejoy TE, Rankin JM, Bierregaard RO, Brown KS, Jr., Emmons LH, Van Der Voort ME. 1984. Ecosystem decay of Amazon forest remnants. Pages 295-325 in: *Extinctions* (Nitecki MH, ed.). University of Chicago Press, Chicago, Illinois.

Lovejoy TE, Bierregaard RO, Rylands AB, Malcolm JR, Quintela CE, Harper LH, Brown KS, Powell AH, Powell GVN, Schubart HOR, Hays MB. 1986. Edge and other effects of isolation on Amazon forest fragments. Pages 257-285 in: *Conservation biology: The science of scarcity of diversity* (Soule ME, ed.). Sinauer Associates, Sunderland, Massachusetts.

Malcolm JR. 1988. Small mammal abundances in isolated and non-isolated primary forest reserves near Manaus, Brazil. *Acta Amazonica* 18: 67-83.

Mendoza E, Fay J, Dirzo R. 2005. A quantitative analysis of forest fragmentation in Los Tuxtlas, southeast Mexico: patterns and implications for conservation. *Revista Chilena de Historia Natural* 78:451-467.

Moore RP, Robinson WD, Lovette IJ, Robinson TR. 2008. Experimental evidence for extreme dispersal limitation in tropical forest birds.

- 582 *Ecology Letters* 11: 960-968.
- 583 Morton ES. 1973. On the evolutionary advantages and disadvantages of fruit
584 eating in tropical birds. *American Naturalist* 107:8-22.
- 585 Nemark WD. 1991. Tropical forest fragmentation and the local extinction of
586 understory birds in the eastern Usambara Mountains, Tanzania.
587 *Conservation Biology* 5:67-78.
- 588 O'Grady JJ, Reed DH, Brook BW, Frankham R. 2004. What are the best
589 correlates of predicted extinction risk? *Biological Conservation*
590 118:513-520.
- 591 Pahl LI, Winter JW, Heinsohn G. 1988. Variation in responses of arboreal
592 marsupials to fragmentation of tropical rainforest in north eastern
593 Australia. *Biological Conservation* 46:71-82
- 594 Patten MA, Gomez de Silva H, Smith-Patten BD. 2010. Long-term changes in
595 the bird community of Palenque, Chiapas, in response to rainforest loss.
596 *Biodiversity and Conservation* 19:21-36.
- 597 Pennington TD, Sarukhan J. 1968. *Arboles Tropicales de Mexico*. Instituto
598 Nacional de Investigaciones Forestales, D.F., Mexico.
- 599 Pimm SL, Jones HH, Diamond J. 1988. On the risk of extinction. *American*
600 *Naturalist* 132:757-785.
- 601 Pimm SL, Raven P, Peterson A, Şekercioğlu Ç, Ehrlich PR. 2006. Human
602 impacts on the rates of recent, present, and future bird extinctions.
603 *Proceedings of the National Academy of Sciences USA*
604 103:10941-10946.
- 605 Powell AH, Powell GVN. 1987. Population dynamics of male euglossine bees in

- 606 Amazonian forest fragments. *Biotropica* 19: 176-179.
- 607 Puebla F, Winker K. 2004. Dieta y dispersión de semillas de dos especies de
608 tangara (*Habia*) en dos tipos de vegetación en Los Tuxtlas, Veracruz,
609 México. *Ornitología Neotropical* 15:53-64.
- 610 Ramos MA. 1983. *Seasonal movements of bird populations at a Neotropical*
611 *study site in southern Veracruz, Mexico*. Ph.D. dissertation. University
612 of Minnesota, Minneapolis.
- 613 Ramos MA. 1988. Eco-evolutionary aspects of bird movements in the
614 northern Neotropical region. Pages 251-293 in: *Acta XIX Congressus*
615 *Internationalis Ornithologici*, Vol. I, (Ouellet H, ed.). University of Ottawa
616 Press, Ottawa, Canada.
- 617 Rappole JH, Warner DW. 1980. Ecological aspects of migrant bird behavior in
618 Veracruz, Mexico. Pages 353-393 in: *Migrant Birds in the Neotropics:*
619 *Ecology, Behavior, Distribution, and Conservation*, (Keast A, Morton ES,
620 eds.). Smithsonian Institution Press, Washington, D. C.
- 621 Rappole JH, Morton ES, Lovejoy TE, III, Ruos JL. 1983. *Nearctic avian migrants*
622 *in the Neotropics*. United States Fish and Wildlife Service, Washington
623 D.C.
- 624 Rappole JH, Powell GVN, Sader SA. 1994. Remote-sensing assessment of
625 tropical habitat availability for a Nearctic migrant: the Wood Thrush.
626 Pages 91-103 in: *Mapping the diversity of nature* (Miller RI, ed.).
627 Chapman and Hall, London.
- 628 Rappole JH, Winker K, Powell GVN. 1998. Migratory bird habitat use in
629 southern Mexico: mist nets versus point counts. *Journal of Field*

- 630 *Ornithology* 69:635-643.
- 631 Remsen JV Jr., Good DA. 1996. Misuse of data from mist-net captures to
- 632 assess relative abundance in bird populations. *Auk* 113: 381-398.
- 633 Robbins CS. 1980. Effect of forest fragmentation on breeding bird populations
- 634 in the piedmont of the mid-Atlantic region. *Atlantic Naturalist* 33:31-36.
- 635 Robinson WD. 1999. Long-term changes in the avifauna of Barro Colorado
- 636 Island, Panama, a tropical forest isolate. *Conservation Biology* 13:85-97.
- 637 Robinson WD, Sherry TW. 2012. Mechanisms of avian population decline and
- 638 species loss in tropical forest fragments. *Journal of Ornithology*
- 639 153:S141-S152.
- 640 Robinson WD, Angehr GR, Robinson TR, Petit LJ, Petit DR, Brawn JD. 2004.
- 641 Distribution of bird diversity in a vulnerable Neotropical landscape.
- 642 *Conservation Biology* 18:510-518.
- 643 Rolstad J. 1991. Consequences of forest fragmentation for the dynamics of
- 644 bird populations: conceptual issues and the evidence. *Biological Journal*
- 645 *of the Linnean Society* 42:149-163.
- 646 Rompré G, Robinson WD, Desrochers A, Angehr G. 2007. Environmental
- 647 correlates of avian diversity in lowland Panama rainforests. *Journal of*
- 648 *Biogeography* 34:802-815.
- 649 Schaldach W, Escalante P. 1997. Lista de Aves. Pages 571-573 in: *Historia*
- 650 *Natural de Los Tuxtlas* (González-Soriano E, Dirzo R, Vogt RC, eds.).
- 651 Universidad Nacional Autonoma de Mexico. D.F., Mexico.
- 652 Sodhi NS, Liow LH, Bazzaz FA. 2004. Avian extinctions from tropical and
- 653 subtropical forests. *Annual Review of Ecology Evolution and*

- 654 *Systematics* 35:323-345.
- 655 Soto M, Gama L. 1997. Climas. Pages 7-23 In: *Historia Natural de Los Tuxtlas*
- 656 (González-Soriano E, Dirzo R, Vogt RC, eds.). Universidad Nacional
- 657 Autonoma de Mexico. D.F., Mexico.
- 658 Stotz DF, Fitzpatrick JW, Parker TA, Moskovits DK. 1996. *Neotropical birds:*
- 659 *ecology and conservation*. University of Chicago Press, Chicago.
- 660 Stouffer PC, Bierregaard RO. 1995. Effects of forest fragmentation on
- 661 understory insectivorous birds. *Ecology* 76:2429-45.
- 662 Stratford JA, Robinson WD. 2005. Gulliver travels to the fragmented tropics:
- 663 geographic variation in mechanisms of avian extinction. *Frontiers in*
- 664 *Ecology and the Environment* 3:85-92.
- 665 Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, Waller
- 666 RW. 2004. Status and trends of amphibian declines and extinctions
- 667 worldwide. *Science* 306:1783-1786.
- 668 Terborgh J, Winter B. 1980. Some causes of extinction. Pages 119-133 In:
- 669 *Conservation biology: an evolutionary-ecological perspective* (Soule ME,
- 670 Wilcox BA, eds.). Sinauer Associates, Sunderland, Massachusetts.
- 671 Tilman D, May RM, Lehman CL, Nowak MA. 1994. Habitat destruction and the
- 672 extinction debt. *Nature* 371:65-66.
- 673 Turner IM. 1996. Species loss in fragments of tropical rain forest: a review of
- 674 the evidence. *Journal of Applied Ecology* 33:200-209.
- 675 Van Houtan KS, Pimm SL, Bierregaard RO, Jr., Lovejoy TE, Stouffer PC. 2006.
- 676 Local extinctions in flocking birds in Amazonian forest fragments.
- 677 *Evolutionary Ecology Research* 8:129-148.

- 678 Van Houtan KS, Pimm SL, Halley JM, Bierregaard RO Jr., Lovejoy TE. 2007.
679 Dispersal of Amazonian birds in continuous and fragmented forest.
680 *Ecology Letters* 10:219-229.
- 681 Vega Rivera JH. 1982. Aspectos biológicos de *Myiobius sulphureipygius* (Aves:
682 Tyrannidae) en el área de Santa Marta region de “Los Tuxtlas,”
683 Veracruz, Mexico. Professional thesis. Universidad Nacional Autónoma
684 de México.
- 685 Vetter D, Hansbauer MM, Végvári Z, Storch I. 2011. Predictors of forest
686 fragmentation sensitivity in Neotropical vertebrates: a quantitative
687 review. *Ecography* 34:1-8.
- 688 Whitman AA, Hagan JM III, Brokaw NVL. 1997. A comparison of two survey
689 techniques used in a subtropical forest. *Condor* 99:955-965.
- 690 Willis EO. 1966. The role of migrant birds at swarms of army ants. *Living Bird*
691 5:187-231.
- 692 Willis EO. 1974. Populations and local extinctions of birds on Barro Colorado
693 Island, Panama. *Ecological Monographs* 44:153-169.
- 694 Willis EO. 1979. The composition of avian communities in remanescent
695 woodlots in southern Brazil. *Papeis Avulsos de Zoolgia* 33:1-25.
- 696 Winker K. 1995. Habitat selection in woodland Nearctic-Neotropic migrants on
697 the Isthmus of Tehuantepec. I. Autumn migration. *Wilson Bulletin*
698 107:26-39.
- 699 Winker K. 1997. Introduction to the birds of Los Tuxtlas. Pages 535-541 in:
700 *Historia Natural de Los Tuxtlas* (González-Soriano E, Dirzo R, Vogt RC,
701 Eds.). Universidad Nacional Autónoma de México. D.F., México.

- 702 Winker K, Rappole JH, Ramos MA. 1990. Population dynamics of the Wood
703 Thrush (*Hylocichla mustelina*) on its wintering grounds in southern
704 Veracruz, Mexico. *Condor* 92:444-460.
- 705 Winker K, Oehlenschläger RJ, Ramos MA, Zink RM, Rappole JH, Warner DW.
706 1992. Avian distribution and abundance records for the Sierra de Los
707 Tuxtles, Veracruz, Mexico. *Wilson Bulletin* 104:699-718.
- 708 Winker K, Escalante P, Rappole JH, Ramos MA, Oehlenschläger RJ, Warner DW.
709 1997. Periodic migration and lowland forest refugia in a sedentary
710 Neotropical bird, Wetmore's bush-tanager. *Conservation Biology*
711 11:692-697.
- 712 Zimmerman BL, Bierregaard RO. 1986. Relevance of the equilibrium theory of
713 island biogeography and species-area relations to conservation with a
714 case from Amazonia. *Journal of Biogeography* 13:133-143.

715

Figure captions

717 Figure 1. Satellite view of Volcán San Martín, the northernmost volcano in the
718 Sierra de Los Tuxtlas, showing the distribution of forests (dark areas). The
719 study area is indicated by the white arrow (image from GoogleEarth, 2005).

720

721 Figure 2. Comparative views of Volcán Santa Marta and San Martín Pajapan in
722 the southern Sierra de Los Tuxtlas from 1973 (top; NASA/Skylab) and 2005
723 (bottom; GoogleEarth) showing extent of deforestation, particularly severe in
724 the lowlands.

725

726 Figure 3. Maps of the study area in the northern lowlands of the Sierra de Los
727 Tuxtlas showing a rough outline of all forests types (dark gray areas) in 1979
728 (top, from Landsat image), in 2005 (bottom, from GoogleEarth), and netting
729 sites (black polygons). Numbers indicate field season(s) site was used and
730 correspond to rows in Table 1.

731

732 Figure S1. Species accumulation curve for a representative year with below
733 average net hours (1992, 12,605 net hours).

734



735

736

737 Fig. 1

738
739



740




741

742

Fig. 2

743

744






Fig. 3

Table 1 (on next page)

Tables 1 through 4

Table 1. Sample effort and periods during eight nonbreeding seasons across three decades in the Sierra de Los Tuxtlas, Veracruz, Mexico.

Nonbreeding season	Net hours	Sampling period
1) 1973-74	33,976	15 Aug-26 May
2) 1974-75	36,512	7 Aug-29 May
3) 1986-87	4,310	17 Nov-16 Jan
4) 1992-93	12,605	5 Sep-15 Nov
5) 1993-94	41,142	25 Aug-20 May
6) 1994-95	22,509	15 Aug-15 Nov
7) 2002-03	8,395	21 Feb-27 Apr
8) 2003-04	2,312	5 Apr-29 Apr

Table 2. Outcomes of regression analyses for 14 species showing changes in abundance (capture rates; captures and rates are given in the Appendix) and those not detected in the later sampling periods. Those *P*-values presented in bold are significant at $\alpha = 0.05$.

Species	<i>F</i>	<i>P</i>	<i>R</i> ²	Last captured
<i>Phaethornis striilgularis</i> ^c	6.337	0.045	0.514	2002-03
<i>Hylomanes momotula</i> ^a	0.210	0.890	0.003	1994-95
<i>Trogon collaris</i> ^b	7.041	0.038	0.540	n/a
<i>Xiphorhynchus flavigaster</i> ^b	6.941	0.039	0.536	n/a
<i>Xenops minutus</i> ^c	7.578	0.033	0.558	2003-04
<i>Glyphorhynchus spirurus</i> ^{c,d}	7.529	0.034	0.557	1974-75
<i>Lepidocolaptes souleyetii</i> ^d	3.265	0.121	0.352	1992-93
<i>Ornithion semiflavum</i> ^d	0.327	0.588	0.052	1994-95
<i>Leptopogon amaurocephalus</i> ^d	2.814	0.144	0.319	1994-95
<i>Onychorhynchus coronatus</i> ^{c,d}	6.861	0.040	0.533	1986-87
<i>Myiobius sulphureipygius</i> ^{c,d}	10.555	0.019	0.629	1994-95
<i>Henicorhina leucosticta</i> ^{c,d}	6.740	0.041	0.529	2003-04
<i>Coereba flaveola</i> ^d	2.164	0.192	0.265	1994-95
<i>Eucometis penicillata</i> ^c	18.725	0.005	0.757	2002-03

- ^a Species captured 1986-1995. See text.
- ^b Species showing an increase in abundance.
- ^c Species showing a significant decline.
- ^d Species not captured in later sampling periods.

Table 3. Species not captured or observed from 1992-2004, seasons captured (from Appendix), presence on the field site in later sampling periods, and comments.

Species	Seasons Captured	Presence	Comments
<i>Micrastur ruficollis</i>	1	Y	observed
<i>Crypturellus boucardi</i>	3	Y	observed
<i>Heliomaster longirostris</i>	1	?	hummingbird
<i>Florisuga mellivora</i>	1	?	hummingbird
<i>Chlorostilbon canivetii</i>	2	?	hummingbird
<i>Hylocharis eliciae</i>	1, 2	?	hummingbird
<i>Chloroceryle aenea</i>	1, 2	Y	small streams
<i>Dryocopus lineatus</i>	2	Y	observed
<i>Synallaxis erythrothorax</i>	2	Y	observed
<i>Taraba major</i>	2	N	forest understory
<i>Formicarius analis</i>	1	N	forest understory
<i>Grallaria guatemalensis</i>	1, 3	N	forest understory
<i>Tityra inquisitor</i>	1	Y	observed, canopy
<i>Cotinga amabilis</i>	1	?	canopy
<i>Schiffornis turdina</i>	1	N	forest understory
<i>Polioptila plumbea</i>	1	Y	observed
<i>Myadestes unicolor</i>	1	Y	highlands
<i>Euphonia affinis</i>	2	?	none
<i>Thraupis abbas</i>	1	Y	observed
<i>Thraupis episcopus</i>	2	Y	observed
<i>Saltator atriceps</i>	1, 2	Y	observed
<i>Molothrus aeneus</i>	1	Y	observed

Table 4. Habitat, foraging preference, elevational range, and position within geographical distribution for 18 species of birds at the Estación de Biología Los Tuxtlas (from Howell and Webb, 1995).

Species	Habitat preference	Foraging guild	Elevational distribution (m)	Geographic distribution
<i>Phaethornis striilularis</i>	forest	nectarivore	0-1500	periphery
<i>Hylomanes momotula</i>	forest	frugivore	0-1500	periphery
<i>Trogon collaris</i>	forest	frugivore	0-2400	core
<i>Xenops minutus</i>	forest	insectivore	0-1000	periphery
<i>Xiphorhynchus flavigaster</i>	forest	insectivore	0-1500	core
<i>Glyphorhynchus spirurus</i> *	forest	insectivore	0-1200	periphery
<i>Lepidocolaptes souleyetii</i>	semi-open	insectivore	0-1500	periphery
<i>Taraba major</i> *	forest	insectivore	0-1600	periphery
<i>Formicarius analis</i> *	forest	insectivore	0-750	periphery
<i>Grallaria guatemalensis</i> *	forest	insectivore	50-3500	core
<i>Ornithion semiflavum</i>	edge	insectivore	0-1500	periphery
<i>Leptopogon amaurocephalus</i>	edge	insectivore	0-1300	periphery
<i>Onychorhynchus coronatus</i>	forest	insectivore	0-1200	periphery
<i>Myiobius sulphureipygius</i>	forest	insectivore	0-1000	periphery
<i>Schiffornis turdina</i> *	forest	frugivore	0-750	periphery
<i>Henicorhina leucosticta</i>	forest	insectivore	0-1300	core
<i>Coereba flaveola</i>	edge	frugivore	0-1000	periphery
<i>Eucometis penicillata</i>	forest	frugivore	0-750	periphery

* Presence/Absence data suggest species is extirpated.