

Effect of salt stress on different tiller positions in rice and the regulatory effect of Prohexadione calcium (#89342)

1

First revision

Guidance from your Editor

Please submit by **13 Jul 2024** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for guidance.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)
1 Rebuttal letter(s)
16 Figure file(s)
3 Table file(s)
1 Raw data file(s)




Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor






 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).





Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).




BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  **Impact and novelty is not assessed.** Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Effect of salt stress on different tiller positions in rice and the regulatory effect of Prohexadione calcium

Rongjun Zhang¹, Dianfeng Zheng^{Corresp., 1, 2, 3}, Naijie Feng^{Corresp., 1, 2, 3}, Linfeng¹, Jinning Ma¹, Xiayi Yuan¹, Junyu Huang¹, Lisha Huang¹

¹ College of Coastal Agricultural Sciences, Guangdong Ocean University, Zhanjiang, Guangdong, China

² South China Center of National Saline-Tolerant Rice Technology Innovation Center, Zhanjiang, Guangdong, China

³ Shenzhen Research Institute of Guangdong Ocean University, Shenzhen, Guangdong, China

Corresponding Authors: Dianfeng Zheng, Naijie Feng

Email address: zhengdf@gdou.edu.cn, fengnj@gdou.edu.cn

Soil salinization has resulted in a significant decrease in crop yields, particularly affecting the production of crops like rice (*Oryza sativa* L.). Prohexadione calcium (Pro-Ca) can enhance crop resilience against failure by managing plant height. However, its impact on various tiller positions during the tillering phase of rice under salt stress remains unknown. This study explores the distinct effects of salt stress on the physiological traits of the main stem and different tiller segments of rice plants, along with the role of Pro-Ca in mitigating salt stress. The findings revealed that under salt stress conditions, the number of tillers and leaves on the main stem decreased significantly in rice. Moreover, the levels of malondialdehyde (MDA) and H₂O₂ in the leaves and stems of each tiller position notably increased. The percentage of tillers experiencing reduction or elevation was higher than that of the main stem compared to the respective control. Application of Pro-Ca through foliar spraying under NaCl stress effectively alleviated the impact of salt stress on the tiller growth of rice during the tillering phase. It also boosted the activities of antioxidant enzymes like superoxide dismutase (SOD) and peroxidase (POD) in the leaves and stems of the tillers. Furthermore, it successfully mitigated the damage inflicted by salt stress on the cell membrane of rice tillers during the tillering phase. The regulatory effect of calcium on cyclic acid was particularly pronounced in alleviating the impact on the tillers under salt stress conditions.

Effect of salt stress on different tiller positions in rice and the regulatory effect of Prohexadione calcium

Rongjun Zhang¹, Dianfeng Zheng^{1, 2, 3*}, Naijie Feng^{1, 2, 3*}, Linfeng¹, Jinning Ma¹, Xiayi Yuan¹, Junyu Huang¹, Lisha Huang¹

¹ College of Coastal Agricultural Sciences, Guangdong Ocean University, Zhanjiang, Guangdong 524088;

² South China Center of National Saline-Tolerant Rice Technology Innovation Center, Zhanjiang, Guangdong 524088;

³ Shenzhen Research Institute of Guangdong Ocean University, Shenzhen, Guangdong 518108;

Corresponding Author:

Dianfeng Zheng^{1, 2, 3*}

College of Coastal Agricultural Sciences, Guangdong Ocean University, Zhanjiang, Guangdong 524088

Email address: zhengdf@gdou.edu.cn;

Naijie Feng^{1, 2, 3*}

College of Coastal Agricultural Sciences, Guangdong Ocean University, Zhanjiang, Guangdong 524088.

Email address: fengnj@gdou.edu.cn.

29

30 **Abstract**

31 Soil salinization has resulted in a significant decrease in crop yields, particularly affecting the
 32 production of crops like rice (*Oryza sativa* L.). Prohexadione calcium (Pro-Ca) can enhance crop
 33 resilience against failure by managing plant height. However, its impact on various tiller
 34 positions during the tillering phase of rice under salt stress remains unknown. This study
 35 explores the distinct effects of salt stress on the physiological traits of the main stem and
 36 different tiller segments of rice plants, along with the role of Pro-Ca in mitigating salt stress. The
 37 findings revealed that under salt stress conditions, the number of tillers and leaves on the main
 38 stem decreased significantly in rice. Moreover, the levels of malondialdehyde (MDA) and H₂O₂
 39 in the leaves and stems of each tiller position notably increased. The percentage of tillers
 40 experiencing reduction or elevation was higher than that of the main stem compared to the
 41 respective control. Application of Pro-Ca through foliar spraying under NaCl stress effectively
 42 alleviated the impact of salt stress on the tiller growth of rice during the tillering phase. It also
 43 boosted the activities of antioxidant enzymes like superoxide dismutase (SOD) and peroxidase
 44 (POD) in the leaves and stems of the tillers. Furthermore, it successfully mitigated the damage
 45 inflicted by salt stress on the cell membrane of rice tillers during the tillering phase. The
 46 regulatory effect of calcium on cyclic acid was particularly pronounced in alleviating the impact
 47 on the tillers under salt stress conditions.

48

49

50

51

52

53

54

55 **Introduction**

56 In recent years, soil salinization has become a global issue due to rising sea levels and expanding
 57 salinized land areas caused by global warming (Ahmed et al., 2021), severely reducing global
 58 crop yields and agricultural production (Munns & Tester, 2008; Dai et al., 2022). Enhancing crop
 59 salt tolerance and effectively addressing the damage from salt stress on yield has emerged as a
 60 critical research focus (And et al., 2011). Salt stress is a key abiotic factor that impacts the
 61 growth of most plants. In general, salt concentrations lead to changes in physiological and
 62 biochemical functions, limiting the growth and development of above-ground plant parts and
 63 root systems (dos Santos et al., 2022). When soil salinity increases, the water potential of the soil
 64 solution decreases below that of the plant root cells. This makes root uptake challenging and
 65 results in osmotic stress. Osmotic stress leads to the closure of plant stomata, hindering CO₂
 66 uptake, weakening photosynthesis, causing nutrient deficiencies. The accumulation of Na⁺ and
 67 Cl⁻ in cells affects mineral uptake and transport, inhibits enzyme activity, and leads to
 68 dehydration of plant cells. Salt stress can also result in the accumulation of reactive oxygen
 69 species (ROS), damaging cellular structures and biomolecules, thus limiting the growth of major
 70 crops like rice. (Hussain et al., 2018). Prolonged exposure to harsh environments has led to the
 71 evolution of a range of salt tolerance mechanisms in crops, including: changes in morphology,
 72 water relations, photosynthesis, hormones, ion distribution and biochemical adaptations (dos
 73 Santos et al., 2022), within certain limits, allow them to competitively obtain water from the soil
 74 and maintain nutrient balances in the body in response to ionic stresses thereby surviving adverse
 75 soil conditions (Hoang et al., 2016).

76 Rice (*Oryza sativa* L.), a moderately salt-sensitive crop (Joseph et al., 2010), is severely
 77 affected by salt stress in its growth and development (Zhang et al., 2012), and this effect varies
 78 according to developmental stage, degree and duration of stress, and variety (Zheng et al., 2023).
 79 It has shown that the effects of salt stress on rice germination and emergence are mainly
 80 characterized by a reduction in germination rate, germination speed, and germination energy,

leading to a reduction in shoot length, root length, and dry weight of rice (Taratima et al., 2022). Salt stress during the seedling period is mainly manifested in the damage to leaves and root system (Chang et al., 2019). Salt stress affects rice tillering mainly by reducing tillering capacity and delaying the reproductive process, and the duration of delay is positively correlated with the degree of salt stress, and primary and secondary tillers being more affected than the main stem. Additionally, salt stress reduces soil fertility and causes nutrient imbalance, and salinity stress inhibits nutrient uptake by the root system, ultimately leading to reduced tillering or tiller death due to nutrient deficits (Ruan et al., 2008). Primary stems and primary tillers contribute more to crop yield than secondary tillers due to asymmetric competitive advantages under stress conditions, and these advantages are associated with increased leaf number. Transportation of water and nutrients between the primary stem and tiller through the vascular bundles at the tiller nodes is essential for tiller development and survival (Yang et al., 2022). Salt stress in the formation of young spikes and spiking and flowering stage of rice is mainly manifested in the following ways: yellowing of leaves, delayed spiking, prolonged spiking period, increase in the number of degradation of glumes, shorter spike lengths, decrease in the number of solid grains, less full grains, more black rotting of roots in the late stage, early senescence, and ultimately affecting the yield of rice (Chang et al., 2019).

Plant growth regulators, as organic compounds with effects on growth and development similar to natural plant hormones, control plant growth by initiating various physiological and metabolic processes (Kaya et al., 2023; Zhao et al., 2023). The formation of the endogenous plant hormone gibberellin requires hydroxylase enzymes to catalyze a series of hydroxylation reactions, and these hydroxylases require 2-ketoglutarate as a coenzyme. Prohexadione calcium (Pro-Ca) imitates the coenzymes' structure and competitively hinders their function, thus impeding the synthesis of active gibberellins. Among these hydroxylation reactions, the reaction pathway for the formation of GA1 is the most sensitive to Pro-Ca, whereas the pathway for the formation of GA4 is not involved in the β -hydroxylation reaction, so that Pro-Ca selectively inhibits the synthesis of gibberellin GA1. GA1 is mainly found in the nutrient organs, controlling

the elongation and growth of stems and leaves, while GA4 is mainly found in the reproductive organs, controlling flower bud differentiation and hot grain development. Pro-Ca is an ideal dwarfing agent because of its strong synthetic activity in inhibiting GA1. Pro-Ca inhibits active gibberellin synthesis while protecting the activity of both surviving gibberellins, so Pro-Ca has dual activity on gibberellin metabolism (Kim et al., 2010; Ilias & Rajapakse, 2005). Pro-Ca has been shown by previous authors to have specific regulatory effects on rice, apple, strawberry, etc. (Kim et al., 2010; Kim et al., 2007; Lee et al., 1998). In previous studies, researchers have explored the risks of salt stress, the characteristics of tillering, and the mechanism of action of Pro-Ca acid. Our own research has shown that Pro-Ca acid can mitigate the harm caused by NaCl to the antioxidant capacity, photosynthetic properties, and cell membranes during the tillering stage in rice (Zhang et al., 2023; Huang et al., 2023). However, further investigations are necessary to understand the varying impacts of salt stress on the main stem and tiller, as well as the regulatory function of Pro-Ca.

In this study, we aimed to investigate the differential effects of salt stress on rice main stems and tillers and the regulatory role of calcium switched acid by comparing the relevant morphology building indexes, antioxidant enzyme activities, membrane damage indexes, and soluble protein contents in leaves and stems of rice main stems, first tillers and second tillers at the tillering stage under different treatments.

Materials and Methods

Materials and reagents

Huanghuazhan (conventional rice) was provided by Longping Seed Co. Ltd (Hunan, China), and Xiangliangyou900 (hybrid rice) was provided by Nianfeng Seed Science and Technology Co. Ltd (Hunan, China).

The original solution of the test regulator 5% Pro-Ca used in this experiment was provided by Sichuan Runer Technology Co. Ltd (Chengdu, Sichuan).

Experimental designs

Full and uniform rice seeds were selected, sterilized with 3% H₂O₂ for 15 min and then washed repeatedly with distilled water, distilled water was added until the seeds were submerged, and the seeds were soaked for 24 h at 30°C, after which they were germinated under dark conditions for 24 h. The experiment was selected to be carried out in the daylight linkage greenhouse of the College of Coastal Agriculture, Guangdong Ocean University, and the germinated seeds were uniformly sown on the rice-planting trays (specifications of 28~30 cm × 58~60 cm), about 5-8 seeds in each hole, and the soil used for seedling was a 3:1 mixture of brick red soil and nutrient soil.

After transplanting cultivation using caliber × bottom diameter × height of 19 × 15 × 18 cm plastic pots, each pot containing 3 kg of sun-dried soil, before transplanting a fixed amount of each pot to add 1 L of water, to be stabilized when the water surface line marking, and regularly replenish water to maintain the water layer. When the seedlings in the seedling tray were three leaves and one heart, the seedlings with consistent growth were selected and transplanted, and the depth of transplanting was about 1.5 cm, with 3 holes in each bucket and 1 plant in each hole. After the end of greening and before tillering, select the evening of sunny weather at about 16:00 to carry out regulator treatment through foliar spraying, about 10ml per pot, to ensure that the front and back of the leaf spraying evenly, in order to ensure its normal absorption. The regulator treatment was followed by a 0.3% salt treatment 48 h later. Tagging and tracking marking of tiller occurrence. Tillers were labeled with secondary and leaf positions. Ensure spatial distance between seedlings at each sampling to prevent competition between individuals due to different spatial size.

The experiment was set up with eight treatments, Xiangliangyou900 variety included four treatments as follows: XCK (distilled water), XS (0.3% NaCl), XPro-Ca (100 mg·L⁻¹ Pro-Ca), XPro-Ca+S (100 mg·L⁻¹ Pro-Ca + 0.3% NaCl), and Huanghuazhan variety included four treatments as follows: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca (100 mg·L⁻¹ Pro-Ca), and ZPro-Ca+S (100 mg·L⁻¹ Pro-Ca + 0.3% NaCl), with three replications per treatment. The leaves

and stems of the main stem, the first tiller, and the second tiller were taken every 7 d (7 d, 14 d, 21 d, 28 d, and 35 d after salt treatment) for the determination of related indexes.

Determination of morphological indices

Morphological indexes such as plant height, root length, and number of tillers were measured directly by using vernier calipers to measure the intersection of stem and root to determine the stem base width, and by using a leaf area meter (YX-1241) to measure the inverted two leaves and inverted three leaves of each tiller position, and by using the conventional drying method, the samples of each part of each treatment were killed for several and a half times for 30 min in an oven at 105°C, and then dried at 80°C to a constant weight and then determined the dry weight.

Determination of antioxidative enzyme activities

At 7 d, 14 d, 21 d, 28 d, and 35 d after NaCl stress, the leaves and stems of 15 rice plants with different tiller positions were rapidly frozen in liquid nitrogen, and then stored in -80°C, respectively. Three replicates of 0.5 g samples (leaves and stems weighed separately) were ground in liquid nitrogen, and then 10 ml of pre-cooled phosphate buffer (0.05 mM PBS, pH 7.8) was added, ground to homogenate, and then centrifuged at 6000 × g for 20 min at 4°C, and the supernatant was aspirated and set aside in a 4°C refrigerator.

3 ml of reaction solution (50 ml PBS pH 6.0 + 28 µL guaiacol + 19 µL H₂O₂ at 30% concentration) was mixed with 40 µL of supernatant. The absorbance was recorded every 30 s for 4 times and the dynamic absorbance was measured at 470 nm using a spectrophotometer (GENESYS 180 UV-Vis, Thermo Sci) to determine the peroxidase (**POD**) activity, 1 unit of enzyme activity for each minute of OD increase of 0.01.

0.1 ml of supernatant was taken and mixed with 2.9 ml of reaction solution (100 ml of PBS pH 7.0 + 0.05 ml of H₂O₂ at 30% concentration) and the absorbance at 240 nm was measured using a spectrophotometer ((GENESYS 180 UV-Vis, Thermo Sci) and recorded every 30 s for

four times. CAT activity was calculated using 0.01 decrease in OD per minute as a unit of enzyme activity (Aebi, 1984).

Superoxide dismutase (SOD) activity was determined using the nitro blue tetrazolium (NBT) method (Giannopolitis & Ries, 1977). 0.1 ml of supernatant was added to 2.9 ml of reaction mixture (2.61 ml MET + 0.097 ml EDTA- Na_2 + 0.097 ml NBT + 0.097 ml riboflavin) and irradiated for 20 min at 4000 lux light at 25°C. At the end of the reaction the absorbance at 560 nm was measured using the solvent in the unilluminated cuvette as a control tube, and the total activity of SOD was also calculated using 50% inhibition of NBT photochemical reduction as one enzyme activity unit (U).

0.1 ml of supernatant was mixed with the reaction solution (2.6 ml EDTA- Na_2 + 0.15 ml AsA + 0.15 ml H_2O_2) and the absorbance at 290 nm was determined by spectrophotometer recording every 30 s for 4 times. A change in absorbance of 0.01 in 1 min is defined as one unit of enzyme activity u, and enzyme activity is expressed as u/g (FW) (Nakano & Asada, 1981).

Determination of membrane damage index

Malondialdehyde (MDA) content was determined by TBA method (Guo et al., 2018). 10 ml of phosphate buffer (0.05 mM PBS, pH 7.8) was added to 0.5 g of sample and ground, then centrifuged at $10,000 \times g$ for 10 min at 4°C. 1 ml of the supernatant was taken and mixed with 2 ml of 0.6% TBA (thiobarbituric acid) in a centrifuge tube. The mixture was boiled in a boiling water bath for 15 min and then centrifuged at $10,000 \times g$ and 25°C for 10 min. The absorbance of the supernatant was measured spectrophotometrically at 450 nm, 532 nm, and 600 nm, respectively. The MDA content was calculated according to Equation: MDA content ($\text{mmol} \cdot \text{g}^{-1}$ FW) = $[6.452 \times (\text{OD}_{532} - \text{OD}_{600}) - 0.559 \times \text{OD}_{450}] \times V_t / (V_s \times W)$, where V_t is the total volume of extract, V_s is the volume of extract used for the assay, and W represents the fresh weight of the sample.

0.5 g of the sample was taken, 5 ml of 0.1% TCA solution was added, ground in liquid nitrogen and centrifuged at $10,000 \times g$ for 10 min. Then, 0.5 ml of the supernatant was added to

0.5 ml of 10 mM PBS buffer and 1 ml of KI solution, and the reaction was carried out in the dark at 28°C for 1 h. The H₂O₂ content of the sample was determined by spectrophotometric (GENESYS 180 UV-Vis, Thermo Sci) at 390 nm to determine the H₂O₂ content by measuring the absorbance (Jessup et al., 2018).

Subcellular localisation staining of plant leaf tissues for O₂⁻ and H₂O₂ was carried out with a slight modification of the method of Romero Puertas et al. (2004).

For histochemical detection of analytical O₂⁻: fresh leaves were washed with distilled water, evacuated, and placed in 10 mL centrifuge tubes containing 0.1% (w/v) nitrogen blue tetrazolium (NBT, pH 7.8) for 24 h. Leaves were rinsed with distilled water, 95% ethanol was added, and then boiled in a boiling water bath for 40 min, destained, and photographed.

Analysis Histochemical detection of H₂O₂: Fresh leaves were washed with distilled water, evacuated, and soaked in 1 mg/L diaminobenzidine 13,3'-tetrahydrochloride (DAB, pH 3.8) in a 10 mL plastic centrifuge tube for 24 h. Leaves were rinsed with distilled water, then boiled in 95% (v/v) ethanol for 40 min, decoloured, and photographed.

Histochemical staining of leaf cell death was referred to Martina Schraudner et al. (1998). The treated leaves were washed with distilled water, and the leaves were immersed in 0.25% (w/v) Evans blue solution for 24 h. The leaves were removed, rinsed with distilled water and placed in boiling anhydrous ethanol:glycerol (9:1) for 30 min, and chlorophyll was removed until the underside of the leaf appeared white and photographed.

Determination of the soluble protein content

To determine the soluble protein content, the method of Bradford (1976) was used with the Caumas Brilliant Blue G-250 staining method, 0.5 g of the sample was added to 10 ml of 0.05 mol/L pre-cooled phosphate buffer (pH 7.8) and ground in liquid nitrogen, and centrifuged at 12,000 × g at 4°C for 20 min, and the supernatant was the crude protein extract. The protein content was determined by adding 1 ml of enzyme solution to 5 ml of koammas brilliant blue solution and then shaking well, and the absorbance value at 595 nm was measured after 2 min of

reaction.

Statistical analyses

Using Excel 2016 for statistical analysis and SPSS 25.0 for further examination, one-way ANOVA and Duncan's method were employed for conducting ANOVA and multiple comparisons. The outcomes were presented as mean (X) \pm standard error (SE). Graphs were created using Origin 2018 software, with distinct lowercase letters denoting significant differences between treatments ($P < 0.05$).

Results

Effect of salt stress on morphological indexes at the tillering stage of rice and regulation by Pro-Ca

From the experimental results, it was found that salt stress negatively affected the tillering ability of both rice varieties (Fig. 1a and b), and the number of tillers decreased by 30.0%~44.43% and 12.52%~33.35% in Xiangliangyou900 and Huanghuazhan, respectively, from the 7th to the 35th days (Fig. 1c and d). The number of main stem leaves of Xiangliangyou900 decreased by 9.09%, 5.72%, 15.00%, and 2.04% on the 14th, 21st, 28th, and 35th days, respectively, and that of Huanghuazhan decreased by 6.90%, 21.05%, 11.11%, 12.20%, and 8.16% on the 7th, 14th, 21st, 28th, and 35th days, respectively, after NaCl stress (Fig. 1e and f).

The plant heights of Xiangliangyou900 and Huanghuazhan were reduced by 4.07%~11.60% and 10.31%~28.10%, respectively, compared with the control from the 7th to the 35th days after salt stress (Table 1). The first tiller length of both varieties was reduced by 2.34% to 29.22% and 13.02% to 38.46%, respectively, compared with the CK from the 7th to the 35th days after salt stress (Table 1). Compared with the main stem and the first tiller, the effect of NaCl stress on the length of the second tiller was more significant, and the length of the second tiller of Xiangliangyou900 and Huanghuazhan was reduced by 12.70%~31.86% and 19.60%~39.38% from the 7th to the 35th d after salt stress, respectively (Table 1). The stem base width of the

main stem of Xiangliangyou900 was reduced by 16.45% to 25.68% from 7th to 35th days after salt stress compared with the control, and that of Huanghuazhan was reduced by 17.41% to 38.99% from 7th to the 35th days after NaCl stress (Table 1). Salt stress also significantly reduced the stem base width of the first and second tillers of both varieties. Compared with the CK treatment, the stem base width of the first tiller was reduced by 16.42%~44.02% and 7.74%~43.86%, and the stem base width of the second tiller was reduced by 13.63%~36.32% and 26.62%~43.46%, respectively, in Xiangliangyou900 and Huanghuazhan from the 7th to the 35th days after the salt stress (Table 1).

The main stem leaf area of Xiangliangyou900 was reduced by 19.83% to 41.70% from the 7th to the 35th days after salt stress compared with the control, and the main stem leaf area of Huanghuazhan was reduced by 29.74% to 45.76% from the 7th to the 35th days (Table 2). Compared with the control, the leaf area of the first tiller decreased by 36.83% to 60.23% and 30.66% to 73.22%, and the leaf area of the second tiller decreased by 30.63% to 43.52% and 43.26% to 65.35%, respectively, in Xiangliangyou900 and Huanghuazhan from the 7th to the 35th days after salt stress (Table 2). In addition, compared with the CK, the root length of Xiangliangyou900 decreased by 25.73% to 48.08% from the 7th to the 35th days after salt stress, and the root length of Huanghuazhan decreased by 15.75% to 34% from the 7th to the 35th days after NaCl stress, which were significant differences (Table 2).

As shown in Table 3, the aboveground dry weight of the main stems of Xiangliangyou900 and Huanghuazhan decreased by 27.53% to 62.62% and 29.77% to 52.86%, respectively, compared with the control from the 7th to the 35th d after NaCl stress. Compared with CK, the dry weight of the first tiller decreased by 9.86% to 63.22% and 30.39% to 65.68%, and the dry weight of the second tiller decreased by 8.44% to 37.46% and 20.99% to 65.03%, respectively, in Xiangliangyou900 and Huanghuazhan from the 7th to the 35th days after salt stress (Table 3). In addition, the root dry weight of Xiangliangyou900 decreased by 34.32% to 70.88% from the 7th to the 35th days after NaCl stress compared with the control, and which of Huanghuazhan decreased by 36.33% to 70.21% (Table 3).

We can see from Fig. 1 that exogenous foliar application of Pro-Ca effectively alleviated the inhibitory effect of NaCl stress on the growth parameters of the two varieties. Foliar application of Pro-Ca under NaCl stress increased the number of tillers by 14.32% to 59.99% and 42.86% to 100.04% in Xiangliangyou 900 and Huanghuazhan, respectively, from the 7th to the 35th days. Compared with S treatment, the number of main stem leaves of both rice varieties increased significantly in Pro-Ca+S treatment, where the number of main stem leaves of Xiangliangyou900 increased by 4.16%, 23.33%, 3.03%, 23.53%, and 6.25%, respectively, and the number of main stem leaves of Huanghuazhan increased by 11.11%, 10.00%, 15.62%, 19.44% and 17.78%, respectively (Fig. 1e and f).

As shown in Table 1, compared with the control, foliar spraying Pro-Ca significantly reduced the plant height of the two rice varieties, in which the plant height of Xiangliangyou900 was reduced by 5.00%~15.00% and Huanghuazhan's plant height was reduced by 3.51%~20.42% from the 7th to the 35th days. The spraying of Pro-Ca reduced the first tiller length of Xiangliangyou900 and Huanghuazhan by 8.77%~30.02% and 9.90%~26.17% from the 7th to the 35th days, respectively, and the second tiller length of the two varieties was reduced by 6.91%~22.69% and 6.52%~38.96% (Table 1). Compared with the S treatment, foliar spraying Pro-Ca under NaCl stress significantly alleviated the stem base width of each tiller position in both rice, in which the main stem basal width, first tiller basal width and second tiller basal width of Xiangliangyou900 were increased by 15.24%~44.07%, 30.87%~45.30%, and 10.58%~48.32%, respectively, from the 7th to the 35th days, the stem base width of each tiller position increased by 23.84%~47.56%, 11.19%~76.13%, and 36.28%~76.96%, respectively (Table 1). Compared with S treatment, foliar spraying Pro-Ca under NaCl stress increased the leaf area of main stem, first tiller leaf area and second tiller leaf area of Xianglaingyou900 by 30.50%~65.47%, 30.11%~165.73%, and 38.19%~115.18% from the 7th to the 35th days, respectively, and the leaf area of each tiller position of Huanghuazhan increased by 19.78%~81.97%, 74.45%~329.32%, and 69.65%~204.06%, respectively (Table 2). The root lengths of Xiangliangyou900 and Huanghuazhan under Pro-Ca+S treatment increased by 13.43%

to 52.29% and 8.37% to 26.09%, respectively, from the 7th to the 35th days compared with that of S treatment, and the differences were significant (Table 2). As shown in Table 3, compared with NaCl, the main stem dry weight, first tiller dry weight and second tiller dry weight of Pro-Ca+S treatment of Xiangliangyou900 increased by 35.75%~137.61%, 2.51%~152.69%, and 28.29%~54.31%, respectively, and the dry weights of each tiller position of Huanghuazhan increased by 21.60%~88.90%, 61.33%~174.87%, and 61.22%~205.01%, respectively. Compared with the S treatment, foliar spraying of Pro-Ca alleviated the suppression of below-ground biomass by NaCl stress, and the root dry weight of Xiangliangyou900 increased by 23.64%~61.74% and that of Huanghuazhan increased by 39.22%~139.70% from the 7th to the 35th days (Table 3).

Effect of salt stress on antioxidant enzymes in rice leaves at each tiller position at tillering stage and regulation by Pro-Ca

Compared with the control, the SOD activity of the main stem leaves of Xiangliangyou900 increased by 6.88% to 31.25% from the 7th to the 35th days after salt treatment, and the SOD activity of the main stem leaves of Huanghuazhan decreased by 10.86% and 9.81% on the 7th and 14th days, respectively, and increased by 13.00% to 24.32% from the 21st to the 35th days (Fig. 2a and b). The SOD activity of the first tiller leaves of Xiangliangyou900 decreased by 12.55% and 9.39% on the 7th and 14th days after salt treatment, respectively, and that of the first tiller leaves of Huanghuazhan decreased by 14.48% on the 7th day, and did not show any significant difference compared with the control in the following days (Fig. 2c and d). Compared with the control, the SOD activity of the second tiller leaves of Xiangliangyou900 was significantly increased by 22.51% on the 7th day but did not change significantly from the 14th to the 35th days after NaCl treatment, however, the SOD activity of the second tiller leaves of Huanghuazhan decreased by 3.17% to 32.22% from the 7th to the 35th days (Fig. 2e and f).

As can be seen from Fig. 3a and b, compared with the control, salt stress reduced the CAT activity of main stem leaves of Xiangliangyou900 by 9.03% to 9.63% from the 7th to the 28th

days, and that of Huanghuazhan main stem leaves by 1.89% to 16.99% from the 14th to the 35th days. Meanwhile, as shown in Fig. 3, NaCl stress reduced the CAT activity of the second tiller leaves of Xiangliangyou900 by 0.50% to 32.15% from the 7th to the 35th days, respectively, and that of Huanghuazhan's first tiller leaves by 3.90% to 8.54% from the 21st to 35th days after salt stress, and that of the second tiller leaves by 0.78% to 5.62%.

NaCl stress reduced the POD activity of main stem leaves of Xiangliangyou900 by 2.83% and 13.67% on the 7th and 14th days, respectively (Fig. 4a), and that of Huanghuazhan by 13.01% and 6.55% , respectively, compared with the control (Fig. 4b). In addition, salt stress reduced the POD activity of the first tiller leaves of Xiangliangyou900 by 10.81% and 8.82% on the 7th and 14th days, and increased it by 5.12% and 24.40% on the 21st and 28th days, respectively, compared with the CK (Fig. 4c). The POD activity of the first tiller leaves of Huanghuazhan was reduced by 5.71% to 10.71% from the 7th to the 21st days and increased by 77.45% and 66.59% at the 28th and 35th days, respectively (Fig. 4d). Under salt stress, the POD activities of the second tiller leaves of Xiangliangyou900 and Huanghuazhan were reduced by 3.69%~44.75% and 4.67%~26.11%, respectively, compared with the control from the 7th to the 35th days (Fig. 4e and f).

The results in Fig. 5a and b showed that NaCl stress increased the APX activity of main stem leaves of Xiangliangyou900 by 9.02% to 89.09% from the 14th to the 35th days, and decreased the APX activity of Huanghuazhan's by 56.24% and 7.31% on the 7th and 14th days, respectively, and increased it by 8.55% to 16.30% from the 21st to 35th days. Compared with the CK treatment, the APX activity of the first tiller leaves of Xiangliangyou900 and Huanghuazhan decreased by 12.80%~54.26% and 3.38%~50.04%, respectively, from the 7th to the 35th days after NaCl stress (Fig. 5c and d), and that of the second tiller leaves of the two varieties decreased by 5.62%~34.90% and 2.74%~33.33%, respectively (Fig. 5e and f).

Under Pro-Ca+S treatment, the SOD activity of main stem leaves of Xiangliangyou900 significantly increased by 3.08% to 18.45% from the 7th to the 35th days, and that of Huanghuazhan was only increased by 4.76% and 0.09% on the 7th and 14th days, respectively

(Fig. 2a and b). The SOD activity of the first tiller leaves of both varieties were increased by 1.93% to 21.78% and 4.14% to 31.35% from the 7th to the 35th days after spraying with Pro-Ca compared to S treatment, respectively (Fig. 2c and d). In addition, it can be seen from Fig. 2e and f that under salt stress, spraying Pro-Ca increased the SOD activity of the second tiller leaves of Xiangliangyou900 by 11.64% and 10.13% on the 28th and 35th days, respectively, and that of Huanghuazhan by 5.16%~23.86% from the 21st to the 35th days.

Compared with salt stress, spraying Pro-Ca before salt stress had no significant effect on the CAT activity of leaves of all tillers of Xiangliangyou900, in which, it increased the CAT activity of leaves of the main stem of Huanghuazhan by 6.18%~17.17% from the 14th to the 28th days, respectively. Compared with the S treatment, spraying Pro-Ca increased the CAT activity of the first tiller leaves of Huanghuazhan by 2.37%~7.12% from the 14th to the 28th days, respectively (Fig. 3).

As can be seen in Fig. 4, the POD activity of the main stem leaves of Xiangtwoyou 900 under Pro-Ca+S treatment increased by 2.38% to 51.79% from the 7th to the 28th days, and that of the main stem leaves of Huanghuazhan increased by 8.27% to 26.63% from the 7th to the 21st days, as compared with that of S treatment. The POD activities of the first tiller and second tiller leaves of Xiangliangyou 900 under Pro-Ca+S treatment increased by 0.45% to 12.84% and 0.83% to 23.74%, respectively, from the 7th to the 35th days (Fig. 4c and e). In addition, the spraying of Pro-Ca increased the POD activity of the first tiller leaves of Huanghuazhan under salt stress by 4.84% to 26.07% from the 7th to the 28th days after salt stress, and that of the second tiller leaves by 9.82% to 34.71% from the 7th to the 35th days (Fig. 4d and f).

Compared with NaCl stress alone, spraying Pro-Ca under NaCl stress increased the APX activity of main stem leaves of Xiangliangyou900 by 8.48% to 38.46% from the 7th to the 35th days (Fig. 5a). The APX activity of main stem leaves of Huanghuazhan increased by 1.10% to 57.35% from the 7th to the 35th days, but not significantly (Fig. 5b). Spraying Pro-Ca under NaCl stress increased the APX activity of the first tiller leaves of Xiangliangyou900 and Huanghuazhan by 6.15%~79.71% and 19.02%~103.47%, respectively, from the 7th to the 35th

days (Fig. 5c and d). The APX activity of the second tiller leaves of Xiangliangyou900 was increased by 2.38%~40.31% from the 14th to the 35th days (Fig. 5e). The activity of the second tiller leaves of Huanghuazhan was increased by 0.15%~51.94% from the 7th to the 28th days (Fig. 5f).

Effect of salt stress on antioxidant enzymes in rice stems at each tiller position at tillering stage and regulation by Pro-Ca

Compared with the CK treatment, the SOD activity of the main stem of Xiangliangyou900 decreased by 7.64% to 30.56% from the 7th to the 35th days after NaCl treatment, and that of the main stem of Huanghuazhan decreased by 1.43% to 13.72% from the 21st to the 35th days (Fig. 6a and b). The SOD activity in the stem of the first tiller of Xiangliangyou900 decreased by 3.81% to 11.08% from the 7th to the 35th days after NaCl treatment, and that in the stem of the second tiller decreased by 1.88% to 32.36% from the 7th to the 28th days (Fig. 6c and e). NaCl stress reduced the SOD activity in the stem of the first tiller of Huanghuazhan by 4.98% to 52.21% from the 21st to the 35th days, and reduced the SOD activity in the stem of the second tiller by 3.27% to 26.61% from the 7th to the 35th days, respectively (Fig. 6d and f).

Fig. 7 shows that NaCl stress reduced the main stem CAT activity of Xiangliangyou900 by 5.85% to 25.60% from the 7th to the 28th days. The CAT activity of Huanghuazhan decreased by 6.67% and 26.94% on the 14th and 35th days, respectively, and increased but not significantly at the 7th, 21st, and 28th days. The CAT activity of the first tiller stems of Xiangliangyou900 decreased by 3.10% to 26.99% from the 21st to the 35th days, and that of Huanghuazhan decreased by 3.14%, 3.27%, and 45.44% at the 7th, 21st, and 35th, respectively (Fig. 7c and d). NaCl stress reduced the CAT activity in the stem of the second tiller of Xiangliangyou900 by 8.69% to 38.70% from the 7th to the 28th days after NaCl stress, and that of Huanghuazhan by 10.25% to 37.52% from the 7th to the 35th days (Fig. 7e and f).

Compared with their respective CKs, the POD activity of main stem of Xiangliangyou900 under NaCl stress increased by 5.02% to 44.88% from the 7th to the 35th days, and that of

Huanghuazhan increased by 7.39% to 11.37% from the 7th to the 21st days, and decreased by 18.49% and 35.13% at the 28th and 35th, respectively (Fig. 8a and b). The POD activity of the first tiller stems of Xiangliangyou900 decreased by 5.51% to 45.94% in 7~35 days under salt stress (Fig. 8c). The POD activity in the stem of the first tiller of Huanghuazhan decreased by 2.64% to 10.67% from the 7th to the 28th days, and increased by 4.11% at the 28th day (Fig. 8d). Compared with the control, salt stress reduced the POD activity in the stem of the second tiller by 2.48% to 25.92% and 2.68% to 25.68% in the stem of the second tiller of Xiangliangyou900 and Huanghuazhan, respectively, from the 7th to the 35th days (Fig. 8e and f).

The main stem APX activity of Xiangliangyou900 decreased by 12.12% to 39.98% from the 7th to the 35th days after NaCl stress, and that of Huanghuazhan decreased by 2.65% to 34.28%, respectively (Fig. 9a and b). The APX activity in the stem of the first and second tillers of both varieties decreased by 2.57% to 49.67% and 5.29% to 64.09%, 21.47% to 63.03% and 21.35% to 34.96%, respectively, from the 7th to the 35th days after salt stress (Fig. 9).

Compared with the S treatment, spraying Pro-Ca under salt stress increased the SOD activity of the main stem of Xiangliangyou900 by 13.66% to 65.88% from the 7th to the 28th days, and that of the main stem of Huanghuazhan by 2.05% to 7.96% from the 21st to the 35th days (Fig. 6a and b), the SOD activities in the stem of the first tiller increased by 6.34%~51.90% and 6.91%~97.69% from the 7th to the 35th days in Xiangliangyou900 and Huanghuazhan, respectively (Fig. 6c and d). Spraying Pro-Ca under salt stress increased the SOD activity of the second tiller stems of Xiangliangyou900 by 2.68% to 123.51% from the 14th to the 35th days, and that of the second tiller stems of Huanghuazhan by 3.27% to 26.61% from the 7th to the 35th days (Fig. 6e and f).

Compared with the S treatment, the CAT activities of main stems sprayed with Pro-Ca Xiangliangyou900 and Huanghuazhan under salt stress increased by 5.72% to 92.16% and 6.04% to 40.09% from the 7th to the 35th days, respectively (Fig. 7a and b). Foliar spraying of Pro-Ca also effectively increased the CAT activity in the first tiller stems of two rice varieties under salt stress, which was increased by 0.83%~30.00% at the 7~28 days in Xiangliangyou900 and by

3.59%~95.57% at the 7~35 days in Huanghuazhan (Fig. 7c and d). CAT activity in the second tiller stems of both varieties was increased by 0.93% to 66.28% and 7.57% to 65.13% under Pro-Ca+S treatment compared to S, respectively (Fig. 7e and f).

Compared with the S treatment alone, spraying Pro-Ca under salt stress increased the main stem POD activity by 5.89% to 51.39% and 4.76% to 33.32% from the 7th to the 35th days in Xiangliangyou900 and Huanghuazhan, respectively. The POD activity in the first and second tiller stems of both varieties increased by 0.18% to 24.07% and 13.49% to 98.03%, 14.51% to 31.29% and 4.15% to 33.22%, respectively, from the 7th to the 35th days (Fig. 8).

Compared with salt stress alone, spraying Pro-Ca under NaCl stress increased the main stem APX activity of Xiangliangyou900 by 11.43% to 87.63% from the 7th to the 35th days, and that of Huanghuazhan by 10.98% to 31.58%, respectively (Fig. 9a and b). It increased the APX activity of the first and second tiller stems of both varieties by 0.85% to 103.18% and 10.74% to 99.20%, 4.11% to 34.69% and 18.49% to 83.24%, respectively, from the 7th to the 35th days (Fig. 9).

Effect of salt stress on membrane damage index in rice leaves at each tiller position at tillering stage and regulation by Pro-Ca

Salt stress significantly increased the MDA content of leaves of two rice varieties compared with the control (Fig. 10). Among them, the MDA contents of main stem, first tiller and second tiller leaves of Xiangliangyou900 increased by 13.04%~54.61%, 0.58%~79.40%, and 3.33%~39.59%, respectively, from the 7th to the 35th days after salt stress (Fig. 10a, c, and e), the MDA content of the leaves of each tiller position of Huanghuazhan increased by 10.62% to 127.93%, 13.25% to 75.94%, and 6.01% to 64.67%, respectively, from the 7th to the 35th days after salt stress (Fig. 10b, d, and f). In addition, as seen in Fig. 11, salt stress increased the H₂O₂ content of the main stem leaves of Xiangliangyou900 and Huanghuazhan by 4.57%~38.51% and 3.88%~21.84% at 7~35 days, and the H₂O₂ content of the first tiller leaves of two kinds of rice by 5.24%~39.80% and 0.44%~41.06%. The H₂O₂ content of the second tiller leaves of the two varieties was

increased by 16.46%~28.88% and 8.37%~58.75% at 7~35 days, respectively (Fig. 11).

Compared with S treatment, foliar spraying of Pro-Ca before salt stress decreased the MDA content of main stem leaves of Xiangliangyou900 and Huanghuazhan by 0.53%~27.59% and 8.65%~25.00%, respectively (Fig. 10a and b), and that of the first tiller leaves of the two varieties under the same conditions by 17.24%~26.80% and 2.39%~29.12%, respectively (Fig. 10c and d), from the 7th to the 35th days. The MDA contents of second tiller leaves of the two varieties under the same conditions were decreased by 0.09%~34.98% and 1.60%~26.93%, respectively (Fig. 10e and f). Spraying Pro-Ca before salt stress reduced the H₂O₂ content of main stem leaves of Xiangliangyou900 and Huanghuazhan by 0.63%~22.87% and 5.91%~42.28%, respectively, from the 7th to the 35th days (Fig. 11a and b). Spraying Pro-Ca reduced the H₂O₂ content of the first tiller leaves of Xiangliangyou900 by 4.03%~31.42% and that of Huanghuazhan by 7.56%~26.04%, respectively at 7~35 days (Fig. 11c and d), and reduced the H₂O₂ content of the second tiller leaves of both rice varieties by 5.49%~29.10% and 8.23%~34.97% (Fig. 11e and f). The staining test revealed that there were more spots on the leaves of both rice varieties under salt stress, and the spots on the tiller leaves were larger in area and darker in color. In contrast, the area of spots on the leaves of the treatments sprayed with Pro-Ca before salt stress decreased and became lighter in color (Fig. 12).

Effect of salt stress on membrane damage index in rice stems at each tiller position at tillering stage and regulation by Pro-Ca

Compared with the control, NaCl stress increased the MDA content of the main stem of Xiangliangyou900 and Huanghuazhan by 14.85%~115.59% and 39.38%~95.94%, respectively (Fig. 13a and b), and that of the first tiller stems of Xiangliangyou900 and Huanghuazhan by 4.98%~90.72% and 9.55%~72.13%, respectively, from the 7th to the 35th days after salt stress (Fig. 13c and d), and the MDA content of the second tiller stems of the two varieties increased by 22.21%~156.61% and 7.98%~61.68%, respectively (Fig. 13e and f). NaCl stress at 0.3% also significantly increased the H₂O₂ content in the stems of both rice varieties (Fig. 14). Among

them, salt stress increased the H_2O_2 content of the main stem of Xiangliangyou900 by 1.28% to 6.12% from the 7th to the 35th days, and increased the H_2O_2 content of the main stem of Huanghuazhan by 7.61% to 24.09% from the 7th to the 28th days (Fig. 14a and b). Salt stress increased the H_2O_2 content of the first tiller stem of Xiangliangyou900 by 17.28%, 14.83%, and 8.49% at the 7th, 14th, and 35th days, respectively, and that of Huanghuazhan by 0.63%~9.16% at 14~35 days (Fig. 14c and d). In addition, NaCl stress increased the H_2O_2 content in the stem of the second tiller of Xiangliangyou900 by 1.37%~18.04% from the 7th to the 35th days, and there was no significant change on the 28th day, that of Huanghuazhan increased the content by 2.67% and 28.36% on the at the 7th and 21st days (Fig. 14e and f).

Compared with S treatment, spraying Pro-Ca before salt stress caused a significant decrease in MDA content, in which the MDA content of the main stems of Xiangliangyou900 and Huanghuazhan decreased by 2.53% to 48.87% and 14.94% to 35.39%, respectively, from the 7th to the 35th days (Fig. 13a and b). The MDA content of the first tiller stems of the two varieties sprayed with Pro-Ca decreased by 3.79%~46.45% and 5.39%~27.28%, respectively, from the 7th to the 35th days after salt stress (Fig. 13c and d), compared with S treatment, spraying Pro-Ca decreased the MDA content of the second tiller stems of Xiangliangyou900 and Huanghuazhan by 1.70%~66.09% and 0.32%~37.09% from the 7th to the 35th days, respectively (Fig. 13e and f). Foliar spraying of Pro-Ca before salt stress reduced H_2O_2 in the main stem of Xiangliangyou900 by 7.07%~12.62% from the 7th to the 28th days, and that of Huanghuazhan by 2.79%~14.74% from the 7th to the 35th days (Fig. 14a and b), and reduced the H_2O_2 content in the stem of the first tiller of Xiangliangyou900 and Huanghuazhan by 9.58%~15.60% and 3.20%~12.45% from the 7th to the 35th days respectively (Fig. 14c and d). Compared with salt stress, spraying Pro-Ca reduced the H_2O_2 content in the stem of the second tiller of the two rice varieties by 1.35%~33.05% and 7.92%~24.15%, respectively, from the 7th to the 35th days (Fig. 14e and f).

Effect of salt stress on soluble protein content in rice leaves at each tiller position at

tillering stage and regulation by Pro-Ca

As can be seen from Fig. 15, the soluble protein content of the main stem leaves of Xiangliangyou900 increased by 0.88% to 2.59% from the 7th to the 28th days after salt stress, and the soluble protein content of the main stem leaves of Huanghuazhan decreased by 1.59% and 2.87% on the 7th and 14th days after salt stress, respectively, and increased by 0.44% to 2.41% from the 21st to 35th d. NaCl stress increased the soluble protein content of the first tiller leaves of Xiangliangyou900 by 0.61% and 5.33% on the 21st and 28th days, respectively, and decreased the soluble protein content of the first tiller leaves of Huanghuazhan by 2.06% to 4.22% from the 7th to the 28th days after NaCl stress (Fig. 15c and d). The soluble protein content of the second tiller leaves of Xiangliangyou 900 increased by 3.27% and 0.45% on the 7th and 21st after NaCl stress, and decreased that of Huanghuazhan by 0.31% to 4.63% from the 7th to the 35th days (Fig. 15e and f).

Compared with the S treatment, the soluble protein content of the main stem leaves of Xiangliangyou900 increased by 0.21% to 4.67% from the 7th to the 35th days in Pro-Ca+S treatment, and that of the main stem leaves of Huanghuazhan increased by 1.28% to 1.90% from the 7th to the 21st days (Fig. 15a and b). In addition, spraying Pro-Ca under salt stress increased the soluble protein content of the first tiller leaves of Xiangliangyou900 by 4.30%, 0.25%, and 1.94% at the 7th , 14th , and 35th, respectively, and that of Huanghuazhan by 0.57% to 4.64% from the 7th to the 35th days (Fig. 15c and d). Foliar spraying of Pro-Ca increased the soluble protein content of the second tiller leaves of Xiangliangyou900 by 0.55% to 4.38% from the 14th to the 35th days and Huanghuazhan by 0.16% to 5.15% from the 7th to the 35th days after salt stress (Fig. 15e and f).

Effect of salt stress on soluble protein content in rice stems at each tiller position at tillering stage and regulation by Pro-Ca

As can be seen from the Fig. 16a and b, the soluble protein content of the main stem of Xiangliangyou900 increased by 1.92% to 9.30% from the 14th to the 35th days after salt stress,

and the soluble protein content of the main stem of Huanghuazhan increased by 1.03% to 7.41% from the 7th to the 28th days. Salt stress reduced the soluble protein content of the first tiller stems of Xiangliangyou900 by 0.31%, 5.92%, and 3.83% at the 7th, 28th, and 35th days, and reduced the soluble protein content of the first tiller stems of Huanghuazhan by 1.48% and 1.12% at the 21st and 35th days, respectively (Fig. 16c and d). The soluble protein content of the second tiller stems of Xiangliangyou900 decreased by 0.88% to 12.45% from the 7th to the 35th days after salt stress, and there was no significant difference at the 14th day, while that of the second tiller stems of Huanghuazhan decreased by 4.81% to 12.49% from the 14th to the 35th days (Fig. 16e and f).

Spraying Pro-Ca under salt stress conditions increased the soluble protein content of the main stems of Xiangliangyou900 and Huanghuazhan by 0.97%~9.13% and 2.05%~15.38%, respectively, from the 7th to the 35th days after NaCl stress (Fig. 16a and b). Compared with the S treatment, the soluble protein content of the first tiller stems of Xiangliangyou900 under Pro-Ca+S treatment increased by 0.90% to 14.89% from the 7th to the 35th days, but there was no significant difference at the 21st day, and that of the first tiller stems of Huanghuazhan under Pro-Ca+S treatment increased by 3.89% to 22.64% from the 7th to the 35th days (Fig. 16c and d). Foliar spraying of Pro-Ca under NaCl stress increased the soluble protein content of the second tiller stems of Xiangliangyou900 and Huanghuazhan by 8.09% to 26.44% and 2.11% to 17.37%, respectively, from the 7th to the 35th days after salt stress (Fig. 16e and f).

Discussion

Salt stress, as a significant abiotic stressor, can greatly inhibit crop growth and development. It has the ability to trigger the production of ROS, an excess of which can harm cellular functions and even result in cell death. ROS also act as secondary messengers, activating molecular processes in response to environmental stresses. Therefore, maintaining an appropriate level of ROS is crucial for plant survival under stress conditions (Lee et al., 2022). However, excessive salt stress can disrupt the balance of internal ions in cells (Zulfiqar & Ashraf, 2021). High

concentrations of salt stress also increase the production of reactive oxygen species (ROS), such as mono-linear oxygen ($^1\text{O}_2$), superoxide (O_2^-), hydrogen peroxide (H_2O_2), and hydroxyl radicals (OH) (Ding et al., 2022). Plants have evolved complex mechanisms to combat salt stress-induced oxidative stress, such as antioxidant enzymes like superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD). These enzymes are crucial for scavenging ROS, and an elevation in ROS levels triggers an upsurge in the activities of SOD, POD, CAT, and other enzymes during salt stress conditions. (Huihui et al., 2020). Salt stress damage to crops has been confirmed in various crops (Kousar et al., 2021; Jung et al., 2017; Feng et al., 2023). Jian et al. (2022) demonstrated that growth rate was reduced after 1 day of 100 mM NaCl stress in rice, and more severe wilting symptoms appeared on the tips of rice plant leaves after 15 days. 'IR29' was particularly damaged, which contained high levels of ROS. The results of this experiment indicated that the main stem leaves exhibited a stronger antioxidant capacity under salt stress due to increased SOD, POD, and APX activities. CAT activity decreased in all tillers under salt stress, while the main stem demonstrated higher salt tolerance compared to the tillers by enhancing SOD, CAT, and POD enzyme activities and experiencing a less significant reduction in APX. It was speculated that this discrepancy could be attributed to a lower level of stress in the main stem and a greater resilience in the main stem in contrast to the tillers. Yang et al. (2022) suggested that since tillers mature later than the main stem, the main stem consistently maintains a growth and development advantage. Prolonged exposure to stress further reinforces this superiority, leading to unequal competition for C and N resources between the main stem and tillers, ultimately diminishing seed yield (Tilley et al., 2017). Xie et al. (2019) observed that salt-tolerant wheat under salt stress conditions had higher activities of CAT, POD and APX thus more unfavourable H_2O_2 accumulation and reduced oxidative damage compared to sensitive varieties. Therefore, it is believed that the leaf tiller co-extension law leads to a pre-eminent advantage of temperature and light resource utilisation in the main stem and the first and second tillers, which contributes to the improvement of antioxidant capacity, nutrient accumulation and thus better growth and resistance.

Foliar spraying of Pro-Ca under salt stress effectively mitigated the damage of salt stress on morphogenesis at the tillering stage of rice, as demonstrated in our previous studies. (Zhang et al., 2023; Zhang et al., 2023; Huang et al., 2023), In line with previous experiments, the results of this study indicated that foliar application of Pro-Ca had a more pronounced impact on tiller development compared to the main stem. Pro-Ca effectively alleviated the oxidative damage caused by salt stress on rice leaves. Foliar spraying of Pro-Ca increased the SOD and APX activities of tiller leaves and stems to a stronger extent than that of the main stems, but the effect on CAT activity of main stem leaves was more pronounced; moreover, Pro-Ca showed a better modulation in the alleviation of membrane damage and in the increase of soluble protein content. On the one hand, compared with the main stem, tillers will be subject to stronger oxidative damage, calcium regulator can increase SOD and APX activities as the first line of defence of the plant against oxidative stress (Shafi et al., 2015), on the other hand, it may be related to the spraying site of Pro-Ca, the treatment of the present experiment was in the seedling stage, and the sampling period was in the tillering stage, and the determined tillers were developed from the axils of the leaves that had been sprayed with Pro-Ca in the previous period, therefore, the regulator may have a better regulatory effect on the site to which it was sprayed. In prior trials, it was discovered that despite not being directly treated with ethylene and chlormequat chloride, oat T1 and T2 tillers exhibited delayed stem growth. This could be attributed to the transition from chlormequat chloride-treated to untreated sections of the plant, a phenomenon also observed in wheat. (Kang et al., 2010; Peltonen-Sainio et al., 2003).

Conclusion

This experiment showed that salt stress hindered the growth of leaves and stems of rice tillers during the tillering stage. Additionally, the application of Pro-Ca through spraying effectively reduced the oxidative damage caused by salt stress on the tillers. Interestingly, the impact on the tillers was more pronounced compared to the main stems under similar conditions. This study offers new perspectives on the varying effects of salt stress on rice tillers and the regulatory role

of Pro-Ca.

Acknowledgments

We are very grateful to all authors for their contributions to this article. And we would like to thank the editor and reviewers for their positive comments.

References

- Aebi, H. 1984. [13] Catalase in vitro. In: Methods in enzymology. Vol. 105. *Academic*
- Ahmed, S., Heo, T., Roy Choudhury, A., Walitang, D., Choi, J., Sa, T. 2021. Accumulation of compatible solutes in rice (*Oryza sativa* L.) cultivars by inoculation of endophytic plant growth promoting bacteria to alleviate salt stress. *Applied Biological Chemistry*, 64, 1-14. <https://doi.org/10.1186/s13765-021-00638-x> (2021).
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), 248-254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3) (1976).
- Chang, J., Cheong, B. E., Natera, S., Roessner, U. 2019. Morphological and metabolic responses to salt stress of rice (*Oryza sativa* L.) cultivars which differ in salinity tolerance. *Plant Physiology and Biochemistry*, 144, 427-435. <https://doi.org/10.1016/j.plaphy.2019.10.017> (2019).
- Dai, L., Li, P., Li, Q., Leng, Y., Zeng, D., Qian, Q. 2022. Integrated multi-omics perspective to strengthen the understanding of salt tolerance in rice. *International Journal of Molecular Sciences*, 23(9), 5236. <https://doi.org/10.3390/ijms23095236> (2022).
- Ding, Y., Pan, Q., Gao, W., Pu, Y., Luo, K., He, B. 2022. Reactive oxygen species-upregulating nanomedicines towards enhanced cancer therapy. *Biomater Sci*, 11(4), 1182-1214. <https://doi.org/10.1039/d2bm01833k> (2022).
- dos Santos, T. B., Ribas, A. F., de Souza, S. G. H., Budzinski, I. G. F., Domingues, D. S. 2022. Physiological Responses to Drought, Salinity, and Heat Stress in Plants: A Review. *Stresses*,

2(1):113-135. <https://doi.org/10.3390/stresses2010009> (2022).

Endo, T., Yamamoto, S., Larrinaga, J. A., Fujiyama, H., Honna, T. 2011. Status and causes of soil salinization of irrigated agricultural lands in Southern Baja California, Mexico. *Applied and Environmental Soil Science*, 2011. <https://doi.org/10.1155/2011/873625> (2011).

Feng, C., Gao, H., Zhou, Y., Jing, Y., Li, S., Yan, Z., Xu, K., Zhou, F., Zhang, W., Yang, X., Hussain, M. Z., Li, H. 2023. Unfolding molecular switches for salt stress resilience in soybean: recent advances and prospects for salt-tolerant smart plant production. *Frontiers in Plant Science*, 14, 1162014. <https://doi.org/10.3389/fpls.2023.1162014> (2023).

Giannopolitis, C. N., Ries, S. K. 1977. Superoxide dismutases: I. Occurrence in higher plants. *Plant Physiology*, 59(2), 309-314. <https://doi.org/10.1104/pp.59.2.309> (1977).

Guo, Y. Y., Yu, H. Y., Yang, M. M., Kong, D. S., Zhang, Y. J. 2018. Effect of drought stress on lipid peroxidation, osmotic adjustment and antioxidant enzyme activity of leaves and roots of *Lycium ruthenicum* Murr. seedling. *Russian Journal of Plant Physiology*, 65(2), 244-250. <https://doi.org/10.1134/S1021443718020127> (2018).

Hoang, T. M. L., Tran, T. N., Nguyen, T. K. T., Williams, B., Wurm, P., Bellairs, S., Mundree, S. 2016. Improvement of salinity stress tolerance in rice: challenges and opportunities. *Agronomy*, 6(4), 54. <https://doi.org/10.3390/agronomy6040054> (2016).

Huang, X. X., Zheng, D., Feng, N., Huang, A., Zhang, R., Meng, F., Jie, Y., Mu, B., Mu, D., Zhou, H. 2023. Effects of prohexadione calcium spraying during the booting stage on panicle traits, yield, and related physiological characteristics of rice under salt stress. *PeerJ*, 2023, 11: e14673. <https://doi.org/10.7717/peerj.14673> (2023).

Huihui, Z., Xin, L., Zisong, X., Yue, W., Zhiyuan, T., Meijun, A., Yuehui, Z., Wenxu, Z., Nan, X., Guangyu, S. 2020. Toxic effects of heavy metals Pb and Cd on mulberry (*Morus alba* L.) seedling leaves: Photosynthetic function and reactive oxygen species (ROS) metabolism responses. *Ecotoxicol Environ Saf.* 195, 110469. <https://doi.org/10.1016/j.ecoenv.2020.110469> (2020).

Hussain, S., Cao, X., Zhong, C., Zhu, L., Khaskheli, M. A., Fiaz, S., Zhang, J., Jin, Q. 2018.

701 Sodium chloride stress during early growth stages altered physiological and growth
 702 characteristics of rice. *Chilean journal of agricultural research*, 78(2), 183-197.
 703 <https://doi.org/10.4067/S0718-58392018000200183> (2018).

704 Ilias, I. F., Rajapakse, N. 2005. Prohexadione-calcium affects growth and flowering of petunia
 705 and impatiens grown under photoselective films. *Scientia Horticulturae*, 106(2), 190-202.
 706 <https://doi.org/10.1016/j.scienta.2005.02.023> (2005).

707 Jessup, W., Dean, R. T., Gebicki, J. M. 1994. [29] Iodometric determination of hydroperoxides
 708 in lipids and proteins[M]//Methods in Enzymology. *New York: Academic Press*, 233, 289-303.
 709 [https://doi.org/10.1016/S0076-6879\(94\)33032-8](https://doi.org/10.1016/S0076-6879(94)33032-8) (1994).

710 Jian, G., Mo, Y., Hu, Y., Huang, Y., Ren, L., Zhang, Y., Hu, H., Zhou, S., Liu, G., Guo J, Ling,
 711 Y. 2022. Variety-specific transcriptional and alternative splicing regulations modulate salt
 712 tolerance in rice from early stage of stress. *Rice*, 15(1), 56. [https://doi.org/10.1186/s12284-022-](https://doi.org/10.1186/s12284-022-00599-9)
 713 [00599-9](https://doi.org/10.1186/s12284-022-00599-9) (2022).

714 Joseph, B., Jini, D., Sujatha, S. 2010. Biological and physiological perspectives of specificity in
 715 abiotic salt stress response from various rice plants. *Asian Journal of Agricultural Science*, 2(3),
 716 99-105.

717 Jung, S., Hütsch, B. W., Schubert, S. 2017. Salt stress reduces kernel number of corn by
 718 inhibiting plasma membrane H⁺-ATPase activity. *Plant Physiology and Biochemistry*, 113, 198-
 719 207. <https://doi.org/10.1016/j.plaphy.2017.02.009> (2017).

720 Kang, S. M., Kim, J. T., Hamayun, M., Hwang, I.C., Khan, A. L., Kim, Y. H., Lee, J. H., Lee, I.
 721 J. 2010. Influence of prohexadione-calcium on growth and gibberellins content of Chinese
 722 cabbage grown in alpine region of South Korea. *Scientia Horticulturae*, 125(2), 88-92.
 723 <https://doi.org/10.1016/j.scienta.2010.02.018> (2010).

724 Kaya, C., Ugurlar, F., Ashraf, M., Ahmad, P. 2023. Salicylic acid interacts with other plant
 725 growth regulators and signal molecules in response to stressful environments in plants. *Plant*
 726 *Physiology and Biochemistry*, 196, 431-443. <https://doi.org/10.1016/j.plaphy.2023.02.006> (2023).

727 Kim, H. Y., Lee, I. J., Hamayun, M., Kim, J. T., Won, J. G., Hwang, I. C., Kim, K. U. 2007.

Effect of prohexadione calcium on growth components and endogenous gibberellins contents of rice (*Oryza sativa* L.). *Journal of Agronomy and Crop Science*, 193(6), 445-451. <https://doi.org/10.1111/j.1439-037X.2007.00280.x> (2007).

Kim, Y. H., Khan, A. L., Hamayun, M., Kim, J. T., Lee, J. H., Hwang, I. C., Yoon, C. S., Lee, I. J. 2010. Effects of prohexadione calcium on growth and gibberellins contents of *Chrysanthemum morifolium* R. cv *Monalisa White*. *Scientia Horticulturae*, 123(3), 423-427. <https://doi.org/10.1016/j.scienta.2009.09.022> (2010).

Kousar, T., Sabir, N., Mushtaq, A., Rizwan, S., Jabeen, U., Bashir, F., Ismail, T., Jakhro, M.I., Shahwani, M. N. 2021. Influence of Silica Gel on Ion Homeostasis in Salt Stressed Wheat Varieties of Balochistan. *Silicon*, 13, 4133-4138. <https://doi.org/10.1007/s12633-020-00706-9> (2021).

Lee, I. J., Foster, K. R., Morgan, P. W. 1998. Photoperiod control of gibberellin levels and flowering in sorghum. *Plant Physiology*, 116(3), 1003-1011. <https://doi.org/10.1104/pp.116.3.1003> (1998).

Lee, J., Hanh Nguyen, H., Park, Y., Lin, J., Hwang, I. 2022, Spatial regulation of RBOHD via AtECA4-mediated recycling and clathrin-mediated endocytosis contributes to ROS accumulation during salt stress response but not flg22-induced immune response. *Plant J*, 109(4), 816-830. <https://doi.org/10.1111/tpj.15593> (2022).

Liu, D., Ma, Y., Rui, M., Lv, X., Chen, R., Chen, X., Wang, Y. 2022. Is high pH the key factor of alkali stress on plant growth and physiology? A case study with Wheat (*Triticum aestivum* L.) seedlings. *Agronomy*, 12(8), 1820. <https://doi.org/10.3390/agronomy12081820> (2022).

Munns, R., Tester, M. 2008. Mechanisms of salinity tolerance. *Annual Review of Plant Biology*, 59, 651-681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911> (2008).

Nakano, Y., Asada, K. 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and cell physiology*, 22(5), 867-880. <https://doi.org/10.1093/oxfordjournals.pcp.a076232> (1981).

Peltonen-Sainio, P., Rajala, A., Simmons, S., Caspers, R., Stuthman, D. D. 2003. Plant growth

755 regulator and daylength effects on preanthesis main shoot and tiller growth in conventional and
756 dwarf oat. *Crop science*, 43(1), 227-233. <https://doi.org/10.2135/cropsci2003.2270> (2003).
757 *press*, 121-126. [https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3) (1984).

758 Rasheed, R., Ashraf, M. A., Parveen, S., Iqbal, M., Hussain, I. 2014. Effect of salt stress on
759 different growth and biochemical attributes in two canola (*Brassica napus* L.) cultivars.
760 *Communications in soil science and plant analysis*, 45(5), 669-679.
761 <https://doi.org/10.1080/00103624.2013.867045> (2014).

762 Romero-Puertas, M. C., Rodríguez-Serrano, M., Corpas, F. J., Gomez, M. D., Del Rio, L. A.,
763 Sandalio, L. M. 2004. Cadmium-induced subcellular accumulation of O₂^{•-} and H₂O₂ in pea
764 leaves. *Plant, Cell & Environment*, 27(9), 1122-1134. <https://doi.org/10.1111/j.1365-3040.2004.01217.x> (2004).

766 Ruan, Y., Hu, Y., Schmidhalter, U. 2008. Insights on the role of tillering in salt tolerance of
767 spring wheat from detillering. *Environmental and experimental botany*, 64(1), 33-42.
768 <https://doi.org/10.1016/j.envexpbot.2008.04.004> (2008).

769 Schraudner, M., Moeder, W., Wiese, C., Camp, W.V., Inzé, D., Langebartels, C., Sandermann, Jr.
770 H. 1998. Ozone-induced oxidative burst in the ozone biomonitor plant, tobacco Bel W3. *The*
771 *Plant Journal*, 16(2), 235-245. <https://doi.org/10.1046/j.1365-313x.1998.00294.x> (1998).

772 Shafi, A., Chauhan, R., Gill, T., Swarnkar, M. K., Sreenivasulu, Y., Kumar, S., Kumar, N.,
773 Shankar, R., Ahuja, P. S., Singh, A. K. 2015. Expression of SOD and APX genes positively
774 regulates secondary cell wall biosynthesis and promotes plant growth and yield in *Arabidopsis*
775 under salt stress. *Plant Mol Biol*, 87(6), 615-31. <https://doi.org/10.1007/s11103-015-0301-6>
776 (2015).

777 Taratima, W., Chomarsa, T., Maneerattanarungroj, P. 2022. Salinity Stress Response of Rice
778 (*Oryza sativa* L. cv. Luem Pua) Calli and Seedlings. *Scientifica*, 2022.
779 <https://doi.org/10.1155/2022/5616683> (2022).

780 Tilley, M. S., Heiniger, R.W., Crozier, C.R. 2019. Tiller initiation and its effects on yield and
781 yield components in winter wheat. *Agronomy Journal*, 111(3), 1323-1332.

782 <https://doi.org/10.2134/agronj2018.07.0469> (2019).

783 Xie, X., He, Z., Chen, N., Tang, Z., Wang, Q., Cai, Y. 2019. The Roles of Environmental Factors
784 in Regulation of Oxidative Stress in Plant. *Biomed Res Int*, 8(2019), 9732325.
785 <https://doi.org/10.1155/2019/9732325> (2019).

786 Xu, Q., Xu, X., Zhao, Y., Jiao, K., Herbert, S. J., Hao, L. 2008. Salicylic acid, hydrogen peroxide
787 and calcium-induced saline tolerance associated with endogenous hydrogen peroxide
788 homeostasis in naked oat seedlings. *Plant Growth Regulation*, 54, 249-259.
789 <https://doi.org/10.1007/s10725-007-9247-2> (2008).

790 Yang, H., Xiao, Y., He, P., Ai, D., Zou, Q., Hu, J., Liu, Q., Huang, X., Zheng, T., Fan, G. 2022.
791 Straw mulch-based no-tillage improves tillering capability of dryland wheat by reducing
792 asymmetric competition between main stem and tillers. *The Crop Journal*, 10(3), 864-878.
793 <https://doi.org/10.1016/j.cj.2021.09.011> (2022).

794 Zhang, Q., Xie, J., Zhu, X., Ma, X., Yang, T., Khan, N. U., Zhang, S., Liu, M., Li, L., Liang, Y.,
795 Pan, Y., Li, D., Li, J., Li, Z., Zhang, H., Zhang, Z. 2023. Natural variation in Tiller Number 1
796 affects its interaction with TIF1 to regulate tillering in rice. *Plant Biotechnology Journal*, 21(5),
797 1044. <https://doi.org/10.1111/pbi.14017> (2023).

798 Zhang, R., Zheng, D., Feng, N., Qiu, Q-S., Zhou, H., Liu, M., Li, Y., Meng, F., Huang, X. X.,
799 Huang, A., Li, Y. 2023. Prohexadione calcium enhances rice growth and tillering under NaCl
800 stress. *PeerJ*, 11:e14804. <https://doi.org/10.7717/peerj.14804> (2023).

801 Zhang, R., Zheng, D., Feng, N., Qiu, Q-S., Zhou, H., Meng, F., Huang, X., Huang, A., Li, Y.
802 2023. Prohexadione-calcium alleviates the leaf and root damage caused by salt stress in rice
803 (*Oryza sativa* L.) at the tillering stage. *PLoS ONE*, 18(3): e0279192.
804 <https://doi.org/10.1371/journal.pone.0279192> (2023).

805 Zhang, Z., Liu, Q., Song, H. X., Rong, X., Abdelbagi, M. I. 2012. Responses of different rice
806 (*Oryza sativa* L.) genotypes to salt stress and relation to carbohydrate metabolism and
807 chlorophyll content. *African Journal of Agricultural Research*, 7(1), 19-27.
808 <https://doi.org/10.5897/AJAR11.834> (2012).

809 Zhao, H., Liu, X., Sun, Y., Liu, J., Waigi, M. G. 2023. Effects and mechanisms of plant growth
 810 regulators on horizontal transfer of antibiotic resistance genes through plasmid-mediated
 811 conjugation. *Chemosphere*, 318, 137997. <https://doi.org/10.1016/j.chemosphere.2023.137997>
 812 (2023).

813 Zheng, C., Liu, C., Liu, L., Tan, Y., Sheng, X., Yu, D., Sun, Z., Sun, X., Chen, J., Yuan, D.,
 814 Duan, M. 2023. Effect of salinity stress on rice yield and grain quality: A meta-analysis.
 815 *European Journal of Agronomy*, 144, 126765. <https://doi.org/10.1016/j.eja.2023.126765> (2023).

816 Zulfiqar, F., Ashraf, M. 2021. Nanoparticles potentially mediate salt stress tolerance in plants.
 817 *Plant Physiol Biochem*. 160:257-268. <https://doi.org/10.1016/j.plaphy.2021.01.028> (2021).

818

Figure 1

Effects of Pro-Ca on rice growth under salt stress.

(a-b) Plant growth. Rice morphology after 7, 14, 21, 28, and 35d of salt stress. Figure (a) shows the form of Xiangliangyou900, and Figure (b) shows Huanghuazhan. From left to right, the plants were treated as follows: CK (distilled water), S (0.3% NaCl), Pro-Ca (100 mg•L⁻¹ Pro-Ca), Pro-Ca+S (100 mg•L⁻¹ Pro-Ca + 0.3% NaCl). (c-d) Tiller numbers of Xiangliangyou900 and Huanghuazhan in the main stem between different treatments. (e-f) Leaf numbers in the main stem. Comparison of tiller numbers and leaf numbers in the main stem between different treatments. Values are means ± SD (*n* = 3) and bars indicate SD. Columns with different letters indicate significant difference at *P* < 0.05 (Duncan's test). Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca (100 mg•L⁻¹ Pro-Ca), XPro-Ca+S (100 mg•L⁻¹ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca (100 mg•L⁻¹ Pro-Ca), ZPro-Ca+S (100 mg•L⁻¹ Pro-Ca + 0.3% NaCl).




Figure 2

Effect of Pro-Ca on SOD activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 3

Effect of Pro-Ca on CAT activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 4

Effect of Pro-Ca on POD activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 5

Effect of Pro-Ca on APX activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 6

Effect of Pro-Ca on SOD activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 7

Effect of Pro-Ca on CAT activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 8

Effect of Pro-Ca on POD activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 9

Effect of Pro-Ca on APX activity of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 10

Effect of Pro-Ca on MDA content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 11

Effect of Pro-Ca on H_2O_2 content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 12

Effect of Pro-Ca on O_2^- , H_2O_2 content and cellular activity of rice main stem leaves at each tiller position under salt stress.

From left to right, the three columns represent the tiller stem, the first node position, and the second tiller position. Each image shows the following treatments applied to the plants from left to right: CK (distilled water), S (0.3% NaCl), Pro-Ca ($100\text{ mg}\cdot\text{L}^{-1}$ Pro-Ca), Pro-Ca+S ($100\text{ mg}\cdot\text{L}^{-1}$ Pro-Ca + 0.3% NaCl). The spot area represents the degree of stress, and the larger staining area indicates that the more severe stress of leaves.




Figure 13

Effect of Pro-Ca on MDA content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 14

Effect of Pro-Ca on H_2O_2 content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 15

Effect of Pro-Ca on the soluble protein content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) leaves under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Figure 16

Effect of Pro-Ca on the soluble protein content of rice main stem (a, b), first tiller (c, d), and second tiller (e, f) stems under salt stress.

The different letters are significant differences according to Duncan's new multiple range test ($P < 0.05$) based on one-way ANOVA. Xiangliangyou900: XCK (distilled water), XS (0.3% NaCl), XPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), XPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl), Huanghuazhan: ZCK (distilled water), ZS (0.3% NaCl), ZPro-Ca ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca), ZPro-Ca+S ($100 \text{ mg} \cdot \text{L}^{-1}$ Pro-Ca + 0.3% NaCl).




Table 1(on next page)

Effects of Pro-Ca on plant height and stem base width of rice main stem, first tiller, and second tiller stems at the tillering stage under salt stress.

Values described are the means \pm SE ($n = 3$). Different letters denote significant difference from Duncan's LSD test ($p < 0.05$).

Table 1

Effects of Pro-Ca on plant height and stem base width of rice main stem, first tiller, and second tiller stems at the tillering stage under salt stress.

Time/ d	Treatment	Plant height/ cm			Stem base width/ mm		
		0	I	II	0	I	II
7	XCK	56.4±0.4a	41.3±0.9a	36.5±1.2a	9.4±0.6b	8.0±0.2a	5.2±0.1b
	XS	52.4±0.9b	29.2±2.7b	27.3±1.9c	7.1±0.2c	5.6±0.4b	4.5±0.3b
	XPro-Ca	48.7±0.9c	28.9±0.4b	31.7±0.7b	11.2±0.5a	8.3±0.2a	6.2±0.2a
	XPro-Ca+S	51.2±0.4b	29.9±2.0b	35.0±1.0ab	9.4±0.3b	7.3±0.6a	6.2±0.3a
	ZCK	63.2±2.7a	42.9±2.7a	45.2±5.0a	9.2±0.4a	7.9±0.5b	6.4±0.1a
	ZS	45.5±1.6c	26.4±1.7c	27.4±1.9b	6.3±0.2b	5.2±0.2c	3.6±0.3b
	ZPro-Ca	51.6±0.6b	34.1±2.3b	31.6±2.3b	9.7±1.2a	8.9±0.1a	6.3±0.1a
	ZPro-Ca+S	56.5±1.3b	36.5±1.2ab	35.2±2.0ab	8.5±0.1a	7.2±0.2b	6.0±0.4a
14	XCK	66.5±2.1a	46.9±1.0a	43.6±1.0a	10.8±0.7a	6.9±0.4a	6.7±0.3a
	XS	58.2±0.2b	45.8±1.6a	29.7±1.5c	8.1±0.2b	5.8±0.1b	4.6±0.5b
	XPro-Ca	56.5±0.7b	41.7±1.0b	33.7±3.9bc	12.4±0.5a	7.3±0.4a	6.7±0.1a
	XPro-Ca+S	52.3±0.3c	32.6±0.9c	37.4±0.6ab	11.7±0.9a	7.7±0.1a	6.8±0.1a
	ZCK	67.3±1.9a	56.3±2.0a	51.8±2.6a	9.9±0.4a	8.0±0.6a	6.8±0.2a
	ZS	53.9±2.5b	43.7±1.0b	36.8±1.8bc	6.5±0.1b	4.6±0.4b	4.3±0.9b
	ZPro-Ca	53.5±1.4b	41.6±2.8b	31.6±2.8c	9.2±0.2a	9.6±0.2a	6.9±0.1a
	ZPro-Ca+S	51.8±3.5b	44.2±2.1b	41.7±0.4b	9.6±0.6a	8.2±0.5a	6.8±0.3a
21	XCK	83.3±1.5a	58.1±5.4a	49.0±0.8a	12.6±0.2a	11.4±0.7a	7.8±0.6a
	XS	73.7±1.2bc	48.0±2.6ab	38.0±1.0c	10.5±0.2b	6.4±0.5c	5.0±0.6b
	XPro-Ca	79.2±1.6ab	43.5±2.8b	44.8±0.7b	12.7±0.3a	9.8±0.8ab	7.6±0.2a
	XPro-Ca+S	71.0±2.6c	41.7±4.8b	41.5±2.1bc	12.1±0.7a	9.2±0.5b	7.4±0.2a
	ZCK	76.0±0.6a	64.0±6.1a	56.6±1.0a	10.5±0.3a	10.0±1.0a	7.9±0.2a
	ZS	68.2±2.2b	55.7±0.3a	41.5±1.5b	7.8±0.3b	5.6±0.5b	5.2±0.2c
	ZPro-Ca	73.3±3.2ab	57.7±3.3a	51.1±2.1a	10.9±0.8a	11.1±0.3a	8.3±0.2a
	ZPro-Ca+S	72.3±0.9ab	41.5±4.3b	50.5±2.6a	9.7±0.3a	8.9±0.6a	7.3±0.1b

10

11 **Table 1 continued**

28	XCK	86.8±1.9a	71.6±3.9a	51.6±2.2a	16.0±0.1b	12.1±1.2a	8.5±0.5a
	XS	83.2±1.5ab	58.8±2.8bc	45.0±1.2bc	11.9±0.5c	8.2±0.7a	5.9±0.6b
	XPro-Ca	81.4±1.7b	62.8±1.3b	48.1±1.6ab	17.6±0.5a	13.2±0.6a	7.9±0.2a
	XPro-Ca+S	80.5±0.4b	52.0±1.4c	42.3±0.6c	15.3±0.5b	11.9±2.6a	7.5±0.0a
	ZCK	88.2±1.4a	73.5±2.1a	61.3±0.9a	14.5±0.5a	10.6±0.8a	8.4±0.1b
	ZS	70.5±0.3c	58.4±1.4c	44.9±2.9b	8.9±0.5c	6.1±0.4b	5.5±0.2c
	ZPro-Ca	76.9±3.2b	65.6±2.0b	57.3±3.5a	12.4±0.5b	10.7±0.3a	8.8±0.3b
	ZPro-Ca+S	74.2±0.7bc	70.3±1.4ab	56.0±3.5a	12.1±0.8b	9.6±0.2a	9.7±0.1a
35	XCK	86.2±2.6ab	70.0±3.9a	58.8±1.0a	19.8±1.0a	13.4±0.9a	9.9±0.5a
	XS	80.3±1.6b	64.4±0.8ab	51.3±0.2bc	15.6±0.4b	9.5±0.6b	7.5±0.3b
	XPro-Ca	87.3±1.3a	63.8±2.4ab	52.8±0.4b	20.0±1.1a	15.8±0.7a	10.0±0.9a
	XPro-Ca+S	82.0±2.0ab	55.8±4.9b	49.2±1.4c	18.9±0.3a	13.4±1.1a	8.3±0.0ab
	ZCK	90.0±0.3a	84.0±2.0a	75.8±2.0a	14.9±0.8a	10.3±0.2b	9.3±0.1a
	ZS	80.2±0.6b	70.3±6.6b	61.0±5.1b	12.3±0.5b	9.5±0.3b	6.8±0.2b
	ZPro-Ca	76.3±0.5bc	68.8±1.9b	61.9±2.8b	15.8±0.5a	13.3±0.9a	9.8±0.5a
	ZPro-Ca+S	75.1±2.4c	69.6±2.1b	63.8±0.2b	15.5±0.3a	10.6±0.2b	9.3±0.1a

12 Values described are the means ± SE (*n* = 3). Different letters denote significant difference from

13 Duncan’s LSD test (*p* < 0.05).

14

15

16

Table 2 (on next page)

Effects of Pro-Ca on root length of rice main stem and leaf area per stem of rice main stem, first tiller, and second tiller stems at the tillering stage under salt stress.

Values are the means \pm SE ($n = 3$). Different letters denote significant difference from Duncan's LSD test ($p < 0.05$).

1 **Table 2**
 2 **Effects of Pro-Ca on root length of rice main stem and leaf area per stem of rice main stem,**
 3 **first tiller, and second tiller stems at the tillering stage under salt stress.**

Time/ d	Treatment	Root length/ cm	Leaf area per stem/ cm ²		
			0	I	II
7	XCK	32.2±0.2a	5370.7±726.7ab	1817.1±158.7b	1297.3±249.1b
	XS	16.7±2.1c	4305.7±149.7b	974.4±196.4c	863.1±169.3b
	XPro-Ca	32.3±0.6a	6739.5±461.0a	2828.0±171.5a	995.5±141.1b
	XPro-Ca+S	25.4±0.6b	6116.9±293.6a	2589.2±117.2a	1857.3±57.1a
	ZCK	28.0±0.4a	4316.7±701.1a	2139.5±320.7a	1812.1±29.7a
	ZS	20.2±0.4c	2503.7±181.5b	573.0±47.9b	628.0±114.9b
	ZPro-Ca	28.3±0.7a	4958.6±249.8a	2751.1±330.2a	1735.9±327.2a
	ZPro-Ca+S	24.1±0.6b	4556.0±449.6a	2459.9±198.9a	1909.4±183.4a
14	XCK	34.0±0.7a	8768.8±265.8ab	4157.0±342.1a	2850.5±19.0a
	XS	23.3±1.0d	5765.6±533.7c	2625.8±151.3b	1610.0±138.4b
	XPro-Ca	30.4±0.1b	9411.7±502.4a	3800.7±95.0a	2544.2±169.6a
	XPro-Ca+S	26.5±0.9c	7524.2±54.7b	3416.4±357.4ab	2659.9±94.3a
	ZCK	31.6±0.5a	4542.4±699.9bc	3800.3±49.7b	2476.3±155.0b
	ZS	24.0±0.5c	3191.6±310.9c	1884.3±108.6c	1335.7±146.5c
	ZPro-Ca	30.4±0.2a	7599.9±557.3a	5933.6±537.9a	4297.4±297.2a
	ZPro-Ca+S	29.1±0.2b	5725.1±198.5b	3424.3±163.0b	2854.3±261.0b
21	XCK	36.9±0.9a	13073.6±31.9b	9086.0±1632.7a	3487.7±242.7b
	XS	26.3±0.3c	8518.4±824.1c	4431.7±473.8b	2343.2±156.9c
	XPro-Ca	35.1±0.7a	16377.4±1268.1a	9106.6±655.3a	4243.2±107.5ab
	XPro-Ca+S	31.1±0.4b	13157.8±1026.7b	7390.5±261.9ab	4539.4±405.0a
	ZCK	31.3±0.3a	11510.1±169.6ab	8480.1±214.0b	4263.7±158.3b
	ZS	25.9±0.3c	6699.3±469.0c	3349.3±104.3c	2419.2±134.3c
	ZPro-Ca	31.9±0.4a	12499.1±591.7a	9743.8±430.3a	5790.4±431.2a
	ZPro-Ca+S	28.1±0.5b	10685.9±115.6b	8323.6±267.9b	5174.6±300.1ab

10 **Table 2 continued**

28	XCK	38.9±0.6a	19356.5±214.9a	10053.9±1078.9a	4877.8±232.1a
	XS	28.9±0.5c	11285.3±1893.9b	4555.5±186.5c	3383.9±246.4b
	XPro-Ca	35.5±0.2b	21134.8±1004.4a	11031.7±344.0a	5339.8±348.8a
	XPro-Ca+S	35.7±0.6b	18674.2±442.4a	7893.0±501.8b	4676.2±44.4a
	ZCK	34.3±1.0a	13441.2±333.7a	8997.6±286.3b	6685.3±605.7a
	ZS	27.6±0.6b	7290.5±219.1c	4316.1±337.7c	3482.6±238.6b
	ZPro-Ca	34.9±0.4a	12883.3±750.9a	9499.3±462.4b	7186.9±142.7a
	ZPro-Ca+S	34.8±0.3a	10919.9±694.5b	10803.5±466.6a	6794.2±293.9a
35	XCK	41.9±0.7a	22136.7±1754.2a	15401.6±3314.5ab	7314.4±1013.4a
	XS	30.0±0.4b	14410.6±860.6b	6125.3±374.9c	4154.9±161.7a
	XPro-Ca	39.7±0.7a	22637.8±649.4a	18446.2±1148.1a	7497.9±1618.8a
	XPro-Ca+S	41.0±1.0a	20386.9±589.8a	11051.6±981.7bc	5948.2±318.4a
	ZCK	36.4±0.6a	17411.4±454.4a	11695.5±378.4b	9364.4±773.9a
	ZS	30.7±0.3b	10297.9±287.1d	8110.1±300.1c	4612.9±130.9b
	ZPro-Ca	38.3±1.2a	14462.8±419.4b	12506.7±509.8b	8107.0±497.7a
	ZPro-Ca+S	37.8±0.5a	12335.2±361.4c	14148.4±387.9a	7825.9±929.4a

11 Values are the means ± SE (*n* = 3). Different letters denote significant difference from Duncan's

12 LSD test (*p* < 0.05).

Table 3(on next page)

Effects of Pro-Ca on dry weight per stem of rice main stem, first tiller, and second tiller stems and root dry weight of rice main stem at the tillering stage under salt stress.

Values are the means \pm SE ($n = 3$). Different letters denote significant difference from Duncan's LSD test ($p < 0.05$).

Table 3

Effects of Pro-Ca on dry weight per stem of rice main stem, first tiller, and second tiller stems and root dry weight of rice main stem at the tillering stage under salt stress.

Time/ d	Treatment	Dry weight per stem/ g			Root dry weight/ g
		0	I	II	
7	XCK	0.470±0.017a	0.168±0.004a	0.103±0.005b	0.195±0.007a
	XS	0.340±0.015b	0.062±0.011b	0.065±0.006c	0.118±0.007d
	XPro-Ca	0.478±0.045a	0.178±0.003a	0.133±0.005a	0.170±0.005b
	XPro-Ca+S	0.466±0.011a	0.156±0.015a	0.094±0.012b	0.146±0.004c
	ZCK	0.426±0.053a	0.168±0.023b	0.129±0.004a	0.154±0.005b
	ZS	0.202±0.005b	0.058±0.004c	0.045±0.008b	0.098±0.002c
	ZPro-Ca	0.392±0.011a	0.212±0.009a	0.143±0.012a	0.189±0.007a
	ZPro-Ca+S	0.365±0.006a	0.159±0.008b	0.138±0.008a	0.139±0.013b
14	XCK	0.626±0.020a	0.295±0.019b	0.198±0.002a	0.268±0.011ab
	XS	0.392±0.040b	0.188±0.004b	0.124±0.006c	0.140±0.000c
	XPro-Ca	0.573±0.028a	0.440±0.074a	0.206±0.014a	0.318±0.053a
	XPro-Ca+S	0.532±0.016a	0.239±0.012b	0.168±0.002b	0.191±0.004bc
	ZCK	0.464±0.017a	0.352±0.033a	0.191±0.013b	0.235±0.008b
	ZS	0.305±0.027b	0.148±0.012c	0.101±0.004c	0.112±0.001c
	ZPro-Ca	0.552±0.026a	0.415±0.032a	0.262±0.010a	0.311±0.010a
	ZPro-Ca+S	0.468±0.050a	0.245±0.007b	0.236±0.024ab	0.229±0.009b
21	XCK	1.145±0.066a	0.382±0.003c	0.199±0.003b	0.405±0.020b
	XS	0.638±0.065b	0.345±0.029c	0.182±0.008b	0.266±0.007d
	XPro-Ca	1.194±0.078a	0.579±0.024a	0.250±0.020a	0.266±0.007d
	XPro-Ca+S	1.249±0.030a	0.460±0.018b	0.259±0.016a	0.266±0.007d
	ZCK	0.988±0.049a	0.650±0.019a	0.291±0.012b	0.331±0.021c
	ZS	0.565±0.005c	0.256±0.014b	0.222±0.020c	0.206±0.009d
	ZPro-Ca	1.008±0.062a	0.681±0.047a	0.367±0.012a	0.596±0.013a
	ZPro-Ca+S	0.791±0.037b	0.598±0.031a	0.359±0.018a	0.493±0.035b

10 **Table 3 continued**

28	XCK	2.160±0.134a	0.943±0.041b	0.344±0.009a	1.469±0.073a
	XS	0.896±0.023c	0.617±0.033c	0.224±0.002b	0.428±0.046d
	XPro-Ca	1.828±0.070b	1.373±0.117a	0.326±0.030a	1.278±0.013b
	XPro-Ca+S	1.571±0.045b	0.633±0.029c	0.346±0.022a	0.601±0.048c
	ZCK	1.387±0.121a	0.767±0.051b	0.376±0.013b	0.954±0.012a
	ZS	0.654±0.019b	0.400±0.008c	0.278±0.011b	0.284±0.011d
	ZPro-Ca	1.467±0.114a	0.948±0.047a	0.689±0.030a	0.695±0.019b
	ZPro-Ca+S	1.235±0.126a	0.908±0.076ab	0.824±0.077a	0.631±0.020c
35	XCK	2.817±0.131a	1.765±0.056a	0.446±0.042ab	1.938±0.081ab
	XS	1.053±0.009c	0.725±0.011b	0.350±0.013b	1.014±0.075c
	XPro-Ca	2.802±0.072a	1.547±0.258a	0.603±0.086a	2.153±0.219a
	XPro-Ca+S	2.502±0.099b	1.432±0.039a	0.449±0.036ab	1.639±0.037b
	ZCK	2.243±0.104ab	1.265±0.139b	0.544±0.076b	1.446±0.059a
	ZS	1.575±0.054c	0.881±0.056c	0.429±0.012b	0.551±0.019c
	ZPro-Ca	2.321±0.144a	1.539±0.024a	1.341±0.052a	0.842±0.079b
	ZPro-Ca+S	1.915±0.121bc	1.421±0.018ab	1.212±0.017a	0.767±0.035b

11 Values are the means ± SE (*n* = 3). Different letters denote significant difference from Duncan's

12 LSD test (*p* < 0.05).

13