

Basis of Single-Seed Formation in Chestnut: Cytomorphological Observations Reveal Ovule Developmental Patterns of *Castanea henryi*

Qi Qiu^{Equal first author, 1, 2}, Xiaoming Tian^{Equal first author, 3}, Guolong Wu^{1, 2}, Juntao Wu^{1, 2}, Deyi Yuan^{1, 2}, Xiaoming Fan^{Corresp. 1, 2}

¹ Key Laboratory of Cultivation and Protection for Non-Wood Forest Trees, Ministry of Education, Central South University of Forestry and Technology, Changsha, Hunan province, China

² Key Laboratory of Non-Wood Forest Products of State Forestry Administration, Central South University of Forestry and Technology, Changsha, Hunan, China

³ Hunan Botanical Garden, Changsha, China

Corresponding Author: Xiaoming Fan
Email address: fan_xiaoming001@163.com

Background. Many plants, including those commonly found in the Fagaceae family, produce more flowers and ovules than mature fruits and seeds, which presenting a survival strategy. In *Castanea henryi*, an ovary contains 16–24 ovules, but only one develops into a seed. The other ovules abort or even produce empty buds, but the reason for this is unknown. Such a strict reproductive screening mechanism is rare in other plants.

Methods. In this study, conventional paraffin embedding and semi-thin sectioning techniques, followed by microscopy, were used for cytological studies of ovule development in *C. henryi*.

Results. Pollination affected not only the process of ovule development, but also the proportion of ovules that formed mature embryo sacs. Approximately 54% of the ovules in the pollinated treatment developed normally, while only 17% of the ovules in the unpollinated treatment developed into mature embryo sacs with a seven-cell, eight-nucleated structure. Failure to form mature embryo sacs and the abnormal divisions of the zygote, respectively, were the reasons for the pre- and post-fertilization ovule failures. Using these results, we elucidated the cellular mechanism underlying single-seed formation in *C. henryi*. Our findings offer insight into seed formation mechanisms for further research in plant reproductive biology of *C. henryi* or *Castanea* Miller species.

Basis of Single-Seed Formation in Chestnut: Cytomorphological Observations Reveal Ovule Developmental Patterns of *Castanea henryi*

Qi Qiu ^{1,2,*}, Xiaoming Tian ^{3,*}, Guolong Wu ^{1,2}, Juntao Wu ^{1,2}, Deyi Yuan ^{1,2}, Xiaoming Fan ^{1,2}

¹ Key Laboratory of Cultivation and Protection for Non-Wood Forest Trees, Ministry of Education, Central South University of Forestry and Technology, Changsha, China

² Key Laboratory of Non-Wood Forest Products of State Forestry Administration, Central South University of Forestry and Technology, Changsha, China

³ Hunan Botanical Garden, Changsha, China

* These authors contributed equally to this work.

Corresponding Author:

Xiaoming Fan ^{1,2}

Central South University of Forestry and Technology, Changsha, Hunan, 410004, China

Email address: fan_xiaoming001@163.com

Abstract

Background. Many plants, including those commonly found in the Fagaceae family, produce more flowers and ovules than mature fruits and seeds, which presenting a survival strategy. In *Castanea henryi*, an ovary contains 16–24 ovules, but only one develops into a seed. The other ovules abort or even produce empty buds, but the reason for this is unknown. Such a strict reproductive screening mechanism is rare in other plants.

Methods. In this study, conventional paraffin embedding and semi-thin sectioning techniques, followed by microscopy, were used for cytological studies of ovule development in *C. henryi*.

Results. Pollination affected not only the process of ovule development, but also the proportion of ovules that formed mature embryo sacs. Approximately 54% of the ovules in the pollinated treatment developed normally, while only 17% of the ovules in the unpollinated treatment developed into mature embryo sacs with a seven-cell, eight-nucleated structure. Failure to form mature embryo sacs and the abnormal divisions of the zygote, respectively, were the reasons for the pre- and post-fertilization ovule failures. Using these results, we elucidated the cellular mechanism underlying single-seed formation in *C. henryi*. Our findings offer insight into seed formation mechanisms for further research in plant reproductive biology of *C. henryi* or *Castanea* Miller species.

Introduction

During the development of angiosperms, many flowers, ovules, and seeds fail to develop into mature fruits and seeds ~~and therefore perish~~. Factors such as the inherent characteristics of the species, number of flowers or ovules, processes of pollination and fertilization, and environmental conditions regulating plant growth are closely associated with this failure in development (Zhang et al., 2020; Severino, 2021). The period of ovule development is ~~likely the most~~ crucial phase in the life cycle of plants as it determines reproductive output and offspring viability. In the early stages of the life cycle, viability selection, characterized by high numbers of ovules and a high rate of developmental failure, plays a pivotal role in shaping the genetic composition of plant populations. Importantly, this viability selection can occur at any stage of ovule development (Hufford & Hamrick, 2003). A certain degree of ovule failure is potentially a result of a viability selection process in which rigorous mate selection and selective fertilization are utilized to maximize reproductive success (Susko, 2006; Hasegawa, Suyama & Seiwa, 2009). However, seed failure has become a central problem in food production (Alqudah, Sharma & Börner, 2021). Moreover, ovule failure is a more effective mechanism than seed failure in reducing the costs of fruiting. As ovule abortion occurs earlier in the developmental sequence than seed abortion, the cost of fruiting is substantially reduced. This early-stage abortion allows plants to reallocate and increase the availability of conserved resources (Calviño, 2014).

The phenomenon of multiple ovules but single-seeded fruiting is common in the Fagaceae family (Boavida et al., 1999; Fan et al., 2015). *Castanea mollissima* has 12–18 ovules in one ovary, but only one develops into a seed; it also has three female flower clusters enveloped by an involucre, or, three nuts in an involucre (Du et al., 2021). Similar phenomena also occur in *Castanea crerta*, *Castanea dentate*, *Castanea pumila*, and *Castanea sativa* (Taylor et al., 2012). As only one seed is present in an involucre, there is usually one ovary per involucre in *Castanea henryi*, which has been used as a model to research selective ovule development. This species is vital in targeting poverty alleviation in China's mountainous regions, and its production has become the dominant industry in some areas (Zhang et al., 2016; Li et al., 2019b). The ovary of *C. henryi* contains 16–24 ovules with axile placentation, typically attached at the inner angle of each carpel. However, only one ovule can maintain development and ultimately form a seed, while the rest abort and become brown and withered (Fan et al. 2015; Qiu et al. 2023). The distribution of ovules that mature into viable, fully developed seeds within the fruit is random. These characteristics allow us to study the selection of viability within the fruit in the early stages of life. In a stringent selection process, mate choice and selective fertilization are employed to maximize reproductive success. Previous studies have shown delayed fertilization in *C. henryi*, and it takes six weeks to form a mature seven-cell eight-nuclear embryo sac structure (Fan et al., 2015). During those six weeks, approximately half of the ovules in the ovary are aborted because of abnormal development. Even if a mature embryo sac is formed, only four to five ovules are successfully fertilized. However, the mechanism for single-seed formation in *C. henryi* remains poorly understood.

In this study, we used microscopic sectioning techniques to reveal, at the cytological level, the **causes** of ovule abortion before and after double fertilization and the characteristics during the formation of a single seed. This research serves to provide baseline knowledge for future investigations of *C. henryi* or Fagaceae species reproduction.

Materials & Methods

Plant materials

The experimental site was located on the western campus of the Central South University of Forestry and Technology, Changsha, Hunan Province, China. The Chinese Chinquapin (*Castanea henryi*) Cultivar Huali 4 (Xiangs-SC-SH-010-2015) was **used**, and the pollinated cultivar was Huali 2 (Xiangs-SC-SH-008-2015). The female inflorescences were collected between June and September 2021 for processing and preservation. **Trees in good growth** condition and relatively uniform strength were selected for the experiment. Pollination and non-pollination treatments were applied to the flowers of the same tree. Three trees were treated, and about 120-150 flowers were sampled from each tree.

Materials treatment

In the *C. henryi* female flower, **stigmata are present but do not have the pollination ability of the parent tree for artificial de-male treatment**. In addition to the female flowers, all male inflorescences were removed, and the female flowers were **bagged**, with a **solid bag mouth**, to avoid the entry of exogenous pollen. A total of 400 female flowers were treated to ensure sufficient material for subsequent sampling. On days 5 to 7 of flowering, when the angle between the stigma and the flower mid-axis was about 30° to 45°, **pollination treatment** was carried out on 200 female flowers in sunny and windless weather, and bags were put on immediately after pollination. The other 200 female flowers were not pollinated. In both treatments, the bags were removed two weeks after pollination when the stigmata of the female flowers turned yellow-brown. Samples were taken weekly after treatment. The two main developmental stages we investigated were the mature flower and the young fruit after fertilization.

Paraffin sectioning

The collected plant material was fixed in an ethanol-acetic acid mixture (3:1, v/v), **pumped**, fixed in an acetic acid mixture for 6 h, and stored in 70 % **ethanol**. ~~The specimens were stored at 4 °C until needed. Using the conventional paraffin section method described by Kapil and Sethi (1963), samples were sectioned with a Leica rotary slicer (RM2235, Leica, Heidelberg, Germany); sections had a thickness of 8 µm after ethanol series dehydration and paraffin embedding.~~ The dewaxed sections were stained with safranin and Fast Green, and the slides were sealed with **neutral glue**. Sections were examined, photographed, and imaged using an optical microscope (BX-53, Olympus, Tokyo, Japan).

Semi-thin sectioning

The samples were fixed in 2.5 % glutaraldehyde in 1× phosphate-buffered **saline**. The samples were then dehydrated with an ethanol gradient series and embedded in EMbed 812 resin. The embedded samples were cut to 2 µm with a glass knife in an ultramicrotome (UC7, Leica) and

stained with a mixture of 0.5 % **crocus** and 1 % methyl violet (1:1, v/v) (Tolivia; Navarro & Tolivia, 1994). Stained sections were viewed and photographed using the BX-51 optical microscope.

Results

Ovaries and nuts

Castanea henryi is a monoecious species, ~~consisting of~~ two types of catkins: ~~the~~ staminate catkins and ~~the~~ bisexual catkins with female flowers ~~which~~ usually occur singly (Fig. 1a). In this species, a ring of green acuminate bracts covers the female flowers, and placental hairs or trichomes in the placenta protect the ovary. The involucre progressively hardens into a burr when the female flower differentiates into the fruit (Fig. 1b). The pistils of *C. henryi* comprise the stigma, style, and ovaries (Fig. 1c). When mature, the **thorns** naturally crack, revealing the nut inside. Mature *C. henryi* nuts ~~were~~ conical in shape and their shells ~~were~~ smooth, hard, reddish brown, or bright brown (Fig. 1d).

Development of the ovule

Regardless of the pollination method (~~self, outcrossing, or natural pollination~~), only one seed is produced. In the early stages, there was no difference in ovule size within the same ovary (Fig. 2a). At 5–6 weeks after pollination, ovules continued to develop with enlargement of the ovary (Fig. 2b–c). At week seven after pollination, the volume of one ovule was larger than that of other ovules, and it was full and round; this ovule has already been fertilized. With the continuous expansion of the ovary, the fertilized ovule continued to develop, showing a milky white color, and the volume was much larger than that of the other ovules. Conversely, the other ovules gradually shrunk and became brown (Fig. 2e–f). The ovaries that were ~~artificially~~ bagged and not pollinated did not have fertilized ovules, with obvious volume expansion at week seven after pollination. All ovules in the ovary gradually browned, atrophied, and finally formed empty buds (Fig. 2g–i).

Double fertilization

We observed that the anatropous ovules of *C. henryi* were enveloped by inner and outer integuments to form a micropyle. The development type of embryo sac was typically polygonal with double integuments and thick nuclear tissue. *C. henryi* had delayed fertilization and immature ovules during blooming. Approximately six weeks after pollination, they developed into mature embryo sacs with typical seven-cell and eight-nuclear structures (Fig. 3a). The egg apparatus comprised an egg cell and two synergids near the micropyle, and the nucleus of the egg cell was inclined toward the chalazal end. The **synergism** was pear-shaped, and its nucleus was inclined toward the micropyle end. The central cell was in the middle of the embryonic sac (Fig. 3a). The antipodal cells were close to the chalazal end and degenerated shortly after the embryo sac matured.

~~The~~ **continuous** section ~~showed that~~ the pollen tube ~~entered~~ a synergid cell through the micropyle and ~~released~~ two sperm cells (Fig. 3b). The other synergid ~~remained~~ intact for a period after fertilization and then gradually degenerated. The sperm cells moved to the egg cells ~~with the help of synergistic cells~~ (Fig. 3c). The sperm nucleus ~~gradually approached~~ the egg nucleus

(Fig. 3d–e), and the sperm nuclei attached to the egg nuclear membrane and began to fuse (Fig. 3f). The chromatin of the male nucleus was then diffused, and a small male nucleolus appeared (Fig. 3g). The sperm and egg cells fused to form zygotes (Fig. 3h); the first division of the zygote was performed after a short dormancy period (Fig. 3i). The other sperm nuclei moved to the polar nucleus in the central cell (Fig. 3j–l), and the sperm nucleus and the nuclear membrane of the polar nucleus were close to each other and began to fuse (Fig. 3m). The chromatin in the sperm nucleus was dispersed from one side to the polar nucleus and gradually diffused (Fig. 3n–o). A male nucleolus was also observed (Fig. 3p). The two nuclei fused to form the primary endosperm nucleus, which was located near the middle of the embryo sac (Fig. 3q). The contact between the sperm and secondary nuclei preceded the contact between the sperm and egg nuclei.

Embryonic development

The egg cells formed zygotes after successful double fertilization. After the dormancy period and multiple divisions, the globular embryo gradually formed at week eight after pollination (Fig. 4a–e). The globular embryos were relatively regular (Fig. 4d–f). The cotyledon primordium then developed on both sides of its top, forming a heart-shaped embryo (Fig. 4gg, h), forming a cotyledon embryo. The final mature embryo and integument grew synchronously and filled the entire ovary. The nuts finally matured at week 20 after pollination.

The embryonic development of *C. henryi* did not form an obvious radicle. The development type of the endosperm was nuclear endosperm. The endosperm first formed in the middle of the ovule and approached the end of the micropyle at the globular embryo stage. At the heart-shaped embryo stage, the endosperm cells began to degrade, moved to both sides of the heart-shaped embryo, and became distributed in the embryo sac in a free state (Fig. 4g, h). The endosperm continuously provided nutrients to the embryo during development; therefore, there was no endosperm when the kernel was mature (Fig. 4i). Even if the zygote divided normally, abortion may occur at all stages of embryonic development in fertile ovules, leading to the formation of empty buds.

Ovule abortion before double fertilization

Abortions in plant seed formation are complex and diverse. Abortion may occur at any stage, ranging from the archesporial cell to embryonic development and maturation. Before fertilization, ovule abortion in *C. henryi* mainly showed normal development of the integument and abnormal development of the embryo sac. In a single ovary, approximately 46 % of the ovules exhibited abnormal embryo sac development (Fig. 5), which can be categorized into three situations: the normal embryo sac cavity was not formed during the development process, and the embryo sac cavity was elongated and narrow (Fig. 6a, b); there was a normal embryo sac cavity, but the structure of the egg apparatus structure was missing or incomplete (Fig. 6c); and embryo sac degeneration occurred. Embryo sac degeneration was manifested as follows: the embryo sac developed to the four-nucleate embryo sac stage, and the four-nucleate embryo sac was degraded (Fig. 6d). At the late stage of embryo sac development, the central cell (Fig. 6e), antipodal cell (Fig. 6f), and egg apparatus (Fig. 6g, h) degenerated. Abortive cells with abnormal

embryo sac development showed degeneration of embryo sac, shrinkage, and deformation of nucellus tissue cells, as well as degeneration and disintegration of nucellus tissue (Fig. 6i).

To investigate the impact of pollination on ovule development, we subjected female chestnut flowers to two treatments: pollination and non-pollination. Compared to pollinated ovules, non-pollinated ovules exhibited relatively delayed development, with a reduction in the number of ovules ~~capable of~~ normal development at each stage. Approximately 54 % of ovules in pollinated ovaries developed into mature embryo sacs with a seven-cell eight-nucleus structure. ~~Even without pollination,~~ about 17 % of ovules in a single ovary had a mature embryo sac structure (Table 1). Pollination partially influences the developmental outcome of the embryo sac.

Abortion of ovules after double fertilization

Not all ovules formed seeds after successful fertilization. In the semi-thin sections, approximately 16 % of the ovules ~~experienced~~ post-fertilization abortion (Fig. 5), which can be classified into three types: a normal proembryo structure could not be formed after fertilization, no normal zygote appeared, and only a clump of flocculent tissue was present (Fig. 7a–c); the zygote formed ~~after fertilization,~~ but it did not split (Fig. 7d–f); the zygote was formed ~~after fertilization,~~ and the development of the zygote stagnated after the first normal division to form the second proembryo (Fig. 7g, h). If the ovule formed a zygote ~~after fertilization,~~ then its endosperm developed normally from the primary endosperm nucleus to the free endosperm nucleus (Fig. 7i, j). Moreover, after the formation of the primary endosperm nucleus followed by a very short dormancy period, the dormancy period was very short, and multiple free endosperm nuclei were formed.

Discussion

The extra production of ovules within flowers is commonly observed in various plant species (Sakai & Kojima, 2009). Each ovule has the potential for fertilization and can develop into a viable offspring embryo (Yu, Jiang & Lin, 2022). However, in *C. henryi*, only one ovule ~~can~~ develop into seeds; the reason for this remains unclear. Therefore, we investigated the anatomical characteristics of abortive ovules. ~~Conclusively,~~ *C. henryi* ovule abortion ~~primarily~~ occurs during three stages of embryo sac development: before double fertilization, during double fertilization, and during zygote division. The single-seed formation in chestnuts results from a stepwise screening mechanism.

The embryo sac has a crucial role in the key process of sexual reproduction in plants. Several factors, including resource allocation (Lee & Bazzaz, 1982), positional effects (Silveira & Fuzessy, 2015), and deposition patterns of callose in ovules (Calviño & García, 2009), impede embryo sac development. However, anatomical observations in chestnut ovules suggest that positional effects do not influence embryo sac abortion. In this study, anatomical observations of pollinated and non-pollinated ovules revealed that pollination affected embryo sac development. Similar phenomena have been observed in various plant species, including Orchidaceae (Mayer et al., 2021), in which pollination plays a crucial role in triggering or regulating embryo sac development and ovule maturation, ultimately enabling fertilization (Sakai, 2007). Similarly, in

Ginkgo biloba, pollination triggers the development of the ovule integument (D'Apice et al., 2021).

On the premise that artificial pollination ensures successful fertilization, we found that the main reason for ovule abortion in chestnut after double fertilization was abnormal zygote development. Similar phenomena have also been observed in Hanfu apples and tetraploid black locusts (Yang et al., 2014). In the latter, the abortion rate at the zygotic stage is as high as 50 % (Jiang et al., 2011). Selective abortion in plants is mostly achieved through programmed cell death, which may lead to ovule development deformity, nuclear tissue degeneration, embryo sac abnormality, zygote stagnation, or developmental interruption (Hauser et al., 2006; Wang et al., 2021). The process of nuclear tissue apoptosis in *Ginkgo biloba* has been verified as programmed cell death (Li et al., 2019a). In the following research, programmed cell death may be considered in the study of aborted ovules in *C. henryi*. Exploration of the molecular mechanisms underlying this process is a promising avenue for future study.

In summary, chestnut undergoes layer-by-layer elimination and selects the highest quality ovules to form seeds for successful reproduction. Reducing ovule development may serve as a resource allocation strategy, especially considering the exceptionally high number of ovules produced in chestnut ovaries. This survival strategy is important for resource management within the chestnut ovary.

Conclusions

In this study, we observed the process of double fertilization and embryo abortion before and after fertilization in *C. henryi* to clarify the cytological mechanism of single-seed formation in chestnut. We found that pollination influenced the developmental outcome of embryo sacs. Therefore, chestnut selects the highest quality ovules to form seeds for reproduction. Reducing ovule development acts as a resource allocation and survival strategy. Elucidation of these mechanisms therefore provides insight into future research into plant resource usage and factors affecting growth, survival, and reproduction.

Additional information and declarations

Funding

This research received funding from the National Key R&D Program of China (Grant No. 2022YFD2200400) and the Natural Science Foundation of Hunan Province (grant no. 2022JJ30997). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
The National Key R&D Program of China: 2022YFD2200400.
The Natural Science Foundation of Hunan Province: 2022JJ30997.

References

Alqudah AM, Sharma R, Börner A. 2021. Insight into the genetic contribution of maximum yield potential, spikelet development and abortion in barley. *Plants, People, Planet* 3:721–736. DOI: [10.1002/ppp3.10203](https://doi.org/10.1002/ppp3.10203).

- Boavida LC, Varela MC, Feijó JA. 1999. Sexual reproduction in the cork oak (*Quercus suber* L.). I. The progamic phase. *Sexual Plant Reproduction* 11:347–353. DOI: [10.1007/s004970050162](https://doi.org/10.1007/s004970050162).
- Calviño A. 2014. Effects of ovule and seed abortion on brood size and fruit costs in the leguminous shrub *Caesalpinia gilliesii* (Wall. ex Hook.) D. Dietr. *Acta Botânica Brasilica* 28:59–67. DOI: [10.1590/S0102-33062014000100006](https://doi.org/10.1590/S0102-33062014000100006).
- Calviño A, García CC. 2009. Distinctive patterns of callose deposition provide an early sign of ovule abortion1 in the andromonoecious shrub *Caesalpinia gilliesii*. *Boletín de la Sociedad Argentina de Botánica* 44:49–55.
- D’Apice G, Moschin S, Araniti F, Nigris S, Di Marzo M, Muto A, Banfi C, Bruno L, Colombo L, Baldan B. 2021. The role of pollination in controlling *Ginkgo biloba* ovule development. *New Phytologist* 232:2353–2368. DOI: [10.1111/NPH.17753](https://doi.org/10.1111/NPH.17753).
- Du B, Zhang Q, Cao Q, Xing Y, Q, L, Fang K. 2021. Morphological observation and protein expression of fertile and abortive ovules in *Castanea mollissima*. *PeerJ* 9:e11756. DOI: [10.7717/PEERJ.11756](https://doi.org/10.7717/PEERJ.11756).
- Fan X, Yuan D, Tang J, Tian X, Zhang L, Zou F, Tan X. 2015. Sporogenesis and gametogenesis in Chinese chinquapin (*Castanea henryi* (Skam) Rehder & Wilson) and their systematic implications. *Trees* 29:1713–1723. DOI: [10.1007/s00468-015-1251-y](https://doi.org/10.1007/s00468-015-1251-y).
- Hasegawa Y, Suyama Y, Seiwa K. 2009. Pollen donor composition during the early phases of reproduction revealed by DNA genotyping of pollen grains and seeds of *Castanea crenata*. *New Phytologist* 182:994–1002. DOI: [10.1111/j.1469-8137.2009.02806.x](https://doi.org/10.1111/j.1469-8137.2009.02806.x).
- Hauser BA, Sun K, Oppenheimer DG, Sage TL. 2006. Changes in mitochondrial membrane potential and accumulation of reactive oxygen species precede ultrastructural changes during ovule abortion. *Planta* 223:492–499. DOI: [10.1007/S00425-005-0107-X](https://doi.org/10.1007/S00425-005-0107-X).
- Hufford KM, Hamrick JL. 2003. Viability selection at three early life stages of the tropical tree, *Platypodium elegans* (Fabaceae, Papilionideae). *Evolution; International Journal of Organic Evolution* 57:518–526. DOI: [10.1111/j.0014-3820.2003.tb01543.x](https://doi.org/10.1111/j.0014-3820.2003.tb01543.x).
- Jiang J, Li Y, He J, Hao C, Wang S, Wen F. 2011. Ovule abortion and its mechanism for tetraploid *Robinia pseudoacacia*. *Scientia Silvae Sinicae* 47:40–45. DOI: [10.11707/j.1001-7488.20110506](https://doi.org/10.11707/j.1001-7488.20110506).
- Kapil RN, Sethi SB. 1963. Development of male and female gametophytes in *Camellia sinensis* (L.) O. Kuntze. *Proceedings of the National Institute of Sciences in India* 29:567–574.
- Lee TD, Bazzaz FA. 1982. Regulation of fruit and seed production in an annual legume, *Cassia fasciculata*. *Ecology* 63:1363–1373. DOI: [10.2307/1938864](https://doi.org/10.2307/1938864).
- Li D, Wu D, Li S, Guo N, Gao J, Sun X, Cai Y. 2019a. Transcriptomic profiling identifies differentially expressed genes associated with programmed cell death of nucellar cells in *Ginkgo biloba* L. *BMC Plant Biology* 19:91. DOI: [10.1186/s12870-019-1671-8](https://doi.org/10.1186/s12870-019-1671-8).
- Li Y, Xiong H, Peng X, Zou F, Yuan D. 2019b. Comparison on leaf anatomical structure and photosynthetic characteristics of grafted seedlings and seedlings in *Castanea henryi*. *Southwest China Journal of Agricultural Sciences* 32:792–797. DOI: [10.16213/j.cnki.scjas.2019.4.014](https://doi.org/10.16213/j.cnki.scjas.2019.4.014).
- Mayer JLS, Scopece G, Barone Lumaga MR, Coiro M, Pinheiro F, Cozzolino S. 2021. Ecological and phylogenetic constraints determine the stage of anthetic ovule development in orchids. *American Journal of Botany* 108:2405–2415. DOI: [10.1002/AJB2.1770](https://doi.org/10.1002/AJB2.1770).
- Qiu Q, Tian X, Wu G, Wu J, Fan X, Yuan D. 2023. Comparative analysis of the transcriptome during single-seed formation of *Castanea henryi*: regulation of starch metabolism and endogenous hormones. *BMC Plant Biology* 23:90. DOI: [10.1186/S12870-023-04102-4](https://doi.org/10.1186/S12870-023-04102-4).
- Sakai S. 2007. A new hypothesis for the evolution of overproduction of ovules: an advantage of selective abortion for females not associated with variation in genetic quality of the resulting seeds. *Evolution; International Journal of Organic Evolution* 61:984–993. DOI: [10.1111/j.1558-5646.2007.00083.x](https://doi.org/10.1111/j.1558-5646.2007.00083.x).
- Sakai S, Kojima T. 2009. Overproduction and selective abortion of ovules based on the order of fertilization revisited. *Journal of Theoretical Biology* 260:430–437. DOI: [10.1016/j.jtbi.2009.06.014](https://doi.org/10.1016/j.jtbi.2009.06.014).
- Severino LS. 2021. Plants make smart decisions in complex environments. *Plant Signaling and Behavior* 16:1970448. DOI: [10.1080/15592324.2021.1970448](https://doi.org/10.1080/15592324.2021.1970448).

- Silveira FAO, Fuzessy LF. 2015. Does successful ovule development depend on its position within the pod? Examples from Neotropical Fabaceae. *Plant Species Biology* 30:285–290. DOI: [10.1111/1442-1984.12057](https://doi.org/10.1111/1442-1984.12057).
- Susko DJ. 2006. Effect of ovule position on patterns of seed maturation and abortion in *Robinia pseudoacacia* (Fabaceae). *Canadian Journal of Botany* 84:1259–1265. DOI: [10.1139/b06-094](https://doi.org/10.1139/b06-094).
- Tolivia J, Navarro A, Tolivia D. 1994. Polychromatic staining of epoxy semithin sections: a new and simple method. *Histochemistry* 101:51–55. DOI: [10.1007/BF00315831](https://doi.org/10.1007/BF00315831).
- Wang H, Zhang S, Qu Y, Gao R, Xiao Y, Wang Z, Zhai R, Yang C, Xu L. 2021. Jasmonic acid and ethylene participate in the gibberellin-induced ovule programmed cell death process in seedless pear ‘1913’ (*Pyrus* hybrid). *International Journal of Molecular Sciences* 22:9844. DOI: [10.3390/IJMS22189844](https://doi.org/10.3390/IJMS22189844).
- Yang X, Ma H, Liu G, Lü D, Qin S, Du G. 2014. Anatomical study on the multi-ovule development and abortion of Hanfu apple. *Journal of Integrative Agriculture* 13:770–777. DOI: [10.1016/S2095-3119\(13\)60409-8](https://doi.org/10.1016/S2095-3119(13)60409-8).
- Yu SX, Jiang YT, Lin WH. 2022. Ovule initiation: the essential step controlling offspring number in *Arabidopsis*. *Journal of Integrative Plant Biology* 64:1469–1486. DOI: [10.1111/JIPB.13314](https://doi.org/10.1111/JIPB.13314).
- Zhang S, Li Y, Li Y, Zhao F, Zeng X. 2020. Insights on seed abortion (endosperm and embryo development failure) from the transcriptome analysis of the wild type plant species *Paeonia lutea*. *Bioinformation* 16:638–651. DOI: [10.6026/97320630016638](https://doi.org/10.6026/97320630016638).
- Zhang X, Yuan D, Zou F, Fan X, Tang J, Zhu Z. 2016. A study on the xenia effect in *Castanea henryi*. *Horticultural Plant Journal* 2:301–308. DOI: [10.1016/j.hpj.2017.02.003](https://doi.org/10.1016/j.hpj.2017.02.003).

Table 1(on next page)

Proportion of normal ovule development in single ovary in two treatments.

Notes: Data are means ($\pm$ S.E) for 20 replicates (N = 20). Statistical comparisons were performed using the *t* test.

1 **Table. 1. Proportion of normal ovule development in single ovary in two treatments.**

Weeks after pollination	Developmental stage	Proportion of normal development ovule in pollinated <i>C. henryi</i>	Proportion of normal development ovule in unpollinated <i>C. henryi</i>	p
1	Archegonium, megasporocyte	100 %	100 %	
2–3	Megasporocyte, meiosis, functional megaspore	100 %	100 %	
4–5	Uninucleate embryos sac, two- to eight-nucleate ES	68 ± 1 %	42 ± 2 %	<0.001***
6	Mature ES	54 ± 2 %	17 ± 1 %	<0.001***
7	Fertilization	20 ± 1 %	0 %	<0.001***

2 **Notes:** Data are means (±S.E) for 20 replicates (N = 20). Statistical comparisons were performed using the *t* test.

3

Figure 1

Inflorescence, pistil and realistic pictures of single seed fruiting of *C. henryi*.

a Bisexual inflorescence. **b** Longitudinal section of female flower. **c** Pistil. **d** Fruit dehiscence state. inv = involucre; sty = style; sti = stigma; tr = trichomes; ova = ovary.




Figure 2

Developmental process of *C. henryi* ovule.

a-c represent the ovary developed at the 4th, 5th and 6th week after pollination, respectively. **d-f** represent the ovary normally developed at the 7th, 8th and 9th week after pollination, respectively. **g-i** represent the ovary (empty bud ovary) that abnormally developed at the 7th, 8th and 9th week after pollination, respectively. The red arrow indicates the fertile ovule. All of the above images are of *C. henryi* viewed under a stereomicroscope after stripping the ovary wall.




Figure 3

Egg apparatus of mature embryo sac and process of double fertilization of *C. henryi*.

a Eight-nucleate embryo sac. **b** Pollen tube entering embryo sac and releasing two sperm cells. **c** Sperm cell approaching egg with help of antipodal cell. **d, e** Sperm nucleus gradually moving closer to egg nucleus. **f, g** Sperm cells and egg cells gradually fusing and male chromatin dispersing into egg nucleus and male nucleoli appeared. **h, i** Zygote. **j-l** Sperm nucleus moving toward polar nucleus. **m** Sperm nucleus attached to polar nucleus and gradually fusing. **n, o** Male chromatin dispersing into polar nucleus. **p** Male nucleus appearing in polar nucleus. **q** Primary endosperm nucleus forming. OI = outer integument; II = inner integument; NU = nucellus tissue; EN = egg nucleus; SC = synergid cell; PN = polar nucleus; SN = sperm nucleus; EC = egg cell; MI = micropylar side; MN = male nucleus; Zy = zygote; PEN = primary endosperm nucleus. **a-q** are paraffin sections double-stained with fuchsin-solid green.




Figure 4

Embryonic development of *C. henryi*.

a-b Zygotic first division. **c** Zygotic second division. **d-f** Globular embryo. **g-h** Gradual formation of heart-shaped embryo. **i** Mature seed kernel. EC = endosperm cell; Ke = kernel. Scale bars = 10µm. **a-h** are paraffin sections double-stained with fuchsin-solid green.




Figure 5

Various factors leading to ovule abortion at different developmental stages.




Figure 6

Ovule abortion before fertilization.

a, b Failure to form embryo sac cavity leads to ovule abortion. **c** Abnormal egg apparatus development. **d** Degenerated four-nucleate embryo sac (arrow). **e** Degenerated central cells in aborted ovules. **f** Degenerated antipodal cells within the abortive ovule. **g, h** Egg apparatus degeneration. **i** Germ cell degeneration and nucellus tissue atrophy. AEA = abnormal egg apparatus; DAC = degenerate antipodal cells; DEA = degenerate egg apparatus. Scale bars: 100 μm (**a**), 50 μm (**b, c**), 10 μm (**d-i**). **a-c** are paraffin sections double-stained with saffron-solid green; **d-i** are semi-thin sections stained with saffron-methyl violet mix.




Figure 7

Ovule abortion after fertilization.

a-c Fertilized ovule does not form a normal proembryo structure. **d-f** Zygote formed after fertilization but did not divide. **g, h** Zygote divided to form a diploid embryo but did not continue to divide. **i, j** Normal division of primary endosperm nucleus to form free endosperm nucleus. PEN = primary endosperm nucleus; FEN = free endosperm nucleus; DE = diploid embryo; Zy = zygote. **a-j** are semi-thin sections stained with a mixture of senna-methyl violet.

