

Clusters explaining the relation between menopause and self-reported periodontal disease: A cross-sectional study (#107178)

1

First submission

Guidance from your Editor

Please submit by **7 Nov 2024** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

3 Figure file(s)

2 Table file(s)

2 Other file(s)



Custom checks

Human participant/human tissue checks



Have you checked the authors [ethical approval statement](#)?



Does the study meet our [article requirements](#)?



Has identifiable info been removed from all files?



Were the experiments necessary and ethical?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  **Impact and novelty is not assessed.** Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Clusters explaining the relation between menopause and self-reported periodontal disease: A cross-sectional study

Hani T. Fadel^{Corresp., 1}, Lujain A. Qarah², Manal O. Alharbi³, Alla Al-Sharif¹, Doaa S. Al-Harkan⁴, Saba Kassim¹, Osama Abu-Hammad⁵, Najla Dar-Odeh⁵

¹ Department of Preventive Dental Sciences, College of Dentistry, Taibah University, AlMadinah AlMunawwarah, Saudi Arabia

² Ministry of Health, AlMadinah AlMunawwarah, Saudi Arabia

³ Islamic University in Madinah, AlMadinah AlMunawwarah, Saudi Arabia

⁴ Department of Oral & Maxillofacial Diagnostic Sciences, College of Dentistry, Taibah University, AlMadinah AlMunawwarah, Saudi Arabia

⁵ School of Dentistry, University of Jordan, Amman, Jordan

Corresponding Author: Hani T. Fadel

Email address: hani.fadel@yahoo.com

Background. Menopause is an important milestone in the women's life continuum and is associated with potentially adverse effects, including those related to oral health. This study assessed self-reported periodontal disease in relation to menopausal status.

Methods. A cross-sectional study involving a convenience sample of female university dental hospital attendees was conducted using a validated, self-administered, self-reported periodontal disease questionnaire. A two-step cluster analysis was used to categorize the participants based on menstrual period (MP) continuity, systemic diseases and age. Differences between clusters were analyzed using chi-square test. **Results.** From 112 included participants, 3 clusters resulted from the analysis: Cluster #1 (37 ± 8 years, no systemic diseases and continued MP), Cluster #2 (40 ± 10 , with systemic diseases and continued MP) and Cluster #3 (54 ± 9 , with systemic diseases and discontinued MP). Cluster #3 had less optimal oral hygiene habits and more missing teeth ($p > 0.05$). Clusters #1 and #2 reported more gingival bleeding, tooth sensitivity and calculus ($p > 0.05$). Cluster #3, on the other hand, presented with more self-reported oral dryness ($p \leq 0.05$). **Conclusions.** Within study limits, clusters of menopausal women with systemic diseases reported high symptoms of periodontal disease that were not significantly different from younger individuals, with the exception of oral dryness.

Clusters explaining the relation between menopause and self-reported periodontal disease: A cross-sectional study

Hani T. Fadel,¹ Lujain A. Qarah,² Manal O. Alharbi,³ Alla Al-Sharif,¹ Doaa S. Al-Harkan,⁴ Saba Kassim,¹ Osama Abu-Hammad⁵ and Najla Dar-Odeh⁵

¹ Department of Preventive Dental Sciences, College of Dentistry, Taibah University, AlMadinah AlMunawwarah, Saudi Arabia

² Ministry of Health, AlMadinah AlMunawwarah, Saudi Arabia

³ Islamic University, AlMadinah AlMunawwarah, Saudi Arabia

⁴ Department of Oral & Maxillofacial Diagnostic Sciences, College of Dentistry, Taibah University, AlMadinah AlMunawwarah, Saudi Arabia

⁵ School of Dentistry, University of Jordan, Amman, Jordan

Corresponding Author:

Hani T. Fadel ¹

Department of Preventive Dental Sciences, College of Dentistry, Taibah University, PO Box 40515, 41511 Madinah, Saudi Arabia

Email address: hani.fadel@yahoo.com

Abstract

Background. Menopause is an important milestone in the women's life continuum and is associated with potentially adverse effects, including those related to oral health. This study assessed self-reported periodontal disease in relation to menopausal status.

Methods. A cross-sectional study involving a convenience sample of female university dental hospital attendees was conducted using a validated, self-administered, self-reported periodontal disease questionnaire. A two-step cluster analysis was used to categorize the participants based on menstrual period (MP) continuity, systemic diseases and age. Differences between clusters were analyzed using chi-square test.

Results. From 112 included participants, 3 clusters resulted from the analysis: Cluster #1 (37±8 years, no systemic diseases and continued MP), Cluster #2 (40±10, with systemic diseases and continued MP) and Cluster #3 (54±9, with systemic diseases and discontinued MP). Cluster #3 had less optimal oral hygiene habits and more missing teeth ($p>0.05$). Clusters #1 and #2 reported more gingival bleeding, tooth sensitivity and calculus ($p>0.05$). Cluster #3, on the other hand, presented with more self-reported oral dryness ($p\leq0.05$).

Conclusions. Within study limits, clusters of menopausal women with systemic diseases reported high symptoms of periodontal disease that were not significantly different from younger individuals, with the exception of oral dryness.

Introduction

Menopause is defined by the World Health Organization (WHO) as “the permanent cessation of menstruation due to loss of ovarian follicular activity” (World Health Organization, 1996). It is diagnosed retrospectively following 12 months of amenorrhea that is not associated with pathology. The term “Menopause” itself means “without estrogen”, and refers to the time at which cyclic ovarian function - known as menstruation – ceases (World Health Organization, 1996). As levels of circulating sex hormones change, several clinical outcomes ensue with a resulting potential impact on emotional state and quality of life.

Within the context of oral health, it has been shown that hormonal fluctuations during the different stages of menopause may lead to certain inflammatory responses in the body (Mariotti, 2005). Notably, these fluctuations and changes increase the sensitivity of the gingiva to microbial dental plaque and calculus, consequently increasing the susceptibility to adverse oral health outcomes such as gingivitis, periodontitis, oral dysesthesia and xerostomia or dry mouth (Mosley, Smith & Dezan, 2015; “Guideline on Oral Health Care for the Pregnant Adolescent,” 2016). Further, the gingival epithelium becomes thinner, atrophic and more prone to inflammatory changes during menopause (Alves et al., 2015). Hormones have been shown to directly and indirectly exert effects on the periodontium. Estrogen and progesterone affect cellular proliferation, differentiation and growth in target tissues, including keratinocytes and fibroblasts in the gingiva (Mariotti & Mawhinney, 2013; Jafri et al., 2015). Therefore, the sudden decrease in estrogen production may be associated with primary osteoporosis, which has an effect on jaw bones (Alves et al., 2015). The reduction in bone mineral density resulting from osteoporosis contributes to periodontal disease progression in menopausal women (Alves et al., 2015). Moreover, Estrogens may interfere with other periodontal tissues such as the gingiva and periodontal ligament, and influence host immune-inflammatory responses (Alves et al., 2015). Some studies have reported a link between osteoporosis of the mandible and the peripheral skeleton with alveolar bone loss of the mandible and tooth loss in menopausal women (Jonasson & Rythén, 2016).

Despite the available evidence, there are no data that have focused on the personal input of women in Northwest Saudia Arabia with regards to their oral health, namely periodontal disease and the possible relation with their menopausal condition. This can help oral health promotion, early treatment and prevention of further complications associated with different stages of menopause. Specifically, this will aid in the establishment of a referral system between gynecologists, general practitioners and dentists that involves regular dental checkups for early detection and intervention, and based on the fact that oral health is an integral component of the overall health.

The objective of this study was thus to assess the self-reported periodontal disease among a selected group of women attending a university outpatient dental clinic in relation to their menopausal status.

Materials & Methods

Study Design

This was an observational cross-sectional analytical study, conducted at the Taibah University College of Dentistry outpatient clinics in AlMadinah AlMunawwarah, Saudi Arabia. Approval from the Taibah University College of Dentistry Research Ethics Committee was obtained prior to commencement of the study, as all set procedural institutional guidelines were followed (Approval no. TUCDREC/20180107/Fadl).

Ethical Considerations

The study was conducted in accordance with the ethical guidelines set by the Declaration of Helsinki (2013). Before recruitment, the purpose of the study was explained to each of the participants and informed consent was obtained. They were also informed that participation was voluntary, with no negative repercussions in terms of quality of offered healthcare due to declining to participate. Participants were also assured that all information will be kept confidential and will only be used for research and educational purposes. In addition, participants were informed whenever necessary of their treatment needs based on the findings of this study.

Study Population and Sample Size Calculation

The study involved a convenience / consecutive sample from all female individuals aged between 22 and 75 years, who were attending the college clinics between November 2018 and July 2019. Children and individuals below 22 years were excluded alongside those with dementia or mental health or radiotherapy. Approximately 13,000 patients (women and men) visit the clinics each year.

Based on the reported prevalence of the outcome “gingivitis” in the general population of 75%, with a desired level of confidence of 95%, and a margin of error of 10, and an estimated population of women attending the clinics annually of 6,500, a sample size of 97 participants was required for the study (Chisquares Inc. Sample Size Calculation for Cross-sectional Studies with Percentage as Outcome.). Incomplete responses were to be excluded from the analysis.

Interview and Questionnaire

Data were collected by means of an interview and a self-administered questionnaire. The interview was conducted in the waiting area and involved explanation of the study, giving consent to participate and filling the questionnaire. The questionnaire took 3-5 minutes to complete and included providing information about age, education level, marital stats, siblings, current health status and social and oral hygiene habits. Participants were also asked about

continuation of their menstrual period (MP). Moreover, a valid and reliable Arabic version of the self-reported periodontal health questionnaire was included (Khader, Alhabashneh & Alhersh, 2014). This section consisted of seventeen close ended dichotomous questions, representing the outcome variables.

Data Analysis

Means and standard deviations were used to describe continuous variables, while frequencies and percentages described categorical variables. For simplicity during data presentation, the different chronic illnesses that the participants were suffering from were grouped under one category, i.e. systemic conditions, since most were considered of inflammatory nature. Similarly, different long-term medications were grouped together, acknowledging their negative effect on salivary secretion. Initial analyses did not reveal any clear patterns when comparing participants with or without systemic conditions or menstrual periods in terms of oral health (data not shown). Accordingly, a 2-step cluster analysis was performed in order to identify hidden patterns and relationships by categorizing the participants based on the available background, demographic and health data. Following a number of attempts, a clustering model of “Good” quality resulted via Silhouette’s measure. The finally selected clustering predictors, in order of importance; were menstrual period continuation, presence of systemic diseases and age. The differences between groups/clusters in terms of quantitative demographics, oral hygiene habits, oral findings and the self-reported periodontal disease questions were analyzed using chi-square test. A p-Value ≤ 0.05 was considered statistically significant. The Statistical Package for Social Sciences (SPSS) software, version 20 (SPSS Inc., Chicago, IL, USA) was used for the analyses.

Results

Of a total of 114 participating female patients, two were excluded due to incomplete data. The remaining 112 returned their completed questionnaires and agreed to participate in the study while awaiting their treatment appointment at the dental clinics. The mean age of the total sample was 40 (± 10) years. Fifty-two percent held a university degree and 66% were employed.

Thirty-six percent of the participants suffered from different systemic conditions such as hypertension and diabetes mellitus. Similarly, 59% were on chronic medication, five of which specifically reporting the use of hormone replacement therapy (HRT). Fourteen percent of the total participants were current smokers, 3 of them smoked cigarettes and the remaining used waterpipes. Twenty percent had already discontinued their menstrual period.

The two-step cluster analysis resulted in 3 clusters, with cluster # 1 (n=65) including women with no systemic conditions and cluster # 3 (n=22) involving those with a completely discontinued menstrual period (Figure 1). Compared to the other two clusters, women in cluster # 3 mostly completed up to high school education ($p < 0.001$), 55% were unemployed ($p < 0.05$) and had the highest number of children ($p < 0.01$) (Table 1). With regards to participants from

cluster # 2, on the other hand, 84% were married ($p > 0.05$) and 80% were on medication ($p < 0.01$) (Table 1).

Cluster # 3 apparently had the least percentage of participants who regularly brushed their teeth (91%), used other tooth cleaning aids (41%) or used mouthwash (41%) compared to the other two clusters, although the differences were not statistically significant ($p > 0.05$) (Figure 2). The same cluster also tended to have the highest prevalence of tooth loss (96%) and replacement of missing teeth (50%) compared to the other clusters (Figure 3). However, the differences were insignificant ($p > 0.05$). With regards to the self-reported periodontal disease (SRPD) questions, the observations of swollen and painful gums, pus, movable teeth, teeth that have changed in position or longer teeth were more reported among individuals in cluster # 3, albeit the lack of statistical significance ($p > 0.05$) (Table 2). The feeling of dry mouth was also significantly higher in cluster # 3 compared to the other 2 clusters ($p < 0.05$) (Table 2). On the other hand, self-reported bleeding gums, tooth sensitivity and calculus deposits were relatively, yet insignificantly; higher in clusters # 1 and # 2 ($p > 0.05$) (Table 2).

Discussion

This study aimed at evaluating self-reported oral health in women in relation to their menopausal status. Women who have not reached menopausal age insignificantly reported more bleeding gums, tooth sensitivity and calculus deposits than their menopausal counterparts. This is in line with findings from larger registries, reporting more oral health problems by younger women (Azofeifa et al., 2014). An explanation could be that younger women are exposed to continuous hormonal changes, with the potential of exaggerated responses of the dental and periodontal tissues to local stimuli (Dar-Odeh et al., 2017). Furthermore, the fact that women in the reproductive age may go through several pregnancies, are minimally aware at the time with regards to their oral health and often avoid dental visits could all contribute to the characteristic adverse oral health findings (Dar-Odeh et al., 2018).

There was an insignificant tendency for older menopausal women with different systemic comorbidities, i.e. Cluster # 3, to observe more painful gums, pus, longer teeth, teeth that are movable, or teeth that have changed their position, all of which known to be related to advanced periodontitis. Recent cohort studies showed that postmenopausal women were more likely to have periodontitis (Park et al., 2023a), and that women with periodontitis are more likely to develop osteoporosis (Choi et al., 2017). Furthermore, it was concluded from previous studies that diabetes mellitus and hypertension are particularly associated with missing teeth among women (Dar-Odeh et al., 2019).

Additional findings from this study confirmed what was mentioned earlier, which showed a tendency for menopausal women to have higher prevalence of missing teeth and tooth replacement. Interestingly, the current study identified relatively poor oral hygiene practices

among menopausal women, which aggravates the risk for dental caries and periodontal disease. Moreover, menopausal women were of low educational backgrounds, mostly unemployed and with more children. These sociodemographic variables could be a factor in predisposing to low awareness to oral hygiene practices and favorable oral health outcomes. Park and co-workers concluded in their nationwide cohort study that good oral hygiene practices are adversely correlated with osteoporotic fractures (Park et al., 2023b).

Menopausal women with systemic comorbidities reported a significantly higher prevalence of dry mouth. Krupa et al. have recently concluded that there was a significant association between menopausal duration and salivary flow rates (Krupa et al., 2023). Older age is associated with polypharmacy which could also predispose to dry mouth (Cannon et al., 2023). On the other hand, hormone replacement therapy (HRT) during menopause is thought to relieve dry mouth symptoms (Wang et al., 2021). However, as low as only five participants reported the use of HRT in the current study.

The relatively small sample size is looked upon as a study limitations, since it may have impacted the extrapolation of the observed findings alongside the convenience sampling that precluded generalizability of the finding. This may dictate the consideration of the current investigation as a pilot study at best. However, the self-reporting of oral health symptoms in menopausal women in this study covers an important aspect, providing unique information that can be utilized for hypothesis generation in future studies. Moreover, the cross-sectional design limits the identification of temporal relationships between menopause and oral health parameters, necessitating the interpretation of findings with caution.

Conclusions

Within limitations of this study, it can be concluded that clusters of menopausal women with systemic diseases reported high symptoms of oral and periodontal disease that were not significantly different from younger individuals, with the exception of oral dryness. Further studies with larger study samples are required to expand on such findings.

References

- Alves RC, Félix SA, Rodriguez-Archilla A, Oliveira P, Brito J, Dos Santos JM. 2015. Relationship between menopause and periodontal disease: a cross-sectional study in a Portuguese population. *International journal of clinical and experimental medicine* 8:11412–11419.
- Azofeifa A, Yeung LF, Alverson CJ, Beltrán-Aguilar E. 2014. Oral health conditions and dental visits among pregnant and nonpregnant women of childbearing age in the United States, National Health and Nutrition Examination Survey, 1999–2004. *Preventing chronic disease* 11:E163. DOI: 10.5888/pcd11.140212.
- Cannon I, Robinson-Barella A, McLellan G, Ramsay SE. 2023. From Drugs to Dry Mouth: A Systematic Review Exploring Oral and Psychological Health Conditions Associated with Dry Mouth in Older Adults with Polypharmacy. *Drugs & aging* 40:307–316. DOI: 10.1007/s40266-023-01017-5.
- Choi J-K, Kim Y-T, Kweon H-I, Park E-C, Choi S-H, Lee J-H. 2017. Effect of periodontitis on the development of osteoporosis: results from a nationwide population-based cohort study (2003–2013). *BMC women’s health* 17:77. DOI: 10.1186/s12905-017-0440-9.
- Dar-Odeh NS, Aleithan FA, Alnazzawi AA, Al-Shayyab MH, Abu-Hammad SO, Abu-Hammad OA. 2017. Factors affecting oral health determinants in female university students: a cross-sectional survey in Saudi Arabia. *International journal of adolescent medicine and health* 32:/j/ijamh.2020.32.issue-1/ijamh-2017-0084/ijamh-2017-0084.xml. DOI: 10.1515/ijamh-2017-0084.
- Dar-Odeh NS, Binsaad S, Gasim R, Haidary R, Afandi A, Hammad S, Hammad O. 2018. Why do Women avoid Dental Visits During Pregnancy? A CrossSectional Survey in Al Madinah, Western Saudi Arabia. *Pesquisa Brasileira em Odontopediatria e Clínica Integrada* 18:1–8. DOI: 10.4034/pboci.2018.181.31.
- Dar-Odeh NS, Borzangy S, Babkair H, Farghal L, Shahin G, Fadhlalmawla S, Alhazmi W, Taher S, Abu-Hammad O. 2019. Association of Dental Caries, Retained Roots, and Missing Teeth with Physical Status, Diabetes Mellitus and Hypertension in Women of the Reproductive Age. *International journal of environmental research and public health* 16. DOI: 10.3390/ijerph16142565.
- Guideline on Oral Health Care for the Pregnant Adolescent. 2016. *Pediatric dentistry* 38:59–66.
- Jafri Z, Bhardwaj A, Sawai M, Sultan N. 2015. Influence of female sex hormones on periodontium: A case series. *Journal of natural science, biology, and medicine* 6:S146-149. DOI: 10.4103/0976-9668.166124.
- Jonasson G, Rythén M. 2016. Alveolar bone loss in osteoporosis: a loaded and cellular affair? *Clinical, cosmetic and investigational dentistry* 8:95–103. DOI: 10.2147/CCIDE.S92774.
- Khader Y, Alhabashneh R, Alhersh F. 2014. Validity of a Self-Reported Periodontal Disease Questionnaire among Jordanians. 4:2–7. DOI: 10.4172/2161-1122.100020.

Krupa NC, Thippeswamy HM, Chandrashekar BR, Thetakala RK. 2023. Impact of Menopausal Duration on Salivary Flow Rate, Tooth Loss, and Oral Health-related Quality of Life in Indian Communities. *Journal of mid-life health* 14:101–106. DOI: 10.4103/jmh.jmh_41_22.

Mariotti AJ. 2005. Estrogen and extracellular matrix influence human gingival fibroblast proliferation and protein production. *Journal of periodontology* 76:1391–1397. DOI: 10.1902/jop.2005.76.8.1391.

Mariotti A, Mawhinney M. 2013. Endocrinology of sex steroid hormones and cell dynamics in the periodontium. *Periodontology 2000* 61:69–88. DOI: 10.1111/j.1600-0757.2011.00424.x.

Mosley JF 2nd, Smith LL, Dezan MD. 2015. An overview of upcoming changes in pregnancy and lactation labeling information. *Pharmacy practice* 13:605. DOI: 10.18549/pharmpract.2015.02.605.

Park K-Y, Kim M-H, Choi S-H, Pang E-K. 2023a. Association of periodontitis with menopause and hormone replacement therapy: a hospital cohort study using a common data model. *Journal of periodontal & implant science* 53:184–193. DOI: 10.5051/jpis.2202480124.

Park J-H, Park M-S, Kim H-J, Lee H, Kim J-W, Song T-J. 2023b. Better oral hygiene is associated with a reduced risk of osteoporotic fracture: a nationwide cohort study. *Frontiers in endocrinology* 14:1253903. DOI: 10.3389/fendo.2023.1253903.

Wang L, Zhu L, Yao Y, Ren Y, Zhang H. 2021. Role of hormone replacement therapy in relieving oral dryness symptoms in postmenopausal women: a case control study. *BMC oral health* 21:615. DOI: 10.1186/s12903-021-01966-6.

World Health Organization. 1996. *Research on the menopause in the 1990s : report of a WHO scientific group*. Geneva, Switzerland: WHO.

World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects. 2013. *JAMA* 310:2191–2194.

Figure 1

An area chart showing the three resulting clusters from the 2-step cluster analysis, with the predictors “Period Continuation”, “Presence of Systemic Diseases” and “Age”.




Figure 2

A bar-chart showing the oral hygiene habits in the total sample (N=112) and the three clusters. No significant differences were observed using chi-square test.




Figure 3

A bar-chart showing the prevalence of self-reported tooth loss and tooth replacement in the total sample (N=112) and the three clusters. No significant differences were observed using chi-square test.




Table 1(on next page)

Demographic characteristics of the total sample (N=112) and the three clusters.

(p-Values in **BOLD** fonts are statistically significant using chi-square test at 0.05). ^a Only 5 (5%) of those on medication are on hormone replacement therapy (HRT). ^b Only 3 (3%) smoke cigarettes, the remaining smokers use water pipe.

Table 1: Demographic characteristics of the total sample (N=112) and the three clusters.
 1

Variable	Total Sample (N=112)	Cluster # 1 (n=65)	Cluster # 2 (n=25)	Cluster # 3 (n=22)	p Value
Marital Status - n (%)					
Not Married	27 (24)	18 (28)	4 (16)	5 (23)	0.502
Married	85 (76)	47 (72)	21 (84)	17 (77)	
Education - n (%)					
High school or lower	54 (48)	26 (40)	12 (48)	20 (91)	0.000
University degree	58 (52)	39 (60)	13 (52)	2 (9)	
Employment - n (%)					
No	38 (34)	22 (34)	4 (16)	12 (55)	0.021
Yes	74 (66)	43 (66)	21 (84)	10 (45)	
Siblings - n (%)					
None	27 (24)	17 (26)	4 (16)	6 (27)	0.001
1 – 2	22 (20)	17 (26)	4 (16)	1 (5)	
3 – 4	33 (29)	19 (29)	12 (48)	2 (9)	
> 4	30 (27)	12 (19)	5 (20)	13 (59)	
Recent Medication - n (%) *					
No	46 (41)	35 (54)	5 (20)	6 (27)	0.005
Yes	66 (59)	30 (46)	20 (80)	16 (73)	
Smoking - n (%) †					
No	96 (86)	53 (82)	22 (88)	21 (96)	0.255
Yes	16 (14)	12 (18)	3 (12)	1 (4)	

(p-Values in **BOLD** fonts are statistically significant using chi-square test at 0.05)
 2

* Only 5 (5%) of those on medication are on HRT
 3

† Only 3 (3%) smoke cigarettes, the remaining smokers use water pipe
 4

5

6

7

Table 2 (on next page)

Positive answers (YES) to the self-reported periodontal disease questionnaire by the total sample (N=112) and the three clusters.

(p-Values in **BOLD** fonts are statistically significant using chi-square test at 0.05)

Table 2: Positive answers (YES) to the self-reported periodontal disease questionnaire by the total sample (N=112) and the three clusters.

Variable	Total Sample (N=112)	Cluster # 1 (n=65)	Cluster # 2 (n=25)	Cluster # 3 (n=22)	p Value
Did you notice any swollen or reddish area in your gums (more than the usual)?	30 (27)	17 (26)	6 (24)	7 (32)	0.820
Do your gums bleed easily?	50 (45)	30 (46)	12 (48)	8 (36)	0.676
Do you feel pain from your gums?	28 (25)	12 (18)	8 (32)	8 (36)	0.161
Does your mouth feel dry?	29 (26)	11 (17)	8 (32)	10 (45)	0.022
Have you noticed an unpleasant smell from your mouth?	35 (31)	24 (37)	5 (20)	6 (27)	0.271
Do you have any abscess or pus in your mouth?	13 (12)	7 (11)	1 (4)	5 (23)	0.128
Does food get trapped between your teeth?	85 (76)	51 (79)	18 (72)	16 (73)	0.755
Do you feel that your teeth move / have moved?	15 (13)	5 (8)	4 (16)	6 (27)	0.060
Have you noticed that your teeth have changed in their position?	23 (20)	12 (18)	5 (20)	6 (27)	0.674
Have you noticed that your teeth have become longer?	18 (16)	8 (12)	4 (16)	6 (27)	0.255
Do you feel any sensitivity in your teeth?	46 (41)	28 (43)	13 (52)	5 (23)	0.111
Do you have any calculus or limestone deposits on your teeth?	60 (54)	33 (51)	16 (64)	11 (50)	0.494
Do you think you have any disease in your gums or tissues that support your teeth (periodontium) or loss of bone that fixes the teeth?	18 (16)	9 (14)	4 (16)	5 (23)	0.618
Has any dentist or hygienist ever told you that you have deep periodontal pockets?	6 (5)	3 (5)	2 (8)	1 (4)	0.801
Have you ever been told that you needed treatment for your gums / periodontal tissues?	16 (14)	7 (11)	3 (12)	6 (27)	0.150
Have you ever received any treatment for your gums / periodontal tissues?	26 (23)	13 (20)	6 (24)	7 (32)	0.522
Have you ever received any surgical treatment for your gums / periodontal tissues?	3 (3)	3 (5)	0 (0)	0 (0)	0.328

(p-Values in **BOLD** fonts are statistically significant using chi-square test at 0.05)

10

11