

Contrast in the density and biomass of fish in a reef system with different fishing intensity in the Mexican Caribbean (#106576)

1

First submission

Guidance from your Editor

Please submit by **31 Oct 2024** for the benefit of the authors (and your token reward) .



Structure and Criteria

Please read the 'Structure and Criteria' page for guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

6 Figure file(s)

2 Raw data file(s)

1 Other file(s)

! Custom checks

Vertebrate animal usage checks



Have you checked the authors [ethical approval statement](#)?



Were the experiments necessary and ethical?



Have you checked our [animal research policies](#)?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. **BASIC REPORTING**
2. **EXPERIMENTAL DESIGN**
3. **VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  **Impact and novelty is not assessed.** Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Contrast in the density and biomass of fish in a reef system with different fishing intensity in the Mexican Caribbean

Jose Manuel Jose Manuel Castro Perez ^{Corresp., 1}, Carmen A Villegas Sánchez ¹, Alejandro Medina Quej ¹, Rigoberto Rosas Luis ¹, Jesús Ernesto Arias González ²

¹ División de Estudios de Posgrado e Investigación, Tecnológico Nacional de México/Instituto Tecnológico de Chetumal, Chetumal, Q.R., Mexico

² Centro de Investigación y Estudios Avanzados del I.P.N-Unidad Mérida, Merida, Yucatán, Mexico

Corresponding Author: Jose Manuel Jose Manuel Castro Perez
Email address: jose.cp@chetumal.tecnm.mx

A wide range of fish species are caught in reef fisheries. However, fishing efforts tend to be highly selective in favor of large species, which generally have low population growth rates, making them more vulnerable to overfishing. When the decline of large predators occurs, fishing efforts start to focus on catching species from lower trophic levels, which can cause a trophic cascade effect. The objective of this research was to detect changes in the density and biomass of fish communities in areas with different fishing intensity in the study area. This study was carried out in Banco Chinchorro in the Mexican Caribbean and analyze the effect of fishing intensity on fish density and biomass, comparing data obtained from visual censuses with dependent information of the fishery. Evidence was found of a relationship between high fishing exploitation and low levels of density and biomass for *Epinephelus striatus*, *E. guttatus* and *Lachnolaimus maximus*. The decline of predators of non-commercial species had no effect on the density and biomass of these species. Both the density and biomass of the fish in the areas studied were influenced by the presence of algae, octocorals, hydrocorals and by variations in the CPUE. This study detected that density and biomass have decreased in some species belonging to the Serranidae and Lutjanidae families in areas with high fishing intensity. On the other hand, little evidence was found that the density and total biomass of families of noncommercially important species increased through the decline of their predators. Finally, although attempts were made to reduce the effect of structural complexity on the density and biomass of fish species, it was found that certain benthic groups and CPUE explained the observed changes in the density and biomass of fish assemblages.

Contrast in the density and biomass of fish in a reef system with different fishing intensity in the Mexican Caribbean.

José M. Castro-Pérez¹, Carmen A. Villegas-Sánchez¹, Alejandro Medina-Quej¹, Rigoberto Rosas-Luis² and Jesús E. Arias-González³.

1. Tecnológico Nacional de México/Instituto Tecnológico de Chetumal, Chetumal Quintana Roo, México.

2. Cátedras CONACyT. Tecnológico Nacional de México /Instituto Tecnológico de Chetumal, Chetumal Quintana Roo, México.

3. Centro de Investigación y Estudios Avanzados del I.P.N-Unidad Mérida, Yucatán, México.

Corresponding Author:

José M. Castro-Pérez¹

Tecnológico Nacional de México/Instituto Tecnológico de Chetumal, Chetumal Quintana Roo, México.

Email address: jose.cp@chetumal.tecnm.mx.

Abstract

A wide range of fish species are caught in reef fisheries. However, fishing efforts tend to be highly selective in favor of large species, which generally have low population growth rates, making them more vulnerable to overfishing. When the decline of large predators occurs, fishing efforts start to focus on catching species from lower trophic levels, which can cause a trophic cascade effect. The objective of this research was to detect changes in the density and biomass of fish communities in areas with different fishing intensity in the study area. This study was carried out in Banco Chinchorro in the Mexican Caribbean and analyze the effect of fishing intensity on fish density and biomass, comparing data obtained from visual censuses with dependent information of the fishery. Evidence was found of a relationship between high fishing exploitation and low levels of density and biomass for *Epinephelus striatus*, *E. guttatus* and *Lachnolaimus maximus*. The decline of predators of non-commercial species had no effect on the density and biomass of these species. Both the density and biomass of the fish in the areas studied were influenced by the presence of algae, octocorals, hydrocorals and by variations in the CPUE. This study detected that density and biomass have decreased in some species belonging to the Serranidae and Lutjanidae families in areas with high fishing intensity. On the other hand, little evidence was found that the density and total biomass of families of noncommercially important species increased through the decline of their predators. Finally, although attempts were made to reduce the effect of structural complexity on the density and biomass of fish species, it was found that certain benthic groups and CPUE explained the observed changes in the density and biomass of fish assemblages.

Keywords: coral reef; artisanal fishing; fish assemblage structure; Banco Chinchorro biosphere reserve; México.

Introduction

The increase in global fishing levels has generated environmental problems that are of public interest (Pauly et al., 2002). There is wide recognition that fish stocks throughout the world are under stress because of overfishing, coastal development, human population growth and climate change (Savo, Morton & Lepofsky, 2017; Russ et al., 2021).

Small-scale fisheries, including those in reef systems, are an important source of livelihood and food security for more than 1 billion people worldwide (Adam et al., 2015; Babcock, Tewfik & Burns-Perez, 2018). In the western Caribbean, more than a million people depend on the integrity and health of the Mesoamerican Reef System (MRS) for their livelihoods. The national economies of four countries (Belize, Guatemala, Honduras, and Mexico) benefit substantially from reef fishery resources and attractiveness as international tourist destinations (Zeller, Graham & Harper, 2011); however, there have been few studies on the effects of fisheries on reef fish communities (Pauly et al., 2002; Adam et al., 2015).

Although a wide range of fish species are caught in reef fisheries, fishing efforts tend to be highly selective in favor of large species, which generally have low rates of population growth, making them more vulnerable to overfishing (Sadovy de Mitcheson et al., 2013). Commercial species often include higher predators, such as serranids (groupers), lutjanids (snappers) and balistids (trigger fish). The decline of these species can increase the abundance of prey, which in turn influences the base of the food chain (Mumby et al., 2006; Ruppert et al., 2017). When the decline of large predators occurs, fishing efforts start to focus on catching species from lower trophic levels, which can cause a trophic cascade effect (Jackson et al., 2001; Mumby et al., 2006, 2012). On the other hand, in recent years, it has been found that coral bleaching in large areas of reefs due to the effect of climate change (Hoegh-Guldberg et al., 2018) and the increase in the frequency or intensity of environmental disturbances such as hurricanes (Knutson et al., 2010) has much more direct effects on fish (Graham et al., 2020) and on benthos (Hughes et al., 2018; Russ et al., 2021) than fishing. The coral reef systems of the Banco Chinchorro Biosphere Reserve (BCBR) support many different species, which are the main component of fisheries in the area. These fisheries are generally small-scale, artisanal and multispecific; however, economic progress, increasing coastal tourism, and an increase in human population have led to greater competition for fishing resources and possible overfishing. Most of the fishermen live in the City of Chetumal, but spend 15 to 30 days in Cayo Centro in the RBBC to carry out their fishing activities. Approximately 41 motor boats powered by an outboard motor operate in the study area. The capture is made daily and freediving equipment is used for it.

The fisheries in the BCBR are closely associated with the extraction of the spiny lobster (*Panulirus argus*) and the queen conch (*Aliger gigas*), although several species of fish are also caught throughout the year, including fishing conducted during spawning aggregations of different species, such as *Lutjanus analis* and *Baliste capriscus* (Castro-Pérez, González & Arias-González, 2011). The federal and state governments of Mexico declared the BCBR a marine protected area on July 19, 1996, with the purpose of conserving and protecting mangrove and coral ecosystems. However, the management program that was established for these purposes was carried out with few scientific studies of the flora and fauna in this complex reef. However, to improve the fishing management plans in marine protected areas, it is necessary to incorporate technical and scientific opinions as well as the knowledge of users and handlers.

Although this study analyzed data obtained from sampling carried out in 2007 and 2008, the resulting information is of great relevance because there is currently no information on the effect of fishing activity on fish communities in the study area, mainly due to how difficult the sampling logistics are to carry out this type of study. For this reason, the present research aims to detect changes in the density and biomass of fish communities in areas with different fishing intensity, so the proposed hypothesis is that fishing reduces the density and biomass of the main fish species in the BCBR.

Materials & methods

Area of study

The BCBR is part of the Mesoamerican Reef System, which covers 1000 km along the coasts of Honduras, Guatemala, Belize and Mexico, and it is located in southeastern Mexico in the state of Quintana Roo, 42 km from the Mahahual coast (18°47' - 18°23' N & 87°14' - 87°27' W). This reef was declared a marine protected area under the category of biosphere reserve in 1996. (Fig. 1A). Oval in shape, the BCBR covers an area of approximately 814.2 km² (45 km long and 18 km wide) and is composed of three cays: *Cayo Norte* (0.4 km²), *Cayo Centro* (5.4 km²), and *Cayo Lobos* (0.1 km²) (Chávez & Hidalgo, 1984; Jordán & Martín, 1987). The reef lagoon covers approximately 553.7 km², with depths ranging from 2 to 12 m and gradually decreases from south to north (González *et al.*, 2003).

Data collection.

Sampling of the fishery.

To identify areas with different fishing intensities in the study area, only the Langosteros del Caribe cooperative of the three that operate there was sampled, due to the complexity of working in all of them at the same time, for this purpose a monthly record was kept of the fish species caught between August 2007 and June 2008. During one week each month, the catches from the boats that arrived at the vessel that received the product were recorded. The weight was measured with a top loading balance with a capacity of 20 kg and an accuracy of 5 g. The fishing locations were recorded by consulting the position with the captain and using a map of the study area divided into 36 fishing quadrats (Fig. 1C). The CPUE was calculated as kilograms of fish caught per fisherman per hour (kg · fisherman⁻¹ · hr⁻¹) for each quadrant.

Sampling of fish communities

Visual censuses were used to record the abundance and size of the reef fish using the linear transect technique proposed by Brock (1954). Five visual censuses (replicas) were carried out at each sampling station, during the same day and time. Within a 50 m transect, the observer recorded the species detected within a distance of 2 m on each side and 5 m in front of the transect. For each observation, the diver recorded the abundance per species and estimated the total length of the fish. Fish lengths were estimated in 1 cm intervals for fish 0 to 10 cm TL and in 5 cm intervals for fish >10 cm TL. Samples were collected in August and September 2008 at 130 stations throughout the study area (Fig. 1B).

Sampling of the benthic groups in the reef habitats

The benthic cover was estimated to evaluate the characteristics of the habitat on the same date and in the sampling stations where the biological data of the reef fishes were obtained. A video camera was used to film the benthic substrate at each sampling station along five linear 50 m transects, with the footage then analyzed by stopping the image at specific time intervals until 40 frames were obtained. In each frame, at a series of 13 marked points (520 points per transect) that were systematically distributed around the monitor, the benthic organisms were identified according to six higher levels, known as morphostructural groups (MSG): scleractinian corals; octocorals; hydrocorals; algae; seagrass; and sponges (Arias-González *et al.*, 2011).

It is important to mention that in the years in which biological sampling was carried out, the Natural Commission for Protected Natural Areas (CONANP) did not grant permission to carry out research activities without collecting or handling specimens of species not considered at risk within the Banco Chinchorro Biosphere Reserve. However, this working group currently has permission (Oficio No. F00.9/DRBBCH/151/2024) to carry out research work.

Data analysis

Areas with different fishing intensities

Castro-Pérez, González & Arias-González (2011) identified seven fishing zones in the BCBR using the CPUE values for fish of commercial importance in 30 fishing quadrants because in 6 six quadrants, no catches were recorded (Fig. 1C). In the present study, to reduce the effect of the structural complexity of the benthos on the density and biomass of the fish, 21 fishing quadrants were used, located mainly on the reef slope and in the southern section of the reef lagoon, which are characterized by high structural complexity. Combining the CPUE data of the 10 commercially important species of the 21 quadrants and the zones identified by Castro-Pérez, González & Arias-González (2011), we classified the area into the following zones: Zone 1 (Quadrants 4, 34 and 36); Zone 2 (Quadrant 5); Zone 5 (Quadrant 25); and Zone 7 (Quadrants 2, 10, 15, 19, 21, 23, 24, 26, 27, 28, 29, 30, 31, 32, 33 and 35). Similarity was calculated using the Bray–Curtis index. The similarity coefficients were used to construct the similarity dendrogram; subsequently, data were permuted 999 times for a distribution to determine ANOSIM's R statistic ($R = 0$ is identical, $R = -1$ or 1 is most divergent), which facilitated the discovery of patterns among the quadrats and, thus, the detection of areas with different fishing intensities. Both analyses were performed with PRIMER 6.0 software (Clarke & Gorley, 2001).

Changes in the density and biomass of fish species due to fishing

The density (individuals/100 m²) and biomass (kilogram/100 m²) of each of the species were estimated to determine changes in these variables due to the fishing of species of commercial and non-commercial importance using data obtained from censuses visuals. The fish biomass was calculated via the exponential function $W = aL^b$, where W is the weight in kilograms, L is the length obtained from the length intervals and a and b are the constants from the length-weight relationship obtained from both Claro & García-Arteaga (1994) and FishBase. The average density and total biomass of commercially and noncommercially important species were compared among areas with different fishing intensities (detected via the similarity dendrogram) by means of a one-way analysis of variance (ANOVA). Subsequently, a multiple range test (Fisher's least significant difference, or LSD) (Zar, 1999) was applied to ascertain which areas

presented differences. The data were log(x)-transformed to fulfill both normality assumptions (Shapiro–Wilk test) and homoscedasticity (Levene’s test) (Zar, 1999). The same statistical tests were used to verify the average density and biomass differences among areas with different exploitation levels for each of the eight species caught with the most frequency in the study area: *Epinephelus striatus*, *Epinephelus guttatus*, *Lachnolaimus maximus*, *Lutjanus analis*, *Lutjanus griseus*, *Mycteroperca bonaci*, *Ocyurus chrysurus*, and *Sphyrna barracuda* (Castro-Pérez, González & Arias-González, 2011). Moreover, differences in the average density and biomass of the noncommercial species were tested among areas of different fishing intensities, with species grouped into twelve families: Acanthuridae, Balistidae, Chaetodontidae, Haemulidae, Holocentridae, Kyphosidae, Labridae, Malacanthidae, Mullidae, Pomacanthidae, Scaridae, and Serranidae.

Relationship of density and biomass of commercially important species with fishing pressure and benthic variables

The fish assemblages found in the different visual transects can vary for at least three reasons: fishing effort, differences in environmental conditions, and random variation (Clua & Legendre, 2008). Two redundancy analyses (RDA) were carried out using the CANOCO v4.5 program (Ter Braak & Smilauer, 2002), which separately related the density and biomass of the eight commercially important fish species with the coverage data for the different benthic morphostructural groups and the CPUE values for the species of greatest commercial interest. These multivariate methods were applied following the gradient length criterion, the species response models pertaining to the environment and the linear-type CPUE data. The density and biomass values were transformed via the Hellinger distance (Legendre & Gallagher, 2001), while the statistical significance was tested by means of Monte Carlo permutations (N = 9999).

Results

Areas with different fishing intensities

Based on the dendrogram using the CPUE values of the most commercially important species, the formation of three groups was detected with a similarity index of 60 % (Fig. 2A). The ANOSIM test provided statistical support to determine that the three groups were different in their composition and abundance ($R = 0.99$; $P < 0.05$). Considering this spatial separation and the distribution of the CPUE values for these quadrants, it was observed that Group 1 included quadrants that mainly belonged to the reef lagoon in the southern section of the study area (Cayo Lobos), which presented high CPUE values. Group 2 was formed by quadrats located on the reef slope, with intermediate CPUE values, while Group 3 generally contained quadrats located on the reef slope, with high CPUE values. In accordance with the above and to detect changes in the density and biomass of the fish species caused by the fishing activity, three areas with different fishing intensities were classified in the BCBR: moderate level fishing in the fore reef (FRMLF); High Level Fore Reef Fishing (FRHLF); and deep-sea fishing in reef lagoons (RLHLF) (Fig. 2B).

Changes in the density and biomass of fish species due to fishing

Analysis of the group of commercially important species among the areas of different fishing intensities revealed that the average density and biomass values showed significant differences (one-way ANOVA, $P < 0.05$). The analysis with the multiple range test showed that the highest values for these variables were recorded in the moderate fishing area and differed (LSD, $P < 0.05$) from the density and biomass of fish found in the high fishing areas (FRHLF and RLHLF), which were similar (LSD, $P > 0.05$) (Fig. 3A). The analysis of the group of noncommercial species did not reveal differences between the density and biomass values for the areas that presented different fishing efforts (one-way ANOVA, $P > 0.05$) (Fig. 3B).

The individual analysis of the eight commercially important species found that *E. guttatus*, *E. striatus* and *L. maximus* showed significant differences in their density and biomass values (one-way ANOVA, $P < 0.05$). The LSD test showed a decrease in the values of these variables in the high fishing areas, which did not show significant differences between them ($P > 0.05$) but differed ($P < 0.05$) from the moderate fishing zone (FRMLF) that showed high values of density and biomass of these species (Fig. 4).

The density and biomass per family of noncommercial species in the areas with different fishing intensities did not show significant differences (one-way ANOVA, $P > 0.05$), except for the Scaridae family (one-way ANOVA, $P < 0.05$). The analysis with the LSD test showed that the lowest values of density and biomass were present in the moderate fishing zone, but it differed ($P < 0.05$) from the zones where the highest catches occurred (FRHLF and RLHLF), which were statistically similar ($P > 0.05$) (Fig. 5).

Relationship of density and biomass of commercially important species with fishing pressure and benthic variables

The results of the redundancy analysis (RDA) showed low correlations (0.32 to 0.45) between the benthic coverage (seagrass, octocorals, hydrocorals and algae) and the CPUE fishing variable with the density and biomass of commercially important fish species. The percentage of variance explained was 78.1 to 88.0.

The arrangement of the density and environmental variables in the RDA undertaken in the present study revealed that along the first axis, the seagrass cover was associated with the quadrats belonging to the FRMLF and RLHLF areas and had a positive relationship with the density of *S. barracuda* and *L. griseus*. The hydrocoral cover was related to the quadrats in the FRHLF area. The octocoral cover and the grouped CPUE quadrats in the FRHLF area were detected along the second axis, with these variables revealing a strong positive association with the density of *O. chrysurus*. Algae cover was linked to the quadrats in the FRMLF area (C4 and C2), presenting a positive relationship with *B. capriscus* and a negative relationship with the density of *L. analis* (Fig. 6A).

Finally, the RDA results for both biomass and environmental variables showed that the first axis separated the quadrats with greater coverage of algae in the left part of the order where the quadrats belonging to the FRMLF area were grouped (C4 and C21), presenting a negative

relationship with *L. analis* biomass. The second axis showed that the octocoral, hydrocoral and CPUE cover was associated with the quadrats mainly pertaining to the FRHLF area, showing a negative relationship among the CPUE results, the hydrocorals and the biomass of the species *E. striatus*, *E. guttatus*, *L. griseus* and *S. barracuda* (Fig. 6B).

Discussion

Areas with different fishing intensities

This study used CPUE values to detect and analyze three areas with different fishing intensities within the BCBR to identify changes in the biomass and density of reef fish. Although the fishing quadrats of the areas were found in structurally complex habitats, their location and type of habitat within the system influenced their fishing intensity. The quadrats of the areas with high intensities of fish exploitation (FRHLF and RLHLF) were located mainly on the reef slope and the southern part of the reef lagoon (*Cayo Lobos*), where a great variety of habitats provide the fishing communities a greater surface area in which to search for the target resource, namely, the spiny lobster. In contrast, the quadrats in the area with the least fishing intensity (FRMLF) were located primarily in the middle and northern part of the reef complex, in which fishing efforts are carried out less frequently.

Changes in the density and biomass of fish species due to fishing

The results of the present study provide evidence that fishing activity has caused detectable changes in the density and biomass of commercially important fish species in the BCBR. The low density and biomass of the commercial species in the areas of high fishing intensity reflect the extraction of top predators, such as species belonging to the Serranidae and Lutjanidae families. This may be related to the selectivity of the fishing equipment used in the reef complex, which usually consists of harpoons (*Castro-Pérez, González & Arias-González, 2011*), via which the fishermen catch the most expensive and in-demand species in local markets (*Mumby et al., 2012; Kadison et al., 2017*). Therefore, the reduction of the species most valuable to fishing communities may be imminent, leading to the exploitation of other less important species, such as scarids, balistids and pomacanthids. Various studies have found that the decrease in the density and biomass of carnivorous and piscivorous species (Serranidae, Lutjanidae and Caranidae) is caused by an increase in fishing pressure (*Friedlander & DeMartini, 2002; Williams et al., 2011; Valdivia, Cox & Bruno, 2017*). However, it cannot be ruled out that other factors could be intervening in the changes in these biological variables, such as the effects of climate change and hurricanes on their habitats (*Graham et al., 2020; Russ et al., 2021*).

Individual analysis of the eight commercially important species found that *E. striatus*, *E. guttatus* and *L. maximus* showed a decrease in density and biomass in areas with high fishing intensity. The latter can be attributed to the fact that these species have a high economic value in local markets (US\$ 7.79 kg⁻¹), and for this reason, they are caught all year round throughout the reef system through selective fishing, although the highest captures of the first two species were in November, December and January (*J. Castro, 2011, pers. comm.*), months in which these species have been documented to aggregate for reproduction (*Aguilar-Perera & Aguilar-Davila, 1996; Sala, Ballesteros & Starr, 2001; Dahlgren et al., 2016*). It has been reported in several locations in the Caribbean that these three species form an essential part of the small-scale fishery (*Sadovy*

de Mitcheson et al., 2013; Sherman et al., 2018). *E. striatus* and *E. guttatus* have been highly exploited due to their reproductive aggregations that facilitate their capture at predictable sites and seasons (*Sadovy De Mitcheson et al., 2013; Cheung et al., 2013; Sherman et al., 2017*), which is why *E. striatus* is currently listed as threatened by the International Union for Conservation of Nature Red List (IUCN), while *E. guttatus* is considered a minor concern. *L. maximus* is a monandric and protogynous hermaphroditic species (*McBride & Johnson, 2007*), characteristics that have led to a decrease in its population due to overfishing (*Ault, Smith & Bohnsack, 2005*); therefore, at the regional level, this condition is classified as vulnerable by the IUCN.

In the Mexican Caribbean, there is little information on the biology and fisheries of many commercially important species; therefore, there is an urgent need to implement fisheries management and regulation strategies for these species. The only federal law concerning commercially important fish species in the study area is for the Nassau grouper *E. striatus*, which is a species that is associated with the closed season for the Red grouper *Epinephelus morio* and other species of grouper from the Gulf of Mexico and Caribbean Sea, which runs from 01 February to 31 March of each year.

The present research found little evidence that the density and total biomass of families of noncommercially important species increased through the elimination of their predators. This may be due to the small number of predatory species that are captured in the BCBR, causing a moderate indirect effect on the prey populations (*Jennings & Polunin, 1997; Roff et al., 2016*).

The differences in the density and biomass values found in the Scaridae family between areas with different fishing intensities may be linked to the size of the organisms that make up the family. These organisms are generally large in size (e.g., *Scarus vetula* and *Sparisoma viride*), causing them to be less susceptible to predation. In this reef system, organisms (*E. striatus*, *M. bonaci*, *L. analis* and *S. barracuda*) that are capable of consuming large prey are eliminated, leaving smaller predatory species (mesopredators) limited in their capacity to prey on larger organisms. This may have caused an increase in the density and biomass of the scarids in the highly fished areas (FRHLF and RLHLF). While some studies have found that families of large-bodied organisms are more vulnerable to fishing than families of smaller organisms (*Friedlander & DeMartini, 2002; Hawkins & Roberts, 2004*), there is little scientific evidence that species of the Scaridae family are being caught in the BCBR. The foregoing is the result of the implementation of awareness programs on the protection of parrotfish by the authorities of the reserve toward fishermen. These species are currently on the list of protected species in the Official Mexican Standard NOM-059-SEMARNAT.

Relationship of density and biomass of commercially important species with fishing pressure and benthic variables

The density and biomass of the main commercially important fish species in the areas with different fishing intensities were mainly influenced by the presence of algae, octocorals, hydrocorals and CPUE. Although some of these benthic groups contributed to the structural complexity of the reefs, the most important component was the scleractinian corals, which showed no association with the biological variables of the fish species. This was because the fishing quadrants were located on reefs with high structural complexity. Although attempts were

made to reduce the effect of structural complexity on the density and biomass of fish species (Luckhurst & Luckhurst, 1978; Chabanet et al., 1997; Darling, 2017), it was found that certain benthic groups and CPUE explained the observed changes in the density and biomass of fish assemblages in the BCBR. The effect of fishing was based on the negative relationship found between CPUE and the density and biomass of highly exploited species such as *E. striatus*, *E. guttatus*, *L. griseus* and *S. barracuda*.

Although there was a detectable effect of fishing on certain commercially important fish species, there was no evidence that this activity affected the structure of the fish community (top-down trophic cascade effect) that could cause a phase change of benthic groups, as found on highly exploited reefs (Jackson et al., 2001; Mumby et al., 2006; Mumby & Steneck, 2008). Castro-Pérez, González & Arias-González (2011) mentioned that reef-fish fishing in this reserve was moderate because the target species was usually the spiny lobster *Panulirus argus* and fish were only occasionally caught. However, it should be considered that various authors have found that certain environmental disturbances are associated with the degradation of reefs that directly affect the structure of the fish community (bottom-up effect) (Graham et al., 2020; Russ et al., 2021), such as ocean acidity (Kleypas & Yates, 2009; Veron, 2011), coral bleaching, (Hughes et al., 2018) and hurricanes (Knutson et al., 2010) caused mainly by climate change. For this reason, it is important to establish management and conservation measures for the main fish species of commercial importance in these protected areas to counteract the effect of these environmental disturbances on these species, since these environmental phenomena cannot be excluded in protected areas (Jones et al., 2004). In a recent study comparing protected and nonprotected reef areas after hurricanes, high fish biomass was found in protected areas, in contrast to areas in which fishing was not regulated (McClure et al., 2020).

Conclusion

In the BCBR throughout the year, the largest fishing exploitation is for the spiny lobster, but in the months when this crustacean begins to be scarce, the fishermen focus their greatest fishing efforts on the main species of commercially important fish. Due to the above, this study has detected evidence that the density and biomass have decreased in some species belonging to the Serranidae and Lutjanidae families in areas with high fishing intensity, therefore, there is an urgent need to implement fisheries management and regulation strategies for these species. On the other hand, little evidence was found that the density and total biomass of families of noncommercially important species increased through the elimination of their predators. This may be due to the small number of predatory species that are captured in the BCBR, causing a moderate indirect effect on the prey populations. Finally, although attempts were made to reduce the effect of structural complexity on the density and biomass of fish species, it was found that certain benthic groups and CPUE explained the observed changes in the density and biomass of fish assemblages in the BCBR.

References

- Adam TC, Burkepille DE, Ruttenberg BI, Paddock MJ. 2015. Herbivory and the resilience of Caribbean coral reefs: knowledge gaps and implications for management. *Marine Ecology Progress Series* 520:1–20 <http://doi.org/10.3354/meps11170>

- 413 Aguilar-Perera A, Aguilar-Dávila W. 1996. A spawning aggregation of Nassau grouper
414 *Epinephelus striatus* (Pisces: Serranidae) in the Mexican Caribbean. *Environmental Biology of*
415 *Fishes* 45:351–361 <https://doi.org/10.1007/BF00002527>
- 416 Arias-González J, Núñez-Lara E, Rodríguez-Zaragoza F, Legendre P. 2011. Reefscape proxies for
417 the conservation of Caribbean coral reef biodiversity. *Ciencias Marinas* 37(1):87–96
418 <https://doi.org/10.7773/cm.v37i1.1746>
- 419 Ault JS, Smith SG, Bohnsack J. 2005. Evaluation of average length as an estimator of exploitation
420 status for the Florida coral-reef fish community. *ICES Journal of Marine Science* 62(3):417–423
421 <https://doi.org/10.1016/j.icesjms.2004.12.001>
- 422 Babcock EA, Tewfik A, Burns-Perez V. 2018. Fish community and single-species indicators
423 provide evidence of unsustainable practices in a multi-gear reef fishery. *Fisheries Research*
424 208:70–85 <https://doi.org/10.1016/j.fishres.2018.07.003>
- 425 Brock VE. 1954. A preliminary report on a method of estimating reef fish populations. The
426 *Journal Wildlife Management* 18(3):297–308 <https://doi.org/10.2307/3797016>
- 427 Castro-Pérez JM, González GA, Arias-González JE. 2011. Characterizing spatial and temporal
428 reef fisheries in Chinchorro Bank Biosphere Reserve, northern Mesoamerican Reef System.
429 *Hidrobiológica* 21(2):197–209
- 430 Chabanet P, Ralambondrainy H, Amanieu M, Faure G, Galzin R. 1997. Relationships between
431 coral reef substrata and fish. *Coral Reefs* 16:93–102 <https://doi.org/10.1007/s003380050063>
- 432 Chávez EA, Hidalgo E. 1984. Spatial structure of benthic communities of Banco Chinchorro,
433 México. *Advances in reef sciences. A join meeting of the Atlantic Reef Committee an The*
434 *International Society for Reef Studies*. Univesity of Miami, pp. 26–28
- 435 Cheung WWL, Sadovy de Mitcheson Y, Braynen MT, Gittens LG. 2013. Are the last remaining
436 Nassau grouper *Epinephelus striatus* fisheries sustainable? Status quo in the Bahamas.
437 *Endangered Species Research* 20:27–39 <https://doi.org/10.3354/esr00472>
- 438 Clarke KR, Gorley RN. 2001. PRIMER v5: *User manual/ tutorial*. PRIMER-E, Plymouth, UK.,
439 192 pp.
- 440 Claro R, García-Arteaga JP. 1994. Crecimiento. In: Claro R, ed. *Ecología de los peces marinos de*
441 *Cuba*. Instituto de Oceanología Academia de Ciencias de Cuba and Centro de Investigaciones de
442 Quintana Roo, México, 321–402
- 443 Clua E, Legendre P. 2008. Shifting dominance among scarid species on reefs representing a
444 gradient of fishing pressure. *Aquatic Living Resources* 21(3):339–348
445 <https://doi.org/10.1051/alr:2008036>
- 446 Dahlgren CP, Buch K, Rechisky E, Hixon MA. 2016. Multiyear tracking of Nassau grouper
447 spawning migrations. *Marine and Coastal Fisheries* 8:522–535
448 <https://doi.org/10.1080/19425120.2016.1223233>
- 449 Darling ES, Graham NAJ, Januchowski-Hartley FA, Nash KL, Pratchett MS, Wilson SK. 2017.
450 Relationships between structural complexity, coral traits, and reef fish assemblages. *Coral Reefs*
451 36:561–575 <https://doi.org/10.1007/s00338-017-1539-z>

- 452 FISHBASE. World Wide Web Electronic Publication. <http://www.fishbase.org>
- 453 Friedlander AM, Demartini EE. 2002. Contrasts in density, size, and biomass of reef fishes
454 between the northwestern and the main Hawaiian Islands: the effects of fishing down apex
455 predators. *Marine Ecology Progress Series* 230:253–264 <https://doi.org/10.3354/meps230253>
- 456 González A, Torruco D, Liceaga A, Ordaz J. 2003. The shallow and deep bathymetry of the Banco
457 Chinchorro reef in the Mexican Caribbean. *Bulletin Marine Science* 73(1):15–22
- 458 Graham NAJ, Robinson JPW, Smith SE, Govinden R, Gendron G, Wilson SK. 2020. Changing
459 role of coral reef marine reserves in a warming climate. *Nature Communications* 11:2000
460 <https://doi.org/10.1038/s41467-020-15863-z>
- 461 Hawkins JP, Roberts CM. 2004. Effects of artisanal fishing on Caribbean coral reefs.
462 *Conservation Biology* 18(1):215–226 <https://doi.org/10.1111/j.1523-1739.2004.00328.x>
- 463 Hoegh-Guldberg O, Kennedy EV, Beyer HL, McClennen C, Possingham HP. 2018. Securing a
464 long-term future for coral reefs. *Trends in Ecology & Evolution* 33(12):936–944
465 <https://doi.org/10.1016/j.tree.2018.09.006>
- 466 Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum, JK,
467 Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs
468 JA, Hoey AS, Hoogenboom M, Lowe RJ, ... Wilson SK. 2018. Spatial and temporal patterns of
469 mass bleaching of corals in the Anthropocene. *Science* 359:80–83
470 <https://doi.org/10.1126/science.aan8048>
- 471 Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Louis W, Botsford LW, Bourque BJ,
472 Bradbury RH, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan, HS,
473 Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ, Warner RR. 2001. Historical overfishing
474 and the recent collapse of coastal ecosystems. *Science* 293:629–638
475 <https://doi.org/10.1126/science.1059199>
- 476 Jennings S, Polunin NVC. 1997. Impacts of predator depletion by fishing on the biomass and
477 diversity of non-target reef fish communities. *Coral Reefs* 16(2): 71–82
478 <https://doi.org/10.1007/s003380050061>
- 479 Jones GP, McCormick MI, Srinivasan M, Eagle JV. 2004. Coral decline threatens fish biodiversity
480 in marine reserves. *The Proceedings of the National Academy of Science (PNAS)* 101(21):8251–
481 8253 <https://doi.org/10.1073/pnas.0401277101>
- 482 Jordán E, Martin E. 1987. Chinchorro: morphology and composition of a Caribbean atoll. *Atoll*
483 *Research Bulletin* 310:1–20 <https://doi.org/10.5479/si.00775630.310.1>
- 484 Kadison E, Brandt M, Nemeth R, Martens J, Blondeau J, Smith T. 2017. Abundance of
485 commercially important reef fish indicates different levels of over-exploitation across shelves of
486 the U.S. Virgin Islands. *PLoS ONE* 12(7):e0180063
487 <https://doi.org/10.1371/journal.pone.0180063>
- 488 Kleypas JA, Yates KK. 2009. Coral Reefs and Ocean Acidification. *Oceanography* 22(4):108–117
489 <http://www.jstor.org/stable/24861028>

- 490 Knutson T, McBride J, Chan J, Emanuel K, Holland G, Landsea C, Held I, Kossin JP, Srivastava
491 AK, Sugi M. 2010. Tropical cyclones and climate change. *Nature Geoscience* 3:157–163
492 <https://doi.org/10.1038/ngeo779>
- 493 Legendre P, Gallagher ED. 2001. Ecologically meaningful transformations for ordination of
494 species data. *Oecologia*, 129(2):271–280 <https://doi.org/10.1007/s004420100716>
- 495 Luckhurst BE, Luckhurst K. 1978. Analysis of the influence of substrate variables on coral reef
496 fish communities. *Marine Biology* 49:317–323 <https://doi.org/10.1007/BF00455026>
- 497 McBride RS, Johnson MR. 2007. Sexual development and reproductive seasonality of hogfish
498 (Labridae: *Lachnolaimus maximus*), an hermaphroditic reef fish. *Journal of Fish Biology*
499 71:1270–1292 <https://doi.org/10.1111/j.1095-8649.2007.01580.x>
- 500 McClure EA, Sievers KT, Abesamis RA, Hoey AS, Alcala AC, Russ GR. 2020. Higher fish
501 biomass inside than outside marine protected areas despite typhoon impacts in a complex
502 reefscape. *Biological Conservation* 241:108354 <https://doi.org/10.1016/j.biocon.2019.108354>
- 503 Mumby PJ, Dahlgren CP, Harborne AR, Kappel CV, Micheli F, Brumbaugh DR, Holmes KE,
504 Mendes JM, Broad K, Sanchirico JN, Buch K, Box S, Stoffle RW, Gill AB. 2006. Fishing,
505 trophic cascades, and the process of grazing on coral reefs. *Science* 311(5757):98–101
506 <https://doi.org/10.1126/science.1121129>
- 507 Mumby PJ, Steneck RS. 2008. Coral reef management and conservation in light of rapidly
508 evolving ecological paradigms. *Trends in Ecology & Evolution* 23:555–563
509 <https://doi.org/10.1016/j.tree.2008.06.011>
- 510 Mumby PJ, Steneck RS, Edwards AJ, Ferrari R, Coleman R, Harborne AR, Gibson JP. 2012.
511 Fishing down a Caribbean food web relaxes trophic cascades. *Marine Ecology Progress Series*
512 445:13–24 <https://doi.org/10.3354/meps09450>
- 513 Pauly D, Christensen V, Gu  nette S, Pitcher TJ, Sumaila UR, Walters CJ, Watson R, Zeller D.
514 2002. Towards sustainability in world fisheries. *Nature* 418:689–695
515 <https://doi.org/10.1038/nature01017>
- 516 Roff G, Doropoulos C, Rogers A, Bozec YM, Krueck NC, Aurellado E, Priest M, Birrell C,
517 Mumby PJ. 2016. The ecological role of sharks on coral reefs. *Trends in Ecology & Evolution*
518 31(5):395–407 <http://dx.doi.org/10.1016/j.tree.2016.02.014>
- 519 Ruppert JLW, Vigliola L, Kulbicki M, Labrosse P, Fortin MJ, Meekan MG. 2017. Human
520 activities as a driver of spatial variation in the trophic structure of fish communities on Pacific
521 coral reefs. *Global Change Biology* 24:e67-79 <https://doi.org/10.1111/gcb.13882>
- 522 Russ GR, Rizzari JR, Abesamis RA, Alcala AC. 2021. Coral cover a stronger driver of reef fish
523 trophic biomass than fishing. *Ecological Applications* 31(1):e02224.
524 <https://doi.org/10.1002/eap.2224>
- 525 Sadovy de Mitcheson Y, Craig MT, Bertoni AA, Carpenter KE, Cheung WWL, Choat JH,
526 Cornish AS, Fennessy ST, Ferreira BP, Heemstra PC, Liu M, Myers RF, Pollard DA, Rhodes
527 KL, Rocha LA, Russell BC, Samoilys MA, Sanciangco J. 2013. Fishing groupers towards
528 extinction: a global assessment of threats and extinction risks in a billion dollar fishery. *Fish and*
529 *Fisheries* 14(2):119–136 <https://doi.org/10.1111/j.1467-2979.2011.00455.x>

- 530 Sala E, Ballesteros E, Starr RM. 2001. Rapid decline of Nassau grouper spawning aggregations in
531 Belize: fishery management and conservation needs. *Fisheries* 26:23–30
532 [https://doi.org/10.1577/1548-8446\(2001\)026<0023:RDONGS>2.0.CO;2](https://doi.org/10.1577/1548-8446(2001)026<0023:RDONGS>2.0.CO;2)
- 533 Savo V, Morton C, Lepofsky D. 2017. Impacts of climate change for coastal fishers and
534 implications for fisheries. *Fish and Fisheries* 18(5):877–889 <https://doi.org/10.1111/faf.12212>
- 535 Sherman KD, King RA, Dahlgren CP, Simpson SD, Stevens JR, Tyler CR. 2017. Historical
536 processes and contemporary anthropogenic activities influence genetic population dynamics of
537 Nassau Grouper (*Epinephelus striatus*) within the Bahamas. *Frontiers in Marine Science* 4:393
538 <https://doi.org/10.3389/fmars.2017.00393>
- 539 Sherman KD, Shultz AD, Dahlgren CP, Thomas C, Brooks E, Brooks A, Brumbaugh DR, Gittens
540 L, Murchie KJ. 2018. Contemporary and emerging fisheries in The Bahamas—Conservation and
541 management challenges, achievements, and future directions. *Fisheries Management and*
542 *Ecology* 25:319–331 <https://doi.org/10.1111/fme.12299>
- 543 Ter Braak CJF, Smilauer P. 2002. *CANOCO reference manual and CanoDraw for Windows*
544 *user's guide: Software for Canonical Community Ordination* (version 4.5). Biometris,
545 Wageningen <https://edepot.wur.nl/405659>
- 546 Valdivia A, Cox CE, Bruno JF. 2017. Predatory fish depletion and recovery potential on
547 Caribbean reefs. *Science Advances* 3(3):e1601303 <https://doi.org/10.1126/sciadv.1601303>
- 548 Veron JEN. 2011. Ocean acidification and coral reefs: An emerging big picture. *Diversity*
549 3(2):262–274 <https://doi.org/10.3390/d3020262>
- 550 Williams ID, Richards BL, Sandin SA, Baum JK, Schroeder RE, Nadon MO, Zgliczynski BJ,
551 Craig P, McIlwain JL, Brainard RE. 2011. Differences in reef fish assemblages between
552 populated and remote reefs spanning multiple archipelagos across the central and western
553 Pacific. *Journal of Marine Science* 1–14 <https://dx.doi.org/10.1155/2011/826234>
- 554 Zar JH. 1999. *Biostatistical analysis*. Prentice-Hall, Englewood Cliffs, pp. 672.
- 555 Zeller D, Graham R, Harper S. 2011. Reconstruction of total marine fisheries catches for Belize,
556 1950-2008. In: Palomares MLD, Pauly D, eds. *Too Precious to Drill: The Marine Biodiversity*
557 *of Belize*. Fisheries Centre, University of British Columbia, 142–151
558 [http://www.seaaroundus.org/doc/publications/books-and-reports/2011/Palomares-and-Pauly-too-](http://www.seaaroundus.org/doc/publications/books-and-reports/2011/Palomares-and-Pauly-too-precious-to-drill-Belize.pdf)
559 [precious-to-drill-Belize.pdf](http://www.seaaroundus.org/doc/publications/books-and-reports/2011/Palomares-and-Pauly-too-precious-to-drill-Belize.pdf)

560

561

Figure 1

(A) Location of the study area. (B) Distribution of the sampling stations for fish and benthic organisms. (C) scheme of the fishing quadrats.




Figure 2

(A) Dendrogram obtained from the Bray–Curtis similarity coefficients and used for identifying quadrats with different fishing intensities based on the CPUE data. (B) Location of the areas with different fishing intensities.

FRMLF = forereef moderate-level fishing; FRHLF = fore-reef high-level fishing; and RLHLF = reef lagoon high-level fishing.





Figure 3

Comparison of the average ($\pm$ SE) density and biomass among areas with different fishing intensities. (A) commercial species. (B) noncommercial species.

The homogeneous groups (LSD test, $P > 0.05$) are shown by the same letters for each treatment (areas): FRMLF = forereef moderate-level fishing; FRHLF = forereef high-level fishing; and RLHLF = reef lagoon high-level fishing.




Figure 4

Comparison of the average ($\pm$ SE) density and biomass per species of commercial importance among areas with different fishing intensities.

The homogeneous groups (LSD test, $p > 0.05$) are shown by the same letters for each treatment (areas): FRMLF = forereef moderate-level fishing; FRHLF = forereef high-level fishing; and RLHLF = reef lagoon high-level fishing.




Figure 5

Comparison of the average ($\pm$ SE) density and biomass of the *Scaridae* family caught among the areas with different fishing intensities.

The homogeneous groups (LSD test, $p > 0.05$) are shown by the same letters for each treatment (areas): FRMLF = forereef moderate-level fishing; FRHLF = forereef high-level fishing; and RLHLF = reef lagoon high-level fishing.




Figure 6

Ordination of the RDA data for the density (A) and biomass (B) of commercial fish with benthic coverage (fishing quadrats) and the variable of fishing (CPUE) in the BCBR. The symbols represent the areas with different fishing intensities: (red circle) FRM

The abbreviations corresponding to the species are the following: *Balcap* (*Balistes capriscus*); *Epigut* (*Epinephelus guttatus*); *Epistr* (*Epinephelus striatus*); *Lacmax* (*Lachnolaimus maximus*); *Lutana* (*Lutjanus analis*); *Lutgri* (*Lutjanus griseus*); *Mycbon* (*Mycteroperca bonaci*); *Ocychr* (*Ocyurus Chrysurus*); and *Sphbar* (*Sphyraena barracuda*). The letter C = fishing quadrat; FRMLF = forereef moderate-level fishing; FRHLF = forereef high-level fishing; and RLHLF = reef lagoon high-level fishing.

