

Identification and biological characterization of pathogen causing sooty blotch of *Ardisia crispa* (Thunb.) A.DC.

Demei Yang^{1†}, Jiangli Luo^{1†}, Ying Zhou^{1*}, Sixuan Zhou², Xiongwei Liu¹, Chang Liu^{1*}

¹ School of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang, Guizhou, 550025, China.

² Institute of Animal Husbandry and Veterinary Sciences of Guizhou Province, Nanming district Guiyang, Guizhou, 550005, China

† Demei Yang and Jiangli Luo are contributed equally to this work.

* Corresponding authors:

Ying Zhou, yingzhou71@sina.com; Guizhou University of Traditional Chinese Medicine, Huaxi District; Guiyang, Guizhou, 550025, China; 0851-88308059.

Chang Liu, 19liuchang@163.com; Guizhou University of Traditional Chinese Medicine, Huaxi District; Guiyang, Guizhou, 550025, China; 0851-88308059.

ABSTRACT

Sooty blotch, a primary leaf disease affecting *Ardisia crispa* (Thunb.) A.DC. (AC), significantly impacts both the yield and quality of this medicinal plant. However, the specific species of pathogenic fungi responsible for this disease and their mechanisms of pathogenesis remain unclear. To elucidate the etiology of sooty blotch, it is essential to investigate effective prevention and treatment methods, and provide a theoretical basis for the effective protection of AC. Initially, The alterations in internal organelles that result in sooty blotch were examined using transmission electron microscopy (TEM) and scanning electron microscopy (SEM). Additionally, differential genes were analyzed using differential display reverse transcriptase-PCR (DDRT-PCR) in both healthy and diseased leaves of AC. Moreover, the pathogenic fungi were separated, purified and identified from leaves infected with sooty blotch of AC, and subsequently, their pathogenicity and biological characteristics were tested. Furthermore, the inhibitory effect of pathogens was measured using the water extract of traditional Chinese medicine, based on the growth rate of the mycelium. The findings indicated that the photosynthesis rate of diseased leaves was slower than that of healthy leaves, as revealed by TEM and SEM analyses. Additionally, DDRT-PCR results demonstrated that the differentially expressed genes primarily

included those related to zinc finger proteins, acyl-CoA-transferases, and chloroplasts. The phylogenetic tree and pathogenicity test results showed that the pathogens causing sooty blotch of AC were *Annulohypoxyylon stygium* and *Diaporthe angelicae*. Microscopic observation revealed that *A. stygium* and *D. angelicae* exhibited distinct microscopic characteristics, and a pH range of 7-10 along with a subdued light environment were more conducive to the growth of pathogens. Additionally, the water extract of *Lonicerae flos* and *Ardisia crispa* (Thunb.) A.DC. had a strong antifungal action on the two pathogens of sooty blotch, and the water extract of *Ardisia crenate* Sims had a better antifungal action on the *D. angelicae*. In this study, *A. stygium* and *D. angelicae* were reported as pathogenic fungi causing sooty blotch for the first time, and affected the photosynthesis of AC leaf, and these study provides a theoretical basis for the diagnosis and prevention of *Ardisia crispa* (Thunb.) A.DC sooty blotch.

Keywords *Ardisia crispa* (Thunb.) A.DC., Sooty blotch, *Annulohypoxyylon stygium*, *Diaporthe angelicae*, biological characteristics, biocontrol mechanism.

INTRODUCTION

Sooty blotch is a widespread fungal disease affecting a variety of plant species, including Chinese medicinal materials, fruit trees, crops, flowers, and so on. A black mold develops on the infected areas of the plant, which impedes photosynthesis and leads to weak growth, early leaf drop, and primarily affects the leaves, and then the disease can also spread to the branches and eventually cause the plant to wilt gradually (Tham et al., 2023). The pathogens of sooty blotch are typically spread through rain or air currents, carried as spores or fragments of mycelium. When these spores or mycelia germinate on substrates containing honeydew, the sooty blotch pathogenic fungi colonize the plant surface. Initially, the affected plant surface may produce black, bituminous material or have a fluffy center. Over time, the mildew spreads and forms a layer that covers the entire leaves, shoots, and fruits. In the advanced stage of the disease, pathogenic fungi produce structures such as ostracods and conidia in the bituminous coal. These structures are the primary source of infection in the second year after overwintering, leading to multiple re-infections (Gleason et al., 2019). This leads to the loss of crops and medicinal plant, the decline of economic output, the degradation of landscape functions of garden plants, and even environmental pollution. The disease is very contagious and becomes more so as it spreads. It has inflicted severe damage on orchards, medicinal bases, and nursery lands (Araujo et al., 2021).

Radix Ardisia, a valued Miao herbal remedy in Guizhou, is known for its laryngeal benefits (Liu et al., 2022). In 2021, Kaihou Jian Spray, containing this herb, reached 880 million yuan in sales, reflecting rising market demand for its base plants. *Ardisia crispa* (Thunb.) A.DC. (AC), the botanical source of *Radix Ardisia*, is a medicinal plant with a history in traditional Chinese medicine (Wu et al., 2025). It contains alkaloids, saponins, and polysaccharides, which provide unique pharmacological effects such as clearing throat, removing phlegm and dampness, dispersing stasis and eliminating swelling (Wu et al., 2025). It's used for treating lung heat, cough, phlegm production, and asthma. AC also has anti-inflammatory and immunomodulatory properties, attracting modern medical research. However, AC is prone to sooty blotch in Guizhou's humid climate, which can lead to black mildew on leaves, hindering photosynthesis and potentially causing plant death. The increasing incidence of this disease negatively impacts yield and quality, causing economic losses.

Sooty blotch, a global issue, hosts various pathogenic fungi. Research shows 37 genera of pathogens are linked to it, with *Zygophiala*, *Microcyclospora*, *Stomiopeltis*, and *Ramichloridium* being some of these (Zhang et al., 2019). These fungi are adaptable and stress-resistant, surviving in different climates and soils. To manage sooty blotch diseases, it's crucial to study these fungi. Understanding their growth, reproduction, transmission, and pathogenicity is key for targeted prevention and control strategies. Currently, chemical pesticides like carbendazim and chlorothalonil are used, but they can harm the environment, lead to resistance, and reduce effectiveness. Biological control, on the other hand, offers pollution-free benefits and sustained control without resistance. It's a vital component for sustainable development. However, there's limited research on biological control of sooty blotch in AC. Comprehensive research is needed to provide new insights and strategies for managing this disease.

In this study, the changes of organelles were utilized to clearly observe the leaves of *A. crispa* under normal and infected conditions through transmission electron microscopy (TEM) and scanning electron microscopy (SEM). To analyze the differentially expressed genes in the leaves of AC using DDRT-PCR at the molecular level, the aim is to understand the response mechanism to sooty blotch. Additionally, typical pathogenic fungi responsible for sooty blotch were isolated and purified to elucidate the types of pathogenic fungi and the pathogenesis of sooty blotch. The microscopic characteristics of the pathogenic fungi were identified using microscopy. The pathogenicity of the isolated fungi was observed, and their morphological and molecular biological characteristics were investigated. Concurrently, the inhibitory effects of eight types of

traditional Chinese medicine on pathogenic fungi were assessed using the mycelium growth rate method, leading to the selection of appropriate antifungal agents. The study elucidates the molecular mechanisms of biological action from a genetic perspective, offering a reference for related research on the prevention and control of sooty blotch. It contributes to enhancing the quality and output of Chinese medicinal materials, and enriches the disease research of *A. crispa* sooty blotch, and facilitates more in-depth subsequent research.

MATERIALS AND METHODS

Materials

Normal and disease leaves of *Ardisia crispa* were collected from Guizhou University of Traditional Chinese Medicine (N.109.437569, E.19.19680), Jie-hong Zhao professor from Guizhou University of Chinese Medicine, identified the infected leaves as sooty blotch leaves. High speed refrigerated centrifuge Sorvall Legend Micro 17R (Thermo Fisher Scientific Co., LTD). Polysaccharide polyphenol plant total RNA extraction kit (Tiangen Biochemical Technology (Beijing) Co., LTD.), agar gel DNA recovery kit (Tiangen Biochemical Technology (Beijing) Co., LTD.), FastKing one-step genome cDNA first strand synthesis premix kit (Tiangen Biochemical Technology (Beijing) Co., LTD.). Fungal genome DNA extraction kit (solerbio, Beijing Solaibao Technology Co., LTD.), 2xTaq PCR MasterMixII (Tengen Company).

TEM and SEM analysis

Transmission electron microscopy (TEM) was performed using the following optimized protocol: the sample was prefixed with 3% glutaraldehyde and then 1% osmium tetroxide. To ensure dehydration, the concentration gradient of the dehydrating agent acetone is 30%→50%→70%→80%→90%→95%→100%. Then, the samples were permeated and embedded using a mixture of dehydrating agent and Epon812 embedding agent. The microtome was used to generate ultrathin sections with a thickness of 60 to 90 nm. The uranium acetate and lead citrate were performed to staining, and the images were gathered and analyzed to observe specific lesions. Scanning electron microscope (SEM): ultra-pure water was washed twice, 5 min each time, and the concentration gradient was 30%→50%→70%→80%→90%→95%→100% after concentration dehydration of ethanol series, and the specific lesions were observed after ion sputtering 10 min each time.

DDRT-PCR detection

The selection process involves several critical factors, including primer length, GC content, annealing temperature, and the avoidance of primer dimer formation. Typically, primer lengths range from 18 to 24 base

pairs to ensure sufficient specificity and binding strength. The GC content should be maintained between 40% and 60% to ensure the stability of the primer and its annealing efficiency. The annealing temperature is a crucial determinant for the effective binding of primers to templates; thus, preliminary experiments are essential to evaluate both the amplification efficiency and specificity of these primers, in order to select those most suitable for experimental requirements. The total RNA was derived from normal and disease leaves according to RNA extraction kit. The first cDNA was synthesized by using a Fastking one-step method, excluding the leading strand synthesis premixed reagent. Using the normal and disease cDNA as template, the PCR reaction system consisted of 1 μ L cDNA, 1 μ L random primers, 1 μ L anchored primers, 10 μ L 2xPCR mix, 7 μ L ddH₂O. The amplification procedures were as follows: 94 °C 5 min, 94 °C 40 s → 42 °C 40 s → 72 °C 90 s (40 cycles), 72 °C extension for 10 min. See anchored primers and random primers sequences in table S1 and table S2.

The PCR samples were loaded onto a 6% polyacrylamide gel, subjected to electrophoresis, and then photographed and recorded using a gel imaging system. The differential bands were excised and placed in a PCR tube containing 100 μ L of enzyme-free sterile water. The mixture was subsequently crushed into a fine powder and incubated in water at 85 °C for 15 minutes. Following centrifugation at 12,000 r/min for 15 minutes, 3 mol/L NaAc (pH 5.2) and 2.5 times the volume of anhydrous ethanol were added to 1/10 the volume of the supernatant to precipitate the DNA. The mixture was centrifuged again for 15 minutes at 12,000 r/min. The precipitate was washed once with 70% ethanol and once with anhydrous ethanol, each time centrifuging for 15 minutes. Finally, the DNA was dried and dissolved in 20 μ L DEPC water. 1 μ L template was used only for re-amplification, and the amplification system used was similar to that described above. The re-amplified DNA fragments were identified by 1% agarose gel electrophoresis. Marker D2000 was invoked as the control, and the electrophoresis conditions were set to 120V voltage, 400 mA current, and 30 minutes duration. The DNA was then recovered using an agarose gel DNA recovery kit and subsequently detected by 1% agarose gel electrophoresis. The recovered DNA was further amplified by PCR using the same a re-amplified DNA fragments were visualised in a 1% agarose gel mplification system as mentioned earlier. The 40 μ L amplification product was then collected for sequencing. The obtained sequences were tested and compared for similarity with different fragments and related genes using BLAST tools. Additionally, the function of gene searching was associated with Ensembl, Gene-Card, and other databases for comprehensive analysis.

Isolation and purification of pathogenic fungi

The infected leaves were washed with clean water and cut into 5 mm x 5 mm sizes, and make sure to remove external contaminants. The retained parts were cleaned with sterile water and then shaken with 75% ethanol for 30 seconds. Afterwards, shaken with 1% NaClO for 6 minutes. It's were washed with sterile water for 2-3 times and then cultured in Sand's solid medium for 5-7 days, followed by purification for 2-3 times.

Molecular identification

Molecular identification encompasses DNA sequence analysis, genome comparison, and the application of molecular markers. Through the integration of these genetic traits, it becomes possible to accurately distinguish and identify various fungal strains. The Fungal Genomic DNA (Soleibao) extraction kit was used to extract DNA, and 1% agarose gel electrophoresis was employed to detect the DNA with photos taken in the gel imaging system. For PCR amplification, 2 μ L DNA was used as the template and mixed with 8.2 μ L water, 10 μ L Mix enzyme, and 0.4 μ L upstream and downstream primers (Table S3). The PCR amplification was performed with the following conditions: initial denaturation at 94 °C for 5 minutes, followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 55 °C for 30 s, and extension at 72 °C for 30 s. Finally, extension was performed at 72 °C for 10 minutes. The PCR products were subsequently sequencing, and the pathogenic fungi gene sequences were analyzed using a BLAST platform in the NCBI database. Based on the identification and screening of reference strains, MEGA11.0 software was used to laborately constructed phylogenetic tree.

Pathogenicity test of pathogenic fungi

The leaf mycelium inoculation method was employed to treat the normal leaves of *A. crispa* *in vitro*, and normal leaves were used in the blank control experiment. The leaves were carefully placed on a sterilized filter paper in a flat plate under aseptic conditions on a clean table. Then, the mycelium was applied onto the leaf surface, and sterile water was added, ensuring that the water level did not exceed half of the leaves. It was cultured at a consistent temperature and humidity of 28 °C for a duration of 12 days. Throughout the culture period, the development of leaf disease was observed and the symptoms and characteristics were carefully recorded and photographed. The mycelium morphology of pathogens was observed under a microscope. It includes characteristics of fungal colonies, the shape and size of spores, and the structure of mycelia.

Biological characteristics of pathogenic fungi

Effect of pH on pathogenic fungi growth

The effects of pH on the growth of pathogenic fungi were investigated using Scharr's culture medium. Eight different pH levels (5, 6, 7, 8, 9, 10, 11 and 12) were ready. To create holes for inoculation, a 5 mm hole punch was utilized to drill holes in the edge of the pathogen colony. The pathogenic fungi cake was then inoculated into the holes, and each group was repeated three times. The cultures were incubated in an incubator at 28 °C. After 6 days of incubation, the colony diameter was monitored using the cross-crossing method.

Effect of light on pathogenic fungi growth

The effects of light on the growth of pathogenic fungi were examined in this study. The 5 mm hole punch was utilized to create holes at the periphery of the pathogenic fungi colony. The pathogenic fungi were cultured on Scharr's culture medium, and three different light conditions were tested: 24 hours of continuous light, 12 hours of light pursued by 12 hours of darkness, and 24 hours of complete darkness. Each treatment group was replicated three times, and the cultures were kept at a constant temperature of 25 °C. After 6 days of incubation, the colony diameter was detected by the cross-crossing method.

Biological control of pathogenic fungi

With water as solvent, 8 kinds of traditional Chinese medicine were extracted by reflux heating extraction method, combined with filtrate, heated and concentrated to 50 mL (1 g/mL), and stored at 4 °C for later use. A 25 mL concentrated Chinese medicine solution was mixed with 225 mL medium (100 mg/mL) and sterilized at 121 °C for 20 min to prepare the medicated medium. Good active cake with a diameter of 4 mm was drilled with a hole punch and placed in the center of the medicated medium. The positive control was 200 µg/mL for carbendazim and the blank control was sterile water. Each Chinese medicine water extract and pathogenic bacteria, positive control and blank control were repeated in three groups. After incubating at a constant temperature of 28 °C for 6 days, the colony diameter was monitored using the cross method. The experiment was conducted three times for each group, and the colony diameter of each group was measured using the cross-hatching method to obtain three sets of data. Subsequently, the average colony diameter and antibacterial rate for each group were calculated. The mean values and antibacterial rates were determined, along with the standard deviation for each dataset. Calculate the inhibition rate $[(\text{colony diameter of the blank group} - \text{colony diameter of the treatment group}) / (\text{colony diameter of the blank group} - 0.4) * 100\%]$. The results were analyzed using SPSS 26.0 statistical software and expressed as mean $\pm$ standard deviation ($\bar{x} \pm s$). One-way

ANOVA was utilized to compare the results among multiple groups.

RESULTS

The results of transmission electron microscope and scanning electron microscope

Under transmission electron microscopy (TEM), the chloroplasts of the normal leaf mesophyll cells were neatly arranged with clear and complete structures, and the chloroplasts contained a few starch granules. The cell membrane was discontinuous, and the cytoplasmic contents were dispersed in most regions (Figure 1A). In the mesophyll cells of the infected leaves, there were few chloroplasts, and most appeared wrinkled, circular in shape, and small in volume. The cytoplasm contained a small number of vacuoles and lipid droplets. Under electron microscopy, large vacuoles were suspected to contain free cytoplasm, and a large number of bacteria were also observed in the triangular area, with discontinuous cell membranes in some regions (Figure 1B).

Under scanning electron microscopy (SEM), the apparent morphology of the front, reverse and section of the normal and infected leaf was uncommon. There were close-arranged and complete upper epidermal cells on the front of healthy leaves, and the pores in the lower epidermis are uniform in size and orderly in arrangement, and most of them are concealed pores on the reverse side of the healthy leaves (Figure 2A). The healthy leaves section showed that the internal structure was transparent, the palisade tissue was arranged neatly, and the spongy tissue and vascular sheath cell tissue were normal. The helical mycelia were noted to be located between the mesophyll cells of the disease leaf (Figure 2B). These mycelia produced haustoria on the mesophyll cells, which greatly enhanced their ability to connect to the cells. Additionally, the mycelium was observed to protrude from the stomata on the opposite sides of the leaves. In the leaf cross-section, spiral advancing mycelia were observed to traverse through the cell space, while also protruding from the stomata on the back side of the leaf.

DDRT-PCR detection results and analysis

Total RNA was removed from both healthy and disease leaves and subjected to 1% agarose gel electrophoresis for detection. The bands observed were distinct, with the 28 S band appearing approximately twice as bright as the 18 S band, which detects a high level of RNA integrity (Figure S1). Three anchoring primers B0327, B0328 and B0329 were combined with 26 random primers B0301 ~ B0326 for PCR amplification, and 199 differential bands were isolated by polyacrylamide gel electrophoresis (Figure S2). 150

different bands were obtained from disease leaves samples, and 49 discrete bands were obtained from healthy leaves samples, with the size ranging from 100 to 2000 bp, which was consistent with the rule of difference display band results. After DNA recovery from 199 different bands by agarose gel electrophoresis (Figure S3), a total of 48 single bands with strong specificity were obtained (Figure S4). The sequencing results were compared using BLAST in the NCBI database, leading to a total of 69 sequences related to the study (Table S4). The gene sequences were annotated using a gene-bank, and the detailed results can be noted in Table S5. The subsequent homology analysis revealed the involvement of six different metabolic pathways in the samples, namely: 1) stress resistance; 2) Photosynthesis; 3) Cell respiration; 4) Cell proliferation; 5) Antioxidant; and 6) Intracellular transport.

Molecular identification results and analysis

A total of 11 pathogenic fungi strains were obtained from infected leaves of AC. Following phenotypic identification, the strains were designated as AC1 to AC11 (Table S6), respectively. The strains were distinguished according to their morphological characteristics and identified through sequencing (Figure 3; Table S7).

Pathogenicity test results and analysis

The disease morphology of *A. stygium* is different in different periods, the mycelium grew up obviously on the third day, a small part of the leaves began to turn yellow on the eighth day, the leaves are yellowish in whole and black spots can be seen all over the mycelium on the 12th day. The disease morphology of *D. angelicae* in different periods is various, the mycelium grew up obviously on the third day, the tip of the leaf turned yellow on the eighth day, the mycelium spread long foliar surface and most of the foliar surface turned yellow on the 12th day (Figure S5).

The morphological characteristics and the sequencing results showed that the pathogenic fungi of sooty blotch were *Annulohypoxylon stygium* and *Diaporthe angelicae* (Figure 4). Microscopic observations revealed distinct microscopic characteristics among diverse pathogenic fungi of sooty blotch. The mycelium of *A. stygium* exhibited a thick wall, large curvature, few branches, and a slender clumpy and separated appearance. On the other hand, the hypha of *D. angelicae* displayed a thin transparent wall, low curvature, abundant branches, and slender clusters (Figure 4).

The phylogenetic tree was analyzed to determine the evolutionary relationships of *A. stygium* (AC10)

(Figure 5 A), *A. stygium* (AC11) (Figure 5 B) and other relevant strains. The topology of phylogenetic trees presents an interesting insight into genetic differences. Notably, AC10 and AC11 were respectively closely related to MG881822.1 and KC343029.1, indicating a common descent. These observations strongly suggest that there are significant genetic differences between AC10, AC11 and their congeners. The results suggest that phylogenetic trees offer a powerful understanding about the evolutionary relationships of pathogenic fungi and their underlying genetic relationships with reference strain.

Results and analysis of biological characteristics of pathogenic fungi

Results and analysis of the effect of pH on the growth of pathogenic fungi

pH testing results revealed that both pathogenic fungi were able to grow within a pH range of 5 to 12 (Table S8). However, the mycelial growth rate of AC10 was significantly slower at pH 6 and below, indicating that an overly acidic environment was not suitable for its growth. Conversely, the mycelial growth rate of AC10 was faster within a pH range of 7 to 12, with no significant difference found between them. This indicates that the pathogenic fungi is more suitable for growth in a neutral or slightly alkaline environment. The growth pattern of AC11 mirrored that of AC10 (Table S9). The colony morphology of both pathogenic fungi on the sixth day of growth is shown in Figure 6A, while the colony diameters at different pH values on the sixth day in Figure 6B-C. If the entire plate is overgrown, the colony diameter is recorded as 8.60 cm.

Results and analysis of the influence of light on the growth of pathogenic fungi

The photoperiod test results revealed distinct effects of different light conditions on the pathogenic fungi. AC10 showed greater growth in full-bright conditions (Table S10), while there was no significant difference in mycelial growth rate between full-darkness and half-light/half-darkness conditions. However, the difference in growth rate was more pronounced compared to maximum light conditions. AC11, on the other hand, was not influenced by light (Table S11). The colony morphology of the two pathogenic fungi on the 6th day of growth and the colony diameter under different light conditions on the 6th day are shown in Figure 7. In the case of overgrowth of the entire plate, a diameter measurement of 8.60 cm was recorded.

Biological control results and analysis

It can be displayed that different Chinese herbal extracts had different inhibitory effects on pathogenic fungi at the same concentration. The results showed that the inhibitory rate of different plant extracts against pathogenic bacteria was significantly different (Table S12; Table 1). For the Pathogenic fungi *A. stygium*,

antifungal rate was *Ardisia crispa* (62.1633 ± 2.8403) > *Lonicerae flos* (54.3133 ± 0.6982) > *Alangium chinense* (Lour.) Harms (36.8733 ± 3.2520) > *Lonicera japonica* Thunb. (33.7200 ± 0.2300) > *Sophora tonkinensis* Gagnep. (32.6433 ± 4.0539) > *Ardisia crenata* Sims (28.2500 ± 1.8644) > *Aletris spicata* (Thunb.) Franch (6.6233 ± 2.0842) > *Polygonum capitatum* Buch-Ham ex D.Don (-13.3167 ± 2.7936). In summary, the water extract of *A. crispa* and *Lonicerae flos* exhibits potent inhibitory effects against *A. stygium*. The inhibitory rates for *Ardisia crenata*, *Lonicera japonica*, *Sophora tonkinensis* and *Alangium chinense* are approximately 30%. The inhibitory effects of *Aletris spicata* are not significant, and *Polygonum capitatum* may even promote their growth.

For the pathogenic fungi *Diaporthe angelicae*, antifungal rate was *Ardisia crispa* (94.5367 ± 3.3270) > *Ardisia crenata* (74.0433 ± 2.3027) > *Lonicerae flos* (49.5633 ± 7.3847) > *Lonicera japonica* (32.5967 ± 2.3460) > *Sophora Tonkinensis* (9.9200 ± 1.5216) > *Alangium chinense* (4.4267 ± 14.5974) > *Aletris spicata* (-0.5467 ± 4.3481) > *Polygonum capitatum* (-34.4300 ± 0.0000). It can be seen that it had a good inhibitory effect on *D. angelicae* (Table S13; Table 2), *Lonicerae flos*, *Ardisia crispa*, *A. crenata* Sims, while *Lonicera japonica* had a inhibitory rate of about 30%, *Sophora tonkinensis* and *Alangium chinense* had no obvious inhibitory effect, and *Aletris spicata* and *Polygonum capitatum* had a slight promoting effect on growth. The antifungal colony morphology of 8 kinds of Chinese medicine aqueous extracts on 2 kinds of pathogenic fungi is illustrated in Figure 8. This study provides a reference for the development of *Ardisia crispa* (Thunb.) A. DC, *Lonicerae flos* and *Ardisia crenata* Sims as green antibacterial drugs for sooty blotch, so as to reduce the dependence on antibacterial drugs and the occurrence of resistance of pathogenic bacteria, and make the cultivation of Chinese medicinal materials develop in a more healthy and sustainable direction.

DISCUSSION

Sooty blotch is a plant disease that is common in tropical and subtropical areas. As a broad concept for sooty blotch, suggesting that it should be collectively referred to as sooty blotch based on the outward forms of diseases caused by pathogenic fungi, rather than just trying to classify it (Wang et al., 2021). The host range of sooty blotch is extensive, including most tropical trees, as well as some economic fruit trees and flower plants. Sooty blotch are widely distributed and occur in warm and humid areas of the world, and most of them are diverse. For example, at least 60 species of coal pollution bacteria have been reported in the United States, at least 12 genera and 30 species in China, at least 25 species of coal pollution bacteria in Spain, 19 species in

Germany, 6 genera and 12 species in Turkey, 7 genera in Norway, 6 genera and 8 species in Poland, 4 genera and 6 species in Montenegro, and 3 genera and 5 species in Serbia and Slovenia (Li et al., 2016), Mauritius, Hawaii, South Africa, Reunion Island, Pakistan, India, South China (Ganesan et al., 2022), and Taiwan. When sooty blotch occurs, the branches and leaves are the focus with a layer of mildew that resembles soot, hence its name. Sooty blotch indirectly impacts the plant's physiological functions by reducing the effective photosynthetic area of the leaves, thereby altering the internal structure or composition of plant tissue. In the case of fruit trees, sooty blotch can also infect the fruits, causing black discoloration and decreasing their quality (Li et al., 2022). The harm caused by sooty blotch cannot be ignored (Liu et al., 2020). This common pathogenic fungi produces black colonies and spots on the surface of host plants, often damaging the leaves, stems, and fruits (Chomnunti et al., 2014). As a consequence, plant leaves may display black sooty mold layer symptoms on their upper surface.

Plant pathomycetes live on the surface of parasitizing plants and causing plant diseases. Pathogenicity tests revealed the presence of *A. stygium* and *D. angelicae* as the two types of pathogenic fungi. Microscopic observation showed distinct microscopic characteristics in these diverse pathogenic fungi. Through the study of their biological characteristics, it was observed that the mycelium growth rate of the *A. stygium*, which causes sooty blotch, was considerably slower at pH 6 and below compared to other groups. This indicates that an overly acidic environment is not suitable for the growth of pathogenic fungi (Li et al., 2022). The difference in growth rate was considerably larger when compared to the growth rate under full-light conditions. Among the approximately 1.0×10^5 species of fungi currently-known, more than 8000 species of fungi can cause diseases on plants (Sang et al., 2020). After plant tissues are adversely affected by pathogenic fungi, the defense enzyme system in the host body will undergo corresponding changes (Rajani et al., 2021). In the incompatibility interaction, local cell necrosis plaques significantly different from those surrounding healthy tissues are constituted at the infected site, that is, Hyper-sensitive reaction (HR) (Li et al., 2022). In affinity interaction, some fungi use stomata or trauma on the host surface to invade and produce some infection structures formed by specialized mycelia. The pathogenic fungi can invade and establish a parasitic relationship with the host, and resulting in plant infection (Mapuranga et al., 2022). In 1883, Frank reported that after the conidium of *Fusicladium* sp. And *Colletotrichum lindemuthianum* germinated on the surface of plants, the tips of the bud tubes expanded to form appressorium (Shang et al., 2021). The production of fungal appressorium directly

affects the molecular recognition of pathogenic fungi and host plants, as well as the invasion of pathogenic fungi and the occurrence of host plant defense response (Dar et al., 2023). The mycelium of many plant parasitic fungi grows on the surface of the host plant, from which collateral branches invade the host cell to absorb nutrients, which is called haustorium. The parasitism of Erysiphales stop and Uredinomycetes sap in plants is characterized by the formation of phagocytes (Polonio et al., 2021). The outer membrane of the haustorium, known as the haustorium outer membrane, serves as the interface between the haustorium and the host cell plasma membrane (Polonio et al., 2021). This change is believed to facilitate the passive diffusion of nutrients from the host cell towards the haustorium (Liu et al., 2021).

Referred to the DDRT-PCR technique using long primers and the above two-step PCR amplification as Enhanced Diff-ferential Display (Song et al., 2023; Christensen et al., 2023). The DDRT-PCR results indicated a more pronounced gene expression in infected *A. crispa* (Thunb.) A.DC. Since Fridovich proposed the theory of biological free radicals, the relationship between the cellular protective enzyme system and plant disease resistance has garnered widespread attention (Hassan et al., 2022). The zinc finger protein, a superfamily protein with a typical domain, was deemed to be associated with the regulation of plant growth, development, and stress adaptation, and plays a crucial role in regulating plant adaptation to stressful environments. Anion channels have the ability to regulate plant growth, development, and nutrient absorption, making them essential for plants to cope with drought, salt stress, aluminum toxicity, and phosphorus deficiency. The regulatory subunit can influence the activity of the catalytic subunit by binding to an effector and altering its conformation. CXE (Carboxylesterase), a supergene family of polymeric proteins, is primarily engaged in the hydrolysis of restricted substances in plants, such as esters, sulfates, and amides. It plays a major role in plant growth and development. In the study of signal transduction of plant immune response, 26 S proteasome plays a role in the regulation of protein degradation by numerous important transducers (Livneh et al., 2016). The discovery of these differential genes preliminarily elucidated the molecular mechanism of sooty blotch.

CONCLUSIONS

This study is the first to report *A. stygium* and *D. angeliae* as pathogenic fungi causing sooty blotch on *A. crispa*. Additionally, *L. flos* and *A. crispa* demonstrated strong antifungal activity against two sooty blotch pathogens. These findings provide a theoretical basis for developing eco-friendly prevention and control strategies for *A. crispa* sooty blotch, while also supporting the sustainable and healthy growth of the *A. crispa*

cultivation industry.

ADDITIONAL INFORMATION AND DECLARATIONS

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of the article. This work was supported by the Science and Technology Fund of Guizhou Province (QKHJC-ZK[2024]general 376); Graduate education Innovation program of Guizhou University of Traditional Chinese Medicine (YCXYKYS2023010); High-level Innovative Talents of Guizhou Province (QianKeHe platform talents-GCC [2023] 047).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

REFERENCES

- Araujo L, Pinto F A M F, de Andrade C C L, Gomes L B, Mituti T, Duarte V. 2021.** Pseudocercospora actinidiae causes sooty spot disease on kiwifruit in Santa Catarina, Brazil. Australasian Plant Disease Notes 16(1):22 DOI 10.1007/s13314-021-00436-w.
- Chomnunti P, Hongsanan S, Aguirre-Hudson B, Tian Q, Peršoh D, Dhami M K, Alias A S, Xu J C, Liu X Z, Stadler M, Hyde K D. 2014.** The sooty moulds. Fungal diversity 66:1-36 DOI

10.1007/s13225-013-0263-4.

- Christensen H, Olsen J E. 2023.** Primer Design: Design of Oligonucleotide PCR Primers and Hybridization Probes. Introduction to Bioinformatics in Microbiology. Cham: Springer International Publishing 89-110 DOI 10.1007/978-3-319-99280-8_5.
- Dar M A, Mahajan A. 2023.** An overview on Alternaria leaf in apple trees in Kashmir region. IJAR 9(6):01-03 DOI 10.22271/allresearch.
- Ganesan S, Singh H S, Biswal D, Kishore K, Samant D, Sahu S, Pattanaik M, Srinivas P, Samuel D K. 2022.** Etiology of sooty blotch disease of Aegle marmelos and its management. Pest Management in Horticultural Ecosystems 28(2):174-183 DOI 10.5958/0974-4541.2022.00057.1.
- Gleason M L, Zhang R, Batzer J C, Guangyu Sun G Y. 2019.** Stealth pathogens: the sooty blotch and flyspeck fungal complex. Annual Review of Phytopathology 57(1):135-164 DOI 10.1146/annurev-phyto-082718-100237.
- Hassan H M. 2022.** On 'Intracellular production of superoxide radical and of hydrogen peroxide by redox active compounds' by H. Moustafa Hassan, Irwin Fridovich. Archives of Biochemistry and Biophysics 726:109256 DOI 10.1016/j.abb.2022.109256.
- Li H Y, Zhang R, Sun G Y. 2016.** Research progress on ectophytic sooty blotch and flyspeck fungi. Mycosystema 35(12):1441-1455 DOI 10.13346/j.mycosystema.160128.
- Li B, Chen Y, Tian S. 2022.** Function of pH-dependent transcription factor PacC in regulating development, pathogenicity, and mycotoxin biosynthesis of phytopathogenic fungi. The FEBS Journal 289(7):1723-1730 DOI 10.1111/febs.15808.
- Li L, Zhu X M, Zhang Y R, Cai Y Y, Wang J Y, Liu M Y, Wang J Y, Bao J D, Lin F C. 2022.** Research on the molecular interaction mechanism between plants and pathogenic fungi. International Journal of Molecular Sciences 23(9):4658 DOI 10.3390/ijms23094658.
- Liu C, Pan J, Yin Z G, Feng T, Zhao J, Dong X, Zhou Y. 2022.** Integrated transcriptome and metabolome analyses revealed regulatory mechanisms of flavonoid biosynthesis in Radix Ardisia. PeerJ 10:e13670 DOI 10.7717/peerj.13670.
- Liu J, Liu M, Tan L, Huai B, Ma X, Pan Q, Zheng P, Wen Y, Zhang Q, Zhao Q, Kang Z, Xiao S. 2021.** AtSTP8, an endoplasmic reticulum-localised monosaccharide transporter from Arabidopsis, is recruited to

the extrahaustorial membrane during powdery mildew infection. *New Phytologist* 230(6):2404-2419 DOI 10.1111/nph.17347.

Liu W Y, Hui C M, Liu G L, Liu G W, Luo X H, Liu Y J. 2020. Influence of sooty blotch on photosynthetic and eco-physiological characteristics of *Phyllostachys nigra* var. *Henonis* 33 (04):160-168 DOI 10.5555/20203477317.

Livneh I, Cohen-Kaplan V, Cohen-Rosenzweig C, Avni N, Ciechanover A. 2016. The life cycle of the 26S proteasome: from birth, through regulation and function, and onto its death. *Cell research* 26(8):869-885 DOI: 10.1038/cr.2016.86

Mapuranga J, Zhang L, Zhang N, Yang W. 2022. The haustorium: The root of biotrophic fungal pathogens. *Frontiers in Plant Science* 13:963705 DOI 10.3389/fpls.2022.963705.

Polonio Á, Pérez-García A, Martínez-Cruz J, Fernández-Ortuño D, Vicente A D. 2021. The haustorium of phytopathogenic fungi: A short overview of a specialized cell of obligate biotrophic plant parasites. *Progress in Botany* 82:337-355 DOI 10.1007/124_2020_45.

Polonio Á, Fernández-Ortuño D, de Vicente A, Pérez-García A. 2021. A haustorial-expressed lytic polysaccharidemonooxygenase from the cucurbit powdery mildew pathogen *Podosphaera xanthii* contributes to the suppression of chitin-triggered immunity. *Molecular plant pathology* 22(5):580-601 DOI 10.1111/mpp.13045.

Rajani P, Rajasekaran C, Vasanthakumari MM, Olsson SB, Ravikanth G, Uma Shaanker R. 2021. Inhibition of plant pathogenic fungi by endophytic *Trichoderma* spp. through mycoparasitism and volatile organic compounds. *Microbiological Research* 242:126595 DOI 10.1016/j.micres.2020.126595.

Shang J, Shang Y, Tang G, Wang C. 2021. Identification of a key G-protein coupled receptor in mediating appressorium formation and fungal virulence against insects. *Science China Life Sciences* 64:466-477 DOI 10.1007/s11427-020-1763-1.

Song T K, Zamri N E M, Ibrahim R, Mohtar J A, Abbas H, Rahman A M A. 2023. Differential display reverse transcription polymerase chain reaction (DDRT-PCR) for grey oyster mushroom samples grown with acoustic sound treatment. *IOP Conference Series: Earth and Environmental Science*. 1216(1):012018 DOI 10.1088/1755-1315/1216/1/012018.

Tham K X, Batzer J C, Gleason M L, Rosli H. 2023. Sooty blotch and flyspeck on mango (*Mangifera indica*

L.) caused by Peltaster-like species in Malaysia. *Crop Protection* 171:106288 DOI 10.1016/j.cropro.2023.106288.

Wang B, Zhang R, Sun G Y. 2021. Research progress of adaptive evolution mechanism of sooty blotch and flyspeck fungi on plant surface niche. *Mycosystema* 40(10):2550-2561 DOI 10.13346/j.mycosystema.210240.

Wu DH, Lyu CG, Zhao D, Yang CG, Liu SQ, Zhu JT, Yang YL, Guo LP, Kang CZ. 2025. *Ardisia Crispae* Radix et Rhizoma: A review of botany, traditional uses, phytochemistry, pharmacology, and toxicology. *J Ethnopharmacol* 338(Pt 3):119093. doi: 10.1016/j.jep.2024.119093.

Zhang J Y, Luo G M, Chai H W, Zhu Z H, Liu P P. 2019. Identification of the pathogen causing gardenia sooty mold disease and screening of bacteriostatic agents. *Chinese Herbal Medicines* 42(05):1018-1022. DOI 10.13863/j.issn1001-4454.2019.05.013.