

Early conch morphology of a gigantic Cretaceous ammonoid, *Pachydesmoceras* (Desmoceratidae) (#112946)

1

First revision

Guidance from your Editor

Please submit by **4 Apr 2025** for the benefit of the authors .



Structure and Criteria

Please read the 'Structure and Criteria' page for guidance.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data.



Image check

Check that figures and images have not been inappropriately manipulated.

If this article is published your review will be made public. You can choose whether to sign your review. If uploading a PDF please remove any identifiable information (if you want to remain anonymous).

Files

Download and review all files from the [materials page](#).

1 Tracked changes manuscript(s)

1 Rebuttal letter(s)

5 Figure file(s)

2 Table file(s)



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. **BASIC REPORTING**
2. **EXPERIMENTAL DESIGN**
3. **VALIDITY OF THE FINDINGS**
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  **Impact and novelty is not assessed.** Meaningful replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Conclusions are well stated, linked to original research question & limited to supporting results.



The best reviewers use these techniques

Tip

Example

Support criticisms with evidence from the text or from other sources

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult. I suggest you have a colleague who is proficient in English and familiar with the subject matter review your manuscript, or contact a professional editing service.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Early conch morphology of a gigantic Cretaceous ammonoid, *Pachydesmoceras* (Desmoceratidae)

Ryoji Wani ^{Corresp. 1}

¹ Faculty of Environment and Information Sciences, Yokohama National University, Yokohama, Japan

Corresponding Author: Ryoji Wani
Email address: wani@ynu.ac.jp

Gigantic ammonoids, with conch diameters exceeding 1 m, remain one of the most enigmatic groups of extinct organisms. Their paleoecology has been the subject of ongoing debate, with some uncertainties arising from preservation biases, especially of an early conch. This study focuses on an exceptionally preserved early conch of the giant Cretaceous ammonoid *Pachydesmoceras denisonianum* from southern India. Conch morphology and the ontogenetic trajectories of constrictions and septal spacings were examined. The results indicate that constrictions were frequently present in the early conch; based on the shell layers observed in the cross-section, these constrictions likely resulted from periods of halted or slowed growth. The common occurrence of constrictions during early ontogeny suggests that *Pachydesmoceras* lifespan may have been longer than previously assumed. Additionally, the ontogenetic patterns of septal spacing do not appear to reflect these growth halts or slowdowns.

Early conch morphology of a gigantic Cretaceous ammonoid,

Pachydesmoceras (Desmoceratidae)

Ryoji Wani¹

¹Faculty of Environment and Information Sciences, Yokohama National University, Yokohama, Japan; wani@ynu.ac.jp

ABSTRACT

Gigantic ammonoids, with conch diameters exceeding 1 m, remain one of the most enigmatic groups of extinct organisms. Their paleoecology has been the subject of ongoing debate, with some uncertainties arising from preservation biases, especially of an early conch. This study focuses on an exceptionally preserved early conch of the giant Cretaceous ammonoid *Pachydesmoceras denisonianum* from southern India. Conch morphology and the ontogenetic trajectories of constrictions and septal spacings were examined. The results indicate that constrictions were frequently present in the early conch; based on the shell layers observed in the cross-section, these constrictions likely resulted from periods of halted or slowed growth. The common occurrence of constrictions during early ontogeny suggests that *Pachydesmoceras*

lifespan may have been longer than previously assumed. Additionally, the ontogenetic patterns of septal spacing do not appear to reflect these growth halts or slowdowns.

INTRODUCTION

Gigantic marine invertebrates evolved across several groups during and after the Cambrian explosion (Klug *et al.*, 2015b). Ammonoids flourished from the Devonian to the end of the Cretaceous period and are a prime example of this trend. During this period, some ammonoid species grew to enormous sizes, with conch diameters exceeding 1 m, and these species are known worldwide (Klug *et al.*, 2015b; Tajika, Nützel & Klug, 2018). One of the most notable examples is the Late Cretaceous *Parapuzosia seppenradensis* (Puzosiinae, Desmoceratidae, Perisphinctina; for higher taxonomy, see Besnosov & Mikhailova, 1991, and Yacobucci, 2015), which is currently recognized as the largest ammonoid species (Ifrim *et al.*, 2021). Another prominent gigantic ammonoid is *Pachydesmoceras* (Puzosiinae, Desmoceratidae, Perisphinctina; Kin & Niedźwiedzki, 2012; Tajika, Nützel & Klug, 2018), which occurred from the upper Albian to upper Turonian in regions across Europe, Africa, Madagascar, India, Japan, and New Zealand (Wright, 1996). The type species of this genus is *Pachydesmoceras denisonianum*, with the lectotype (designated by Matsumoto, 1987) originating from the northeast of the village of Odiyam in the Ariyalur region of southern India.

The inner whorls of gigantic ammonoids are often preserved poorly, primarily because the earliest whorls tend to be dissolved during diagenesis (Maeda, 1987; Maeda & Seilacher, 1996; Maeda et al., 2010; Wani & Gupta, 2015). Ifrim et al. (2021) investigated the ontogeny of the largest ammonoid, *Parapuzosia seppenradensis*, including the early conch morphology (approximately 110 mm in conch diameter). In contrast, the internal conch morphology of *Pachydesmoceras* has rarely been documented, with *P. pachydiscoide* and *P. kossmati* being the exceptions (Matsumoto, 1988; Kennedy & Klinger, 2014).

During our fieldwork in the Ariyalur area of southern India (a collaborative fieldwork conducted with Dr. K. Ayyasami from the Geological Survey of India, authorized under permit no. 874/2002), an exceptionally preserved early conch of *Pachydesmoceras denisonianum* (the type species of *Pachydesmoceras*) was collected from a horizon nearly identical to that of its lectotype, close to the type locality. In this study, the outer and internal morphology of this specimen were examined to identify the early conch morphology of a gigantic ammonoid that has rarely been recognized in previous studies.

MATERIALS

A single gigantic ammonoid specimen was discovered in the Karai Formation (Uttatur Group), located approximately 4 km southwest of Odiyam village (11°13'01"N, 78°59'32"E) in the

Ariyalur area (for detailed geological information, see *Sundaram & Rao, 1986; Sundaram et al., 2001*). The ammonoid, with a conch diameter of approximately 0.7 m, was unfortunately fragmented into several pieces, preventing the collection of the complete specimen. Only the nearly intact innermost part of the conch was retrieved in the field. Based on the conch morphology of the broken fragments and the large conch diameter, this specimen was identified as *Pachydesmoceras denisonianum*. The geological age of the specimen was considered to be late Albian–early Cenomanian, based on associated species from nearby localities, including *Mariella bergeri*, *Puzosia compressa*, and *Mortoniceras* spp.

The preserved outermost whorl shows little to no outer shell layer, likely due to dissolution and/or peeling caused by the adjacent broken whorl. Consequently, the presence and prominence of ribs on the shell surface of the preserved outermost whorl could not be accurately assessed.

The specimen examined in this study is deposited in the Mikasa City Museum (MCM), Hokkaido, Japan, with the registered number (MCM-W2145). Additionally, for the comparison of conch morphology, the morphological data presented in *Matsumoto (1988)* were employed.

METHODS

To observe the outer conch shape of the early conch, the specimen was first blackened using

colloidal graphite and then whitened with ammonium chloride. Conch terminology follows that of Klug *et al.* (2015a).

The examined specimen was subsequently polished along its median plane (plane of symmetry) using silicon carbide powder. Constrictions were observed on the median plane, and the spacing between successive constrictions was measured. These spacings are the rotational angle between two consecutive constrictions at the ventral positions. The center of rotation was defined as the center of the approximated logarithmic spiral. The relationship between the measured constriction spacings and conch diameter was examined throughout early ontogeny. Additionally, cross-sectional observations of the shell structure, particularly around the constrictions, were made.

The septal spacing between successive septa was also measured on the median plane. These spacings were defined as the rotational angle between two consecutive septa at the positions where the septum intersects with the siphuncle. The relationship between the measured septal spacings and conch diameter was examined throughout early ontogeny.

EARLY CONCH MORPHOLOGY OF *PACHYDESMOCERAS DENISONIANUM*

Outer morphology of early conch

The early conch of the gigantic *Pachydesmoceras denisonianum* (109 mm in conch diameter)

exhibits a discoidal shape ($B/D = 0.38$), is weakly compressed ($B/H = 0.91$), and has a moderate whorl expansion ratio ($WER = 1.90$) and a very wide and evolute umbilicus ($U/H = 0.78$) (Table 1; Fig. 1). The whorl is ovoid in cross-section, with a rounded venter, convex flanks, rounded umbilical shoulder, and steep umbilical wall.

Constrictions are frequently observed, with 10 distinct constrictions identifiable on the preserved outermost whorl (Fig. 1A; Table S1). All constrictions are concave, prosiradiate, and project forward on the venter, exhibiting nearly uniform widths and depths from the umbilical seams to the venters. Due to the dissolution or peeling of the outer shell layer caused by the adjacent broken whorl, the presence and intensity of other ornamentations, such as ribs, remain uncertain.

Internal morphology of early conch

The earliest whorl is dissolved, and only approximately 1.5 whorls are preserved, with the smallest preserved conch diameter measuring 32 mm (Fig. 2). Thus, the earliest conch morphology, such as the ammonitella and initial chamber, cannot be observed or evaluated. The early conch follows a logarithmic spiral, with a moderately embracing imprint zone rate ($IZR = 0.29$; Table 1).

Twenty constrictions, including the 10 that are discernible on the outer morphology, are

recognized at the median plane (Figs. 1, 2; Table S1). The conch diameters corresponding to the first and second constrictions are not measurable due to the dissolution of the earliest whorl. The smallest measurable conch diameter with a constriction is 33 mm (corresponding to the third constriction; Fig. 2; Table S1). Due to the dissolution of the earliest whorl, it remains unclear whether constrictions exist before the first observed constriction (Fig. 2). The rotational angles between successive constrictions of the 20 observed constrictions in the specimen ranged from 22° to 82° (average = 42.3°, standard deviation = 13.29°; Figs. 1–3; Table S1).

Two well-preserved constrictions reveal that ribs are positioned adapically to the constrictions in the cross sections (Fig. 2). Additionally, the outer shell layer at the constrictions is distinctly oblique to the shell surface. This feature is especially noticeable in the adoral parts of the constrictions (Fig. 2B, C). In contrast, the inner shell layer remains continuous, even across the constrictions.

The preserved first septum was located at a conch diameter of 38 mm (phragmocone diameter without the body chamber), and 25 septa were recognized in this specimen (Fig. 2; Table S1). The rotational angles between successive septa ranged from 18° to 26° (average = 21.9°, standard deviation = 1.93°; Figs. 2, 3; Table S1).

DISCUSSION

Intraspecific and interspecific comparison of early conch morphology

Based on the measurements of *Pachydesmoceras denisonianum*, including the lectotype and specimen examined in this study (Table 1), intraspecific comparisons were made regarding the conch morphology at different conch diameters (Fig. 4). The graph between umbilicus width (U) and conch diameter (D) indicates that the U/D ratio tended to increase with growth, reflecting an enlarging umbilicus relative to the conch diameter (Fig. 4D). In contrast, the graphs of morphological ratios for whorl width (W/H and W/D; Fig. 4E, F) show greater variation in larger specimens, although the **sample size** is limited. These trends suggest that whorl width (W) exhibited more variation than other morphological parameters. This variability in whorl width may be attributed to the inflation of body chambers in *Pachydesmoceras* as it matured (Matsumoto, 1988), or to diagenetic deformation in larger specimens (Maeda et al., 2010; Klug et al., 2015b).

A relatively smaller conch of *P. denisonianum* (166 mm in diameter; Table 1) was reported by Collignon (1961, pl. 8), collected from the Cenomanian of Madagascar. This specimen displays crowded ribs of unequal length, with some of the longer ribs accompanied by indistinct constrictions (Matsumoto, 1988). In the specimen examined in this study, ribs are not visible on the shell surface, likely due to the preservation of the outer shell layer. Given the relationship between constrictions and ribs observed in Collignon's specimen (1961), a similar

relationship is likely present in the examined specimen.

At a later stage in *P. denisonianum*, Matsumoto (1988) noted that constrictions became less distinct and may only appear as shallow furrows along some of the longer ribs. Considering this observation alongside the early conch morphology identified in this study, it can be inferred that the frequency and prominence of constrictions would decrease with growth in *P. denisonianum*.

Early conch morphology has been recognized in *P. pachydiscoide* and *P. kossmati* (Matsumoto, 1988; Kennedy & Klinger, 2014). According to Matsumoto (1988) and Kennedy & Klinger (2014), early conchs of both species exhibit frequent constrictions associated with ribs, which later become less distinct. In contrast, the ribs become more prominent, thicker, and coarser in the later stages. In the mature stage, the ribs gradually weaken, resulting in a nearly smooth conch (Matsumoto, 1988). From the early to later stages, these ontogenetic trends in shell ornamentation are similar to those observed in *P. denisonianum*. Therefore, observing the early conch in the type species of the genus *Pachydesmoceras* is crucial for interpreting the taxonomy and phylogenetic relationships of this and related genera.

Implication for shell growth from constrictions and associated ribs

The observation of the cross section of the examined specimen revealed that ribs are located just

adapically to the constrictions (Fig. 2). The nearly identical width and depth of each constriction (Fig. 1) suggests that the ribs at these positions extend along the constrictions, resulting in “long ribs.” Similarly, in the relatively smaller specimen from the Cenomanian of Madagascar (Collignon, 1961), some long ribs are accompanied by indistinct constrictions. Matsumoto (1988) noted that long ribs begin to appear at least by the middle stage in *Pachydesmoceras*. In later stages, these long ribs are spaced at gradually broadening intervals, with shorter ribs intercalated. However, based on the current observations, long ribs may appear as early as the initial stages, at least when the conch diameter reaches 33 mm.

An examination of the well-preserved constrictions revealed that the outer shell layer at these constrictions is distinctly oblique to the shell surface, which is especially noticeable in the adoral parts of the constrictions (Fig. 2). The discontinuous shell layers across the constrictions and ribs suggest a transition in growth phases at these points, potentially indicating a growth halt or slowdown. Similar distinct shell layers have been reported by Bucher *et al.* (1996), who studied the shell layers associated with conch ornamentation called megastria. This thick, continuous line extends around the flanks and venter of an ammonoid conch. Their study observed discontinuous outer shell layers and continuous inner shell layers at the megastria, concluding that such discontinuities represented growth halts. Based on these observations, constrictions have also been hypothesized to be associated with growth halts or slowdowns

(Arkell, Kummel & Wright, 1957; Kulicki, 1974; Kennedy & Cobban, 1976; Obata et al., 1978; Westermann, 1990; Bucher et al., 1996; Bucher, 1997; Klug et al., 2015a; De Baets, Landman & Tanabe, 2015; Uryu, 2015). The findings in this study align with this hypothesis. The repeated occurrence of constrictions in the early conch (at least 20 constrictions up to a conch diameter of 109 mm) suggests that the examined specimen experienced several growth halts or slowdowns during early ontogeny. As constrictions become less frequent in the later stages of *Pachydesmoceras*, growth halts, or slowdowns may decrease with growth. However, the frequency of constrictions could vary between specimens or species. Since only a single specimen was analyzed in this study, further investigation with additional specimens is necessary to determine whether the frequency of these growth halts or slowdowns are species-dependent, genus-dependent, or environmentally influenced (e.g., seawater temperature, chemical composition of seawater, nutritional condition, and oxygen condition).

Constrictions are commonly observed on the conchs of *Pachydesmoceras* and other ammonoids, in contrast to Cretaceous nautiloids from the same region, which do not exhibit constrictions except a nepionic constriction formed at hatching (Blanford, 1862; Stoliczka, 1863–1866; Wani & Ayyasami, 2009; Wani, Kurihara & Ayyasami, 2011). This difference may be attributed to growth halts or slowdowns in ammonoids, as opposed to nautiloids, which lack such features.

Given the repeated halts or slowdowns in shell growth during the early ontogeny and the large conch diameter of *P. denisonianum*, it is likely that their lifespan was longer than previously assumed. However, accurately estimating their life duration remains challenging at present.

Ontogenetic trajectory of septal spacing

The measured septal spacings of the early conch in *Pachydesmoceras denisonianum* reveal a nearly stable pattern with a slightly increasing trend (standard deviation = 1.93°) (Fig. 3B). The precise correlation between the timing of apertural and septal formation remains unclear. However, *Matsumoto (1988)* reported that the body chamber length of the lectotype is 240° . Based on this, the apertural position at each septum was estimated in the present study, assuming that the body chamber length remained constant at 240° throughout the ontogeny of the examined specimen (Fig. 3C; Table S1). Fig. 3C illustrates that, despite a decrease in constriction spacings (conch diameter ~ 70 mm), the septal spacing continues to exhibit a stable increasing trend. Therefore, the comparison of graphs for constriction and septal spacings (Fig. 3C) indicates no clear relationship between graphs for constriction and septal spacings, suggesting that constrictions and septal spacings are not directly linked in *Pachydesmoceras*. This lack of connection implies that growth halts or slowdowns were not recorded in the

ontogenetic trajectories of septal spacings, at least in *Pachydesmoceras*.

The ontogenetic trajectories of *Pachydesmoceras denisonianum* were compared with those of other ammonoids from the subfamilies Puzosiinae (e.g., *Puzosia* sp., from the Turonian of the Ariyalur area, southern India) and Desmocerotinae (e.g., *Desmoceras latidorsatum* var. *media*, from the Albian of the Mahajanga area, Madagascar), both within the family Desmocerotidae (Takai et al., 2022; Nishino et al., 2024; Fig. 5). Although the comparable conch diameters among the examined three taxa are limited, the observed ranges of septal spacings in *Pachydesmoceras* fall within those of *Desmoceras* (especially those with phragmocone diameters >1 mm). The slope of the slightly increasing trend in *Pachydesmoceras* is almost parallel to that of *Puzosia*. These suggest that the septal spacings of *Pachydesmoceras* share characteristics with both taxa. These trends in the Puzosiinae and Desmocerotinae are similar to those observed in other Cretaceous Perisphinctina but differ from those seen in Phylloceratina and Lytoceratina from the Cretaceous period (Arai & Wani, 2012; Iwasaki, Iwasaki & Wani, 2020; Kawakami, Uchida & Wani, 2022; Takai et al., 2022; Kawakami & Wani, 2023; Nishino et al., 2024).

Growth halts in ammonoids, which may be accompanied by constrictions, likely result from the covariance between isometric or allometric growth of the aperture and ornamentation (Bucher, 1997) and/or the balance between conch and soft part growth. The results of this work

indicate that when the growth balance between the conch and soft parts in *Pachydesmoceras* was disrupted, this imbalance may have been compensated not by changes in septal spacing, but by modifications in the apertural shape, leading to the formation of constrictions. Whether this phenomenon was common across ammonoids remains uncertain, and therefore, the relationship between the ontogenetic trajectories of constrictions and septa in various taxa across different geological ages warrants further investigation in future studies. This would provide a deeper understanding of how growth patterns and environmental factors influenced the morphological evolution of ammonoids over time.

ACKNOWLEDGEMENTS

I sincerely thank the Director General of the Geological Survey of India (Kolkata), K. Ayyasami, and S. Anantharaman for their kind help during fieldwork and fossil sampling in the Ariyalur area, southern India. Without their assistance, the fieldwork would not have been possible. I also thank A. Tajika and D. Aiba for their support during literature reviews and E. Yazykova and K. De Baets as well as the editor (V. Brygadyrenko) for their valuable and thoughtful comments on an earlier draft of the manuscript.

REFERENCES

- Arai K, Wani, R. 2012.** Variable growth modes in Late Cretaceous ammonoids: implications for diverse early life histories. *Journal of Paleontology* **86**: 258–267.
- Arkell WJ, Kummel B, Wright CW. 1957.** Mesozoic Ammonoidea. In: Moore RC, ed. *Treatise on Invertebrate Paleontology. Part L, Mollusca 4*. Lawrence: Geological Society of America and University of Kansas Press, 80-465.
- Besnosov NV, Mikhailova IA. 1991.** Higher taxa of Jurassic and Cretaceous Ammonitida. *Paleontological Journal* **25** (4): 1–19.
- Blanford HF. 1862.** On the Cretaceous and other rocks of the South Arcot and Trichinopoly districts, Madras. *Memoirs of the Geological Survey of India* **4**: 1–217.
- Bucher H. 1997.** Caractères périodiques et modes de croissance des ammonites: comparaison avec les gastéropodes. *Geobios* **20**: 85–99.
- Bucher H, Landman NH, Klokak SM, Guex J. 1996.** Mode and rate of growth in ammonoids. In: Landman NH, Tanabe K, Davis RA, eds. *Ammonoid Paleobiology*. New York: Plenum Press, 407–461.
- Collignon M. 1961.** Ammonites néocrétacées du Menabe (Madagascar). 7.– Les Desmoceratidae. *Annales Géologiques de Madagascar* **31**: 1–115.
- De Baets K, Landman NH, Tanabe K. 2015.** Ammonoid embryonic development. In: Klug C, Korn D, De Baets K, Kruta I, Mapes RH, eds. *Ammonoid Paleobiology, From*

Anatomy to Ecology. Amsterdam: Springer, 113–205.

Ifrim C, Stinnesbeck W, González González A H, Schorndorf N, Gale AS. 2021. Ontogeny, evolution and palaeogeographic distribution of the world’s largest ammonite *Parapuzosia (P.) seppenradensis* (Landois, 1895). *PLoS ONE* 16(11), e0258510.

Iwasaki T, Iwasaki Y, Wani R. 2020. Polymorphism in Late Cretaceous phylloceratid ammonoids: evidence from ontogenetic trajectories of septal spacing. *Papers in Palaeontology* 6: 155–172.

Kawakami Y, Uchida N, Wani R. 2022. Ontogenetic trajectories of septal spacing and conch shape in the Late Cretaceous gaudryceratid ammonoids: implications for their post-embryonic palaeoecology. *Palaeontology* 65: e12587.

Kawakami Y, Wani R. 2023. Stepwise growth changes in early post-embryonic stages among Cretaceous tetragonitid ammonoids. *Paläontologische Zeitschrift* 2023: <https://doi.org/10.1007/s12542-023-00650-0>.

Kennedy WJ, Cobban WA. 1976. Aspects of ammonite biology, biogeography, and biostratigraphy. *Special Papers in Palaeontology* 17: 1–94.

Kennedy WJ, Klinger HC. 2014. Cretaceous faunas from Zululand and Natal, South Africa. *Valdedorsella, Pseudohaploceras, Puzosia, Bhimautes, Pachydesmoceras, Parapuzosia (Austiniceras) and P. (Parapuzosia)* of the ammonite subfamily

Puzosiinae Spath, 1922. *African Natural History* **10**: 1–46.

Kin A, Niedźwiedzki R. 2012. First record of the puzosiine ammonite genus *Pachydesmoceras* from the Middle and Upper Turonian of Poland. *Cretaceous Research* **33**: 15–20.

Klug C, De Baets K, Kröger B, Bell MA, Korn D, Payne J L. 2015b. Normal giants? Temporal and latitudinal shifts of Palaeozoic marine invertebrate gigantism and global change. *Lethaia* **48**: 267–288.

Klug C, Korn D, Landman NH, Tanabe K, De Baets K, Naglik C. 2015a. Describing ammonoid conchs. In: Klug C, Korn D, De Baets K, Kruta I, Mapes RH, eds. *Ammonoid Paleobiology, From Anatomy to Ecology*. Amsterdam: Springer, 3–24.

Kulicki C. 1974. Remarks on the embryogeny and postembryonal development of ammonites. *Acta Palaeontologica Polonica* **19**: 201–224.

Maeda H. 1987. Taphonomy of ammonites from the Cretaceous Yezo Group in the Tappu area, northwestern Hokkaido, Japan. *Transactions and Proceedings of the Paleontological Society of Japan, New Series* **148**: 285–305.

Maeda H, Kumagae T, Matsuoka H, Yamazaki Y. 2010. Taphonomy of large *Canadoceras* (ammonoid) shells in the Upper Cretaceous Series in South Sakhalin, Russia. *Paleontological Research* **14**: 56–68.

Maeda H, Seilacher A. 1996. Ammonoid taphonomy. In: Landman NH, Tanabe K, Davis RA,

eds. *Ammonoid Paleobiology*. New York: Plenum Press, 543–578.

Matsumoto T. 1987. Note on *Pachydesmoceras*, a Cretaceous ammonite genus. *Proceedings of the Japan Academy, Series B* **63**: 5–8.

Matsumoto T. 1988. A monograph of the Puzosiidae (Ammonoidea) from the Cretaceous of Hokkaido. *Palaeontological Society of Japan, Special Papers* **30**: 1–179.

Nishino Y, Komazaki K, Arai M, Hattori A, Uoya Y, Iida T, Wani R. 2024. Covariable changes of septal spacing and conch shape during early ontogeny: a common characteristic between Perisphinctina and Ancyloceratina (Ammonoidea, Cephalopoda). *Journal of Paleontology* **98**: 102–114, <https://doi.org/10.1017/jpa.2023.96>.

Obata I, Futakami M, Kawashita Y, Takahashi T. 1978. Apertural features in some Cretaceous ammonites from Hokkaido. *Bulletin of the National Science Museum, Series C, Geology & Paleontology* **4**: 139–155.

Stoliczka F. 1863–1866. The fossil Cephalopoda of the Cretaceous rocks of southern India. Ammonoidea, with revision of the Nautilidae. *Memoirs of the Geological survey of India, Palaeontologia Indica* **3**: (1), 41–56 (1863); (2–5), 57–106 (1864); (6–9), 107–154 (1865); (10–13), 155–216 (1866).

Sundaram R, Henderson RA, Ayyasami K, Stilwekk JD. 2001. A lithostratigraphic revision

and palaeoenvironmental assessment of the Cretaceous System exposed in the onshore
Cauvery Basin, southern India. *Cretaceous Research* **22**: 743–762.

Sundaram R, Rao PS. 1986. Lithostratigraphy of Cretaceous and Palaeocene rocks of
Tiruchirappalli District, Tamil Nadu, south India. *Records of the Geological Survey of
India* **116**: 9–23.

Tajika A, Nützel A, Klug C. 2018. The old and the new plankton: ecological replacement of
associations of mollusc plankton and giant filter feeders after the Cretaceous? *PeerJ*
6:e4219; DOI 10.7717/peerj.4219.

Takai F, Matsukuma S, Hirose K, Yamazaki T, Aiba D, Wani R. 2022. Conservative
ontogenetic trajectories of septal spacing during the post-embryonic stage in Cretaceous
ammonoids of the subfamily Desmoceratinae. *Lethaia* **55**: 1–12. doi:
10.18261/let.55.2.2.

Urgy S. 2015. Theoretical Modelling of the Molluscan Shell: What has been Learned From the
Comparison Among Molluscan Taxa? In: Klug C, Korn D, De Baets K, Kruta I,
Mapes RH, eds. *Ammonoid Paleobiology, From Anatomy to Ecology*. Amsterdam:
Springer, 207–251.

Wani R, Ayyasami K. 2009. Ontogenetic change and intra-specific variation of shell
morphology in the Cretaceous nautiloid (Cephalopoda, Mollusca) *Eutrephoceras*

clementinum (d'Orbigny, 1840) from the Ariyalur area, southern India. *Journal of Paleontology* **83**: 365–378.

Wani R, Gupta NS. 2015. Ammonoid taphonomy. In: Klug C, Korn D, De Baets K, Kruta I, Mapes RH, eds. *Ammonoid Paleobiology, From macroevolution to paleogeography*. Amsterdam: Springer, 555–597.

Wani R, Kurihara K, Ayyasami K. 2011. Large hatchling size in Cretaceous nautiloids persists across the end-Cretaceous mass extinction: new data of Hercoglossidae hatchlings. *Cretaceous Research* **32**: 618–622.

Westermann GEG. 1990. New developments in ecology of Jurassic–Cretaceous ammonoids. In: Pallini G, Cecca F, Cresta S, Santantonio M, eds. *Atti del secondo convegno internazionale. Fossili, Evoluzione, Ambiente, Pergola, 1987*. Ostra Vetere, Italy: Tectnostampa, 459–478.

Wright CW. 1996. *Treatise on Invertebrate Paleontology part L, Mollusca 4 Revised. Volume 4*. Lawrence: Geological Society of America and University of Kansas Press.

Yabe H. 1914. Ein neuer Ammonitenfund aus der Trigonina Sandstein Gruppe von Provinz Tosa. *Science Reports of the Tohoku Imperial University, 2nd series* **1**, 71–74.

Yacobucci MM. 2015. Macroeolution and paleobiogeography of Jurassic-Cretaceous ammonoids. In: Klug C, Korn D, De Baets K, Kruta I, Mapes RH, eds. *Ammonoid*

Paleobiology, From Macroevolution to Paleogeography. Amsterdam: Springer, 189–
228.

Figure and table captions

Figure 1. Photographs of the examined specimen, MCM-W2145. (A) Left lateral view; (B, C) ventral views; (D) right lateral view. Stars are only shown in the left-side photograph, indicating the ventral positions of constrictions that can be recognized in the preserved outer whorl. Scale bar is 10 mm.

Figure 2. Cross section of the examined specimen. (A) Photograph of the median section of the examined specimen; (B, C) enlarged photographs of constrictions and ribs shown in black squares and its schematic drawings. The exterior is toward the top of the photos. Black arrow, the preserved smallest conch; white arrows, the ventral position of constrictions; star, the smallest position of constriction that can be recognized in the preserved outer whorl; circle, the center of the approximated logarithmic spiral.

Figure 3. Ontogenetic trajectories of constriction and septal spacing. (A) Graph of constriction spacing through early ontogeny; (B) graph of septal spacing through early ontogeny; (C) comparison between constriction and septal spacings. The apertural position at each septum was estimated under the assumption that the body chamber length remained 240° throughout

ontogeny, consistent with the measurements of the lectotype (likely an adult macroconch;
Matsumoto, 1988).

Figure 4. Measurements of *Pachydesmoceras denisonianum*. (A) Conch diameter/umbilicus
width relationship; (B) conch diameter/whorl height relationship; (C) conch diameter/whorl
width relationship; (D) umbilicus width/conch diameter ratio (U/D); (E) whorl width/whorl
height ratio (W/H); (F) whorl width/conch diameter ratio (W/D). Solid circle, measurements of
lectotype.

**Figure 5. Comparison of ontogenetic trajectories of septal spacings in the subfamilies
Puzosiinae and Desmoceratinae.** *Puzosia* and *Desmoceras* data are from *Nishino et al. (2024)*
and *Takai et al. (2022)*, respectively.

Table 1. Morphological data of *Pachydesmoceras denisonianum*. Measurements except the
examined specimen are from *Matsumoto (1988)*. D, conch diameter; d, conch diameter at 180°
adapically from D; U, umbilical width; H, whorl height; h, whorl height at 180° adapically from
H; W, whorl width; ah, aperture height.

Table S1. Raw data of *Pachydesmoceras denisonianum* conch morphology.

Figure 1

Figure 1. Photographs of the examined specimen , MCM-W2145.

(A) Left lateral view; (B, C) ventral views; (D) right lateral view. Stars are only shown in the left-side photograph, indicating the ventral positions of constrictions that can be recognized in the preserved outer whorl. Scale bar is 10 mm.



Figure 2

Figure 2. Cross section of the examined specimen.

(A) Photograph of the median section of the examined specimen; (B, C) enlarged photographs of constrictions and ribs shown in black squares and its schematic drawings. The exterior is toward the top of the photos. Black arrow, the preserved smallest conch; white arrows, the ventral position of constrictions; star, the smallest position of constriction that can be recognized in the preserved outer whorl ; circle, the center of the approximated logarithmic spiral .




Figure 3

Figure 3. Ontogenetic trajectories of constriction and septal spacing.

(A) Graph of constriction spacing through early ontogeny; (B) graph of septal spacing through early ontogeny ; (C) comparison between constriction and septal spacings. The apertural position at each septum was estimated under the assumption that the body chamber length remained 240° throughout ontogeny, consistent with the measurements of the lectotype (likely an adult macroconch; *Matsumoto, 1988*).




Figure 4

Measurements of *Pachydesmoceras denisonianum* .

(A) Conch diameter/umbilicus width relationship; (B) conch diameter/whorl height relationship; (C) conch diameter/whorl width relationship; (D) umbilicus width/conch diameter ratio (U/D); (E) whorl width/whorl height ratio (W/H); (F) whorl width/conch diameter ratio (W/D). Solid circle, measurements of lectotype.




Figure 5

Comparison of ontogenetic trajectories of septal spacings in the subfamilies Puzosiinae and Desmocerotinae.

Puzosia and *Desmoceras* data are from *Nishino et al. (2024)* and *Takai et al. (2022)* , respectively.




Table 1 (on next page)

Table 1. Morphological data of *Pachydesmoceras denisonianum*.

Measurements except the examined specimen are from *Matsumoto (1988)*. D, conch diameter; d, conch diameter at 180° adapically from D; U, umbilical width; H, whorl height; h, whorl height at 180° adapically from H; W, whorl width; ah, aperture height.

specimen	D (mm)	d (mm)	U (mm)	H (mm)	h (mm)	W (mm)	ah (mm)	U/D	U/H	W/D	W/H	H/h	whorl expansion rate (WER; $(D/d)^2$)	imprint zone rate (IZR; $(H-ah)/H$)	reference
Lectotype, GSI. 208	995		410	345	208	300		0.41	1.19	0.30	0.87	1.44			Stoliczka (1865)
GSJ. F3469	460		156	179	136	195		0.34	0.87	0.42	1.09	1.43			Matsumoto (1988)
Yabe, 1914, pl. 12	446		142	188		160		0.32	0.76	0.36	0.85				Yabe (1914)
YKC. 610612	365		120	143	131	183		0.33	0.84	0.50	0.93	1.40			Matsumoto (1988)
MNHN. 3750	166		52	67	45	65		0.31	0.78	0.39	0.97	1.43			Collignon (1961)
MCM-W2145	109	79	35	45	31	41	32	0.32	0.78	0.38	0.91	1.45	1.90	0.29	this study