



Morphological, phytochemical, and pharmacological properties of the genus *Tamarix* in Kazakhstan species: a review

Arailym Daulbayeva¹, Gulnara Kadyrbayeva¹, Kaldanay Kozhanova¹, Shazlin Shaharudin², Nurgali Rakhymbayev¹, Zoya Allambergenova¹, Rabiga Anarbayeva³, Urziya Alimova¹, Aigerim Kantureyeva¹, Ainash Baidullaeva¹, Vladimir An¹ and Bakkonat Kydyrbai¹

¹ School of Pharmacy, S.D. Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan

² School of Health Sciences, Universiti Sains Malaysia, Kota Bharu, Malaysia

³ Department of Medicine Technology and Pharmacognosy, South Kazakhstan Medical Academy, Shymkent, Kazakhstan

ABSTRACT

The tree-like plants of the genus *Tamarix* belong to the Tamaricaceae family and predominantly grow in subtropical areas, steppes, and saline soils. These plants exhibit various pharmacological properties, including antioxidant, antidiabetic, anti-inflammatory, and antimicrobial characteristics. The current flora of Kazakhstan also includes 13 *Tamarix* species. Despite the diversity of the *Tamarix* taxon, the species discovered in the region have received limited investigation. Therefore, this review analyzed Kazakhstan and international-related scientific studies concerning the morphological, phytochemical, and pharmacological properties of 13 species within the *Tamarix* genus. The plants' biological features and potential applications were also comprehensively analyzed. Consequently, this review significantly contributed to botanical science and practical pharmacology. Conservation and sustainable employment of the *Tamarix* species for medical purposes could also be developed.

Submitted 21 February 2025
Accepted 19 August 2025
Published 30 September 2025

Corresponding authors
Gulnara Kadyrbayeva,
chilnara_k@mail.ru
Shazlin Shaharudin, shazlin@usm.my

Academic editor
Mansour Sobeh

Additional Information and
Declarations can be found on
page 21

DOI 10.7717/peerj.20059

© Copyright
2025 Daulbayeva et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Agricultural Science, Plant Science, Taxonomy, Pharmacology

Keywords Antioxidant, Antidiabetic, Anti-inflammatory, Antimicrobial, Taxonomy

INTRODUCTION

Tamarix L. is a genus of deciduous shrubs and small trees belonging to the family *Tamaricaceae*, widely distributed across arid regions of Eurasia and Africa (Bencherif *et al.*, 2020). These plants typically inhabit saline soils, river floodplains, and desert environments, exhibiting high salt tolerance and a remarkable ability to stabilize sandy substrates.

The taxonomy of the genus *Tamarix* is considered highly complex, as its species are morphologically similar, which complicates accurate identification and leads to discrepancies among taxonomists. Historically, up to 200 species and intraspecific taxa of *Tamarix* have been described. However, a revision by B. Baum (1978) reduced this number to 54 species and 15 infraspecific forms (Venturella, Baum & Mandracchia, 2007). More recent data have clarified the taxonomic structure of the genus: according to the online resource *Plants of the World Online* (Royal Botanic Gardens Kew, 2024), the genus

currently includes 73 accepted species. Such discrepancies are primarily due to a high number of synonyms and varying interpretations of species boundaries by different authors. For instance, the Russian digital flora reference *Plantarium* considers *Tamarix karelinii* Bunge to be a hybrid (*T. hispida* Willd. \times *T. ramosissima* Ledeb.) ([Plantarium, 2025](#)), whereas *Royal Botanic Gardens Kew* (2024) recognizes *T. karelinii* as a distinct species. This highlights the importance of utilizing up-to-date taxonomic databases and incorporating regional floristic sources when clarifying species nomenclature.

In Kazakhstan, *Tamarix* species are predominantly found in desert and semi-desert regions, including the Caspian Lowland, the Aral Sea basin, the Moyynkum and Balkhash deserts, and the river valleys of southern regions. Among them, *T. ramosissima*, *T. hispida*, and *T. laxa* are the most widespread and ecologically significant species ([Ile-Balkhash, 2025](#); [The Ferns Database, 2025](#); [Breckle, Wucherer & Dimeyeva, 2012](#)).

Notably, the ecological, medicinal, and economic aspects of the *Tamarix* genus underscore its significance. The *Tamarix* species have also demonstrated resilience in extreme conditions, including high soil salinity and drought. This observation suggests that the plants are valuable subjects for soil stabilization and erosion prevention-related studies. Furthermore, the properties of this species are highly suitable for ecological projects aimed at combating desertification and restoring lands surrounding the Aral Sea ([Ashirbekov, 2013](#); [Daryo.uz, 2024](#)).

Tamarix species have long attracted scientific interest due to their ecological adaptability and rich phytochemical profile. These plants thrive in arid and saline environments and play an essential role in stabilizing desert ecosystems. Their ability to produce a wide range of secondary metabolites, including polyphenols, flavonoids, and tannins, underlies their demonstrated antioxidant, anti-inflammatory, and antimicrobial activities ([Bikbulatova & Korul'kina, 2001](#); [Baktybayeva et al., 2019](#)).

Taxonomically, the genus *Tamarix* is considered complex due to overlapping morphological traits and frequent hybridization among species, which complicates accurate species identification and classification ([Sheidai et al., 2018](#)). This complexity is especially relevant in Kazakhstan, where several *Tamarix* species share similar habitats and morphological characteristics.

In Kazakhstan, *Tamarix* plants are also valued for their medicinal applications. Local populations use the bark, leaves, and flowers of *T. ramosissima* and related species to treat digestive disorders, hematological conditions, and infections. Ethnomedical practices across Central Asia, India, and the Middle East describe *Tamarix*-based remedies for ailments such as jaundice, tuberculosis, menstrual irregularities, and various skin conditions ([Ibn Sino, 1996](#); [Amirdovlat, 1990](#); [Zakhidov, 1991](#); [Gemedzhieva, Sultanova & Abilov, 2013](#)). These uses highlight the ethnopharmacological significance of the genus in traditional medicine systems.

Studies on the chemical composition and bioactivity of *Tamarix* species in Kazakhstan have confirmed the presence of valuable bioactive compounds, yet few have combined these data with taxonomic assessments ([Abilov & Sultanova, 2018](#)). One of the earliest systematic efforts to classify *Tamarix* was made by Joseph Franz Frein in 1903, laying the foundation for future botanical studies on the genus ([Villar et al., 2014](#)). Consequently,

scientists have advanced their complex structure comprehension of the genus arising from the development of molecular phylogenetics.

Various sources and research methodologies reported differing numbers of plant species in the *Tamarix* genus. Villar, Alonso & Crespo (2022) suggested a taxonomic revision of the genus *Tamarix* into seven species, of which three were native to the Iberian Peninsula and the Balearic Islands. This outcome updated the data of Flora Iberica published in 1993. Sheidai et al. (2018) revealed that the genus *Tamarix* comprised 54 and 90 species, with Asia and the eastern Mediterranean as the primary speciation centers. Akhiani et al. (2019) documented a new species (*T. humboldtiana*) in southern Iran. The information confirmed the existence of various species in the region. Several studies also noted approximately 70 species of *Tamarix* that adapted to hot and dry environments (Krylenko, Krylenko & Krylenko, 2020).

Pollen investigations of various *Tamarix* species in Iran revealed diagnostic signs that aided taxonomic identification (Shagholi, Keshavarzi & Sheidai, 2020). This finding presented significant diversity, which complicated species separation. The palynological characteristics of the genus *Tamarix* in Israel were also described in “Pollen Morphology of the Genus *Tamarix* in Israel”, complicating taxonomic clarification of the genus based on palynological attributes (Eshel et al., 2019).

Molecular approaches are vital in distinguishing the taxonomy and phylogeny of the *Tamarix* genus, such as inter-simple sequence repeat (ISSR), random amplified polymorphic (RAPD), deoxyribonucleic acid (DNA), internal transcribed spacer (ITS) sequences, single nucleotide polymorphism (SNP) markers, DNA barcoding, and microsatellites. Hence, incorporating genetic markers and DNA analysis can improve three main processes: (i) accurate species categorization, (ii) hybridization identification, and (iii) practical examination of evolutionary relationships within the genus (Sheidai et al., 2018; Arianmanesh et al., 2015; Lee, Gaskin & Kim, 2019; Terrones et al., 2022). Comprehensive chloroplast genome-related studies can also further expand the understanding of the taxonomic correlations among the species (Wang, Cao & Wei, 2023).

Previous reviews on the genus *Tamarix*, such as those by Bahramsoltani et al. (2020) and Li et al. (2024), have primarily focused on the traditional uses, phytochemical composition, and pharmacological activity of *Tamarix* species at the global level. In contrast, the present review is centered on 13 *Tamarix* species native to Kazakhstan. A comparative analysis of their taxonomic status is presented based on both the international database *Plants of the World Online* (POWO) and regional data relevant to Kazakhstan and Central Asia. The review also integrates ethnopharmacological information on the use of *Tamarix* in Kazakhstan and addresses ecological aspects related to their practical application, including afforestation practices in the Aralkum region.

Therefore, this review aims to summarize the pharmacological potential of these species by examining their morphological characteristics, geographic distribution, phytochemical composition, and known pharmacological properties.

Thus, this study represents an interdisciplinary review that combines taxonomic, ecological, phytochemical, and pharmacological data on *Tamarix* species found in

Kazakhstan. It expands the existing body of knowledge and provides a scientific basis for the sustainable use and conservation of these species.

SURVEY METHODOLOGY

This review conducted a literature search across four scientific databases: (i) Web of Science, (ii) Scopus, (iii) SciFinder, and (iv) PubMed. The keywords employed were “*Tamarix*”, “*Tamarix* morphology”, “*Tamarix* phytochemical compounds”, “*Tamarix* pharmacological properties”, and “*Tamarix* habitats”. The search covered publications from 1951 to 2025, and only peer-reviewed articles and academic sources in English, Russian, and Kazakh were considered.

Only studies focusing on the morphological, phytochemical, and pharmacological properties of *Tamarix* L. plants in the flora of Kazakhstan were included. Additionally, the review incorporated data from authoritative regional sources, such as the *Flora of Kazakhstan*, academic textbooks, and the Plantarium website (<https://www.plantarium.ru>), which provides high-quality taxonomic and ethnobotanical information.

Morphological characteristics of plants in the genus *Tamarix*

The genus *Tamarix* encompasses multiple species exhibiting significant morphological variations. These plants in the genus have adapted to moist and frequently saline soils found in arid and semi-arid climates. Generally, *Tamarix* plants manifest as shrubs or trees characterized by leaves that have evolved into scales. The flowers of these plants are collected in dense inflorescences measuring between five cm and 10 cm, blooming during spring, summer, and fall. *Tamarix* seeds also possess plumose tips, facilitating dispersal through wind and water (Akhami et al., 2019; Eshel et al., 2019).

Several characteristic features of *Tamarix* have been demonstrated in its morphological data, such as leaf shape with arrangement, floral organ number with structure, and pollen grain attributes (Sheidai et al., 2018; Eshel et al., 2019). Although a considerable degree of hybridization and phenotypic plasticity complicating identification can occur, this information has enabled species categorization (Arianmanesh et al., 2016; Souddi & Bendi Djelloul, 2020). Numerous species in the genus *Tamarix* also possess synonyms that complicate their systematics, necessitating reevaluation during identification and classification (Villar, Alonso & Crespo, 2022; Gorshkova, 1949; Zieliński, 1994; Assadi, Ramak Maassoumi & Khatamsaz, 1989). Table 1 lists the morphological attributes of the 13 *Tamarix* species (two hybrids) native to Kazakhstan. These species include *T. laxa* Willd., *T. meyeri* Boiss., *T. elongata* Ledeb., *T. passerinoides* Delile., *T. hispida* Willd., *T. leptostachys* Bunge, *T. arceuthoides* Bunge, *T. hohenackeri* Bunge, *T. gracilis* Willd., *T. ramosissima* Ledeb., *T. litwinowii* Gorschk., *T. karelinii* Bunge, hybrid *T. hispida* Willd. × *T. ramosissima* Ledeb., *T. eversmannii* Presl., hybrid *T. leptostachys* Bunge × *T. ramosissima* Ledeb.

Geographical distribution

The plants within the *Tamarix* genus are correlated through a complex taxonomic network. These plants are among the most studied and widespread genera worldwide. *Tamarix*

Table 1 Morphological feature summary of *Tamarix* plants in Kazakhstan.

Species	Morphology	Reference
1. <i>Tamarix litwinowii</i> Gorschek	<p>Shrub height up to 3 metres.</p> <p>Bark matt, light brown or grey, smooth.</p> <p>Leaves: Leaves of growing shoots: ovoid-elongated, 1–2 mm, acuminate, descending. The leaves of the annual branches are broad, long, acuminate, slightly descending, and protruding.</p> <p>Flowers: 4-membered during spring flowering, sometimes 5-membered during autumn flowering, small, less than 5 mm in diameter. It usually blooms twice a year.</p> <p>Capsule: narrow, 3 mm long, 1 mm wide, approximately 3–4 times longer than the calyx.</p> <p>The anthers are rather blunt.</p>	<p><i>Pavlov (1963); Kostina, Barabanshchikova & Pavlova (2020); Flora of Turkmenistan (1951)</i></p>
2. <i>Tamarix laxa</i> Willd.	<p>Branched bush, 2–3 metres high, large, branches spreading, glabrous, green and bluish. Phanerophyte.</p> <p>Bark: old branches grey; young branches short, reddish-brown, grey-cherry, or grey, brittle.</p> <p>Leaves are erect, up to 2 mm, ovate, acuminate at the apex, narrowed to the base, yellowish-green; Racemes are 1–3 cm long, 4–7 mm wide, sparse, almost sessile, and appear early in spring on 1-year-old branches.</p> <p>Flowers are pink, in dense racemes (4x1 cm), in turn collected in apical panicles on short peduncles. 4-membered during spring flowering and 5-membered during autumn.</p> <p>Capsule: narrow, 3–4 mm long, about 1 mm wide, 5–6 times longer than the calyx.</p> <p>The anthers are heart-shaped, slightly acuminate or obtuse, dark purplish.</p>	<p><i>Pavlov (1963); Flora of Turkmenistan (1951); Afanasyev (2012); Chinese Plant Names (2024); Stankov & Taliev (1949); Baum (1978); Golub Valentin (2008); Shang & Lowry II (2007); Chepinoga et al. (2024); Fungi.su (2024)</i></p>
3. <i>Tamarix meyeri</i> Boiss.	<p>Shrub to 2–4 (8) metres tall, green or bluish-glabrous, with grey, reddish-brown, or brownish-grey bark.</p> <p>Bark: old trunks brownish-grey, bark of year-old branches purplish, yellowish, or greenish.</p> <p>Leaves: annual branches lanceolate, 3–4 mm long with a semi-stemmed base, brown, obtuse, or slightly acuminate, slightly curved inward, slightly broadened at the base, strongly descending.</p> <p>Flowers: 4-membered; bracts narrow-long; lower: blunted with a short truncated blunt end; upper: slightly acuminate.</p> <p>Capsule: 5–7 mm long, twice as long as the calyx.</p> <p>Pollen grains preprolate, acute.</p>	<p><i>Souddi & Bendi Djelloul (2020); Pavlov (1963); Flora of Turkmenistan (1951); Stankov & Taliev (1949); Baum (1978); Venturella, Gargano & Mandracchia (2012)</i></p>

(continued on next page)

Table 1 (continued)

Species	Morphology	Reference
4. <i>Tamarix elongata</i> Ledeb.	<p>Shrub height 1–3 metres. shrub with blackish brown bare bark.</p> <p>The bark of old trunks is grey; that of the year-old branches is greyish-yellow, pale, or reddish-brown.</p> <p>Leaves in the lower part of branches large, broadly linear, lanceolate or oblong-lanceolate, 4 to 9 mm long, 1–3 mm wide, acute, first leaves broad, somewhat heart-shaped, thick or fleshy.</p> <p>Flowers are 4-membered; calyx with triangular or ovate, acuminate; membranous lobes along the edge; corolla is large.</p> <p>Capsule: 4–6 mm long, 2.5–4 times longer than the calyx.</p> <p>Anthers: reddish or orange-yellow, ovoid-conical capsule.</p>	<p>Pavlov (1963); Stankov & Taliev (1949); Baum (1978); Shang & Lowry II (2007)</p>
5. <i>Tamarix passerinoides</i> Delile. Synonyms: <i>Tamarix aucheriana</i> (Decne. ex Walp.) B.R. Baum, <i>Tamarix macrocarpa</i> (Ehrenb.) Bunge	<p>Shrub height 1-2 metres, low spreading tree with bark from reddish-brown to dark purple, young parts are pubescent.</p> <p>The bark of old branches is dark brown-grey, with a purple-grayish tint.</p> <p>Leaves: strongly auricular, bifurcated, 1.5 to 2.5 mm long, somewhat ovate with spreading tip, 1-2.5 mm long (on old branches 3-3.5 mm long), 1-1.5 mm wide with punctate glands.</p> <p>Flowers are five-lobed, pinkish to purple-pink. Inflorescences are simple or loosely folded, usually at the beginning of flowering. The racemes are 2 to 5 cm long and 8 to 10 mm wide.</p> <p>Capsule: large, 10–12 mm long, with three leathery valves.</p> <p>The apical heart-shaped anthers are reddish coloured and acuminate. The pollen grains were prolate. Tectum: Medium reticulate.</p>	<p>Shagholi, Keshavarzi & Sheidai (2020); Arianmanesh et al. (2016); Gorshkova (1949); Zieliński (1994); Assadi, Ramak Maassoumi & Khatamsaz (1989); Pavlov (1963); Flora of Turkmenistan (1951); Baum (1978); Qaiser (1983); De Martis, Loi & Polo (1984); Elkordy & Faried (2017); Qaiser & Perveen (2004)</p>
6. <i>Tamarix gracilis</i> Willd. Synonyms: <i>Tamarix affinis</i> Bunge, <i>Tamarix polystachya</i> Ledeb.	<p>Shrub height of 1.5–4 metres. Shrub tree with brown to blackish-brown bark, young parts completely glabrous or with papillae.</p> <p>The bark is greyish-green or brownish-chestnut, with grey-pale cork spots on the leaf axils and along the stem.</p> <p>The leaves are oval-lanceolate and acute. Flowering racemes are thin, 2.5 to 5 cm long, sparse, and on bare stalks. sessile, with a narrow base, much longer than wide, 1–3 mm long.</p> <p>Capsule: 4–6(8) mm long, nearly four times longer than the calyx.</p> <p>Spring flowers are 4-membered; summer flowers are 4–5-membered, up to 5 mm in diameter; calyx lobes are ovoid, green, with a serrated translucent edge, obtuse, rarely acuminate, and without a keel.</p> <p>Pollen: unit of distribution: monads. Size: medium. Sculpture: mesh. Number of apertures: 3. Aperture position: equatorial. Aperture type: slit.</p>	<p>Shagholi, Keshavarzi & Sheidai (2020); Gorshkova (1949); Zieliński (1994); Assadi, Ramak Maassoumi & Khatamsaz (1989); Pavlov (1963); Stankov & Taliev (1949); Baum (1978); Shang & Lowry II (2007)</p>

(continued on next page)

Table 1 (continued)

Species	Morphology	Reference
7. <i>Tamarix hispida</i> Willd.	<p>Shrub height of 1.5–5 metres. A small tree with reddish brown bark, usually very hairy to almost bare, at least with papillae-like brush top bark of old branches and trunks reddish-grey; annual branches and growing shoots with reddish or ochre-grey bark densely covered with short trichomes.</p> <p>Leaves are sessile, cordate-cylindrical at the base, 1.5–2.25 mm long. Leaves of annual branches narrowly lanceolate, acuminate, leaves of green shoots ovate-lanceolate, 0.52 mm long, acute, with a point curved inward, semiclasping, pubescent.</p> <p>Flowers: 5-membered: calyx 1–1.3 mm long, with broad, obtuse, or pointed lobes, without keel, with a broad, transparent, serrated edge.</p> <p>Capsule: 4–5 mm long, 4–5 times longer than the calyx.</p> <p>The anthers are rather blunt.</p>	<p>Arianmanesh et al. (2016); Pavlov (1963); Flora of Turkmenistan (1951); Stankov & Taliev (1949); Baum (1978); Shang & Lowry II (2007)</p>
8. <i>Tamarix leptostachys</i> Bunge	<p>Shrub height of 1–5 metres shrub or very small tree, tall, with brown bark; young parts glabrous, sometimes with sparse fibrous-papillate leaves. The bark of the old branches is reddish-grey; annuals are grey-purple or reddish.</p> <p>Leaves are sessile with a narrow base, 2–3 mm long. The younger leaves are somewhat ribbed, 2–4 mm long, 0.5–2 mm wide, and acute.</p> <p>The flowers are five-petalled and pink. The petals are almost free, ovate or tubular-ovate, acute to obtuse, finely serrated along the edge, 0.5–0.75 mm long, 0.25–0.4 mm wide.</p> <p>Capsule: elongate-pyramidal, about 1.8 mm long, twice as long as the calyx.</p> <p>Pollen grains to prolate. Tectum: Finely reticulate.</p> <p>Pollen grains with columellae</p>	<p>Pavlov (1963); Baum (1978); Shang & Lowry II (2007); Qaiser (1983); Qaiser & Perveen (2004); Plantarium plant Determinant Online (2024)</p>
9. <i>Tamarix arceuthoides</i> Bunge	<p>Shrub height of 1–4 metres. Very small tree, 2–3 (-4) m tall, with reddish-brownish-pinkish bark, densely branched and glabrous.</p> <p>The bark of the trunks and old branches is grey, the bark of young woody ones is reddish-brown, dark purplish, or brownish-brown, and the annual shoots are emerald green, rarely syzy-green, glabrous, thin, straight and ascending.</p> <p>The leaves are sessile with a narrow base, usually with rough papillate edges, 1–2.5 mm long. Racemes are 1.5 to 5 cm long and 3 to 4 mm wide.</p> <p>Flowers 5-membered; calyx 0.5 to 0.7 mm long; bracts as long as the peduncle or slightly longer; entirely herbaceous; from linear to triangular to narrowly ovate; acuminate at the apex.</p> <p>Capsule: 4–6 times longer than the calyx lobes appressed to the capsule.</p> <p>The pollen grains were prolate. Tectum: Finely reticulate.</p> <p>The anthers are cordate, yellowish.</p>	<p>Akhani et al. (2019); Shaghali, Keshavarzi & Sheidai (2020); Arianmanesh et al. (2016); Pavlov (1963); Baum (1978); Shang & Lowry II (2007); Qaiser (1983); Qaiser & Perveen (2004); Plantarium plant Determinant Online (2024); Zhang et al. (2018); Ovchinnikov (1981)</p>

(continued on next page)

Table 1 (continued)

Species	Morphology	Reference
<p>10. <i>Tamarix hohenackeri</i> Bunge. Synonyms: <i>Tamarix rosea</i> Bunge <i>Tamarix smyrnensis</i> Auct.</p>	<p>Shrub height 1-3 m. Bushes with cranberry bark and linear, blunt, slightly curved.</p> <p>The bark of the old branches is greyish-brown; bark of the year old shoots is dark purple.</p> <p>Leaves linear or ovate-lanceolate, 1 to 5 mm long, long acuminate, acuminate, pointed, with the tip curved inward, semi-stemmed, without keel; racemes lateral, 1 to 8 cm long.</p> <p>Flowers are 4-membered, peduncles are long, calyxes, and flower brushes are yellowish elongated, 4 to 10 cm long and 5 to 6 mm wide, dense, and slightly widened at the base.</p> <p>Capsule: 4–5 mm long, three times shorter than the ovary.</p> <p>The anthers are blunt.</p>	<p>Gorshkova (1949); Zieliński (1994); Assadi, Ramak Maassoumi & Khatamsaz (1989); Pavlov (1963); Kostina, Barabanshchikova & Pavlova (2020); <i>Flora of Turkmenistan</i> (1951); Stankov & Taliev (1949); Shang & Lowry II (2007).</p>
<p>11. <i>Tamarix ramosissima</i> Ledeb. Synonyms: <i>Tamarix altaica</i> Nied. <i>Tamarix odessana</i> Steven ex Bunge <i>Tamarix pallasii</i> Desv. <i>Tamarix pentandra</i> Pall.</p>	<p>Shrub height 1–5 m. Shrub or low-growing tree. Phanerophyte, hyperhalophyte</p> <p>The cork of old trunks and branches is dark grey; the bark of year old shoots is reddish or orange-reddish.</p> <p>Leaves with small sessile green or glaucous scale-shaped leaves that secrete salt on their surface.</p> <p>The flowers are small, with a calyx and a pink corolla of four to five petals. Flowers are collected in large clusters. The bracts are equal to the pedicel.</p> <p>Capsule: trigonous-pyramidal, 3–5 mm long, 3–4 times longer than the calyx.</p> <p>Pollen grains to subprolate, microreticulate.</p> <p>The anthers are rather blunt.</p>	<p>Akhani et al. (2019); Shagholi, Keshavarzi & Sheidai (2020); Arianmanesh et al. (2016); Gorshkova (1949); Zieliński (1994); Assadi, Ramak Maassoumi & Khatamsaz (1989); Pavlov (1963); Kostina, Barabanshchikova & Pavlova (2020); <i>Flora of Turkmenistan</i> (1951); Afanasyev (2012); Shang & Lowry II (2007); Qaiser (1983); Qaiser & Perveen (2004); Zhang et al. (2018); Heywood et al. (2007); Crins (1989); Gaskin & Schaal (2003); Gaskin et al. (2004); Britannica (2019).</p>
<p>12. <i>Tamarix karelinii</i> Bunge Hybrid <i>Tamarix hispida</i> Willd. × <i>Tamarix ramosissima</i> Ledeb. Synonyms: <i>Tamarix hispida</i> var. <i>karelinii</i> (Bunge) B. R. Baum</p>	<p>Shrub height 2–4 m.</p> <p>The bark of the trunks is grey; bark of annual branches are grey-purple; branches glabrous, rarely slightly rough.</p> <p>The leaves are simple, sessile, ovoid, covered with salt glands, 1–2 mm long, 0.5–1 mm wide, acute.</p> <p>Flowers are pink to dark pink, five-lobed, open or semi-open, almost sessile; the calyx is approximately 1 mm long, with rounded, obtuse, membranous lobes along the edge.</p> <p>Capsule: 5–6 mm long, 5–6 times longer than the calyx.</p> <p>Pollen grains to subprolate. Tectum: Medium -coarse reticulate. Anthers with short pointed tips.</p>	<p>Gorshkova (1949); Zieliński (1994); Assadi, Ramak Maassoumi & Khatamsaz (1989); Pavlov (1963); Kostina, Barabanshchikova & Pavlova (2020); Golub Valentin (2008); Qaiser (1983); Qaiser & Perveen (2004); <i>Plantarium plant Determinant Online</i> (2024)</p>
<p>13. <i>Tamarix eversmannii</i> Presl. Hybrid <i>Tamarix leptostachys</i> Bunge × <i>Tamarix ramosissima</i> Ledeb.</p>	<p>Shrub height 2–3 m.</p> <p>The bark of old trunks is reddish-grey; the bark of one-year-old shoots is reddish-brown.</p> <p>Leaves are ovate-lanceolate or ovate-oblong, 1.5–4 mm long, rounded at the base, descending, acute at the apex, usually bent inward; racemes apical, 1–3(4) cm long, 2–3 mm wide, sparse, in a sparse, spreading panicle.</p> <p>Flowers are 5-membered; calyx 1 mm long with triangular-ovate, pointed membranous lobes at the edge.</p> <p>Pollen grains are pointed; styles 3(4), twice as short as the ovary.</p>	<p>Pavlov (1963)</p>

species can also adapt to different ecological conditions, and these plants can thrive in the arid and semi-arid regions of Eurasia. One considerable example involves the primary centers of speciation for the genus *Tamarix* in Central Asia (Kazakhstan, Afghanistan, Iran, and Turkmenistan) (Sheidai et al., 2018). These species have also adapted to saline soils and wet conditions along rivers and oases across Middle Eastern regions (Iran, Iraq, and Saudi Arabia). Similarly, eastern European *Tamarix* plants grow along rivers and in salt marshes within Ukraine and southern Russia. These species of the genus *Tamarix* are also discovered along the coasts, rivers, and lakes of North Africa, indicating high adaptability to arid environments and saline soils (Terzoli et al., 2013; Zhang et al., 2003).

Tamarix species (such as *T. ramosissima* and *T. chinensis*) were initially introduced to North America in the 19th century. Nevertheless, these plants exhibit invasive characteristics in the southwestern states of the United States (Arizona, California, Nevada, New Mexico, and Texas). The species thrives along rivers and reservoirs, modifying ecosystems and competing with native flora (Abby Brownell, 2013; iNaturalist contributors, 2024e). In Kazakhstan, *Tamarix* species play a significant role in desert and semi-desert ecosystems. They are primarily distributed in the Caspian Lowland, the Aral Sea region, the Moyynkum and Balkhash deserts, and the river valleys of southern Kazakhstan, where they form dense thickets in floodplain forests (tugai) and on saline soils. The most common and widely distributed species are *T. ramosissima* Ledeb., *T. hispida* Willd., and *T. laxa* Willd. (Ile-Balkhash, 2025). These three taxa constitute the dominant component of tamarisk thickets across much of their range, from the western deserts to the southeastern regions of the country.


In contrast, other species have more localized distributions. For example, *T. arceuthoides* Bunge is known from southwestern Kazakhstan (Caspian region) (The Ferns Database, 2025), while *T. aralensis* Bunge is primarily confined to the Aral Sea basin. Such narrowly distributed species, often restricted to drying lakebeds or specific desert zones, may be considered potentially vulnerable under changing environmental conditions.

Regional shifts in *Tamarix* species dominance have also been observed. In the desiccated zones of the Aral Sea, *T. laxa* and *T. elongata* dominate older desiccation sites (since the 1960s), *T. hispida* prevails in areas desiccated in the 1980s, and *T. ramosissima* is more common in river deltas (Breckle, Wucherer & Dimeyeva, 2012). This species in particular demonstrates a remarkable capacity to adapt to arid conditions and saline soils (Sheidai et al., 2018).

Figure 1 illustrates the habitat map of the *Tamarix* species in Kazakhstan. Table 2 presents the global and regional distribution of *Tamarix* species in Kazakhstan, structured according to the floristic zoning system proposed by Rachkovskaya, Khramtsov & Volkova (2003).

Phytochemical compounds

The compositions of the varying *Tamarix* species have been significantly investigated. One prominent example includes biologically active compounds in specific species, such as *T. hohenackeri* Bunge and *T. ramosissima* Ledeb. Consequently, all assessed species have shown the presence of flavonoids, phenolic acids, and triterpenoids. Certain species



Source: Ian Macky PAT maps, Public Domain (https://ian.macky.net/pat/map/kz/kz_blu.gif)

Figure 1 Geographical distribution of *Tamarix* species in Kazakhstan.

Full-size DOI: 10.7717/peerj.20059/fig-1

have also revealed other classes of phytochemical compounds, including coumarins, anthraquinones, tannins, tetraterpenes, and steroids (Bikbulatova & Korul'kina, 2001; Xing et al., 2014; Bahramsoltani et al., 2020; Umbetova et al., 2004a; Sultanova et al., 2002; Sultanova et al., 2004b; Utkin, 1966; Ren et al., 2019; Hong et al., 2018; Sultanova et al., 2001; Umbetova et al., 2005; Zhang & Tu, 1994; Sultanova et al., 2004a; Orabi et al., 2013; Xiao et al., 2013; Umbetova et al., 2004b; Sultanova et al., 2013; Yao et al., 2017). This data suggests that the genus *Tamarix* is rich in phytochemical compounds and exhibits various biological activities (antioxidant, anti-inflammatory, and antimicrobial properties). Nevertheless, identifying the compounds necessitates assessing various plant parts using diverse extraction methods to enhance the yield of particular phytochemical compounds. The following subsections delineate the primary chemical compounds in these plants.

Flavonoids

Various *Tamarix* species in Kazakhstan and China have presented flavonoids and their derivatives, such as *T. hispida* Willd., *T. hohenackeri* Bunge, *T. laxa* Willd., *T. ramosissima* Ledeb., and *T. elongata* Ledeb. The primary flavonoids indentified in this process include quercetin, kaempferol with its derivatives, and flavonoids associated with the genus *Tamarix* (tamarixetin and its derivatives) (Bikbulatova & Korul'kina, 2001; Xing et al., 2014; Umbetova et al., 2004a; Sultanova et al., 2002; Sultanova et al., 2004b; Utkin, 1966; Ren et al., 2019; Hong et al., 2018; Sultanova et al., 2001; Umbetova et al., 2005; Zhang & Tu, 1994). Interestingly, a distinctive compound isolated from *T. ramosissima* Ledeb. has also been reported as, 8-(1-(1-(3,4-dihydroxyphenyl)ethyl)-3',4'-dimethoxy-3,5,7-trihydroxyflavone (ramosissimin) (Hong et al., 2018).

Table 2 Geographical distribution of *Tamarix* L.

No	Species	Geographical distribution around the world	Geographical distribution in Kazakhstan
1	<i>Tamarix litwinowii</i>	Central Asia, Eastern Iran, Northern Afghanistan, Turkmenistan (Pavlov, 1963)	Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan (Ile-Balkhash, 2025)
2	<i>Tamarix laxa</i> Willd.	Turkmenistan, Caucasus, Central Asia, Iran, Afghanistan, Western China, Mongolia (Pavlov, 1963)	Semipalatinsk, near the Caspian, Aktobe, Embi, Turgen, Western and Eastern Melkosop. Zaisan, Northern Ust-Urt, Mangyshlak, Priaral, Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan, Balkhash, Alakol (Pavlov, 1963)
3	<i>Tamarix meyeri</i> Boiss.	Turkmenistan, Caucasus, Central Asia, Iran, Minor Asia, Syria, Egypt, Tajikistan, Afghanistan (Pavlov, 1963)	Mangyshlak (Pavlov, 1963)
4	<i>Tamarix elongata</i> Ledeb	Russia, Kazakhstan, Turkmenistan, Uzbekistan, Central Asia, Western China, southern Mongolia (Pavlov, 1963)	Tobyl, Yeshim, Embi, Turg., Western Melkosop. Zaisan, Northern Ust-Urt, Mangyshlak, Aral, Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan, Balkhash, Alakol (Pavlov, 1963)
5	<i>Tamarix passerinoides</i> Delile.	Turkmenistan, Iran, Asia Minor, Arabia, Egypt, Algerian Sahara, Libya, Sinai (Gorshkova, 1949 ; Zieliński, 1994 ; Assadi, Ramak Maassoumi & Khatamsaz, 1989)	South Usturt (Pavlov, 1963)
6	<i>Tamarix gracilis</i> Willd.	Western Siberia, Western China, Mongolia, Kazakhstan, Tajikistan, Turkmenistan, Turkey (Gorshkova, 1949 ; Zieliński, 1994 ; Assadi, Ramak Maassoumi & Khatamsaz, 1989 ; Pavlov, 1963)	Tobyl Eshim, Irtysh, Semey, near the Caspian, Mugodzhar, Emb, Turk, Western and Eastern Melkosop, Northern Ustyurt, Aral, Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan (Yany Kurgan station) (Pavlov, 1963)
7	<i>Tamarix hispida</i> Willd.	Central Asia, Afghanistan, China, Mongolia, Kazakhstan, Russia, Uzbekistan, Turkmenistan, Afghanistan (Pavlov, 1963)	near the Caspian, Embi, Turg., Western and Eastern Melkosop, Northern Ustyurt, Aral, Kyzylorda, Turkestan (Pavlov, 1963)
8	<i>Tamarix leptostachys</i> Bunge	Central Asia (northern part), Western China, Mongolia (Pavlov, 1963)	Embi, Turg., Zaysan, Aral, Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan, Balkhash, Alakol (Pavlov, 1963)

(continued on next page)

Table 2 (continued)

No	Species	Geographical distribution around the world	Geographical distribution in Kazakhstan
9	<i>Tamarix arceuthoides</i> Bunge	Central Asia (Tien Shan, Pamir-Alai, Kopetdag), Iran, Iraq, Afghanistan, Western China, Mongolia, Kazakhstan, Turkmenistan, Uzbekistan, Kyrgyzstan, Pakistan, Kashmir (Pavlov, 1963)	Muyun-Kum, Balkhash, Dzungen Alatau, Zail. Kung. Alatau, Kirg Alatau, Karat, west of the TSh (Pavlov, 1963)
10	<i>Tamarix hohenackeri</i> Bunge	Caucasus, Central Asia, Northern Iran, Mongolia, Armenia, Azerbaijan, China (Gansu, Jiangxi, Nei-Mongol, Zizhiqiu, Ningxia-Hui, Qinghai and Xinjiang), Crimea, Kazakhstan, Republic of Georgia, Russia, Turkey, Greece (Bikbulatova & Korul'kina, 2001 ; Xing et al., 2014 , Shishkin, 1949 ; Tropicos, 2024 ; EUNIS, 2018)	Kyzylorda, Betpakdala, Muyun-Kum, Kyzylkum, Turkestan, Balkhash, Alakol, Dzungen Alatau, Zail. Kung. Alatau, Chu-II mountains (Pavlov, 1963)
11	<i>Tamarix ramosissima</i> Ledeb.	Caucasus, Central Asia, Iran, Iraq, Asia Minor, Afghanistan, China, Mongolia, Ukraine, Kazakhstan, Tajikistan, Uzbekistan, Kyrgyzstan, Azerbaijan, Turkmenistan, Tibet, China, Korea, Armenia, Georgia, Pakistan, Qatar, Turkey, Ukraine, Uzbekistan (Gorshkova, 1949 ; Zieliński, 1994 ; Assadi, Ramak Maassoumi & Khatamsaz, 1989 ; Kaiser, 1983 ; Shishkin, 1949 ; Gaskin & Schaal, 2002 ; CABI International, 2024)	Tobyl Eshim, Semipalatinsk, Prikaspi, Aktobe, Zaysan, Northern Ust-Urt, Mugodzh, Embi, Turg, Western and Eastern Melkosop, Betpakdala, Muyun-Kum, Balkhash-Alakol, Southern Ust-Urt, Kyzylkum, Turkestan, Dzungen Alatau, Chu-II Mountains, Karat (Pavlov, 1963)
12	<i>Tamarix karelinii</i> Bunge (Hybrid <i>Tamarix hispida</i> Willd. × <i>Tamarix ramosissima</i> Ledeb.)	Central Asia, Iran, Mongolia (Gorshkova, 1949 ; Assadi, Ramak Maassoumi & Khatamsaz, 1989)	Turg., Kyzylorda, Betpakdala, Balkhash-Alakol, Turkestan, Zail. Kung. Alatau, Karatau (Pavlov, 1963)
13	<i>Tamarix eversmannii</i> Presl. (Hybrid <i>Tamarix leptostachys</i> Bunge × <i>Tamarix ramosissima</i> Ledeb.)	Southeast Europe (Pavlov, 1963)	near the Caspian, Aral, Kyzylorda, Muyun-Kum, Balkhash-Alakol (Pavlov, 1963)

Phenolic acids

Considering the diverse biological activities of the genus *Tamarix*, phenolic acids serve as essential constituents in its phytochemical profiles. Previous studies asserted that phenolic acids were extracted from the aerial components of three species in Kazakhstan and China: (i) *T. hispida* Willd., (ii) *T. hohenackeri* Bunge, and (iii) *T. ramosissima* Ledeb. ([Bikbulatova & Korul'kina, 2001](#); [Xing et al., 2014](#); [Ren et al., 2019](#); [Sultanova et al., 2004a](#); [Xiao et al., 2013](#)). These studies identified the presence of gallic, ellagic, ferulic, caffeic,




Figure 2 The chemical compounds of *T. ramosissima* Ledeb.

Full-size [DOI: 10.7717/peerj.20059/fig-2](https://doi.org/10.7717/peerj.20059/fig-2)

and protocatechuic acids within the various components with antioxidant and anti-inflammatory effects.

Triterpenoids

Triterpenes are abundant in *Tamarix* plants within Kazakhstan and China, such as *T. hispida* Willd., *T. hohenackeri* Bunge, *T. laxa* Willd., *T. ramosissima* Ledeb., and *T. elongata* Ledeb. (Xing et al., 2014; Sultanova et al., 2004b; Umbetova et al., 2004b; Sultanova et al., 2013). Specifically, a triterpenoid compound (isotamarixen) extracted from *T. hispida* Willd is a potential novel compound exhibiting antioxidant, anti-inflammatory, anti-tumor, and neuroprotective effects (Sultanova et al., 2004b).

Other phytochemicals

A range of phytochemical constituents have been identified in *Tamarix* species, including alkaloids, coumarins, anthraquinone glycosides, lactams, tannins, tetraterpenes, and phytosterols (Xing et al., 2014; Orabi et al., 2013; Umbetova et al., 2004b). These compounds exhibit diverse pharmacological activities. Notably, the lactam derivative (*E*)-3-(4-hydroxy-3-methoxybenzylidene)-4-(4-hydroxy-3-methoxyphenyl) pyrrolidine-2-one (commonly referred to as tamaractam) was isolated from the branches and leaves of *T. ramosissima* Ledeb. (Yao et al., 2017). Therefore, phytochemicals demonstrate significant biodiversity, offering several potential applications in pharmaceuticals and medicine due to their complex interactions with biological systems. Figs. 2, 3, 4 and 5 provide an overview of the chemical compounds in the *Tamarix* genus.

Primary metabolites

Although most phytochemical investigations on *Tamarix* have focused on secondary metabolites, several studies have also explored the presence of primary metabolites,




Figure 3 Chemical compounds *T. Hohenackeri* Bunge.

Full-size [DOI: 10.7717/peerj.20059/fig-3](https://doi.org/10.7717/peerj.20059/fig-3)




Figure 4 The chemical compounds of *T. Hispida* Willd.

Full-size [DOI: 10.7717/peerj.20059/fig-4](https://doi.org/10.7717/peerj.20059/fig-4)




Figure 5 The chemical compounds of *T. Laxa* Willd. And *T. elongata* Ledeb.

Full-size [DOI: 10.7717/peerj.20059/fig-5](https://doi.org/10.7717/peerj.20059/fig-5)

including amino acids, carbohydrates, vitamins, fatty acids, and minerals. These components not only play crucial physiological roles but also contribute to the adaptive potential of *Tamarix* species under extreme environmental conditions (Abilov & Sultanova, 2018).

Amino acids

The amino acid composition of *Tamarix* species—particularly *T. hohenackeri* Bunge and *T. ramosissima* Ledeb.—includes tryptophan, cysteine, threonine, glutamic acid, methionine, asparagine, glutamine, and arginine. Additional amino acids such as aspartic acid, serine, glycine, alanine, proline, tyrosine, valine, histidine, isoleucine, leucine, phenylalanine, and lysine have been identified in *T. ramosissima* (Abilov & Sultanova, 2018).

Kazakhstani studies further report the presence of 18 free amino acids in *T. elongata* Ledeb., *T. laxa* Willd., *T. ramosissima* Ledeb., and *T. hispida* Willd., with alanine, proline, leucine, and serine being predominant. The amino acid content was notably higher in samples collected from saline soils, suggesting a possible correlation with environmental salinity stress (Abilov & Sultanova, 2018).

Carbohydrates

The carbohydrate profiles of *T. elongata*, *T. laxa*, *T. ramosissima*, and *T. hispida* include monosaccharides such as glucose, galactose, rhamnose, and arabinose. Another study reported the presence of glucose, arabinose, xylose, galactose, and sucrose in *T. hohenackeri* and *T. ramosissima* (Abilov & Sultanova, 2018).

Vitamins

Vitamin analysis of *T. elongata*, *T. laxa*, *T. ramosissima*, and *T. hispida* revealed low levels of vitamin C, ranging from 0.003% to 0.046%. Vitamin E content in samples collected in the Almaty region ranged between 0.054% and 0.086% (Abilov & Sultanova, 2018).

Fatty acids

Fatty acid profiling of *T. elongata*, *T. laxa*, *T. ramosissima*, and *T. hispida* revealed a consistent qualitative composition across species, with 11 major saturated and unsaturated fatty acids identified. These include myristic, pentadecanoic, palmitic, palmitoleic, stearic, oleic, linoleic, eicosenoic, eicosadienoic, eicosatrienoic, and arachidonic acids. Quantitative variations among species were also noted (Abilov & Sultanova, 2018).

Mineral composition

Ash residue analysis of *Tamarix* species revealed the presence of essential macro- and microelements such as potassium, sodium, calcium, magnesium, copper, iron, zinc, cobalt, lead, nickel, and manganese. Sodium, potassium, and calcium were the most abundant macroelements (Abilov & Sultanova, 2018).

Pharmacological attributes

Tamarix plants possess multiple biologically active attributes owing to their diverse chemical composition, including flavonoids, tannins, alkaloids, and essential oils. Various

Table 3 Pharmacological activity of the genus *Tamarix* of the flora of Kazakhstan.

No	Pharmacological activity	Species of plant	Part of the plant	Extragent	The mechanism of action	Compound(s)	Ref
1	Antioxidant activity	<i>Tamarix laxa</i> Willd., <i>Tamarix hispida</i> Willd., <i>Tamarix ramosissima</i> Ledeb., <i>Tamarix elongata</i> Ledeb.	Aerial parts	Ethanol, chloroform, ethyl acetate	Free radical scavenging	Ursolic acid, isotamarixen, pentacyclic triterpenoid, hispidulin, cirsimaritin and isorhamnetin	Ren et al. (2019) ; Sultanova et al. (2001) ; Sultanova et al. (2004a) ; Sultanova et al. (2013) ; Bukhari et al. (2021)
		<i>Tamarix ramosissima</i> Ledeb.		Ethanol (1:5)	Normalising the effects of oxidative stress in diabetes mellitus.	caffeic acid, luteol, kaempferol chlorogenic acid, caffeic acid, rutoside	Rotaru et al. (2018)
2	Antibacterial activity	<i>Tamarix elongata</i> Ledeb, <i>Tamarix arceuthoides</i> Bunge, <i>Tamarix ramosissima</i> Ledeb, <i>Tamarix macrocarpa</i> , <i>Tamarix ramosissima</i>	Leaves	Ethanol, chloroform and n-hexane extracts	Suppression of bacterial colony growth, Inhibition of biofilms	gallic acid, isoferulic acid, kaempferol and quercetin, hispidulin, cirsimaritin and isoramnetin, polyphenolic compounds	Ren et al. (2019) ; Sultanova et al. (2001) ; Rotaru et al. (2018) ; Zazharskyi et al. (2020) ; Al-Shamma, Kadhum & Al-Hiti (2005) ; Akya et al. (2014) ; Rotaru & Varut (2020)
3	Antimitogenic and chemosensitising activity	<i>Tamarix passerinoides</i> Delile, <i>Tamarix aucheriana</i> <i>Tamarix elongata</i> Ledeb	Aerial parts	Methanol Petroleum ether	Induction of apoptosis, cell cycle alteration	Syringic acid, methyl ferulate	Abaza et al. (2013) , Abaza et al. (2016)
4	Antidiabetic activity	<i>Tamarix hispida</i> Willd.	Aerial parts	Ethanol	Inhibition of enzymes associated with glucose metabolism	Hydrolysable tannins containing residues of gallic acid and its derivatives	Zhumagaliyeva et al. (2021)
5	Anti-leishmanial activity	<i>Tamarix ramosissima</i> Ledeb	Not specified	Chloroform, hydroalcoholic extract	Suppression of the viability of parasitic protozoa	Not specified	Qadir et al. (2014)
6	Anti-inflammatory action	<i>Tamarix hohenackeri</i> Bunge	Not specified	Not specified	Reduction in the production of inflammatory cytokines	Not specified	Chen et al. (2018)
7	Inhibition of ACE and platelet aggregation	<i>Tamarix hohenackeri</i> Bunge	Not specified	Ethyl acetate	Inhibition of angiotensin-converting enzyme and reduction of platelet aggregation	Flavonoids and phenolic compounds	Xing et al. (2014)
8	Inhibitory effect on α -glucosidase activity	<i>Tamarix hispida</i> Willd.	Aerial parts	Ethanol (50%)	Inhibition of α -glucosidase enzyme activity	Isotamarixen	Sultanova et al. (2013)
9	Antiproliferative	<i>Tamarix ramosissima</i> Ledeb <i>Tamarix elongata</i> Ledeb	Aerial parts	Ethanol Ligh petrol	Induces apoptosis in rheumatoid arthritis fibroblast-like synoviocytes, reducing the viability of synovial fibroblasts	Ramosissimin, methylferulate	Hong et al. (2018) , Abaza et al. (2013)
10	Antithrombolytic activity	<i>Tamarix arceuthoides</i>		n-hexane	Platelet destruction		Bukhari et al. (2021)

Tamarix species have also revealed significant antioxidant, antibacterial, anti-inflammatory, and antidiabetic activities (see [Table 3](#)).

Antioxidant properties

Antioxidant effectiveness is associated with considerable polyphenolic contents (such as flavonoids and tannins), which are known for their ability to neutralize free radicals and protect cells from oxidative stress. Significant examples include *T. laxa*, *T. elongata*,

T. hispida, and *T. ramosissima*, possessing antioxidant attributes arising from terpenoid compounds (ursolic acid and isotamarixene) for capturing free radicals and preventing cell and DNA damage. The 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical capture method also presents that *T. ramosissima* extracts containing hispidulin and circymaritin exhibit potent antioxidant potential (Bahramsoltani et al., 2020). Furthermore, an inhibitory effect on α -glycosidase activity and significant antioxidant properties have been denoted in triterpenoid isotamarixene (3- α -[3, 4-dihydroxy-trans-cinnamoyloxy]-D-friedoolean-14-en-28-oic acids) within a 50% *T. hispida* ethanol extract (Sultanova et al., 2002; Ren et al., 2019; Li et al., 2024; Karker et al., 2024a).

The N-hexane extract of *T. arceuthoides* demonstrates protective effects on DNA from oxidative damage caused by H₂O₂, indicating notable antioxidant potential. This outcome signifies that the active compounds in the extract can inhibit or diminish oxidative damage to cellular structures. Multiple studies have also reported the antioxidant activities of other plant species for the genus *Tamarix*. This necessity has been attributed to the plant's high levels of phenolic compounds and flavonoids. Consequently, these studies have stated that phenolic acids and flavonoids can effectively protect against oxidative stress, positioning them as promising candidates for medical applications to treat oxidative process-related diseases (neurodegenerative disorders and cardiovascular diseases). The data have successfully confirmed the significant potential of the *Tamarix* genus in developing natural antioxidant agents (Umbetova et al., 2004b; Sultanova et al., 2013; Yao et al., 2017; Bukhari et al., 2021; Zazharskyi et al., 2020; Gupta, Dubey & Kumar, 2016; Karker et al., 2024b; Zar Kalai et al., 2024; Samejo et al., 2023; Esma et al., 2023; Iqbal et al., 2022; Nisar et al., 2023).

Antibacterial activity

Different extracts of *Tamarix ramosissima* have shown varying antibacterial potential depending on the plant part and extraction solvent. A water–acetone leaf extract demonstrated *in vitro* activity against 15 microbial strains, with minimum inhibitory concentrations (MICs) ranging from 12.5 to 100 μ g/mL and antifungal effects observed at 400 μ g/mL. In contrast, the ethanolic bark extract was tested against seven bacterial and four fungal strains, showing MICs of 5–10 mg/mL and minimum bactericidal concentrations (MBCs) of 10–25 mg/mL. The bark extract exhibited stronger antibacterial activity against Gram-positive bacteria compared to Gram-negative ones (Bahramsoltani et al., 2020; Yusufoglu & Alqasoumi, 2011; Abo-Dola et al., 2015). The observed antibacterial effects may be attributed to the individual action or synergistic interactions of these identified compounds. Another example is the effective growth inhibition of various bacterial and fungal pathogens, such as *Corynebacterium diphtheriae* and *Aspergillus niger*, observed during *in vitro* assays using ethyl acetate and water-acetone extracts of *T. ramosissima*. These results highlight the potential these extracts as natural antimicrobial agents for combating infectious diseases (Rotaru & Varut, 2020).

Zazharskyi et al. (2020) argued that *T. elongata* extracts inhibited bacterial growth, including antibiotic-resistant pathogens. Based on the findings reported by Bukhari et al. (2021) the *n*-hexane extract of *T. arceuthoides* exhibited notable antibacterial activity,

forming inhibition zones of 27.25 ± 0.37 mm against *Micrococcus spp.*, 14.75 ± 0.28 mm against *Staphylococcus aureus*, and 14.25 ± 0.24 mm against *Escherichia coli*. For comparison, the standard antibiotic erythromycin produced inhibition zones of 25.35 ± 0.26 mm, 30.25 ± 0.23 mm, and 34.75 ± 0.34 mm against the same bacterial strains, respectively. Although the extract showed a larger inhibition zone than erythromycin specifically against *Micrococcus spp.*, its overall efficacy was lower against the other tested strains.

In addition to antibacterial effects, *T. arceuthoides* extracts also demonstrated antibiofilm activity. The chloroform extract inhibited biofilm formation by *S. aureus* and *E. coli* by 47.36% and 45.53%, respectively, while the *n*-hexane extract produced slightly lower inhibition values of 45.92% for *S. aureus* and 38.58% for *E. coli*. In comparison, ciprofloxacin, used as a standard control, achieved significantly higher inhibition rates of 77.57% and 82.65%, respectively. These findings indicate that although *T. arceuthoides* extracts possess promising antibacterial and antibiofilm potential, their efficacy remains lower than that of standard antibiotics when assessed under controlled experimental conditions (Bukhari et al., 2021).

The antimicrobial activity of *T. macrocarpa* ethanol extract has been effectively evaluated, and it is active against *S. aureus*, *Bacillus subtilis*, and *Candida albicans*. This plant contains a high concentration of phenolic and polyphenolic compounds (gallic acid, isoferulic acid, campferol, and quercetin) based on the phytochemical analysis (Al-Shamma, Kadhun & Al-Hiti, 2005). Several studies have also computed a significant relationship between the antimicrobial properties of the genus *Tamarix* and the necessity of developing strategies to address pathogenic microorganisms regarding antibiotic-resistant forms. These studies have indicated that biologically active compounds in plants (flavonoids, phenolic acids, and tannins) demonstrate antimicrobial activity against bacteria and fungi. The antimicrobial activity of *Tamarix* is also influenced by its geographical origin and growing conditions, necessitating study to identify the most effective sources for pharmaceutical applications. Thus, studies concerning *Tamarix* species' antimicrobial properties persist, presenting new opportunities for drug development to treat infectious diseases and applications within the pharmaceutical industry (Paul et al., 2022; El-Amier et al., 2021; Dawood, Al-Daody & Al-Takay, 2021; Romeilah et al., 2021; AbouZid & Sleem, 2011; Al-Omar et al., 2020; Boulaaba et al., 2015; Lefahal et al., 2010).

Antimitogenic and chemosensitizing activity

Abaza et al. (2013) indicated that syringic acid and methyl ferulate isolated from methanolic *T. passerinoides* Delile and *T. aucheriana* extracts could be adjuvant agents in colorectal cancer management (Lefahal et al., 2010; Al-Judaibi, 2020). The studies explained that the primary actions of these compounds included inhibiting tumor cell growth, inducing apoptosis, and enhancing cell sensitivity to chemotherapeutic agents. A positive correlation between syringic acid and chemotherapy efficiency was also reported. Peydayesh et al. (2020) investigated the inhibitory effect of silver nanoparticles from *T. hispida* extracts on the cyclin D1 protein expression in human cancer cell lines. The study described the methodology based on biosynthesized silver nanoparticles for evaluating the anti-cancer

effect concerning molecular mechanisms, cell cycle regulation, and cancer cell proliferation. This method offered an alternative approach to nanoparticle production that eliminated chemicals commonly employed in nanomaterial synthesis. Therefore, the biological synthesis method could reduce toxicity and impart additional biologically active properties to nanoparticles due to compounds present in extracts ([Aljubouri & Alobaidi, 2023](#)).

Numerous studies on the *Tamarix* genus have garnered scientific interest regarding its potential anti-cancer properties. These studies demonstrate that *Tamarix* extracts are active against various cancer cell types, including breast, lung, and colon cancers. The anti-tumor effects observed in laboratory conditions also indicate a growth reduction of cultured malignant cell tumors, suggesting that these plants are promising candidates for pre-clinical studies ([Abaza et al., 2013](#); [Abaza et al., 2016](#); [Lefahal et al., 2010](#); [Al-Judaibi, 2020](#); [Peydayesh et al., 2020](#); [Nicolini et al., 2014](#); [Abdel-Tawab et al., 2019](#); [Boulaaba et al., 2013](#)).

Antidiabetic activity

Given the increasing prevalence of diabetes (particularly type 2 diabetes) involving insulin resistance and metabolic disorders, studies regarding the antidiabetic properties of the genus *Tamarix* have attracted interest from the scientific community. Plants within this genus usually contain various biologically active compounds that can positively influence glucose and lipid metabolisms. Specifically, *T. hispida* possesses tannins inhibiting the enzyme α -glucosidase, preventing a rapid increase in blood sugar levels ([AbouZid & Sleem, 2011](#)). Specific antidiabetic property-related studies of extracts from several *Tamarix* genera have also demonstrated their capacity to reduce blood glucose levels in experimental animals. This outcome is attributed to improved insulin sensitivity and increased glucose absorption capacity of tissues ([Alnuqaydan & Rah, 2021](#); [Alnuqaydan et al., 2022](#); [Fayed et al., 2023](#); [Al Sobeai, 2018](#); [Sehrawat & Sultana, 2006](#); [Ullah et al., 2017](#); [Hebi & Eddouks, 2017](#); [Mahfoudhi et al., 2016](#)).

Anti-inflammatory action

[Chen et al. \(2018\)](#) assessed the anti-inflammatory properties of *T. hohenackeri* extracts. The study identified potential neuroinflammation inhibitors from four *T. hohenackeri* extracts while examining their possible mechanisms of action. Consequently, reduced inflammatory response was observed due to the flavonoids and phenolic acids reducing the cytokine production in lipopolysaccharide-activated microglia. Previous studies also examined the anti-inflammatory properties of other *Tamarix* species ([Zhumagaliyeva et al., 2021](#); [Chen et al., 2018](#); [Yusufoglu & Alqasoumi, 2011](#); [Fayez et al., 2024](#); [Hamed et al., 2024](#)). These studies indicated that the main mechanisms of their action involved inhibiting inflammatory mediators and lowering inflammatory cells' activity. The *Tamarix* plant extracts were also reported as adjunctive treatments (inflammatory conditions) and aided formulation of phytotherapeutic agents (anti-inflammatory and antioxidant properties) owing to their capacity to suppress the activity of pro-inflammatory cytokines and enzymes while reducing oxidative stress,

Inhibition of ACE and platelet aggregation

[Xing et al. \(2014\)](#) investigated bioactive compounds isolated from *T. hohenackeri* Bunge a species traditionally used in folk medicine and known for its radioprotective properties. The study suggested that flavonoids and phenolic compounds present in the extract exhibited inhibitory effects on angiotensin-converting enzyme (ACE) and platelet aggregation, indicating potential cardiovascular benefits. In another study, [Bukhari et al. \(2021\)](#) evaluated the thrombolytic activity of *T. arceuthoides* *in vitro*. The experiment involved incubating human blood clots with plant extracts at 37 °C and measuring the percentage of clot lysis. Streptokinase served as the positive control, showing 70.83 ± 0.84% clot lysis. Among the tested extracts, the n-hexane extract of *T. arceuthoides* demonstrated thrombolytic activity of 19.72%, whereas the chloroform extract exhibited 18.57%. Although the activity was lower compared to the standard, these findings suggest the presence of bioactive compounds with potential clot-dissolving properties that warrant further investigation.

Antileishmanial activity

[Akya et al. \(2020\)](#) investigated the antileishmanial effects of hydroalcoholic and chloroformic extracts of *Tamarix ramosissima* against *Leishmania major* and *Leishmania tropica* promastigote forms. The hydroalcoholic extract reduced parasite viability at all tested concentrations, with an IC₅₀ value of 0.1 mg/mL for both species. The chloroformic extract showed moderate inhibitory effects, with a measured IC₅₀ of 50 mg/mL. These findings indicate that *T. ramosissima* exhibits high antileishmanial activity and holds promise as a natural source for developing novel agents against leishmaniasis ([Akya et al., 2020](#)).

Antiproliferative and apoptosis-inducing effects

The tamaractam and ramosissimin compounds isolated from *T. ramosissima* induce apoptosis while inhibiting the growth of synoviocytes associated with rheumatoid arthritis ([Umbetova et al., 2004a](#); [Hong et al., 2018](#); [Yao et al., 2017](#)).

CONCLUSIONS

The study of 13 *Tamarix* species found in Kazakhstan has shown that species such as *T. ramosissima*, *T. hispida*, and *T. laxa* have been more extensively studied compared to less explored species like *T. eversmannii*, *T. meyeri*, and *T. arceuthoides*. Further research is needed to conduct a detailed analysis of their morphological, molecular phylogenetic, and chemotaxonomic characteristics. Among the identified biological activities, the most promising ones include antioxidant, antibacterial, anti-inflammatory, antidiabetic, and anticancer properties. The antioxidant activity is attributed to the high content of polyphenols and flavonoids, making *Tamarix* a potential source of natural antioxidants for medical and food applications. The antibacterial activity, particularly pronounced in the bark and leaf extracts of *T. ramosissima*, suggests potential for developing novel natural antimicrobial agents. The anti-inflammatory and antidiabetic properties indicate

the potential of *Tamarix* for developing phytopharmaceuticals, while specific compounds such as tamarixetin and tamaractam may be utilized in cancer therapy.

Potential areas of application include pharmacology, agriculture, and the food industry. In medicine, *Tamarix* can be used to develop treatments for inflammatory and metabolic diseases. In agriculture, its high tolerance to drought and salinity makes it a valuable resource for soil stabilization and land restoration. In the food industry, *Tamarix* extracts may serve as natural antioxidants and functional additives.

To resolve controversial taxonomic issues, it is recommended to use molecular genetic methods (ISSR, RAPD, ITS, SNP markers) to clarify phylogenetic relationships and distinguish closely related species, apply DNA barcoding to identify hybrid forms, conduct comprehensive morphological and anatomical studies, including palynomorphological analysis, and perform chemotaxonomic analysis based on secondary metabolite profiles. The development of a unified database integrating molecular, morphological, and chemical characteristics of *Tamarix* species in Kazakhstan will enable the standardization of their identification and classification criteria. Further research will not only help eliminate existing gaps in the study of these plants but also expand their practical applications in various scientific and industrial fields.

It is also important to highlight that *T. androssowii*, one of the species distributed in Kazakhstan, is listed in the Red Book of Kazakhstan (2014) as a rare and endangered species. This fact underscores the dual importance of the genus *Tamarix*—not only for the development of phytopharmaceuticals and other practical applications but also for biodiversity conservation and sustainable use of plant resources in Kazakhstan's arid and semi-arid ecosystems.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Arailym Daulbayeva performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Gulnara Kadyrbayeva conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Kaldanay Kozhanova conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Shazlin Shaharudin analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Nurgali Rakhymbayev performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

- Zoya Allamberganova analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Rabiga Anarbayeva analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Urziya Alimova analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Aigerim Kantureyeva analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Ainash Baidullaeva analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Vladimir An analyzed the data, authored or reviewed drafts of the article, and approved the final draft.
- Bakkonat Kydyrbai analyzed the data, authored or reviewed drafts of the article, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

This is a literature review.

REFERENCES

- Abaza MS, Afzal M, Al-Attiah RJ, Guleri R. 2016.** Methylferulate from *Tamarix aucheriana* inhibits growth and enhances chemosensitivity of human colorectal cancer cells: possible mechanism of action. *BMC Complementary and Alternative Medicine* **16**(1):384 DOI 10.1186/s12906-016-1358-8.
- Abaza MS, Al-Attiah R, Bhardwaj R, Abbadi G, Koyippally M, Afzal M. 2013.** Syringic acid from *Tamarix aucheriana* possesses antimitogenic and chemo-sensitizing activities in human colorectal cancer cells. *Pharmaceutical Biology* **51**(9):1110–1124 DOI 10.3109/13880209.2013.781194.
- Abby B. 2013.** Tamarix Spread, Establishment, and Removal. Available at <https://www.lakeforest.edu/news/tamarix-spread-establishment-and-removal> (accessed on 030 May 2024).
- Abdel-Tawab FM, Shebl SA, Alsenosy NK, Abdou HS. 2019.** Assessment of anticarcinogenic effect of *Tamarix nilotica* on human lung cancer cell line. *Egyptian Journal of Genetics and Cytology* **48**(1):107–119.
- Abilov Z, Sultanova N. 2018.** *Chemical composition and biological activity of plants of the genus Tamarix*. Almaty: Kazakh University, 254.
- Abo-Dola M, Lutfi M, Bakhiet A, Mohamed A. 2015.** Anti-inflammatory, analgesic, antipyretic and the membrane-stabilizing effects of *Tamarix aphylla* ethanolic extract. *European Journal of Medicinal Plants* **5**(4):341–348 DOI 10.9734/EJMP/2015/13888.
- AbouZid S, Sleem A. 2011.** Hepatoprotective and antioxidant activities of *Tamarix nilotica* flowers. *Pharmaceutical Biology* **49**(4):392–395 DOI 10.3109/13880209.2010.518971.

- Afanasyev IA. 2012. The study of the nature of salt production by the tamarix plant growing in the Astrakhan region // Modern high-tech technologies. 5–8. Available at <https://top-technologies.ru/ru/article/view?id=30749> (accessed on 14 April 2024).
- Akhani H, Samadi N, Noormohammadi A, Borsch T. 2019. A new species of *Tamarix* (Tamaricaceae) from Hormozgan Province. S Iran, supported by morphology and molecular phylogenetics. *Willdenowia* 49:127–139 DOI 10.3372/wi.49.49113.
- Akya A, Chegene Lorestani R, Zhaleh H, Nemati Zargarani F, Ghadiri K, Rostamian M. 2020. Effect of *Vigna radiata*, *Tamarix ramosissima* and *Carthamus lanatus* extracts on *Leishmania major* and *Leishmania tropica*: an *in vitro* study. *Chinese Herbal Medicines* 12(2):171–177 DOI 10.1016/j.chmed.2019.12.006.
- Akya A, Mojarab M, Farshchian M, Ahmadi K. 2014. The effect of stem bark extracts of *Tamarix ramosissima* shrub on *Pseudomonas aeruginosa* isolates. *Scientific Journal of Kurdistan University of Medical Sciences* 19:128–134.
- Al-Judaibi AA. 2020. *Tamarix arabica* and *Salvadora persica* as antibacterial agents. *AIMS Microbiology* 6(2):121 DOI 10.3934/microbiol.2020008.
- Al-Omar MS, Mohammed HA, Mohammed SA, Abd-Elmoniem E, Kandil YI, Eldeeb HM, Chigurupati S, Sulaiman GM, Al-Khurayyif HK, Almansour BS, Suryavamshi PM, Khan RA. 2020. Anti-microbial, anti-oxidant, and α -amylase inhibitory activity of traditionally-used medicinal herbs: a comparative analyses of pharmacology, and phytoconstituents of regional halophytic plants’ diaspora. *Molecules* 25(22):5457 DOI 10.3390/molecules25225457.
- Al-Shamma AA, Kadhum EJ, Al-Hiti MA. 2005. Antimicrobial activity and phytochemical investigation of *Tamarix Macrocarpa* (Ehrenb.) bge wildly grown in Iraq. *Medical Plants* 29.1:99–104.
- Al Sobeai SM. 2018. Anticancer, cytotoxic effect of *Tamarix aphylla*, and antibacterial screening efficiency against multidrug-resistant human pathogens. *Asian Journal of Pharmaceutical and Clinical Research* 11(11):241–246 DOI 10.22159/ajpcr.2018.v11i11.27309.
- Aljubouri SA, Alobaidi KH. 2023. Assessment of the Antivirulence Potential of *Tamarix aphylla* Ethanolic Extract against Multidrug-Resistant *Streptococcus mutans* Isolated from Iraqi Patients. *The Scientific World Journal* 2023(1):6662262 DOI 10.1155/2023/6662262.
- Alnuqaydan AM, Almutary AG, Alshehri OY, Henidi HA, Alajlan AM, Al Tamim A, Alowafeer A, Rather MY, Rah B. 2022. Evaluation of the cytotoxic activity of *Tamarix articulata* and its anticancer potential in prostate cancer cells. *Journal of Applied Pharmaceutical Science* 12(02):089–108 DOI 10.7324/JAPS.2021.120209.
- Alnuqaydan AM, Rah B. 2021. *Tamarix articulata* Inhibits cell proliferation, promotes cell death mechanisms and triggers G0/G1 cell cycle arrest in hepatocellular carcinoma cells. *Food Technology and Biotechnology* 59(2):162–173 DOI 10.17113/ftb.59.02.21.6904.
- Amirdovlat A. 1990. *Unnecessary for the ignorant*. M: Science. Moscow: Nauka.

- Arianmanesh R, Mehregan I, Assadi M, Nejadstari T. 2015.** SEM studies on morphology and function of salt glands in the genus *Tamarix* L.(Tamaricaceae) from Iran. *Asian Journal of Biological and Life Sciences* **4**(3):212–216.
- Arianmanesh R, Mehregan I, Assadi M, Nejadstari T. 2016.** Comparative Morphology of the Genus *Tamarix* (Tamaricaceae) in Iran. *International Letters of Natural Sciences* **60**:1–12 DOI [10.56431/p-6s8gxp](https://doi.org/10.56431/p-6s8gxp).
- Ashirbekov U. 2013.** Experience of afforestation on the dried-up bottom of the Aral Sea. *Problems of Desert Development* **68**:1–5.
- Assadi M, Ramak Maassoumi AA, Khatamsaz M. 1989.** *Flora of Iran*. Tehran: Ministry of Agriculture, Islamic Republic of Iran, 1–151.
- Bahramsoltani R, Kalkhorani M, Zaidi SMA, Farzaei MH, Rahimi R. 2020.** The genus *Tamarix*: traditional uses, phytochemistry, and pharmacology. *Journal of Ethnopharmacology* **246**:112245 DOI [10.1016/j.jep.2019.112245](https://doi.org/10.1016/j.jep.2019.112245).
- Baktybayeva LK, Namaz ER, Kalzhan KM, Berlin KD, Umbetiyarova LB. 2019.** Pharmacological properties of endemic plants growing in the steppes of Kazakhstan. DOI [10.26577/EJE.2019.V58.I1.08](https://doi.org/10.26577/EJE.2019.V58.I1.08).
- Baum B. 1978.** *The genus Tamarix*. Jerusalem: Israel Academy of Sciences and Humanities.
- Bencherif K, Trodi F, Hamidi M, Dalpè Y, Hadj-Sahraoui AL. 2020.** Biological overview and adaptability strategies of *Tamarix* plants, *T. articulata* and *T. gallica* to abiotic stress. In: *Springer eBooks*. New York: Springer, 401–433 DOI [10.1007/978-981-15-9380-2_14](https://doi.org/10.1007/978-981-15-9380-2_14).
- Bikbulatova TN, Korul’kina LM. 2001.** Composition of *Tamarix hokenakeri* and *T. ramosissima*. *Chemistry of Natural Compounds* **37**:216–218 DOI [10.1023/A:1012501620145](https://doi.org/10.1023/A:1012501620145).
- Boulaaba M, Snoussi M, Saada M, Mkadmini K, Smaoui A, Abdelly C, Ksouri R. 2015.** Antimicrobial activities and phytochemical analysis of *Tamarix gallica* extracts. *Industrial Crops and Products* **76**:1114–1122 DOI [10.1016/j.indcrop.2015.08.020](https://doi.org/10.1016/j.indcrop.2015.08.020).
- Boulaaba M, Tsolmon S, Ksouri R, Han J, Kawada K, Smaoui A, Abdelly C, Isoda H. 2013.** Anticancer effect of *Tamarix gallica* extracts on human colon cancer cells involves Erk1/2 and p38 action on G 2/M cell cycle arrest. *Cytotechnology* **65**(6):927–936 DOI [10.1007/s10616-013-9564-4](https://doi.org/10.1007/s10616-013-9564-4).
- Breckle S-W, Wucherer W, Dimeyeva LA. 2012.** Vegetation of the aralkum. In: Veste, Breckle, Wucherer, eds. *Aralkum –A man-made desert: the desiccated floor of the Aral Sea (Central Asia)*. Cham: Springer, 215–230 DOI [10.1007/978-3-642-21117-1_9](https://doi.org/10.1007/978-3-642-21117-1_9).
- Britannica. 2019.** The Editors of Encyclopaedia. “tamarisk”. Encyclopedia Britannica. Available at <https://www.britannica.com/plant/tamarisk> (accessed on 30 May 2024).
- Bukhari SA, Mustafa G, Bashir S, Anjum F. 2021.** *In vitro* biological evaluation and DNA damage protection activities of cotoneaster afghanicus G.Klotz and *Tamarix arceuthoides* bunge extracts. *Pakistan Journal of Botany* **53**(1):197–202 DOI [10.30848/PJB2021-1\(15\)](https://doi.org/10.30848/PJB2021-1(15)).
- CABI International. 2024.** *Tamarix ramosissima* (saltcedar) [original text by Pasiecznik N]. CABI Invasive Species Compendium. Wallingford, United Kingdom: CAB

International. Available at <https://www.cabi.org/ISC/datasheet/52503> (accessed on 30 May 2024).

- Chen R, Yang Y, Xu J, Pan Y, Zhang W, Xing Y, Ni H, Sun Y, Hou Y, Li N. 2018.** *Tamarix hohenackeri* Bunge exerts anti-inflammatory effects on lipopolysaccharide-activated microglia *in vitro*. *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology* **40**:10–19 DOI [10.1016/j.phymed.2017.12.035](https://doi.org/10.1016/j.phymed.2017.12.035).
- Chepinoga VV, Barkalov VYu, Ebel AL, Knyazev MS, Baikov KS, Bobrov AA, Chkalov AV, Doronkin VM, Efimov PG, Friesen NV, German DA, Gontcharov AA, Grabovskaya-Borodina AE, Gureyeva II, Ivanenko YA, Kechaykin AA, Korobkov AA, Korolyuk EA, Kosachev PA, Kupriyanov AN, Luferov AN, Melnikov DG, Mikhailova MA, Nikiforova OD, Orlova LV, Ovchinnikova SV, Pinzhenina EA, Poliakova TA, Shekhovstsova IN, Shipunov AB, Shmakov AI, Smirnov SV, Tkach N, Troshkina VI, Tupitsyna NN, Vasjukov VM, Vlasova NV, Verkhovzina AV, Anenkhonov OA, Efremov AN, Glazunov VA, Khoreva MG, Kiseleva TI, Krestov PV, Kryukova MV, Kuzmin IV, Lashchinskiy NN, Pospelov IN, Pospelova EB, Zolotareva NV, Sennikov AN. 2024.** Checklist of vascular plants of Asian Russia. *Botanica Pacifica. A Journal of Plant Science and Conservation* **13**(Special issue):3–310 First published online 07 Nov 2024 DOI [10.17581/bp.2024.13S01](https://doi.org/10.17581/bp.2024.13S01).
- Chinese Plant Names. 2024.** *Tamarix laxa* Willd. Available at http://www.efloras.org/florataxon.aspx?flora_id=3&taxon_id=200014309 (accessed on 14 May 2024).
- Crins W. 1989.** The Tamaricaceae in the southeastern United States. *Journal of the Arnold Arboretum* **70**:403–405 DOI [10.5962/bhl.part.19790](https://doi.org/10.5962/bhl.part.19790).
- Daryo.uz. 2024.** Kazakhstan launches saxaul planting campaign in Aral Sea Region. Available at <https://daryo.uz/en/2024/04/08/kazakhstan-launches-saxaul-planting-campaign-in-aral-sea-region>.
- Dawood NAS, Al-Daody AC, Al-Takay TK. 2021.** *HPLC-UV analysis and antioxidant activities of phenolic compounds from bark of Tamarix articulata vahl shrubs grown in mosul province*. Bristol: IOP Publishing, 012046.
- De Martis B, Loi MC, Polo MB. 1984.** Il genere *Tamarix* L. (Tamaricaceae) in Sardegna. *Webbia* **37**(2):211–235 DOI [10.1080/00837792.1984.10670275](https://doi.org/10.1080/00837792.1984.10670275).
- El-Amier YA, Soufan W, Almutairi KF, Zaghloul NS, Abd-ElGawad AM. 2021.** Proximate composition, bioactive compounds, and antioxidant potential of wild halophytes grown in coastal salt marsh habitats. *Molecules* **27**(1):28 DOI [10.3390/molecules27010028](https://doi.org/10.3390/molecules27010028).
- Elkordy A, Faried A. 2017.** Pollen morphology and numerical analysis of *Tamarix* L. (Tamaricaceae) in Egypt and its systematic implication. *Bangladesh Journal of Plant Taxonomy* **24**(1):91–105 DOI [10.3329/bjpt.v24i1.33036](https://doi.org/10.3329/bjpt.v24i1.33036).
- Eshel A, Alekperov C, Langgut D, Oren I. 2019.** Pollen morphology of the genus *Tamarix* in Israel. In: Gul B, Böer B, Khan M, Clösener-Godt M, Hameed A, eds. *Sabkha Ecosystems. Tasks for vegetation science*, vol 49. Cham: Springer DOI [10.1007/978-3-030-04417-6_29](https://doi.org/10.1007/978-3-030-04417-6_29).
- Esma ATK, Rabah A, Djamila B, Mohamed SB, Chawki B, Ramazan E. 2023.** *In vitro* assessment of antioxidant, neuroprotective, anti-urease and anti-tyrosinase

- capacities of *Tamarix africana* leaves extracts. *Journal of Traditional Chinese Medicine* **43**(2):252.
- EUNIS. 2018. *Myricaria germanica* (L.) Desv. Available at <https://eunis.eea.europa.eu/species-names-result.jsp> (accessed on 30 May 2024).
- Fayed MA, Bakr RO, Yosri N, Khalifa SA, El-Seedi HR, DI Hamdan, Refaey MS. 2023. Chemical profiling and cytotoxic potential of the n-butanol fraction of *Tamarix nilotica* flowers. *BMC Complementary Medicine and Therapies* **23**(1):169 DOI [10.1186/s12906-023-03989-8](https://doi.org/10.1186/s12906-023-03989-8).
- Fayez N, Khalil W, Abdel-Sattar E, Abdel-Fattah AFM. 2024. Involvement of TNF α , IL-1 β , COX-2 and NO in the anti-inflammatory activity of *Tamarix aphylla* in Wistar albino rats: an *in-vivo* and *in-vitro* study. *BMC Complementary Medicine and Therapies* **24**(1):57 DOI [10.1186/s12906-024-04359-8](https://doi.org/10.1186/s12906-024-04359-8).
- Flora of Turkmenistan. 1951. The front cover. Alexander Downlow. *Science* **5**:271.
- Fungi.su. 2024. Loose tamarisk, (*Tamarix laxa* Willd.). Available at https://fungi.su/articles.php?article_id=1917 (accessed on 14 May 2018).
- Gaskin JF, Ghahremani-nejad F, Zhang D, Londo JP. 2004. A Systematic Overview of Frankeniaceae and Tamaricaceae from Nuclear rDNA and Plastid Sequence Data. *Annals of the Missouri Botanical Garden* **91**(3):401–409.
- Gaskin JF, Schaal BA. 2002. Hybrid *Tamarix* widespread in US invasion and undetected in native Asian range. *Proceedings of the National Academy of Sciences of the United States of America* **99**(17):11256–11259 DOI [10.1073/pnas.132403299](https://doi.org/10.1073/pnas.132403299).
- Gaskin JF, Schaal BA. 2003. Molecular phylogenetic investigation of U.S. invasive *Tamarix*. *Systematic Botany* **28**(1):86–95.
- Gemedzhieva NG, Sultanova NA, Abilov JA. 2013. Assessment of the raw material base of *R. Tamarix* L. species from the Southern Balkhash region. *Problems of Botany in Southern Siberia and Mongolia* **12**:12–15.
- Golub Valentin. 2008. The desert vegetation communities of the Lower Volga Valley. *Feddes Repertorium* **105**:499–515 DOI [10.1002/fedr.19941050716](https://doi.org/10.1002/fedr.19941050716).
- Gorshkova SG. 1949. The genus Grebenshchik is *Tamarix* L. Flora of the USSR. M.; L.: publishing House of the USSR Academy of Sciences, 15. 300–301.
- Gupta D, Dubey J, Kumar M. 2016. Phytochemical analysis and antimicrobial activity of some medicinal plants against selected common human pathogenic microorganisms. *Asian Pacific Journal of Tropical Disease* **6**(1):15–20 DOI [10.1016/S2222-1808\(15\)60978-1](https://doi.org/10.1016/S2222-1808(15)60978-1).
- Hamed MM, El-Wakil EA, Abdel-Aziz MS, El Shahed ZA, Abdel-Hady H. 2024. UPLC-ESI-MS Analysis of Secondary Metabolites from *Tamarix Aphylla* With Its In-Vitro Anti-Inflammatory and Anti-Microbial Activities. *Egyptian Journal of Chemistry* **67**(9):1–12 DOI [10.21608/ejchem.2024.271590.9403](https://doi.org/10.21608/ejchem.2024.271590.9403).
- Hebi M, Eddouks M. 2017. Hypolipidemic activity of *Tamarix articulata* Vahl. in diabetic rats. *Journal of Integrative Medicine* **15**(6):476–482 DOI [10.1016/S2095-4964\(17\)60361-3](https://doi.org/10.1016/S2095-4964(17)60361-3).
- Heywood VH, Brummitt RK, Culham A, Seberg O. 2007. *Flowering plant families of the world* (Vol. 88, p. 68). Ontario: Firefly books.

- Hong Z, Li J, Biswas S, Jiang CS, Huang Y, Sun T, Niu Y, Yu JQ, Li WQ, Yao Y. 2018.** Ramosissimin, a new flavonol from *Tararix ramosissima*, induces apoptosis in rheumatoid arthritis fibroblast-like synoviocytes. *Die Pharmazie* **73**(3):169–173 DOI 10.1691/ph.2018.7890.
- Ibn Sino AA. 1996.** The canon of medical science. Volume II. *Tashkent* **1996**:294.
- Ile-Balkhash. 2025.** Vegetation and soils of the Balkhash region. Available at <https://ile-balkhash.kz/priroda/rastitelnost-i-pochvy> (accessed on 15 April 2024).
- iNaturalist contributors, iNaturalist. 2024.** iNaturalist research-grade observations. iNaturalist.org. Available at <https://www.gbif.org/occurrence/4121250392> (accessed on 30 May 2024).
- Iqbal A, Begum N, Rabbi F, Akhtar N, Rahman KU, Khan WM, Shah Z. 2022.** In-vitro antimicrobial, antioxidant and enzyme inhibitory activities of fixed oil extracted from stem bark of *Tamarix aphylla*. *Pharmaceutical Chemistry Journal* **56**:1116–1122 DOI 10.1007/s11094-022-02762-4.
- Karker M, Oueslati S, Falleh H, Msaada K, Legault J, Abdelly C, Pichetted A, Ksouri R. 2024a.** Phytochemical investigation, antioxidant, anti-inflammatory and cytotoxic activities of Tunisian medicinal *Tamarix africana* Poir. *International Journal of Environmental Health Research* **34**(5):2366–2377 DOI 10.1080/09603123.2023.2249424.
- Karker M, Oueslati S, Falleh H, Msaada K, Legault J, Abdelly C, Pichette A, Ksouri R. 2024b.** Phytochemical investigation, antioxidant, anti-inflammatory and cytotoxic activities of Tunisian medicinal *Tamarix africana* Poir. *International Journal of Environmental Health Research* **34**(5):2366–2377 DOI 10.1080/09603123.2023.2249424.
- Kostina MV, Barabanshchikova NS, Pavlova IV. 2020.** Structural and lithological features of the shoot systems of species of the genus *Tamarix* L. (Tamaricaceae), which determine the adaptation of these species in the Moscow region. Bulletin of the Moscow Society of Nature Testers. *Department of Biology* **125**(6):21–32.
- Krylenko S, Krylenko V, Krylenko M. 2020.** On possibility of using *Tamarix* as indicator of the relief dynamics. *Bulletin of Science and Practice* **6**:57–69 DOI 10.33619/2414-2948/60/06.
- Lee SR, Gaskin JF, Kim YD. 2019.** Molecular diagnosis for a *Tamarix* species from two reclaimed lands along the Yellow Sea in Korea inferred from genome wide SNP markers. *Journal of Systematics and Evolution* **57**(3):247–255 DOI 10.1111/jse.12432.
- Lefahal M, Benahmed M, Louaar S, Zallagui A, Duddeck H, Akkal S. 2010.** Antimicrobial activity of *Tamarix gallica* L. extracts and isolated flavonoids. *Advances in Natural and Applied Sciences* **4**(3):289–293.
- Li F, Xie W, Ding X, Xu K, Fu X. 2024.** Phytochemical and pharmacological properties of the genus *Tamarix*: a comprehensive review. *Archives of Pharmacal Research* **47**(5):410–441.
- Mahfoudhi A, Grosso C, Gonçalves RF, Khelifi E, Hammami S, Achour S, Trabelsi-Ayadi M, Valente P, Andrade PB, Mighri Z. 2016.** Evaluation of antioxidant, anticholinesterase, and antidiabetic potential of dry leaves and stems in *Tamarix aphylla* growing wild in Tunisia. *Chemistry & Biodiversity* **13**(12):1747–1755 DOI 10.1002/cbdv.201600157.

- Nicolini F, Burmistrova O, Marrero MT, Torres F, Hernández C, Quintana J, Estevez F. 2014. Induction of G2/M phase arrest and apoptosis by the flavonoid tamarixetin on human leukemia cells. *Molecular Carcinogenesis* 53(12):939–950 DOI 10.1002/mc.22055.
- Nisar J, SM Ali Shah, Ayaz S, Akram M, Rashid A, Mustafa I, Nisar Z. 2023. In vitro comparative evaluation of *Tamarix gallica* extracts for antioxidant and antidiabetic activity. *Experimental Biology and Medicine* 248(3):253–262 DOI 10.1177/15353702221139208.
- Orabi KY, Abaza MS, El Sayed KA, Elnagar AY, Al-Attiyah R, Guleri RP. 2013. Selective growth inhibition of human malignant melanoma cells by syringic acid-derived proteasome inhibitors. *Cancer Cell International* 13(1):82 DOI 10.1186/1475-2867-13-82.
- Ovchinnikov PN. 1981. Flora of the Tajik SSR. —L. Izd. Nauka, volume VI. 594–596.
- Paul S, Naher K, Yeasmin S, Acharrya RN, Golder M, Ahmed KS, Hossain H, Dev S, Das AK. 2022. Phytochemicals, antioxidant, and chemopreventive potential of *Tamarix Indica* leaf extracts. *Journal of Herbs, Spices & Medicinal Plants* 28(2):143–159 DOI 10.1080/10496475.2021.2015506.
- Pavlov NV. 1963. *Flora of Kazakhstan*. Alma-Ata, 6. Kazakh SSR, Alma-Ata, 72–73.
- Peydayesh M, Raisi M, Kaykavousi K, Gharaghani MA, Abdollahpour-Alitappeh M, Mosazade F, Seifalianf A, Khatami M. 2020. The inhibitory effect of *Tamarix hispida* mediated silver nanoparticles on Cyclin D1 protein expression of human cancer cells line. *Inorganic and Nano-Metal Chemistry* 50(11):1144–1149 DOI 10.1080/24701556.2020.1735432.
- Plantarium. 2025. *Tamarix karelinii* Bunge. Available at <https://www.plantarium.ru/page/view/item/37282.html> (accessed on 15 April 2025).
- Plantarium plant Determinant Online. 2024. Available at <https://www.plantarium.ru/page/view/item/37287.html> (accessed on 14 May 2024).
- Qadir MI, Abbas K, Hamayun R, Ali M. 2014. Analgesic, anti-inflammatory and anti-pyretic activities of aqueous ethanolic extract of *Tamarix aphylla* L.(Saltcedar) in mice. *Pakistan Journal of Pharmaceutical Sciences* 27(6):1985–1988.
- Qaiser M. 1983. The genus tamarix (Tamaricaceae) in Pakistan. *The Iranian Journal of Botany* 2(1):21–68.
- Qaiser M, Perveen A. 2004. Pollen flora of Pakistan-XXXVII. Tamaricaceae. *Pakistan Journal of Botany* 36:1–18.
- Rachkovskaya EI, Khramtsov VN, Volkova EA. 2003. *Botanical Geography of Kazakhstan and Middle Asia (within the desert region)*. St. Petersburg: Russian Academy of Sciences.
- Ren X, Bao Y, Zhu Y, Liu S, Peng Z, Zhang Y, Zhou G. 2019. Isorhamnetin, Hispidulin, and Cirsimaritin Identified in *Tamarix ramosissima* Barks from Southern Xinjiang and Their Antioxidant and Antimicrobial Activities. *Molecules* 24(3):390 DOI 10.3390/molecules24030390.
- Romeilah RM, El-Beltagi HS, Shalaby EA, Younes KM, Hani EL, Rajendrasozhan S, Mohamed H. 2021. Antioxidant and cytotoxic activities of *Artemisia monosperma*

- L. and *Tamarix aphylla* L. essential oils. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* **49**(1):12233–12233 DOI 10.15835/nbha49112233.
- Rotaru LT, Varut RENATA. 2020. Antibacterial property and molecular docking studies of *Tamarix ramosissima* plant extract. *Journal of Science and Arts* **20**(1):135–142.
- Rotaru LT, Varut RM, Covei MB, Costache II, Novac M, Nicolaescu O, Florescu C, Petrica A, Kostici R, Ciobanu D. 2018. Determination of antioxidant components and activity of *Tamarix ramosissima* comparative with *vaccinium myrtillus* on streptozotocin-diabetic mice. *Revista de Chimie* **69**(7):1860–1865 DOI 10.37358/RC.18.7.6432.
- Royal Botanic Gardens Kew. 2024. Plants of the World Online. Available at <https://powo.science.kew.org/>.
- Samejo MQ, Bughio SH, Memon S, Khan H, Memon GZ, Memon NN. 2023. Phenolic acid profiling and antioxidant activity of *Tamarix dioica* leaves, stem and flowers. *Pakistan Journal of Scientific & Industrial Research Series B: Biological Sciences* **66**(2):120–124.
- Sehrawat A, Sultana S. 2006. Evaluation of possible mechanisms of protective role of *Tamarix gallica* against DEN initiated and 2-AAF promoted hepatocarcinogenesis in male Wistar rats. *Life Sciences* **79**(15):1456–1465 DOI 10.1016/j.lfs.2006.04.009.
- Shagholi T, Keshavarzi M, Sheidai M. 2020. Pollen morphology of some *Tamarix* (Tamaricaceae) species of Iran. *Acta Botanica Hungarica* **62**:151–161 DOI 10.1556/034.62.2020.1-2.9.
- Shang C-B, Lowry II P. 2007. Araliaceae, flora of China. *Flora of China* **13**:59–65.
- Sheidai M, Teymoori H, Noormohammadi Z, Mehrabian AR, Koohdar F, Ghasemzadeh-Baraki S. 2018. Species delimitation in the genus *Tamarix*: morphological and molecular data. *Phytotaxa* **343**(2):101–115 DOI 10.11646/PHYTOTAXA.343.2.1.
- Shishkin BK. 1949. *Flora of the USSR, volume 15: malvales, parietales, myrtiflorae*. Leningrad: Akademia Nauk, 565.
- Souddi M, Bendi Djelloul BG. 2020. Phylogenetic relationships of *Tamarix* species (Tamaricaceae) based on I.T.S sequences: a bioinformatic simulation for Algeria. *International Journal of Agriculture and Biosciences* **9**(5):226–231.
- Stankov SS, Taliev VI. 1949. The determinant of higher plants of the European part of the USSR. –Ed. reprint. and an additional one. Moscow: Soviet Science.
- Sultanova NA, Abilov ZA, Omurkamzinova VB, Chaudri IM. 2002. Flavonoids of the aerial part of *Tamarix hispida*. DOI 10.1023/A:1015706520986.
- Sultanova NA, Abilov ZA, Shul’Ts EE, Omurkamzinova VB. 2004a. Biologically active compounds from *Tamarix hispida*. II. DOI 10.1023/B:CONC.0000033946.96188.86.
- Sultanova NA, Abilov ZA, Umbetova AK, Choudhary MI. 2013. Biologically active Terpenoids from *Tamarix* species. *Eurasian Chemico-Technological Journal* **15**(3):219–226 DOI 10.18321/ectj225.
- Sultanova N, Makhmoor T, Abilov ZA, Parween Z, Omurkamzinova VB, ur Rahman A, Choudhary MI. 2001. Antioxidant and antimicrobial activities of *Tamarix ramosissima*. *Journal of Ethnopharmacology* **78**:201–205 DOI 10.1016/s0378-8741(01)00354-3.

- Sultanova N, Makhmoor T, Yasin A, Abilov ZA, Omurkamzinova VB, Atta-ur Rahman, Choudhary MI. 2004b. Isotamarixen - a new antioxidant and prolyl endopeptidase-inhibiting triterpenoid from *Tamarix hispida*. *Planta Medica* 70(1):65–67 DOI 10.1055/s-2004-815458.
- Terrones A, van der Bank M, Moreno J, Juan A. 2022. DNA barcodes and microsatellites: how they complement for species identification in the complex genus *Tamarix* (Tamaricaceae). *Journal of Systematics and Evolution* 60(5):1140–1157 DOI 10.1111/jse.12830.
- Terzoli S, Cattani E, Sabatti M, Valentini R, Zilberstein A, Kuzminsky E. 2013. Primer Note: A novel set of EST-SSR markers in *Tamarix*: a resource to characterize this genus. *Silvae Genetica* 62:104–109 DOI 10.1515/sg-2013-0013.
- The Ferns Database. 2025. *Tamarix arceuthoides*. Available at <https://temperate.theferns.info/viewtropical.php?id=Tamarix+arceuthoides> (accessed on 15 April 2025).
- Tropicos. 2024. Missouri Botanical Garden. Available at <https://tropicos.org> (accessed on 30 May 2024).
- Ullah R, Tariq SA, Khan N, Sharif N, ZU Din, Mansoor K. 2017. Antihyperglycemic effect of methanol extract of *Tamarix aphylla* L. Karst (Saltcedar) in streptozocin–nicotinamide induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine* 7(7):619–623 DOI 10.1016/j.apjtb.2017.06.005.
- Umbetova A, Choudhary M, Sultanova N, Burasheva G, Abilov Z. 2005. Flavonoids of Plants from the Genus *Tamarix*. *Chemistry of Natural Compounds* 41(6):628–630 DOI 10.1007/s10600-006-0023-8.
- Umbetova AK, Esirkegenova SZ, Chaudri IM, Omurkamizinov VB, Abilov ZA. 2004a. Flavonoids of plants from the genus *Tamarix*. *Chemistry of Natural Compounds* 40(3):297–298 DOI 10.1023/B:CONC.0000039149.63358.b2.
- Umbetova AK, Esirkegenova SZ, Chaudri IM, Omurkamizinov VB, Abilov ZA. 2004b. Triterpenoids from plants of the genus *Tamarix*. *Chemistry of Natural Compounds* 40(3):332–335 DOI 10.1007/s10600-006-0113-7.
- Utkin LM. 1966. The structure of tamarixin. *Chem Nat Compd* 2:128–130 DOI 10.1007/BF01134230.
- Venturella G, Baum B, Mandracchia G. 2007. The genus *Tamarix* (Tamaricaceae) in Sicily: First contribution. *Flora Mediterranea* 17:25–46.
- Venturella G, Gargano ML, Mandracchia G. 2012. First record of *Tamarix meyeri* (Tamaricaceae) for western Europe. *Plant Biosystems - an International Journal Dealing with All Aspects of Plant Biology* 146(2):484–489 DOI 10.1080/11263504.2011.629010.
- Villar JL, Alonso MÁ, Crespo MB. 2022. Synopsis of the genus *Tamarix* (Tamaricaceae) in the Iberian Peninsula and Balearic Islands. *Anales del Jardín Botánico de Madrid* 79(2):e132 DOI 10.3989/ajbm.2639.
- Villar JL, Juan A, Alonso MÁ, Crespo MB. 2014. Type specimens of *Tamarix* (Tamaricaceae) described by Josef Franz Freyn in. *Phytotaxa* 172(3):289–292 DOI 10.11646/phytotaxa.172.3.11.

- Wang X, Cao Q, Wei Y. 2023. The complete chloroplast genome sequence of *Tamarix arceuthoides* Bunge and *Tamarix ramosissima* Ledeb, (Tamaricaceae). *Mitochondrial DNA. Part B, Resources* 8(5):541–545 DOI 10.1080/23802359.2023.2209215.
- Xiao LP, Shi ZJ, Xu F, Sun RC. 2013. Characterization of lignins isolated with alkaline ethanol from the hydrothermal pretreated *Tamarix ramosissima*. *BioEnergy Research* 6:519–532 DOI 10.1007/s12155-012-9266-3.
- Xing Y, Liao J, Tang Y, Zhang P, Tan C, Ni H, Wu X, Li N, Jia X. 2014. ACE and platelet aggregation inhibitors from *Tamarix hohenackeri* Bunge (host plant of Herba Cistanches) growing in Xinjiang. *Pharmacognosy Magazine* 10(38):111–117 DOI 10.4103/0973-1296.131021.
- Yao Y, Jiang CS, Sun N, Li WQ, Niu Y, Han HQ, Miao ZH, Zhao XX, Zhao J, Li J. 2017. Tamaractam, a new bioactive lactam from *Tamarix ramosissima*, induces apoptosis in rheumatoid arthritis fibroblast-like synoviocytes. *Molecules (Basel, Switzerland)* 22(1):96 DOI 10.3390/molecules22010096.
- Yusufoglu HS, Alqasoumi SI. 2011. *Anti-inflammatory and wound healing activities of herbal gel containing an antioxidant Tamarix aphylla leaf extract*. Faisalabad, Pakistan: Asian Network for Scientific Information.
- Zakhidov K. 1991. *The treasury of healing remedies*. Dushanbe: Irfon.
- Zar Kalai F, Oueslati S, Dakhlaoui S, Hammami M, Msaada K, Ksouri R. 2024. Chemical profiling of maceration and decoction of *Tamarix gallica* L. organs and *in vitro* biological properties. *International Journal of Environmental Health Research* 34(6):2517–2528 DOI 10.1080/09603123.2023.2256679.
- Zazharskyi VV, Davydenko PO, Kulishenko O, Borovik IV, Zazharska NM, Brygadyrenko V. 2020. *Antibacterial and fungicidal activities of ethanol extracts of 38 species of plants*. Dnipro, Ukraine: Regulatory Mechanisms in Biosystems DOI 10.15421/012037.
- Zhang JW, D’Rozario A, Duan SM, Wang XY, Liang XQ, Pan BR. 2018. Epidermal characters of *Tamarix* L.(Tamaricaceae) from Northwest China and their taxonomic and palaeogeographic implications. *Journal of Palaeogeography* 7(2):179–196 DOI 10.1016/j.jop.2018.01.003.
- Zhang Y, Tu P. 1994. Chemical constituents of *Tamarix ramosissima*. *Chinese Traditional and Herbal Drugs* 25(1):22–24.
- Zhang DY, Yang WK, Pan BR, Yin LK. 2003. Characters of *Tamarix hispida* Willd communities and its ecological & physiological adaptation. *Journal of Desert Research* 23(4):446.
- Zhumagaliyeva S, Amanzholkyzy A, Sultanova N, Abilov Z. 2021. Use of ultrasound for extraction of biologically active substances from *tamarix hispida* willd. *Chemistry of Plant Raw Material* 3:283–289 DOI 10.14258/jcprm.2021038695.
- Zieliński J. 1994. *Tamarix* L.–Pp. 30–34. In: Browicz K, ed. *Chorology of trees and shrubs in south-west Asia and adjacent regions* 10. Ponzan: Bogucki Wydawnictwo Naukowe.