

# Relationship between triterpenoid anticancer drug resistance, autophagy, and caspase-1 in adult T-cell leukemia

Tsukasa Nakanishi, Yuan Song, Cuiying He, Duo Wang, Kentaro Morita, Junichi Tsukada, Tamotsu Kanazawa, Yasuhiro Yoshida

We previously reported that the inflammasome inhibitor cucurbitacin D (CuD) induces apoptosis in human leukemia cell lines. Here, we investigated the effects of CuD and a B cell lymphoma extra large (Bcl-xL) inhibitor on autophagy in peripheral blood lymphocytes (PBL) isolated from adult T cell leukemia (ATL) patients. CuD induced PBL death in patients but not in healthy donors. This effect was not significantly inhibited by treatment with rapamycin or 3-methyladenine (3-MA). The Bcl-xL inhibitor Z36 induced death in primary cells from ATL patients including that induced by CuD treatment, effects that were partly inhibited by 3-MA. Similarly, cell death induced by the steroid prednisone was enhanced in the presence of Z36. A western blot analysis revealed that Z36 also promoted CuD-induced poly ADP ribose polymerase cleavage. Interestingly, the effects of CuD and Z36 were attenuated in primary ATL patient cells obtained upon recurrence after umbilical cord blood transplantation, as compared to those obtained before chemotherapy. Furthermore, cells from this patient expressed high level of caspase-1, and treatment with caspase-1 inhibitor enhanced CuD-induced cell death. Taken together, these results suggest that rescue from resistance to steroid drugs can enhance chemotherapy, and that caspase-1 is a good marker for drug resistance in ATL patients.

Relationship between triterpenoid anticancer drug resistance, autophagy, and capase-1 in adult  
T-cell leukemia

Tsukasa Nakanishi<sup>1,2</sup>, Yuan Song<sup>1,3,\*</sup>, Cuiying He<sup>1,\*</sup>, Duo Wang<sup>1,\*</sup>, Kentaro Morita<sup>1</sup>,  
Junichi Tsukada<sup>2</sup>, Tamotsu Kanazawa<sup>1</sup>, and Yasuhiro Yoshida<sup>1</sup>

<sup>1</sup>Department of Immunology and Parasitology; <sup>2</sup>Department of Hematology, University of  
Occupational and Environmental Health, Japan, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-  
8555, Japan.

<sup>3</sup> Department of Clinical Laboratory, the Fourth Hospital of Hebei Medical University, No. 169  
Tian Shan Street, Shijiazhuang, 050035, China.

Corresponding author: Yasuhiro Yoshida, PhD

Tel: +81-93-691-7431; Fax: +81-93-602-4488;

E-mail: [freude@med.uoeh-u.ac.jp](mailto:freude@med.uoeh-u.ac.jp)

17 Running title: Cucurbitacin D-induced cell death in ATL cells

18

19 Word count; 4014 with references (references 1068)

# **ABSTRACT**

We previously reported that the inflammasome inhibitor cucurbitacin D (CuD) induces apoptosis in human leukemia cell lines. Here, we investigated the effects of CuD and a B cell lymphoma extra large (Bcl-xL) inhibitor on autophagy in peripheral blood lymphocytes (PBL) isolated from adult T cell leukemia (ATL) patients. CuD induced PBL death in patients but not in healthy donors. This effect was not significantly inhibited by treatment with rapamycin or 3-methyladenine (3-MA). The Bcl-xL inhibitor Z36 induced death in primary cells from ATL patients including that induced by CuD treatment, effects that were partly inhibited by 3-MA. Similarly, cell death induced by the steroid prednisone was enhanced in the presence of Z36. A western blot analysis revealed that Z36 also promoted CuD-induced poly ADP ribose polymerase cleavage. Interestingly, the effects of CuD and Z36 were attenuated in primary ATL patient cells obtained upon recurrence after umbilical cord blood transplantation, as compared to those obtained before chemotherapy. Furthermore, cells from this patient expressed high level of caspase-1, and treatment with caspase-1 inhibitor enhanced CuD-induced cell death. Taken together, these results suggest that rescue from resistance to steroid drugs can enhance chemotherapy, and that caspase-1 is a good marker for drug resistance in ATL patients.

# INTRODUCTION

Adult T cell leukemia (ATL) is an aggressive and malignant blood disease. There are many ATL patients in the Kyushu area of Japan who have very poor prognosis, even with intensive chemotherapy (Iwanaga et al. 2012) owing to rapid disease progression and frequent infectious complications. The median survival of patients in Japan is six months (Bazarbachi et al. 2004). Several drugs including synthetic glucocorticoids are used in chemotherapy; however, the treatment is not completely effective due to the development of drug resistance.

Cucurbitacin (Cu) is a type of cucurbit belonging to the triterpenoid group of molecules with a steroid structure (Fig. 1A). We previously reported that CuD has antitumorigenic activity (Takahashi et al. 2009) and showed that it acts as a proteasome inhibitor and induces apoptosis in leukemia cell lines with differential sensitivity (Ding et al. 2011). Glucocorticoid steroid hormones are universally used in the treatment of acute lymphoblastic leukemia, but the development of resistance is associated with poor prognosis (Den Boer et al. 2003). However, the mechanistic basis for the development of resistance to steroid-type molecules by ATL cells is not well understood.

Autophagy is an evolutionarily conserved mechanism for the degradation of cellular waste components and is critical for the maintenance of homeostasis (Puleston & Simon 2014).

Defective autophagy is associated with diseases such as cancer; it has been implicated in tumor development and survival in blood diseases (Bosnjak et al. 2014; Nencioni et al. 2013; Pierdominici et al. 2014). Some studies have reported the induction of cell death via inhibition of autophagy in treatments against T cell leukemia (Jia et al. 1997; Nahimana et al. 2009; Wang et al. 2014), but there are no similar reports for ATL, and the extent to which autophagy is involved in the induction in chemically induced cell death in ATL is unclear.

During the preparation of this communication, Paugh et al. reported caspase-1-related glucocorticoid resistance in leukemia cells (Paugh et al. 2015). In this study, we investigated the antitumorigenic effects of another steroid-structured agent, CuD, in primary ATL cells with respect to autophagy involvement and drug resistance.

# **MATERIALS AND METHODS**

## **Reagents and antibodies**

CuD was purchased from Nacalai Tesque (Kyoto, Japan) and dissolved in 10% ethanol to a stock concentration of 1 mg/mL. Z36 was obtained from Enzo Life Sciences (Farmingdale, NY). The autophagy inhibitor 3-MA was purchased from Sigma (St. Louis, MO) and was diluted in phosphate-buffered saline. The anti-poly ADP ribose polymerase (PARP) antibody (9542S) was purchased from Cell Signaling Technology (Danvers, MA). The  $\beta$ -actin (A5441) antibody was obtained from Sigma; the anti-microtubule-associated protein 1A/1B-light chain 3 (LC3)antibody (M186-3) was purchased from MBL (Nagoya, Japan). Antibodies against  $\alpha$ -tubulin (H-300; sc-5546) and caspase-1 (M-20; sc-514) were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Caspase-1 inhibitor (#400010) was obtained from Calbiochem (San Diego, CA).

## **Preparation of human peripheral blood lymphocytes (PBLs)**

PBLs were obtained from ATL patients or healthy donors following the ethical guidelines of University Occupational and Environmental Health, Japan (No H23-061); donor information is summarized in Table1. Cells were harvested three times from ATL patient number 1: once

before chemotherapy; after three rounds of THP-COP chemotherapy [30 mg/m<sup>2</sup>pirarubicin (THP-ADR) for 1 day, 600 mg/m<sup>2</sup> cyclophosphamide for 1 day, 1 mg/m<sup>2</sup>vincristine for 1 day, and 30 mg/m<sup>2</sup> prednisolone for 5 days] but before umbilical cord blood transplantation; and at the time of recurrence 96 days after the transplantation. Cells were isolated using Lymphoprep (Axis Shield, Oslo, Norway) according to the manufacturer's protocol. Briefly, 6 ml of blood were overlaid on 3 ml of Lymphoprep reagent and centrifuged at  $600 \times g$  for 30 min. After removing the PBL layer, cells were washed with PBS, resuspended in medium, and counted. Prepared cells were cultured as described below.

## **Proliferation assay**

Cell proliferation was assessed with the Cell Titer-Glo luminescent cell viability assay (Promega, Madison, WI) as previously described (Ding et al. 2011). Briefly, cells were seeded in 96-well plates at  $1 \times 10^4$ /well, and then treated with Z36 and/or CuD in the presence or absence of 3-MA. A 10- $\mu$ l volume of assay reagent was added to each well, and fluorescence was measured with a luminometer (Luminescencer-JNR-II; ATTO, Tokyo, Japan).

## **Western blotting**

Cells were lysed with radioimmunoprecipitation assay buffer (RIPA) (Yoshida et al. 2009) to



obtain whole-cell extracts. Equivalent amounts of protein (10  $\mu$ g) were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, then transferred to and immobilized on nitrocellulose membranes (Amersham, Buckinghamshire, UK) that were probed with the appropriate primary and secondary antibodies. Protein bands were detected using the ImmunoStar LD detection system (Wako, Osaka, Japan), and the signal intensity was quantified using ImageJ software (National Institutes of Health, Bethesda, MD). Expression levels of target proteins were normalized to that of  $\beta$ -actin or  $\alpha$ -tubulin.

# **Statistical analysis**

Results are expressed as mean  $\pm$  SD and inter-group differences were evaluated by analysis of variance with Scheffe's post-hoc analysis. A P value  $< 0.05$  was considered statistically significant.

# RESULTS

## CuD induces death in primary cells from ATL patients

We examined the effects of CuD on primary ATL patient cells. The cell viability assay confirmed that CuD was not toxic to PBLs from healthy donors (Fig. 1B), as we have previously reported (Ding et al. 2011). Activated PBL cells by mitogen were also unaffected by CuD (Supplemental Fig. S1A). In contrast, CuD markedly reduced the viability of primary ATL patient cells (Fig. 1C). To examine the involvement of autophagy, we treated cells with rapamycin, a mammalian target of rapamycin inhibitor and autophagy inducer, and found that it had no effect on CuD-induced cell death (Fig. 1C). Similarly, treatment with the autophagy inhibitor 3-MA did not affect CuD-induced antitumorigenic activity in primary ATL patient cells, while 3-MA treatment alone slightly decreased cell viability (Fig. 1D).

## Z36 inhibits proliferation and enhances CuD-induced cell death in primary ATL patient cells

The effects of the B cell lymphoma (Bcl)-2 inhibitor Z36 on ATL patient cells before chemotherapy were examined. Low concentrations (3 and 5  $\mu$ M) of Z36 were not toxic to PBLs from healthy donors; however, cell viability was decreased at a high concentration (7.5  $\mu$ M) (Fig.

2A and Supplemental Fig. S2B). Application of Z36 inhibited the proliferation of ATL patient cells in a dose-dependent manner (Fig. 2B, from Patient 1; other data shown in Supplemental Fig. S2A) and enhanced CuD-induced cell death, which was partly abrogated by treatment with 3-MA (Fig. 2C and Supplemental Fig. S3). These results were confirmed in 10 patients and three healthy controls, although only a representative case is shown. Z36 had similar effects on prednisolone-treated ATL patient cells (Fig. 2D), suggesting that cell death induced by steroid-like agents acts via a different mechanism than Z36-induced cell death.

### **Autophagy negatively regulates CuD- or Z36-induced PARP cleavage**

We examined the levels of cleaved PARP and LC3-II proteins by western blotting, and found that Z36 induced PARP cleavage in primary ATL patient cells, contrary to previous reports (Lin et al. 2009), while Z36 treatment increased LC3-II expression in these cells, as previously reported (Fig. 3A). Treatment with the autophagy inhibitor 3-MA reduced levels of cleaved PARP proteins induced by Z36 and CuD (Fig. 3A). Co-treatment with CuD and Z36 did not increase LC3-II expression relative to cells treated with Z36 only (Fig. 3B).

### **CuD and Z36 have limited effects in primary ATL patient cells after disease recurrence**

As expected, treatment with CuD (0.19  $\mu$ M) and/or Z36 (7.5  $\mu$ M) and/or 3-MA (10 mM) had no

149 effect on primary ATL patient cell viability after chemotherapy, since there were no leukemia  
 150 cells (Fig. 4A). Furthermore, PARP cleavage was not induced by CuD and/or Z36 in these cells  
 151 after chemotherapy (Fig. 4B, middle four lanes). Interestingly, the effects of CuD and Z36 were  
 152 attenuated in primary ATL patient cells upon recurrence after umbilical cord blood  
 153 transplantation (Fig. 4B, right four lanes) as compared to the post-chemotherapy state (Fig. 4B,  
 154 C). Increased caspase-1 levels have been linked to glucocorticoid resistance in leukemia cells  
 155 (Paugh et al. 2015). We found here that caspase-1 levels were increased after chemotherapy,  
 156 implying the development of drug resistance. Consistent with this supposition, the effects of  
 157 combined CuD and Z36 treatment on primary ATL patient cells were enhanced in the presence  
 158 of a caspase-1 inhibitor (Fig. 4E).

# DISCUSSION

There is growing interest in natural products as a source of new drugs for cancer therapy. We previously reported that one such product, CuD, could induce human T-cell leukemia cell death in vitro via proteasome inhibition (Ding et al. 2011). In the present study, we demonstrated that CuD inhibits the proliferation of primary ATL patient cells. This is the first evidence of an inhibitory effect by a member of Cucurbitaceae on tumor cells. On the other hand, CuD showed no toxicity to peripheral blood mononuclear cells from healthy donors or primary cells from mouse (Song et al. 2013), and it was in fact found to enhance PBL proliferation in the current study. Conversely, CuD strongly induced primary ATL patient cell death, as determined by the cell viability assay and by direct counting following CuD treatment (Supplemental Fig. S4). As expected, primary ATL patient cells after chemotherapy were unaffected by CuD or Z36 treatment, but cells obtained upon recurrence showed reduced responsiveness to CuD and Z36 after umbilical cord transplantation as compared to those obtained from the first occurrence of the disease. Given that recurrence after transplantation is associated with poor prognosis (Hishizawa et al. 2010), our results suggest that these cells developed drug resistance. Caspase-1 level is a marker for glucocorticoid (e.g., prednisolone) resistance in leukemia cells (Paugh et al. 2015); indeed, we observed an upregulation of caspase-1 expression in primary ATL patient cells after recurrence. However, caspase-1 level was unexpectedly elevated after transplantation,

178 suggesting that drug resistance was induced at this stage.

179     Given that drug resistance influences chemotherapeutic trials, developing novel treatment  
180 strategies and identifying new target molecules represent an ongoing challenge. The role of  
181 autophagy in hematological malignancies has been widely investigated (Macintosh & Ryan  
182 2013). Autophagy inhibitors have been shown to overcome cell resistance to anti-cancer drugs  
183 (Carew et al. 2007); autophagy induction is required to overcome glucocorticoid resistance in  
184 acute lymphoblastic leukemia cells (Bonapace et al. 2010). Bortezomib is a proteasome inhibitor  
185 and apoptosis inducer (Pei et al. 2003) that is used clinically for the treatment of multiple  
186 myeloma (Sonneveld et al. 2012). It was previously shown that bortezomib treatment in the  
187 presence of autophagy inhibitors enhances apoptosis (Hoang et al. 2009; Jia et al. 2012). This  
188 finding is consistent with our observation that the effects of the proteasome inhibitor CuD were  
189 enhanced in the presence of 3-MA. On the other hand, autophagy has been shown to enhance  
190 apoptosis induction in blood malignancies (Fang et al. 2012), which is similar to our results  
191 obtained by co-treatment with Z36, a Bcl-xL inhibitor and autophagy-specific inducer in HeLa  
192 cells or THP-1 cells (Casalino-Matsuda et al. 2015). Indeed, we found that Z36 treatment  
193 increased primary ATL patient cell death by inducing not only LC3-II expression but also PARP  
194 cleavage, suggesting that it functions as an apoptosis inducer in ATL cells, which has not been  
195 previously reported. Z36 was only found to suppress PBL proliferation at a concentration of 3

196  $\mu\text{M}$  or greater, implying that lymphoid cells are more sensitive to Z36 than healthy donor cells.

197 In this scenario, autophagy protects against chemically induced apoptosis.

198 In conclusion, we described the distinct effects of two Bcl-xL inhibitors, which were

199 abolished in the presence of a caspase-1 inhibitor. Based on these findings, acquired drug

200 resistance should be considered in the context of the type of cell death that is induced by the

201 levels of specific molecules such as caspase-1.

202 \*Authorship

203 Yuan Song, Cuiying He, Duo Wang contributed equally to this paper.

204



# REFERENCES

- Bazarbachi A, Ghez D, Lepelletier Y, Nasr R, de The H, El-Sabban ME, and Hermine O. 2004. New therapeutic approaches for adult T-cell leukaemia. *The lancet oncology* 5:664-672. 10.1016/S1470-2045(04)01608-0
- Bonapace L, Bornhauser BC, Schmitz M, Cario G, Ziegler U, Niggli FK, Schafer BW, Schrappe M, Stanulla M, and Bourquin JP. 2010. Induction of autophagy-dependent necroptosis is required for childhood acute lymphoblastic leukemia cells to overcome glucocorticoid resistance. *The Journal of clinical investigation* 120:1310-1323. 10.1172/JCI39987
- Bosnjak M, Ristic B, Arsikin K, Mircic A, Suzin-Zivkovic V, Perovic V, Bogdanovic A, Paunovic V, Markovic I, Bumbasirevic V, Trajkovic V, and Harhaji-Trajkovic L. 2014. Inhibition of mTOR-dependent autophagy sensitizes leukemic cells to cytarabine-induced apoptotic death. *PloS one* 9:e94374. 10.1371/journal.pone.0094374
- Carew JS, Nawrocki ST, Kahue CN, Zhang H, Yang C, Chung L, Houghton JA, Huang P, Giles FJ, and Cleveland JL. 2007. Targeting autophagy augments the anticancer activity of the histone deacetylase inhibitor SAHA to overcome Bcr-Abl-mediated drug resistance. *Blood* 110:313-322. 10.1182/blood-2006-10-050260
- Casalino-Matsuda SM, Nair A, Beitel GJ, Gates KL, and Sporn PH. 2015. Hypercapnia Inhibits

Autophagy and Bacterial Killing in Human Macrophages by Increasing Expression of Bcl-2 and Bcl-xL. *J Immunol* 194:5388-5396. 10.4049/jimmunol.1500150

Den Boer ML, Harms DO, Pieters R, Kazemier KM, Gobel U, Korholz D, Graubner U, Haas RJ, Jorch N, Spaar HJ, Kaspers GJ, Kamps WA, Van der Does-Van den Berg A, Van Wering ER, Veerman AJ, and Janka-Schaub GE. 2003. Patient stratification based on prednisolone-vincristine-asparaginase resistance profiles in children with acute lymphoblastic leukemia. *J Clin Oncol* 21:3262-3268. 10.1200/JCO.2003.11.031

Ding N, Yamashita U, Matsuoka H, Sugiura T, Tsukada J, Noguchi J, and Yoshida Y. 2011. Apoptosis induction through proteasome inhibitory activity of cucurbitacin D in human T-cell leukemia. *Cancer* 117:2735-2746. 10.1002/cncr.25711

Fang J, Rhyasen G, Bolanos L, Rasch C, Varney M, Wunderlich M, Goyama S, Jansen G, Cloos J, Rigolino C, Cortelezzi A, Mulloy JC, Oliva EN, Cuzzola M, and Starczynowski DT. 2012. Cytotoxic effects of bortezomib in myelodysplastic syndrome/acute myeloid leukemia depend on autophagy-mediated lysosomal degradation of TRAF6 and repression of PSMA1. *Blood* 120:858-867. 10.1182/blood-2012-02-407999

Hishizawa M, Kanda J, Utsunomiya A, Taniguchi S, Eto T, Moriuchi Y, Tanosaki R, Kawano F, Miyazaki Y, Masuda M, Nagafuji K, Hara M, Takanashi M, Kai S, Atsuta Y, Suzuki R, Kawase T, Matsuo K, Nagamura-Inoue T, Kato S, Sakamaki H, Morishima Y, Okamura

- J, Ichinohe T, and Uchiyama T. 2010. Transplantation of allogeneic hematopoietic stem cells for adult T-cell leukemia: a nationwide retrospective study. *Blood* 116:1369-1376. 10.1182/blood-2009-10-247510
- Hoang B, Benavides A, Shi Y, Frost P, and Lichtenstein A. 2009. Effect of autophagy on multiple myeloma cell viability. *Molecular cancer therapeutics* 8:1974-1984. 10.1158/1535-7163.MCT-08-1177
- Iwanaga M, Watanabe T, and Yamaguchi K. 2012. Adult T-cell leukemia: a review of epidemiological evidence. *Frontiers in microbiology* 3:322. 10.3389/fmicb.2012.00322
- Jia L, Dourmashkin RR, Allen PD, Gray AB, Newland AC, and Kelsey SM. 1997. Inhibition of autophagy abrogates tumour necrosis factor alpha induced apoptosis in human T-lymphoblastic leukaemic cells. *British journal of haematology* 98:673-685.
- Jia L, Gopinathan G, Sukumar JT, and Gribben JG. 2012. Blocking autophagy prevents bortezomib-induced NF-kappaB activation by reducing I-kappaBalpha degradation in lymphoma cells. *PloS one* 7:e32584. 10.1371/journal.pone.0032584
- Lin J, Zheng Z, Li Y, Yu W, Zhong W, Tian S, Zhao F, Ren X, Xiao J, Wang N, Liu S, Wang L, Sheng F, Chen Y, Jin C, Li S, and Xia B. 2009. A novel Bcl-XL inhibitor Z36 that induces autophagic cell death in Hela cells. *Autophagy* 5:314-320.
- Macintosh RL, and Ryan KM. 2013. Autophagy in tumour cell death. *Seminars in cancer*

259 *biology* 23:344-351. 10.1016/j.semcancer.2013.05.006

260 Nahimana A, Attinger A, Aubry D, Greaney P, Ireson C, Thougard AV, Tjornelund J, Dawson

261 KM, Dupuis M, and Duchosal MA. 2009. The NAD biosynthesis inhibitor APO866 has

262 potent antitumor activity against hematologic malignancies. *Blood* 113:3276-3286.

263 10.1182/blood-2008-08-173369

264 Nencioni A, Cea M, Montecucco F, Longo VD, Patrone F, Carella AM, Holyoake TL, and

265 Helgason GV. 2013. Autophagy in blood cancers: biological role and therapeutic

266 implications. *Haematologica* 98:1335-1343. 10.3324/haematol.2012.079061

267 Paugh SW, Bonten EJ, Savic D, Ramsey LB, Thierfelder WE, Gurung P, Malireddi RK, Actis M,

268 Mayasundari A, Min J, Coss DR, Lauder milk LT, Panetta JC, McCorkle JR, Fan Y,

269 Crews KR, Stocco G, Wilkinson MR, Ferreira AM, Cheng C, Yang W, Karol SE,

270 Fernandez CA, Diouf B, Smith C, Hicks JK, Zanut A, Giordanengo A, Crona D, Bianchi

271 JJ, Holmfeldt L, Mullighan CG, den Boer ML, Pieters R, Jeha S, Dunwell TL, Latif F,

272 Bhojwani D, Carroll WL, Pui CH, Myers RM, Guy RK, Kanneganti TD, Relling MV,

273 and Evans WE. 2015. NALP3 inflammasome upregulation and CASP1 cleavage of the

274 glucocorticoid receptor cause glucocorticoid resistance in leukemia cells. *Nat Genet*

275 47:607-614. 10.1038/ng.3283

276 Pei XY, Dai Y, and Grant S. 2003. The proteasome inhibitor bortezomib promotes mitochondrial

injury and apoptosis induced by the small molecule Bcl-2 inhibitor HA14-1 in multiple myeloma cells. *Leukemia* 17:2036-2045. 10.1038/sj.leu.2403109

Pierdominici M, Barbati C, Vomero M, Locatelli SL, Carlo-Stella C, Ortona E, and Malorni W. 2014. Autophagy as a pathogenic mechanism and drug target in lymphoproliferative disorders. *FASEB journal* 28:524-535. 10.1096/fj.13-235655

Puleston DJ, and Simon AK. 2014. Autophagy in the immune system. *Immunology* 141:1-8. 10.1111/imm.12165

Song Y, Ding N, Kanazawa T, Yamashita U, and Yoshida Y. 2013. Cucurbitacin D is a new inflammasome activator in macrophages. *International immunopharmacology* 17:1044-1050. 10.1016/j.intimp.2013.10.003

Sonneveld P, Schmidt-Wolf IG, van der Holt B, El Jarari L, Bertsch U, Salwender H, Zweegman S, Vellenga E, Broyl A, Blau IW, Weisel KC, Wittebol S, Bos GM, Stevens-Kroef M, Scheid C, Pfreundschuh M, Hose D, Jauch A, van der Velde H, Raymakers R, Schaafsma MR, Kersten MJ, van Marwijk-Kooy M, Duehrsen U, Lindemann W, Wijermans PW, Lokhorst HM, and Goldschmidt HM. 2012. Bortezomib induction and maintenance treatment in patients with newly diagnosed multiple myeloma: results of the randomized phase III HOVON-65/ GMMG-HD4 trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 30:2946-2955. 10.1200/JCO.2011.39.6820

295 Takahashi N, Yoshida Y, Sugiura T, Matsuno K, Fujino A, and Yamashita U. 2009.

296 Cucurbitacin D isolated from *Trichosanthes kirilowii* induces apoptosis in human

297 hepatocellular carcinoma cells in vitro. *International immunopharmacology* 9:508-513.

298 10.1016/j.intimp.2009.01.006

299 Wang W, Zhou J, Shi J, Zhang Y, Liu S, Liu Y, and Zheng D. 2014. Human T-cell leukemia

300 virus type 1 Tax-deregulated autophagy pathway and c-FLIP expression contribute to

301 resistance against death receptor-mediated apoptosis. *J Virol* 88:2786-2798.

302 10.1128/JVI.03025-13

303 Yoshida Y, Liu J, Sugiura T, Ishidao T, Ueno S, Yanagita H, Fueta Y, Kunugita N, Hori H, and

304 Yamashita U. 2009. The indoor air pollutant 2-ethyl-hexanol activates CD4 cells.

305 *Chemico-biological interactions* 177:137-141. 10.1016/j.cbi.2008.10.053

306

307

# Figure legends

**Figure 1. CuD induces primary ATL patient cell death.** (A) Structures of CuD and prednisolone. (B, C) PBLs were freshly prepared from healthy donor (B) or ATL patients (C). Cells ( $1 \times 10^4$ /well) were treated with indicated concentrations of CuD or vehicle in the presence or absence of rapamycin (100  $\mu$ M) for 24 h. (D) PBLs ( $1 \times 10^4$ /well) from ATL patient 1 were treated with CuD (0.19  $\mu$ M) in the presence or absence of 3-MA (10  $\mu$ M) for indicated periods. Cell viability was assessed with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. \* $P < 0.05$  vs. vehicle without CuD treatment at each time point.

**Figure 2. Z36 inhibits proliferation and enhances CuD-induced death in primary ATL patient cells.** (A–D) PBLs ( $1 \times 10^4$ /well) from healthy donors (A) or ATL patients (B–D) were treated with indicated concentrations of Z36 or dimethyl sulfoxide (DMSO, a control solvent for Z36) for 12 or 24 h, or CuD (0.19  $\mu$ M) or prednisolone for 24 h.. (C) PBLs from ATL patients were pre-treated with 10 mM 3-MA or vehicle for 2 h, and then exposed to 0.19  $\mu$ M CuD, Z36 (7.5  $\mu$ M), or vehicle (Et-OH, 0.01%) for 24 h. (D) \* $P < 0.05$  vs. vehicle at each time point. Cell viability was determined using the Cell Titer-Glo assay. Results are expressed as mean activity

relative to controls and SD from quadruplicate cultures. \*P< 0.05 vs. vehicle; <sup>†</sup>P < 0.05 vs. treatment with CuD only; <sup>#</sup>P < 0.05 vs. CuD and Z36 co-treatment.

**Figure 3. Autophagy negatively regulates CuD- or Z36-induced PARP cleavage in ATL patient cells.** (A, B) PBLs ( $2 \times 10^5$ /well) from ATL patients were treated with CuD (0.19  $\mu$ M), Z36 (7.5  $\mu$ M), or vehicle in the presence or absence of 3-MA (10 mM) for 24 h. Whole cell extracts were used for western blot analysis of cleaved PARP (A ) and LC3-II (B) expression;  $\beta$ -actin or tubulin was used as a loading control. ET, ethanol; DM, DMSO.

**Figure 4. CuD and Z36 show limited effects on ATL patient-derived primary cells upon disease recurrence.** (A, C) PBLs ( $1 \times 10^4$ /well) from ATL patients after chemotherapy (A) or from ATL patients upon recurrence after umbilical cord blood transplantation (C) were treated with Z36 (7.5  $\mu$ M) or vehicle for 24 h. Cell viability was determined with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. (B) PBLs ( $1 \times 10^4$ /well) from a patient before therapy (left four lanes), after umbilical cord blood transplantation (middle four lanes), and after recurrence (right four lanes) were treated with CuD (0.19  $\mu$ M) and/or Z36 (7.5  $\mu$ M) or with vehicle for 24 h. Whole cell extracts were used for western blot analysis of cleaved PARP;  $\beta$ -actin was used as a loading control. M,



molecular marker. (D) Whole cell extracts from a patient before therapy (ori), after umbilical cord blood transplantation (Ch), and after recurrence (Rec) were analyzed for caspase-1 expression by western blotting;  $\beta$ -actin was used as a control. (E) ATL patient cells ( $1 \times 10^4$ /well) obtained upon recurrence after umbilical cord blood transplantation were treated with CuD (0.19  $\mu$ M) and Z36 (7.5  $\mu$ M) or vehicle for 24 h in the presence or absence of capase-1 inhibitor (20  $\mu$ M). Cell viability was determined with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. \* $P < 0.05$  vs. vehicle; # $P < 0.05$  vs. CuD and Z36 co-treatment.

# Supplementary Figure 1

PBL from a heathy donor ( $1 \times 10^4$ ) were treated with several concentrations of CuD or Z36, and stimulated with LPS (1  $\mu$ g/ml; left panel) or ConA (5  $\mu$ g/ml; right panel).

# Supplementary Figure 2

Peripheral blood lymphocytes (PBLs) from ATL, T cell lymphoma patients, or healthy donors were treated with Z36 at the indicated concentrations for 24 h. The viability of cells was determined and expressed as the mean activity  $\pm$  SD of quadruplicate wells. \* $P < 0.05$  vs. vehicle.

362 Supplementary Figure 3

363 Peripheral blood lymphocytes (PBLs) from ATL patients were treated with CuD (0.19  $\mu$ M)  
 364 and/or Z36 (7.5  $\mu$ M) for 24 h. The viability of cells was determined and expressed as the mean  
 365 activity  $\pm$  SD of quadruplicate wells. \*P < 0.05 vs. vehicle.

366

367 Supplementary Figure 4

368 PBLs ( $1 \times 10^4$ /well) from ATL patients were treated with 3-MA (10 mM) or PBS for 24 h,  
 369 and/or CuD (0.19  $\mu$ M). \*P < 0.05 vs. vehicle. Cell viability was determined using the Cell Titer-  
 370 Glo assay. Results are expressed as mean count relative to controls and SD from quadruplicate  
 371 cultures. \*P < 0.05 vs. vehicle; #P < 0.05 vs. treatment with CuD only.

372

# **Table 1**(on next page)

Characteristics of ATL cell lines, ATL patients, and healthy donors

PBLs were obtained from ATL patients or healthy donors as following ethics in University of Occupational and Environmental Health, Japan. Donor information is summarized in Table 1.

**Table 1.** Characteristics of ATL cell lines, ATL patients, and healthy donors

	Disease	Sex	Age (years)	HTLV-1	Tax	Lymphoid cells (%)	CuD treatment (inhibition to Ct, %)	CuD + Z36 treatment (inhibition to Ct, %)
Healthy donor 1	None	M	29	—	—			
Healthy donor 2	None	F	31	—	—			
Healthy donor 3	None	F	27	—	—			
Healthy donor 4	None	F	37	—	—			
Patient 1	Acute ATL	M	68	+	+	14	54	71
	Before chemo							
	Acute ATL				n.e.	2	0	8
	After chemo							
	Acute ATL							
Patient 2	Recurrence after	M	61		n.e.	14	2	43
	umbilical cord blood							
	transplantation							
Patient 2	Acute ATL	M	61	+	+	65	24	62
Patient 3	Acute ATL	F	75	+	+	8	38	60
Patient 4	Acute ATL	F	75	+	+	11	7	52
Patient 5	Acute ATL	M	64	+	n.e.	2	10	22
Patient 6	Acute ATL	M	82	+	n.e.	28	16	46
Patient 7	Chronic ATL	M	52	+	+	41	55	94
Patient 8	Mycosis fungoides	M	63	—	—	14	34	70
Patient 9	T-cell lymphoma	M	63	—	—	7	0	92
Patient 10	Acute ATL	F	65	+	n.e.	94	15	66


M, male; F, female; \*, information not available; n.e., not examined.

# 1


CuD induces primary ATL patient cell death.

(A) Structures of CuD and prednisolone. (B, C) PBLs were freshly prepared from healthy donor (B) or ATL patients (C). Cells ( $1 \times 10^4$  /well) were treated with indicated concentrations of CuD or vehicle in the presence or absence of rapamycin (100  $\mu$ M) for 24 h. (D) PBLs ( $1 \times 10^4$  /well) from ATL patient 1 were treated with CuD (0.19  $\mu$ M) in the presence or absence of 3-MA (10  $\mu$ M) for indicated periods. Cell viability was assessed with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. \*P< 0.05 vs. vehicle without CuD treatment at each time point.


**A**




**B Healthy donor**



**C ATL Patient**



**D ATL Patient**




# 2


Z36 inhibits proliferation and enhances CuD-induced death in primary ATL patient cells.

(A-D) PBLs ( $1 \times 10^4$  /well) from healthy donors (A) or ATL patients (B-D) were treated with indicated concentrations of Z36 or dimethyl sulfoxide (DMSO, a control solvent for Z36) for 12 or 24 h, or CuD (0.19  $\mu$ M) or prednisolone for 24 h.. (C) PBLs from ATL patients were pre-treated with 10 mM 3-MA or vehicle for 2 h, and then exposed to 0.19  $\mu$ M CuD, Z36 (7.5  $\mu$ M), or vehicle (Et-OH, 0.01%) for 24 h. (D) \*P< 0.05 vs. vehicle at each time point. Cell viability was determined using the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. \*P< 0.05 vs. vehicle; ¶ P < 0.05 vs. treatment with CuD only; # P < 0.05 vs. CuD and Z36 co-treatment.


**A Healthy donor**




**B ATL Patient**



**C ATL Patient**



**D ATL Patient**






### 3

Autophagy negatively regulates CuD- or Z36-induced PARP cleavage in ATL patient cells.

(A, B) PBLs ( $2 \times 10^5$ /well) from ATL patients were treated with CuD ( $0.19 \mu\text{M}$ ), Z36 ( $7.5 \mu\text{M}$ ), or vehicle in the presence or absence of 3-MA ( $10 \text{ mM}$ ) for 24 h. Whole cell extracts were used for western blot analysis of cleaved PARP (A ) and LC3-II (B) expression;  $\beta$ -actin or tubulin was used as a loading control. ET, ethanol; DM, DMSO. [p]




# 4


CuD and Z36 show limited effects on ATL patient-derived primary cells upon disease recurrence.

(A, C) PBLs ( $1 \times 10^4$  /well) from ATL patients after chemotherapy (A) or from ATL patients upon recurrence after umbilical cord blood transplantation (C) were treated with Z36 (7.5  $\mu$ M) or vehicle for 24 h. Cell viability was determined with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. (B) PBLs ( $1 \times 10^4$  /well) from a patient before therapy (left four lanes), after umbilical cord blood transplantation (middle four lanes), and after recurrence (right four lanes) were treated with CuD (0.19  $\mu$ M) and/or Z36 (7.5  $\mu$ M) or with vehicle for 24 h. Whole cell extracts were used for western blot analysis of cleaved PARP;  $\beta$ -actin was used as a loading control. M, molecular marker. (D) Whole cell extracts from a patient before therapy (ori), after umbilical cord blood transplantation (Ch), and after recurrence (Rec) were analyzed for caspase-1 expression by western blotting;  $\beta$ -actin was used as a control. (E) ATL patient cells ( $1 \times 10^4$  /well) obtained upon recurrence after umbilical cord blood transplantation were treated with CuD (0.19  $\mu$ M) and Z36 (7.5  $\mu$ M) or vehicle for 24 h in the presence or absence of capase-1 inhibitor (20  $\mu$ M). Cell viability was determined with the Cell Titer-Glo assay. Results are expressed as mean activity relative to controls and SD from quadruplicate cultures. \*P < 0.05 vs. vehicle; # P < 0.05 vs. CuD and Z36 co-treatment.


# **A ATL Patient after chemo**




# **C Recurrence after the umbilical cord blood transplant**



# **B**



# **D**



# **E**

