

# Behavioral flexibility in an invasive bird is independent of other behaviors

**Corina J Logan** Corresp. 1, 2

<sup>1</sup> Department of Zoology, University of Cambridge, Cambridge, United Kingdom

<sup>2</sup> SAGE Center for the Study of the Mind, University of California, Santa Barbara, Santa Barbara, California, United States

Corresponding Author: Corina J Logan  
Email address: cl417@cam.ac.uk

Behavioral flexibility is considered important for a species to adapt to environmental change. Behavioral flexibility relates to problem solving ability and speed in unpredictable ways, which leaves an open question of whether behavioral flexibility instead varies with differences in other behaviors, such as neophobia or exploration. If present, such correlations would mask which behavior causes individuals to vary. I investigated whether behavioral flexibility (reversal learning) performances were linked with other behaviors in great-tailed grackles, an invasive bird. I found that behavioral flexibility did not significantly correlate with neophobia, exploration, risk aversion, persistence, or motor diversity. This suggests that great-tailed grackle performance in behavioral flexibility tasks reflects a distinct source of individual variation. Maintaining multiple distinct sources of individual variation, and particularly variation in behavioral flexibility, may be a mechanism for coping with the diversity of novel elements in their environments and facilitating this species' invasion success.

# **Behavioral flexibility in an invasive bird is independent of other behaviors**

Corina J. Logan

SAGE Center for the Study of the Mind, University of California, Santa Barbara, CA 93106

U.S.A.

Current address: Department of Zoology, University of Cambridge, Cambridge CB2 3EJ U.K.,

cl417@cam.ac.uk

## **BACKGROUND**

Behavioral flexibility, defined here as changing preferences according to changing circumstances based on learning (Logan 2016a,b), is considered a key factor involved in a species' ability to adapt to environmental change (Lefebvre et al. 1997, Griffin & Guez 2014, Buckner 2015, Chow et al. 2016). However, it is not known how behavioral flexibility works: is it a problem solving ability or does it arise because of links with other behaviors such as neophilia and exploration (see review in Griffin 2016)? The few studies that have investigated whether behavioral flexibility relates to problem solving ability and speed have found that these traits do not covary in predictable ways (Boogert et al. 2011, Griffin et al. 2013, Isden et al. 2013, Logan 2016a, Bebus et al. 2016). This leaves an open question of whether behavioral flexibility varies with differences in other behaviors such as exploration, neophobia, risk aversion, persistence, and motor diversity (the number of different motor actions used to attempt to solve a novel problem). There is debate about whether differences in behavior among individuals are linked to suites of correlated behaviors or whether individual behaviors, such as behavioral flexibility, can vary independently (Coppens et al. 2010, Cole et al. 2011, Sih & Del Giudice 2012). Reversal learning, a measure of behavioral flexibility, is predicted to fall on the fast-slow behavioral type

continuum where fast individuals are exploratory, risk seeking, and persistent with poor accuracy because of the speed with which they solve problems, whereas slow individuals are neophobic, risk averse, and more accurately solve problems (Figure 1; Sih & Del Giudice 2012). Accordingly, slow individuals should be more behaviorally flexible because they might be less impulsive and spend more time sampling the environment (Sih & Del Giudice 2012). If present, these types of behavioral correlations could mask whether differences between individuals in their behavioral flexibility performances are caused by behavioral flexibility or a correlated behavior (Herrmann et al. 2010, Thornton & Lukas 2012, Seed et al. 2012).

Results from the few studies that investigated the relationship between behavioral flexibility and other behaviors are equivocal. There are a variety of ways to measure behavioral flexibility in an experimental context and all involve allowing an individual to learn about a task, which then changes after the individual becomes proficient. Individuals that adapt their behavior to these changing circumstances are considered to exhibit behavioral flexibility. Paradigms testing behavioral flexibility include tasks such as a multi-access box (Auersperg et al., 2011; Manrique, Völter & Call, 2013), water tubes (Logan et al. 2014, Logan 2016a), and episodic-like memory and future planning experiments (Clayton & Dickinson, 1998; Dally, Emery & Clayton, 2006; Raby et al., 2007), however the most widely used measure is reversal learning (e.g., Bond et al. 2007, Tebbich et al. 2010, Boogert et al. 2011). Reversal learning involves learning to associate one option with a reward, which subsequently becomes incorrect when the reward is moved to a different option, thus forcing the individual to reverse their preference to consistently obtain the reward. Switching preferences is also thought to require inhibition to refrain from choosing the previously rewarded response (Manrique et al. 2013, Griffin & Guez 2014, Liu et al. 2016, but see Homberg et al. 2007), which could be related to a propensity to be more

neophobic and less exploratory (Sih & Del Giudice 2012), and/or a propensity to continue to sample the environment (Logan 2016a). Consistent with predictions, black-capped chickadees (*Poecile atricapillus*) and great tits (*Parus major*) that were more flexible (faster to reverse a previously learned preference) were slower to explore (Verbeek et al. 1994, Guillette et al. 2011), and great tits that were more flexible (reversal learning) were more neophobic (Verbeek et al. 1994). Two studies provided evidence inconsistent with the predictions that behavioral flexibility will positively correlate with neophobia and negatively with exploration: there were no correlations between behavioral flexibility (reversal learning) and neophobia or exploration in Florida scrub jays (*Aphelocoma coerulescens*; Bebus et al. 2016), and also no correlations with activity or boldness in wild cavies (*Cavia aperea*; Guenther et al. 2014).

Innovativeness, considered a subcategory of behavioral flexibility, was linked with the number of motor actions used to try to solve a novel problem, but not with persistence or neophobia in several bird species (Griffin et al. 2014, Diquelou et al. 2016, Griffin & Diquelou 2015). Innovativeness, defined as inventing new behaviors to solve novel problems or using existing behaviors in new ways (Griffin & Guez 2014), is distinct from behavioral flexibility. For example, great-tailed grackles exhibit behavioral flexibility in two tests involving reversal learning, showing that they are among the fastest bird species to both learn an initial preference and to reverse this preference (Logan 2016a). However, grackles are not particularly inventive when it comes to creating new behaviors to solve novel problems: they did not successfully innovate string pulling or stick tool use, which are behaviors that many other bird species engage in (Logan 2016b). While it is unknown how motor diversity interacts directly with behavioral flexibility, the prediction is that these traits will positively correlate because increasing the

number of motor actions attempted could increase the probability and speed of finding a successful solution to a novel problem (Diquelou et al. 2016).

To determine whether behavioral flexibility is related to a variety of behaviors in one species, I investigated great-tailed grackles (*Quiscalus mexicanus*, family Icteridae, hereafter referred to as grackles), a generalist forager (Skutch 1954, Johnson & Peer 2001, Wehtje 2003) that is behaviorally flexible (Logan 2016a). Grackles are a native invasive species (Peer 2011): they have expanded their range north from Central America into North America by over 5500% over the course of 120 years following the expansion of human modified environments, which is their preferred habitat (Wehtje 2003). Behavioral flexibility is hypothesized to be a mechanism involved in successful species invasions (Sol & Lefebvre 2000). While grackles could have expanded their range simply because of an increase in their suitable habitat, species differences in traits that facilitate adapting to environmental change, such as diet, are additionally implicated (Blackburn et al. 2009). Behavioral flexibility is one such trait and a better understanding of how it works could have implications for managing species invasions. To better understand behavioral flexibility, I tested the hypothesis that individual variation in behavioral flexibility correlates with variation in other behaviors on the fast-slow continuum.

I predicted that individuals that were more behaviorally flexible would also be the most neophobic and risk averse, the least persistent and exploratory, and use more motor actions. I quantified grackles' activity levels (exploration) when placed in a novel environment and also measured the amount of time spent in the riskiest sections of the aviary (risk aversion). I measured grackles' neophobic reactions to a novel object next to a food dish in comparison with controls where only a food dish was present. Persistence and motor diversity were measured from videos of a stick tool use experiment (Logan 2016b), where no bird successfully invented

stick tool use. Therefore, birds were never rewarded for their actions, which is important when measuring persistence because a food reward could differentially influence persistence across individuals: those who are better at the task would receive more food rewards, which might increase their persistence in future trials.

## **METHODS**

### **Ethics**

This research was conducted in accordance with the following permits: U.S. Fish and Wildlife Service (scientific collecting permit number MB76700A-0), U.S. Geological Survey Bird Banding Laboratory (federal bird banding permit number 23872), California Department of Fish and Wildlife (scientific collecting permit number SC-12306), and the Institutional Animal Care and Use Committee at the University of California Santa Barbara (IACUC protocol numbers 860 and 860.1).

### **Subjects**

Eight adult great-tailed grackles (4 females and 4 males) were caught in the wild and held for 2-3 months in aviaries before being released back to the wild (see Logan 2016a for full details). Half of the birds were caught in Santa Barbara, California in September 2014 and released in December 2014 (Tequila, Margarita, Cerveza, and Michelada; batch 1) and the other half were caught in January 2015 and released in March 2015 (Refresco, Horchata, Batido, Jugo; batch 2).

### **Study Set Up**

Grackles were housed individually in aviaries (183 cm high by 119 cm wide by 236 cm long) at the University of California Santa Barbara. Grackles had *ad libitum* access to water at all times, and unrestricted amounts of food (Mazuri® Small Bird Food) for a minimum of 20 hrs per day. On testing days, their main diet was removed for up to 4 hrs while they participated in experiments and could eat bread or peanuts if successful. Apparatuses were placed on tables (60 cm wide by 39 cm long) and rolled into each aviary for sessions (approximately 20 min per session), which were visually isolated from other grackles and video recorded with a Nikon D5100 camera on a tripod. Experimenters stood just outside the aviary door and in full view of the grackles during the persistence and motor diversity sessions, which did not interfere with behavior (i.e., they readily interacted with the apparatus) because they were habituated to humans in the wild and in the aviary.

## Statistical Analyses

Data were analyzed in R 3.2.1 (R Core Team 2015). For those tests that involved p-values, a result was considered statistically significant when  $p < 0.05$ . When multiple p-values were obtained for one experiment, a Bonferroni-Holm correction was applied to avoid obtaining false positive results simply by conducting many tests on the same data.

## Data Accessibility

Data are available at the KNB Data Repository (Logan 2016c; [https://knb.ecoinformatics.org/#view/corina\\_logan.18.13](https://knb.ecoinformatics.org/#view/corina_logan.18.13)). Behavioral flexibility data were previously published and are available at KNB (Logan 2016d).

## Videos

Clips of videos from each experiment are available at: <https://youtu.be/4efJ2afatus>.

## Exploration and Risk Aversion

The exploration session lasted 60 min, starting 30 min after a wild bird's release into the aviary, a novel environment. The grackles' previous experience was always the same: they were trapped, blood was collected, and colored rings put on their legs; they were transported to the aviary in a cat carrier in a car, biometrics were taken, and then they were released into the aviary where they were singly housed, given food and water, and the camera was set up outside their door. The camera was restarted every 20 min, otherwise experimenters were out of visual and auditory contact when recording.

Exploration is measured in a number of different ways and I chose two measures for the purposes of this study, which have been used to measure exploration in other species: the amount of activity in a novel environment (exploration; e.g., Verbeek et al. 1994, Fox et al. 2009) and the amount of time spent in the risky areas of the novel environment (risk aversion; e.g., Lynn & Brown 2009, Lerman et al. 2012, Jolles et al. 2014). It is unclear whether these measure the same behavior because both have not been measured in a single study. Therefore, I conducted a Spearman's rank correlation test to determine whether they are distinct behaviors.

**Exploration:** Videos were coded by Linnea Palmstrom to determine how often birds moved between sections of the aviary. The aviary was invisibly sectioned into 5 areas with sections 1 and 5 in the upper half of the aviary, where the perches were located, and sections 2-4 on the ground (Figure 2). Their exploration score was the number of times they moved from one aviary section to another over the course of the 60 min session, which was an appropriate time



period (Montiglio et al. 2010) because an individual's activity level was ranked similarly regardless of whether activity occurred in the first 5 min or over the whole session (Spearman's rank correlation:  $S=31.43$ ,  $p=0.10$ ,  $\rho=0.63$ ,  $n=8$ ).

**Risk aversion:** I (invisibly) divided the aviary into safer versus riskier sections (Figure 2) and used the percentage of time spent in sections 4 and 5 as a measure of risk aversion. The rear of the aviary was considered less risky because it was the farthest from the door where the camera and other equipment were visible, while the ground and the perches in the front of the aviary were more risky because these sections were next to the door and walking on the ground is more dangerous than flying. Food and water were placed on the ground near the door. The aviary was covered in tarpaulins on three sides (both sides and rear), while the side with the door (front) and the ceiling were wire mesh that the bird could see through.

## Neophobia

The neophobia sessions began on a grackle's sixth day in the aviary and involved three 10-min trials with trials 1 and 3 serving as a way to quantify food motivation by placing a food bowl alone on the table, while trial 2 had a novel object 2 cm to the right of the food bowl (as in Boogert et al. 2006). There were 2 min between trials. In all trials, the food bowl contained 1/4 of a peanut and the latency to land on the table and to feed was recorded as well as which object was approached first in trial 2 (the food bowl or the novel object). Three novel objects were presented in random order to each bird: a GoPro camera inside its clear waterproof case, a stone dropping training apparatus (see Logan 2016a), and a colored U-tube apparatus (see Logan et al. 2016). The stone dropping training apparatus was a clear acrylic box (8.8 cm tall by 18 cm wide by 11 cm deep) with a clear acrylic tube (9 cm tall, outer diameter=5 cm) on top. The colored U-

tube apparatus was a box (8cm tall by 40 cm wide by 30 cm deep) with a wooden frame covered in cardboard and a clear acrylic top covered by colored paper. Two clear acrylic tubes (both 17 cm tall, one with an outer diameter of 5.1 cm and the other 2.5 cm) protruded from the center of the box and were marked with colored tape at the top. If a grackle did not come to the table within the 10 min period it received a trial duration of 601 seconds. The neophobia tests were conducted on three consecutive days, with one novel object presented to the bird on each day.

Data were analyzed using the latency to land on the table rather than the latency to feed because birds came to the table more often than they ate the food. The data were not normally distributed (Anderson-Darling normality test: GoPro:  $A=3.08$ ,  $p<0.001$ ; stone dropping apparatus:  $A=2.76$ ,  $p<0.001$ ; U-tube:  $A=2.46$ ,  $p<0.001$ ). Therefore, non-parametric paired Wilcoxon signed rank tests with continuity corrections were conducted to determine whether latencies in control trials (averaged) differed from novel object trials. Neophobia scores were obtained for each novel object and summed for an overall score per individual. Scores were calculated by subtracting the latency to land on the table during the novel object trials (trial 2) from the average latency during control trials (trials 1 and 3). Positive scores indicate less neophobia while negative scores indicate more neophobia. Repeatability of individual neophobic responses across contexts was measured using Spearman's rank correlations to determine whether grackles maintained similar neophobia ranks with each of the three novel objects.

## **Persistence and Motor Diversity**

Persistence and motor diversity were calculated as in Griffin & Diquelou (2015). Persistence was calculated as the attempt rate: the number of times a bird came to the table or interacted with (touched) the apparatus or stick across 21 trials of a stick tool use experiment (105 min/bird).

Motor diversity was calculated by counting the number of different motor actions (described in Table 3) performed per individual across the 21 trials of the experiment. Videos were watched from trials 1 to 21 and behaviors from the ethogram (Table 3) were coded at their first observation.

The stick tool use experiment involved an apparatus with a wooden base and rear with clear cast acrylic walls providing a narrow gap at the front and top of the apparatus to insert a stick and retrieve a piece of bread. Birds were given 21 5 min trials to innovate tool use: first, 3 trials with the stick placed on the table next to the apparatus, then 3 trials with the stick inserted into the apparatus, and finally 15 trials with the stick inserted in the apparatus and tool use demonstrated by the human experimenter.

### **Measure of Behavioral Flexibility**

These grackles were previously tested on reversal learning of a color discrimination task consisting of a gold tube and a silver tube placed on the table at the same time with one color containing hidden food and the opportunity to make only one choice per trial (Logan 2016a). Grackles initially learned to search for food hidden in the gold tube and, once proficient, the food was switched to the silver tube and the number of trials required to reach proficiency was assessed. Behavioral flexibility scores were calculated as the number of trials to reverse a color preference minus the number of trials needed to initially learn the color preference (Table 4). Proficiency in the initial discrimination and reversal was demonstrated if individuals chose correctly in at least 17 of the most recent 20 trials with at least 8 or 9 trials correct per set of 10. I then investigated whether relationships between individual variation in behavioral flexibility and exploration, risk aversion, neophobia, persistence, and motor diversity conformed to predictions.

229

## 230 **General Analyses**

231 I determined whether behavioral flexibility (response variable: behavioral flexibility score)  
 232 negatively correlated with exploration and persistence while examining whether batch had an  
 233 effect (explanatory variables) using a Generalized Linear Model (GLM; MCMCglmm function,  
 234 MCMCglmm package; Hadfield 2014) with a Poisson distribution and log link using 13000  
 235 iterations with a thinning interval of 10 and a burnin of 3000. The GLM showed acceptable  
 236 convergence (lag time autocorrelation values were  $<0.01$ ; Hadfield 2010). Risk aversion and  
 237 motor diversity were excluded from the analysis because they significantly covaried with  
 238 exploration and persistence, respectively. A Spearman's rank correlation was used to investigate  
 239 the relationship between behavioral flexibility and neophobia because residuals were not  
 240 normally distributed.

241       Given the small sample size ( $n=7$  for behavioral flexibility scores), I conducted a further  
 242 analysis to determine whether GLM results were likely to be reliable given the data (Burnham &  
 243 Anderson 2002). I compared the Akaike weights (range: 0-1, the sum of all model weights  
 244 equals 1; Akaike 1981) between the test model (above) and a null model (behavioral flexibility  
 245 score as the response variable and 1 as the explanatory variable) using the dredge function in the  
 246 MuMIN package (Bates et al. 2011). If the best fitting model has a high Akaike weight ( $>0.89$ ;  
 247 Burnham & Anderson 2002), then it indicates that the results are likely given the data. The null  
 248 model was strongly supported with an Akaike weight of 0.92, thus indicating the results are  
 249 reliable even with a small sample size.

250

## 251 **Interobserver Reliability**

Linnea Palmstrom coded the exploration/risk aversion videos, I coded neophobia videos, and Katherine Lister coded persistence and I coded motor diversity from videos of a tool use experiment (in Logan 2016b). To measure interobserver reliability, I randomly chose 21% of the videos using [www.random.org](http://www.random.org) and had a coder who was naïve to the hypotheses (Katharina Brecht) recode their exploration (from which measures of risk aversion are calculated), persistence, and neophobia. I randomly chose 3 of the 8 birds using [www.random.org](http://www.random.org) and had Katharina recode their motor diversity (36% of the videos). A higher percentage of motor diversity videos were recoded because agreement determinations were based on the total number of motor actions per bird, which required watching all videos for an individual. There was agreement between Katharina and all other observers for each study: exploration (intraclass correlation coefficient (ICC)=0.998, 95% confidence intervals (CI) =0.98-1.00), neophobia (ICC=0.87, 95% CI=0.67-0.95), persistence (land on table: ICC=0.79, 95% CI=0.49-0.93; interact with apparatus: ICC=1.00, 95% CI=0.999-1.00; interact with stick: ICC=1.00, 95% CI=NA), and motor diversity (ICC=0.71, 95% CI=0.54-0.82; ICCs calculated using R package: irr, function: icc, Gamer et al. 2012).

## RESULTS

### Exploration and Risk Aversion

Exploration and risk aversion were significantly positively correlated, indicating that these two variables actually measure the same behavior (Spearman's rank correlation:  $S=8.55$ ,  $p=0.002$ ,  $\rho=0.90$ ,  $n=8$ ). Therefore, I used exploration to represent this behavior in further analyses (see General Analyses section).

**Exploration:** Grackles varied in how many times they changed sections across the 60 min session (0-163), with Refresco having no section changes and Jugo having the most (Table 1). They also varied in the total number of sections they visited during the session (1-5; Table 1).

**Risk aversion:** Grackles varied in how much time they spent in the riskiest sections of the aviary with Batido spending the most time and Refresco the least (Table 1). All grackles (except Refresco) moved through other sections of the aviary and they varied in how much time they spent in sections 4 and 5 (Table 1).

## Neophobia

There were no significant differences between the latency to land on the table in controls (pre [trial 1] or post [trial 3] novel object trials) versus novel object trials (trial 2) (Wilcoxon tests with Bonferroni-Holm corrected p-values: GoPro: *trials 1-2*  $V=21$ ,  $p=1.00$ , 95% confidence interval=-283-267, *trials 2-3*  $V=8$ ,  $p=1.00$ , 90% CI=-427-277.5; stone dropping apparatus: *trials 1-2*  $V=7$ ,  $p=1.00$ , 80% CI=-369-338, *trials 2-3*  $V=0$ ,  $p=0.54$ , 80% CI=-455-(-41); U-tube: *trials 1-2*  $V=1$ ,  $p=0.88$ , 80% CI=-481-(-85), *trials 2-3*  $V=1$ ,  $p=0.88$ , 80% CI=-507-(-190.5)). Refresco and Margarita were overall less neophobic than the other grackles, and Horchata was the most neophobic (Table 2). There were many trials in which the bird did not come to the table. However, this did not usually appear to be due to neophobia because it happened in many control trials as well as novel object trials, indicating that it might have been due to a lack of motivation to eat or explore the object.

There was no individual repeatability of neophobia scores across contexts (Spearman's rank correlation with Bonferroni-Holm corrected p-values: GoPro vs. stone dropping apparatus:

296  $S=79.21$ ,  $p=1.00$ ,  $\rho=-0.41$ , GoPro vs. U-tube:  $S=56.00$ ,  $p=1.00$ ,  $\rho=0.00$ ; U-tube vs. stone  
 297 dropping apparatus:  $S=20.68$ ,  $p=0.88$ ,  $\rho=0.63$ ).

298

### 299 **Persistence and Motor Diversity**

300 Grackles varied in the number of motor actions they used (1-15) and in how persistent they were  
 301 (18-4047 total number of times a bird landed on the table, touched the apparatus, or touched the  
 302 stick; Table 4). A post-hoc analysis indicated that these two variables were significantly  
 303 positively correlated, and thus both measured the same behavior (Spearman's rank correlation:  
 304  $S=8.55$ ,  $p=0.002$ ,  $\rho=0.90$ ,  $n=8$ ). Therefore, only persistence was used in further analyses (see  
 305 General Analyses section).

306

### 307 **Does behavioral flexibility positively correlate with motor diversity and risk aversion, and** 308 **negatively with exploration and persistence?**

309 Birds that were more flexible (i.e., faster to reverse a preference: number of trials to reverse a  
 310 preference minus the number of trials to initially learn the preference) did not have higher  
 311 exploration scores, they were not more persistent, and there were no batch effects (Figure 3,  
 312 Supplementary Table S1).

313 Risk aversion and motor diversity significantly covaried with exploration and persistence,  
 314 respectively, and these relationships were investigated further. I confirmed that the relationship  
 315 between these variables and behavioral flexibility was the same as their collinear variables with  
 316 an additional GLM. This GLM was the same as above, except the explanatory factors were  
 317 motor diversity, risk aversion and batch. As above, flexibility did not correlate with risk aversion  
 318 or motor diversity (Figure 3, Supplementary Table S2).

319

# 320 Does behavioral flexibility positively correlate with neophobia?

321 Grackles that were more flexible (i.e., faster to reverse a preference) did not have lower  
 322 neophobia scores, which would indicate more neophobia. There was no significant correlation  
 323 between behavioral flexibility scores and neophobia (Figure 3; Spearman's rank correlation:  
 324  $S=92$ ,  $p=0.12$ ,  $\rho=-0.65$ ).

325

## 326 DISCUSSION

### 327 Exploration and Risk Aversion

328 The exploration and risk aversion scores significantly correlated with each other, indicating they  
 329 measured the same behavior. While risk aversion scores could have been confounded by the  
 330 placement of food and water in a risky section, which might attract birds to this area, they spent  
 331 only 0-14% of their time in the section with the food and water (section 4). This indicates that  
 332 they behaved more according to the prediction that this section would be treated as risky even  
 333 when an attractor was present.

334 I question whether the measure of exploration actually measured exploration in this  
 335 species. A bird that is stressed tends to fly back and forth in an aviary, which is not an indicator  
 336 of exploration, but would be interpreted as such according to the section change measure of  
 337 exploration. In this study, Jugo mostly flew back and forth between the perches near the top of  
 338 the aviary while looking up and out of the aviary and not attending to the environment within the  
 339 aviary. In contrast, Horchata also had many section changes, but she usually walked calmly on  
 340 the ground, thus perhaps in her case this measure of exploration was appropriate. Therefore, at  
 341 the species level, activity levels are likely not a good indicator of exploration behavior. Indeed, a



distinction is made between forced exploration, where an individual is placed in a novel environment, and voluntary exploration, where an individual in a familiar environment is provided with the opportunity to enter a novel environment (Guenther et al. 2014). A study on wild guinea pigs (*Cavia aperea*) found that these two variations of exploration measure different behaviors: forced and voluntary exploration activity did not correlate in juveniles or adults (Guenther et al. 2014).

Voluntary exploration would likely be a more accurate measure of actual exploratory behavior in grackles, which could also involve voluntary exploration of a novel object in a familiar environment. Such a measure is also called a neophilia test where a novel object is placed in a familiar aviary in the presence of (but not next to) their regular food source to determine how soon the bird approaches and interacts with the apparatus and for how long (as in Mettke-Hoffman et al. 2002). This kind of test would also likely more directly relate to how grackles have expanded their range so rapidly: rather than exploring novel environments, grackles are more likely to have successfully expanded their range by exploring novel objects. Grackles have not necessarily needed to adapt to novel environments during their range expansion because it coincided with an increase in their suitable (human-managed) habitat (Wehtje 2003). Exploration is more likely to have played a role in exploiting novel objects in their environment because humans throw away products that may be novel to grackles (e.g., egg cartons, yogurt cups) and design new potential food sources (e.g., dumpsters) where food is not necessarily obvious, therefore the objects must be explored to determine whether they contain food.

## Neophobia

Grackles were not generally neophobic because no significant differences were found between controls and novel object trials in the latency to land on the table. Indeed, the GoPro camera, which was also the smallest of the novel objects, appeared to attract their attention more than the food. Comparing grackles with other species that have been tested using a similar design, it appears that they are less neophobic than starlings (*Sturnus vulgaris*; Boogert et al. 2006), blue tits (*Cyanistes caeruleus*; Herborn et al. 2010), Japanese quail (*Coturnix japonica*; Zimmer et al. 2013), Chimango caracaras (*Milvago chimango*; Biondi et al. 2010), European greenfinches (*Carduelis chloris*; Herborn et al. 2011), Indian mynas (Sol et al. 2012, Griffin & Diquelou 2015), and mountain chickadees (*Poecile gambeli*; Kozlovsky et al. 2015), and that they have similar levels of neophobia as noisy miners (*Manorina melanocephala*; Griffin & Diquelou 2015) and a different group of mountain chickadees (Fox et al. 2009).

### **Persistence and Motor Diversity**

The persistence and motor diversity scores significantly correlated with each other, indicating they measured the same behavior. This suggests that the longer a bird persists in attempting to solve a task, the more likely it is to use a wider variety of motor actions. Therefore, it is likely that individuals that used few motor actions would likely have used more if they were perhaps more motivated to interact with the task. Measuring persistence and motor diversity in a variety of contexts could address this potential issue and clarify whether these variables actually do covary on a task that all individuals persist on. These results are different from findings using a similar experimental design on Indian mynas and noisy miners where motor diversity differed between species, but persistence did not, thus indicating these are two separate behaviors (Griffin & Diquelou 2015). However, birds in Griffin & Diquelou (2015) could receive food rewards

from the apparatus if successful (i.e., at the end of each bout of persistence) and such positive reinforcement for persisting could have increased persistence for successful individuals in subsequent trials.

# **Behavioral Flexibility**

Contrary to predictions, behavioral flexibility did not correlate with exploration, neophobia, risk aversion, persistence, or motor diversity. The small sample size might have limited my ability to detect significant correlations, however the behavior of the models suggested this was not the case. It is perhaps not surprising that behavioral flexibility did not correlate with neophobia (the only behavior I was able to obtain repeatability measures from) because neophobia was not expressed consistently across contexts. This could indicate a further source of individual variation in grackles or it could result from the inability of the method to accurately measure neophobia in this species. The latter highlights the importance of conducting repeatability tests when attempting to understand how two variables correlate because two unpredictable variables (behavioral flexibility and neophobia) would not likely correlate with each other. These results are similar to results from Florida scrub jays where behavioral flexibility (reversal learning) did not correlate with neophobia or exploration (Bebus et al. 2016). The few studies that investigate the relationship between behavioral flexibility and other behaviors either show relationships opposite to predictions (Verbeek et al. 1994, Guillette et al. 2011) or show no correlations (this study, Bebus et al. 2016). One prediction was supported in only one test: reversal learning speed negatively correlated with neophobia (Verbeek et al. 1994). This accumulating evidence suggests the need to reconsider the basis for hypotheses linking other behaviors with behavioral flexibility.

# CONCLUSIONS

Traditionally, behavioral flexibility is thought of as a cognitive ability (see review in Shettleworth 2010) and is considered as such in hypotheses linking it with other behaviors (Sih & Del Giudice 2012, Guenther et al. 2013). However, mixed results, with none conforming to predictions, from grackles, keas and New Caledonian crows question this assumption. Grackles lacked correlations between behavioral flexibility and problem solving ability and speed, and individuals that were behaviorally flexible in one type of test were not necessarily flexible in a different type of test (Logan 2016a). The more exploratory keas were more behaviorally flexible on a multi-access box and faster to innovate new solutions to novel problems than New Caledonian crows (Auersperg et al. 2011). These mixed results indicate a need to look beyond cognitive and behavioral measures that might correlate with behavioral flexibility and investigate relationships with factors such as physiology and genetics. For example, grackles that are in better phenotypic condition (e.g., have better immunity) might have the capacity to be more behaviorally flexible than individuals in worse phenotypic condition. Non-behavioral, non-cognitive individual factors have yet to be measured in relation to behavioral flexibility. Considering behavioral flexibility in this more integrated way could allow experimenters to manipulate this elusive trait to understand what it is and how it works.

Though the sample size is small, these results provide further support that behavioral flexibility represents a distinct axis of individual variation in behavior. Behaviors that do not correlate with each other are suggested to represent “inherent individual differences” in each of the traits measured (Cole et al. 2011, p. 495). For example, great tit problem solving ability did not correlate with body condition, neophobia, or exploration; therefore problem solving was

considered its own behavior that varies across individuals rather than varying due to links with other individual traits (Cole et al. 2011). The methods used to measure neophobia and exploration in grackles might not have accurately represented these behaviors, therefore further investigations using different methods that are validated measures of these behaviors in grackles should be explored before entirely ruling out correlations with behavioral flexibility. Previous research on grackles and other species has shown that behavioral flexibility is independent from innovativeness (Logan 2016b), problem solving ability and speed (Boogert et al. 2011, Isden et al. 2013, Logan 2016a), or that it negatively correlates with problem solving speed (Griffin et al. 2013). The majority of evidence so far indicates that individual variation in behavioral flexibility is not confounded with other behaviors, although two alternative hypotheses cannot yet be ruled out: the behaviors might not have been measured with enough consistency across studies to directly compare the results, or the behaviors are not repeatable enough within individuals to reliably covary with each other. Further research is needed to distinguish which hypothesis is supported.

It could be adaptive for invasive species, such as the grackle, to maintain many independent axes of individual variation and, in particular, variation in behavioral flexibility. Indeed, Western bluebirds rely on existing intrapopulation variation when expanding their range (Duckworth 2008). While Western bluebirds rely on variation in dispersal strategies, grackles may rely on maintaining individual variation in behavioral flexibility, which could allow them to more quickly adapt to changing or unpredictable environments.

## FUTURE DIRECTIONS

Future research investigating neophobia, exploration, persistence and motor diversity in this species would benefit from a larger sample size, replicability of results from multiple groups, and finding measures that are repeatable within individuals to determine the reliability of these conclusions. Investigating each variable using multiple methods will facilitate an understanding of which methods actually measure the behaviors of interest.

## ACKNOWLEDGEMENTS

I am grateful to Linnea Palmstrom for coding the exploration videos; Katherine Lister for coding the persistence videos; Katharina Brecht for coding all videos for interobserver reliability analyses; Luisa Bergeron, Christin Palmstrom, Linnea Palmstrom, and Michelle Gertsvolf for trapping and aviary assistance; Steve Rothstein for use of the aviaries; Joe Jablonski for making the cast acrylic apparatuses; Jill Zachary and Kathy Frye at Santa Barbara City Parks and Recreation for use of the Andree Clark Bird Refuge and East Beach Park; Estelle Sandhaus and Chris Briggs at the Santa Barbara Zoo for access to wild grackles; Dieter Lukas, Ted Bergstrom, and Carl Bergstrom for discussions; and Alecia Carter for analysis and manuscript feedback.

## REFERENCES

- Akaike H. 1981.** Likelihood of a model and information criteria. *Journal of Econometrics* **16**:3-14 doi:10.1016/0304-4076(81)90071-3
- Auersperg AMI, von Bayern AMP, Gajdon GK, Huber L, Kacelnik A. 2011.** Flexibility in problem solving and tool use of kea and New Caledonian crows in a multi access box paradigm. *PLoS ONE* **6**:e20231 doi:10.1371/journal.pone.0020231

- 478 **Bates D, Maechler M, Bolker B. 2011.** lme4: linear mixed-effects models using S4 classes. R  
479 package version 0.999375-42. Available at [http:// CRAN.R-project.org/ package=lme4](http://CRAN.R-project.org/package=lme4)  
480 (accessed 12 January 2016).
- 481 **Bebus SE, Small TW, Jones BC, Elderbrock EK, Schoech SJ. 2016.** Associative learning is  
482 inversely related to reversal learning and varies with nestling corticosterone exposure.  
483 *Animal Behaviour* **111**:251-260 doi:10.1016/j.anbehav.2015.10.027.
- 484 **Biondi LM, Bó MS, Vassallo AI. 2010.** Inter-individual and age differences in exploration,  
485 neophobia and problem-solving ability in a Neotropical raptor (*Milvago chimango*).  
486 *Animal Cognition* **13**:701-710 doi:10.1007/s10071-010-0319-8
- 487 **Blackburn TM, Cassey P, Lockwood JL. 2009.** The role of species traits in the establishment  
488 success of exotic birds. *Global Change Biology* **15**:2852–2860 doi:10.1111/j.1365-  
489 2486.2008.01841.x
- 490 **Boogert NJ, Reader, SM, Laland KN. 2006.** The relation between social rank, neophobia and  
491 individual learning in starlings. *Animal Behaviour* **72**:1229-1239  
492 doi:10.1016/j.anbehav.2006.02.021
- 493 **Boogert NJ, Anderson RC, Peters S, Searcy WA, Nowicki S. 2011.** Song repertoire size in  
494 male song sparrows correlates with detour reaching, but not with other cognitive  
495 measures. *Animal Behaviour* **81**:1209-1216 doi:10.1016/j.anbehav.2011.03.004
- 496 **Buckner C. 2013.** A property cluster theory of cognition. *Philosophical Psychology* **28**:307-336  
497 doi: 10.1080/09515089.2013.843274
- 498 **Burnham KP, Anderson DR. 2002.** *Model selection and multimodel inference: a practical*  
499 *information- theoretic approach*, 2nd edn. New York, NY: Springer.

- 500 **Chow PKY, Lea SE, Leaver LA. 2016.** How practice makes perfect: the role of persistence,  
501 flexibility and learning in problem-solving efficiency. *Animal Behaviour* **112**:273-283  
502 doi: 10.1016/j.anbehav.2015.11.014
- 503 **Clayton NS, Dickinson A. 1998.** Episodic-like memory during cache recovery by scrub jays.  
504 *Nature* **395**:272-274 doi:10.1038/26216
- 505 **Cole EF, Cram DL, Quinn JL. 2011.** Individual variation in spontaneous problem-solving  
506 performance among wild great tits. *Animal Behaviour* **81**:491-498  
507 doi:10.1016/j.anbehav.2010.11.025
- 508 **Dally JM, Emery NJ, Clayton NS. 2006.** Food-caching western scrub-jays keep track of who  
509 was watching when. *Science* **312**:1662-1665 doi:10.1126/science.1126539
- 510 **Diquelou MC, Griffin AS, Sol D. 2016.** The role of motor diversity in foraging innovations: a  
511 cross-species comparison in urban birds. *Behavioral Ecology* **27**:584-591  
512 doi:10.1093/beheco/arv190
- 513 **Duckworth RA. 2008.** Adaptive dispersal strategies and the dynamics of a range expansion. *The*  
514 *American Naturalist* **172**:S4-S17 doi:10.1086/588289
- 515 **Fox RA, Ladage LD, Roth TC, Pravosudov VV. 2009.** Behavioural profile predicts dominance  
516 status in mountain chickadees, *Poecile gambeli*. *Animal Behaviour* **77**:1441-1448  
517 doi:10.1016/j.anbehav.2009.02.022
- 518 **Gamer M, Lemon J, Fellows I, Singh P. 2012.** Various coefficients of interrater reliability and  
519 agreement. R Package. URL: <http://www.r-project.org> Accessed April 2016.
- 520 **Griffin AS. 2016.** Innova- tiveness as an emergent property: a new alignment of comparative  
521 and experimental research on animal innovation. *Philosophical Transactions of the Royal*  
522 *Society B* **371**:20150544 doi:10.1098/rstb.2015.0544



523 **Griffin AS, Diquelou MC. 2015.** Innovative problem solving in birds: a cross-species  
 524 comparison of two highly successful passerines. *Animal Behaviour* **100**:84-94  
 525 10.1016/j.anbehav.2014.11.012

526 **Griffin AS, Diquelou M, Perea M. 2014.** Innovative problem solving in birds: a key role of  
 527 motor diversity. *Animal Behaviour* **92**:221-227 doi: 10.1016/j.anbehav.2014.04.009

528 **Griffin AS, Guez D. 2014.** Innovation and problem solving: a review of common mechanisms.  
 529 *Behavioural Processes* **109**:121-134 doi:10.1016/j.beproc.2014.08.027

530 **Griffin AS, Guez D, Lermite F, Patience M. 2013.** Tracking changing environments:  
 531 Innovators are fast, but not flexible learners. *PLoS ONE* **8**:e84907  
 532 doi:10.1371/journal.pone.0084907

533 **Guenther A, Brust V, Dersen M, Trillmich F. 2014.** Learning and personality types are related  
 534 in caviés (*Cavia aperea*). *Journal of Comparative Psychology* **128**:74-81  
 535 doi:10.1037/a0033678

536 **Guenther A, Finkemeier MA, Trillmich F. 2014.** The ontogeny of personality in the wild  
 537 guinea pig. *Animal Behaviour* **90**:131-139 doi:10.1016/j.anbehav.2014.01.032

538 **Guillette LM, Reddon AR, Hoeschele M, Sturdy CB. 2011.** Sometimes slower is better: slow-  
 539 exploring birds are more sensitive to changes in a vocal discrimination task. *Proceedings*  
 540 *of the Royal Society of London B* **278**:767-773 doi:10.1098/rspb.2010.1669

541 **Hadfield J. 2010.** MCMCglmm: Markov chain Monte Carlo methods for generalised linear  
 542 mixed models.  
 543 [http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=E395FF7367BF580D916E465F](http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=E395FF7367BF580D916E465F55302C60?doi=10.1.1.160.5098&rep=rep1&type=pdf)  
 544 [55302C60?doi=10.1.1.160.5098&rep=rep1&type=pdf](http://citeseerx.ist.psu.edu/viewdoc/download;jsessionid=E395FF7367BF580D916E465F55302C60?doi=10.1.1.160.5098&rep=rep1&type=pdf) Accessed 1 Jan 2016

545 **Hadfield J. 2014.** MCMCglmm: MCMC generalised linear mixed models. R package.  
 546 <https://cran.r-project.org/web/packages/MCMCglmm/index.html> Accessed 1 Jan 2016

547 **Herborn KA, Macleod R, Miles WT, Schofield AN, Alexander L, Arnold KE. 2010.**  
 548 Personality in captivity reflects personality in the wild. *Animal Behaviour* **79**:835-843.  
 549 doi:10.1016/j.anbehav.2009.12.026

550 **Herborn KA, Coffey J, Larcombe SD, Alexander L, Arnold KE. 2011.** Oxidative profile  
 551 varies with personality in European greenfinches. *Journal of Experimental Biology*  
 552 **214**:1732-1739 doi:10.1242/jeb.051383

553 **Herrmann E, Hernández-Lloreda MV, Call J, Hare B, Tomasello M. 2010.** The structure of  
 554 individual differences in the cognitive abilities of children and chimpanzees.  
 555 *Psychological Science* **21**:102-110 doi:10.1177/0956797609356511

556 **Homberg JR, Pattij T, Janssen MC, Ronken E, De Boer SF, Schoffelmeer AN, Cuppen E.**  
 557 **2007.** Serotonin transporter deficiency in rats improves inhibitory control but not  
 558 behavioural flexibility. *European Journal of Neuroscience* **26**:2066–2073.

559 **Isden J, Panayi C, Dingle C, Madden J. 2013.** Performance in cognitive and problem-solving  
 560 tasks in male spotted bowerbirds does not correlate with mating success. *Animal*  
 561 *Behaviour* **86**:829-838 doi:10.1016/j.anbehav.2013.07.024

562 **Johnson K, Peer BD. 2001.** Great-tailed grackle (*Quiscalus mexicanus*). In *The birds of North*  
 563 *America online* (ed. A Poole). Ithaca, NY: Cornell Lab of Ornithology.

564 **Jolles JW, Fleetwood-Wilson A, Nakayama S, Stumpe MC, Johnstone RA, Manica A. 2014.**  
 565 The role of previous social experience on risk-taking and leadership in three-spined  
 566 sticklebacks. *Behavioral Ecology* **25**:1395-1401 doi:10.1093/beheco/aru146

- 567 **Kozlovsky DY, Branch CL, Pravosudov VV. 2015.** Problem-solving ability and response to  
568 novelty in mountain chickadees (*Poecile gambeli*) from different elevations. *Behavioral*  
569 *Ecology and Sociobiology* **69**:635-643 doi:10.1007/s00265-015-1874-4
- 570 **Lefebvre L, Whittle P, Lascaris E, Finkelstein A. 1997.** Feeding innovations and forebrain  
571 size in birds. *Animal Behaviour* **53**:549-560 doi:10.1006/anbe.1996.0330
- 572 **Lerman SB, Warren PS, Gan H, Shochat E 2012.** Linking foraging decisions to residential  
573 yard bird composition. PLoS ONE 7:e43497. doi:10.1371/ journal.pone.0043497
- 574 **Liu Y, Day LB, Summers K, Burmeister SS. 2016.** Learning to learn: advanced behavioural  
575 flexibility in a poison frog. *Animal Behaviour* **111**:167-172.  
576 doi:10.1016/j.anbehav.2015.10.018
- 577 **Logan CJ. 2016a.** Behavioral flexibility and problem solving in an invasive bird. *PeerJ* 3:e1975.  
578 doi:10.7717/peerj.1975
- 579 **Logan CJ. 2016b (in press).** How far will a behaviourally flexible invasive bird go to innovate?  
580 *Royal Society Open Science*. doi:10.1098/rsos.160247
- 581 **Logan C. 2016c.** Great-tailed grackle neophobia, persistence, exploration, risk aversion, motor  
582 diversity, interobserver reliability data and R code, Santa Barbara, CA USA 2014-2015.  
583 KNB Data Repository. [https://knb.ecoinformatics.org/#view/corina\\_logan.18.13](https://knb.ecoinformatics.org/#view/corina_logan.18.13)
- 584 **Logan C. 2016d.** Great-tailed grackle behavioral flexibility and problem solving experiments,  
585 Santa Barbara, CA USA 2014-2015. KNB Data Repository. doi:10.5063/F1319SVV  
586 <https://knb.ecoinformatics.org/#view/doi:10.5063/F1319SVV>
- 587 **Logan CJ, Harvey B, Schlinger BA, Rensel M. 2016.** Western scrub-jays do not appear to  
588 attend to functionality in Aesop's Fable experiments. *PeerJ* 4:e1707 doi:  
589 10.7717/peerj.1707.

- 590 **Logan CJ, Jelbert SA, Breen AJ, Gray RD, Taylor AH. 2014.** Modifications to the Aesop's
- 591 Fable paradigm change performances in New Caledonian crows. *PLOS ONE* **9**:e103049
- 592 doi:10.1371/journal.pone.0103049
- 593 **Lynn DA, Brown GR. 2009.** The ontogeny of exploratory behavior in male and female
- 594 adolescent rats (*Rattus norvegicus*). *Developmental Psychobiology* **51**:513-520.
- 595 doi:10.1002/dev.20386
- 596 **Manrique HM, Völter CJ, Call J. 2013.** Repeated innovation in great apes. *Animal Behaviour*
- 597 **85**:195-202 doi:10.1016/j.anbehav.2012.10.026
- 598 **Montiglio PO, Garant D, Thomas D, Réale D. 2010.** Individual variation in temporal activity
- 599 patterns in open-field tests. *Animal Behaviour* **80**:905-912
- 600 doi:10.1016/j.anbehav.2010.08.014
- 601 **Overington SE, Cauchard L, Cote KA, Lefebvre L. 2011.** Innovative foraging behaviour in
- 602 birds: What characterizes an innovator? *Behavioural Processes* **87**:274-285
- 603 doi:10.1016/j.beproc.2011.06.002
- 604 **Peer BD. 2011.** Invasion of the emperor's grackle. *Ardeola* **58**:405-409
- 605 doi:10.13157/arla.58.2.2011.405
- 606 **R Core Team. 2015.** *R: A language and environment for statistical computing*. R Foundation for
- 607 Statistical Computing, Vienna, Austria. <https://www.R-project.org>. Accessed 1 Jan 2016.
- 608 **Raby CR, Alexis DM, Dickinson A, Clayton NS. 2007.** Planning for the future by western
- 609 scrub-jays. *Nature* **445**:919-921 doi:10.1038/nature05575
- 610 **Seed A, Seddon E, Greene B, Call J. 2012.** Chimpanzee 'folk physics': bringing failures into
- 611 focus. *Philosophical Transactions of the Royal Society B* **367**:2743-2752
- 612 doi:10.1098/rstb.2012.0222

- 613 **Sih A, Del Giudice M. 2012.** Linking behavioural syndromes and cognition: a behavioural  
614 ecology perspective. *Philosophical Transactions of the Royal Society B* **367**:2762-2772  
615 doi: 10.1098/rstb.2012.0216
- 616 **Shettleworth SJ. 2010.** Cognition, Evolution, and Behavior (2nd ed). Oxford University Press.
- 617 **Skutch AF. 1954.** *Life histories of Central American birds: families Fringillidae, Thraupidae,*  
618 *Icteridae, Parulidae and Coerebidae (No. 598.29728 S629).* Berkeley, CA: Cooper  
619 Ornithological Society.
- 620 **Sol D, Griffin AS, Bartomeus I. 2012.** Consumer and motor innovation in the common myna:  
621 the role of motivation and emotional responses. *Animal Behaviour* **83**:179-188  
622 doi:10.1016/j.anbehav.2011.10.024
- 623 **Sol D, Lefebvre L. 2000.** Behavioural flexibility predicts invasion success in birds introduced to  
624 New Zealand. *Oikos* **90**:599–605 doi:10.1034/j.1600- 0706.2000.900317.x
- 625 **Thornton A, Lukas D. 2012.** Individual variation in cognitive performance: developmental and  
626 evolutionary perspectives. *Philosophical Transactions of the Royal Society B* **367**:2773-  
627 2783 doi:10.1098/rstb.2012.0214
- 628 **Verbeek ME, Drent PJ, Wiepkema PR. 1994.** Consistent individual differences in early  
629 exploratory behaviour of male great tits. *Animal Behaviour* **48**:1113-1121  
630 doi:10.1006/anbe.1994.1344
- 631 **Wehtje W. 2003.** The range expansion of the great-tailed grackle (*Quiscalus mexicanus* Gmelin)  
632 in North America since 1880. *Journal of Biogeography* **30**:1593-1607 doi:  
633 10.1046/j.1365-2699.2003.00970.x
- 634 **Zimmer C, Boogert NJ, Spencer KA. 2013.** Developmental programming: cumulative effects  
635 of increased pre-hatching corticosterone levels and post-hatching unpredictable food

636 availability on physiology and behaviour in adulthood. *Hormones and Behavior* **64**:494-  
 637 500 doi: 10.1016/j.yhbeh.2013.07.002

# **Table 1**(on next page)

## Exploration and risk aversion results

The percentage of time spent in each aviary section, their risk aversion score (percent time spent in the riskiest sections of the aviary; sections 4 and 5) and their exploration score (total number of section changes).

1 Table 1. Exploration and risk aversion results: the percentage of time spent in each aviary  
 2 section, their risk aversion score (percent time spent in the riskiest sections of the aviary; sections  
 3 4 and 5) and their exploration score (total number of section changes).

Bird	Aviary Section					Risk Aversion Score (% time in risky sections)	Exploration Score (section changes)
	1	2	3	4	5		
Tequila	94	0.4	0.5	6	0	6	16
Margarita	96	0	0.1	4	0	4	5
Cerveza	95	3	0	2	0	2	8
Michelada	92	0.06	0	6	2	8	19
Horchata	47	35	5	14	0	14	145
Refresco	100	0	0	0	0	0	0
Batido	44	0.6	0	0	55	55	30
Jugo	73	12	2	3	11	14	163

4



## **Table 2**(on next page)

### Neophobia results

Neophobia scores for each novel object and an overall score for each individual. Neophobia score calculations: the latency to land on the table in controls (trials 1 and 3 averaged) minus the latency in the novel object condition (trial 2) for each object type (GoPro camera, stone dropping apparatus, and U-tube apparatus), and summed across object types for the overall neophobia score (positive=less neophobic [bold text], negative=more neophobic).

1 Table 2. Neophobia scores for each novel object and an overall score for each individual.  
 2 Neophobia score calculations: the latency to land on the table in controls (trials 1 and 3  
 3 averaged) minus the latency in the novel object condition (trial 2) for each object type (GoPro  
 4 camera, stone dropping apparatus, and U-tube apparatus), and summed across object types for  
 5 the overall neophobia score (positive=less neophobic [bold text], negative=more neophobic).

Bird	GoPro	Stone dropping apparatus	U-tube	Neophobia score
Tequila	7	-444.5	-156.5	-594
Margarita	<b>20</b>	0	0	<b>20</b>
Cerveza	-182	<b>167.5</b>	-42.5	-57
Michelada	0	0	-228	-228
Horchata	-580	-1	-277.5	-858.5
Refresco	<b>1</b>	<b>148.5</b>	<b>1</b>	<b>150.5</b>
Batido	<b>187</b>	-275.5	-541	-629.5
Jugo	<b>338</b>	-227.5	-373.5	-263

6

# **Table 3**(on next page)

Motor diversity ethogram

Description of motor actions used while presented with a stick tool use task (techniques 1, 2, 4, 5, 13 and 14 are from Griffin & Diquelou (2015) who refer to ‘gape’ as ‘lever’).

- 1 Table 3. Description of motor actions used while presented with a stick tool use task (techniques  
2 1, 2, 4, 5, 13 and 14 are from Griffin & Diquelou 2015 who refer to ‘gape’ as ‘lever’).

<b>Technique</b>	<b>Description</b>	<b>Body part</b>
1. Vertical peck	Pecks vertically to the horizontal surface of the apparatus with beak open or closed	Bill
2. Horizontal peck	Pecks horizontally to the vertical edges of the apparatus with beak open or closed	
3. Upside Down Peck	Pecks horizontally to the vertical edges of the apparatus while standing on top of the apparatus, thus the head is upside down	
4. Vertical push	Makes closed beak contact with the horizontal surfaces of the apparatus and slides beak vertically along the surface	
5. Grab apparatus	The apparatus is held between the two mandibles	
6. Grab stick	The stick is held between the two mandibles	
7. Pull stick	The stick is held between the two mandibles and pulled	
8. Push stick	The stick is held between the two mandibles and pushed	
9. Move stick	The stick is moved from inside to outside of the apparatus	
10. Manipulate Stick	Manipulate stick inside apparatus	
11. Carry stick away	The stick is held in the beak as the bird flies away from the table	
12. Throw stick	The stick is tossed to the side	
13. Gape	The closed beak is placed under the edge, in an opening, or on a surface of the apparatus and then opened	
14. Gape upside-down	Same as gape but the head is upside-down (or at least 45 degrees from complete upside-down position)	
15. Stand	Stands on top of the apparatus	Feet (or bill)
16. Step	Places one foot on the apparatus	
17. Tips apparatus	Tips apparatus over after standing on top and flying off or by grabbing with bill and pulling over	

3

# **Table 4**(on next page)

Persistence and motor diversity results, and behavioral flexibility scores

Persistence (the total number of times a bird landed on the table, touched the apparatus, or touched the stick), motor diversity (the total number of motor actions used), and behavioral flexibility scores (number of trials to reverse a preference minus the number of trials to initially learn the preference; from Logan 2016a) per bird (- = did not complete this experiment).

1 Table 4. Persistence (the total number of interactions with the table, apparatus, and stick), motor  
 2 diversity (the total number of motor actions used), and behavioral flexibility scores (number of  
 3 trials to reverse a preference minus the number of trials to initially learn the preference; from  
 4 Logan 2016a) per bird (- = did not complete this experiment).

<b>Bird</b>	<b>Sex</b>	<b>Persistence</b>	<b>Motor diversity</b>	<b>Behavioral flexibility score</b>
Tequila	M	175	6	70
Margarita	F	72	5	70
Cerveza	F	81	2	60
Michelada	F	18	1	30
Horchata	F	145	8	100
Refresco	M	1114	14	50
Batido	M	4047	15	-
Jugo	M	197	6	40

5

**Figure 1**(on next page)

Predictions about how behavioral flexibility will relate to other behaviors

Predictions are adapted from Table 1 in Sih and Del Giudice (2012).

## **More behaviorally flexible** (fast reversal learning)

Generally low speed-high accuracy  
 Slow behavioral type  
 Collects more information  
 Slower to learn new activities  
 Faster to learn to avoid  
 Less impulsive  
 Risk averse  
 Slow exploring  
 Cautious  
 Rarely exposed to novel environments

## **Less behaviorally flexible** (slow reversal learning)


Generally high speed-low accuracy  
 Fast behavioral type  
 Collects less information  
 Faster to learn new activities  
 Slower to learn to avoid  
 More impulsive  
 Risk seeking  
 Fast exploring  
 Bold  
 Commonly exposed to novel environments



## Figure 2 (on next page)

### Aviary diagram

The aviary was invisibly sectioned (dotted lines) from least (1) to most risky (4 and 5). The camera was positioned outside a door at the front of the aviary. Food and water bowls were on the ground at the front of the aviary (grey circles) and perches were located in all upper corners (purple lines).



# **Figure 3**(on next page)

Behavioral flexibility scores in relation to other behaviors

The relationship between behavioral flexibility scores and exploration (A, total number of aviary section changes), risk aversion (B, percentage of time spent in risky aviary sections), persistence (C, total number of interactions with the table, apparatus, and stick), motor diversity (D, total number of different motor actions used), and neophobia (E, latency to land on table during controls minus latency to land next to a novel object) (n=7 grackles).

