

Temporal overlaps of feral cats with prey and competitors in primary and human-altered habitats on Bohol Island, Philippines

Vlastimil Bogdan¹, Tomáš Jůnek^{Corresp., 1}, Pavla Jůnková Vymyslická¹

¹ Department of Ecology / Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic

Corresponding Author: Tomáš Jůnek
Email address: tom_junek@yahoo.com

The vertebrate fauna of the Philippines, known for its diversity and high proportion of endemic species, comprises mainly small- to medium-sized forms with a few large exceptions. As with other tropical ecosystems, the major threats to wildlife are habitat loss, hunting and invasive species, of which the feral cat (*Felis catus*) is considered the most damaging. Our camera-trapping study focused on a terrestrial vertebrate species inventory on Bohol Island and tempo-spatial co-occurrences of feral cats with their prey and competitors. The survey took place in the Rajah Sikatuna Protected Landscape, and we examined primary rainforest, its border with agricultural land, and rural areas in the vicinity of villages. Altogether, over 2,885 trap days we captured 30 species of vertebrates – 10 mammals (including *Sus philippensis*), 19 birds and one reptile, *Varanus cumingi*. We trapped 81.8% of expected vertebrates. Based on the number of events, the most frequent native species was the barred rail (*Gallirallus torquatus*). The highest overlap in diel activity between cats and potential prey was recorded with rodents in rural areas ($\Delta=0.62$); the lowest was in the same habitat with ground-dwelling birds ($\Delta=0.40$). Cat activity was not recorded inside the rainforest; in other habitats their diel activity pattern differed. The cats' activity declined in daylight in the proximity of humans, while it peaked at the transition zone between rainforest and fields. Both rodents and ground-dwelling birds exhibited a shift in activity levels between sites where cats were present or absent. Rodents tend to become active by day in cat-free habitats. No cats' temporal response to co-occurrences of civets (*Paradoxurus hermaphroditus* and *Viverra zibetha*) was found but cats in diel activity avoided domestic dogs (*Canis lupus familiaris*). Our first insight into the ecology of this invasive predator in the Philippines revealed an avoidance of homogeneous primary rainforest and a tendency to forage close to human settlements in heterogeneous habitats. A detailed further investigation of the composition of the cat's diet, as well as ranging pattern, is still needed.

Title

Temporal overlaps of feral cats with prey and competitors in primary and human-altered habitats on Bohol Island, Philippines

List of Authors

Vlastimil Bogdan¹, Tomáš Jůnek¹, Pavla Jůnková Vymyslická¹

¹*Department of Ecology, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6 – Suchbátka, Czech Republic,*

Corresponding Author

Tomáš Jůnek,

Lindnerova 448/8, 180 00 Prague, Czech Republic

E-mail address: tom_junek@yahoo.com

Abstract

The vertebrate fauna of the Philippines, known for its diversity and high proportion of endemic species, comprises mainly small- to medium-sized forms with a few large exceptions. As with other tropical ecosystems, the major threats to wildlife are habitat loss, hunting and invasive species, of which the feral cat (*Felis catus*) is considered the most damaging. Our camera-trapping study focused on a terrestrial vertebrate species inventory on Bohol Island and temporal co-occurrences of feral cats with their prey and competitors. The survey took place in the Rajah Sikatuna Protected Landscape, and we examined primary rainforest, its border with agricultural land, and rural areas in the vicinity of villages. Altogether, over 2,885 trap days we captured 30 species of vertebrates – 10 mammals (including *Sus philippensis*), 19 birds and one reptile, *Varanus cumingi*. We trapped 81.8% of expected vertebrates. Based on the number of events, the most frequent native species was the barred rail (*Gallirallus torquatus*). The highest overlap in diel activity between cats and potential prey was recorded with rodents in rural areas ($\Delta=0.62$); the lowest was in the same habitat with ground-dwelling birds ($\Delta=0.40$). Cat activity was not recorded inside the rainforest; in other habitats their diel activity pattern differed. The cats' activity declined in daylight in the proximity of humans, while it peaked at the transition zone between rainforest and fields. Both rodents and ground-dwelling birds exhibited a shift in activity levels between sites where cats were present or absent. Rodents tend to become active by day in cat-free habitats. No cats' temporal response to co-occurrences of civets (*Paradoxurus hermaphroditus* and *Viverra zibetha*) was found but cats in diel activity avoided domestic dogs (*Canis lupus familiaris*). Our first insight into the ecology of this invasive predator in the

Philippines revealed an avoidance of homogeneous primary rainforest and a tendency to forage close to human settlements in heterogeneous habitats. A detailed further investigation of the composition of the cat's diet, as well as ranging pattern, is still needed.

Introduction

The Philippine Archipelago is considered a global biodiversity hotspot, known for its high proportion of endemic species (Ambal *et al.* 2012). The terrestrial vertebrate taxa, which primarily encompass small to medium sized species, inhabit more than 7100 islands. These species include at least 213 mammals (Heaney *et al.* 2010), 674 birds (Lapage 2015), 270 reptiles and 111 amphibians (BREO 2015).

Similar to other oceanic islands, the predominantly small fauna of the Philippines suffers from the presence of competing invasive species, such as *Rattus* spp., and the feral cat (*Felis catus*). The cat is listed as the most widespread and probably most damaging of the four carnivores included on the list of the 100 worst invasive species (Lowe *et al.* 2000). At least 175 vertebrates are threatened or have been driven to extinction by feral cats on at least 120 islands (Medina *et al.* 2011). Meta-analysis has revealed that the negative impact of feral cats is largest for insular endemic mammals, and is exacerbated by the presence of invasive cat prey species such as mice (*Mus musculus*) or rabbits (*Oryctolagus cuniculus*) (Nogales *et al.* 2013). The cat is widely kept as a pet by people throughout the Philippines and can be found foraging in every habitat (Duffy & Capece 2012). Despite the general prevalence of cats in the Philippine landscape, there is a noticeable lack of knowledge regarding the cat's impact on the biodiversity of this archipelago.

Cats feed on a wide range of animals, from arthropods, reptiles and birds to mammals the size of a rabbit (Pearre, Maass & Maass 1998). In Australia alone, with a variety of animals of

similar size such as those found in the Philippines, 400 prey species consumed by cats have been recorded (Doherty *et al.* 2015). In the Philippines, members of the orders Chiroptera and Rodentia are the most numerous mammalian species (Heaney *et al.* 2010). A wide range of terrestrial and arboreal rodents with body mass ranging from the 15-g *Musseromys* spp. to the 2.6-kg *Phloeomys* spp. risk predation by cats. Only adult individuals of *Phloeomys* and *Hystrix pumila* (Heaney *et al.* 2010) exceed the potential prey dimensions. According to size and niche, members of the Tupaiidea (treeshrews), Erinaceidae (moonrats) and Soricidae family (shrews) should be listed as mammalian prey for cats. Similarly, the smallest Philippine primate, *Tarsius syrichta*, which inhabits Bohol and other islands of the Mindanao faunal region, can be included (MacKinnon & MacKinnon 1980).

On Bohol Island (3,269 km²), as on the other Philippine islands, bats and rodents dominate among local mammals. The small mammalian fauna consists of one insectivorous species and nine species of rodents, including the introduced *Mus musculus*, *Rattus rattus*, *Rattus norvegicus*, *Rattus tanezumi* and *Rattus exulans* (Heaney *et al.* 2010). The avifauna of Bohol numbers 235 species, with Passeriformes forming the largest sub-group at 83 species. Bohol is also home to 14 ground-dwelling bird species inhabiting the woody or bushy inland habitats potentially affected by cats (Kennedy 2000).

Along with dogs (*Canis lupus familiaris*), possible competitors of cats on Bohol include two mammalian carnivores, Asian palm civet (*Paradoxurus hermaphroditus*) and Malayan civet (*Viverra zibetha*) (Heaney *et al.* 2010) and two reptile species: yellow-headed water monitor (*Varanus cumingi*) and reticulated python (*Python reticulatus*) (BREO 2015). To our knowledge, no predation between cat and civets has been published.

The timing of activity of mammalian predators is a well discussed topic (e.g. Palomares & Caro 1999; Tambling *et al.* 2015). Time-stamped records from camera traps allow for detailed insights into the time budget and temporal coexistence of animals across trophic guilds, seasons, etc. (Rowcliffe *et al.* 2014), and recent camera trapping studies have successfully examined overlaps in diel activity patterns (Ridout & Linkie 2009), confirming significant activity overlap between carnivores and their preferred prey (Harmsen *et al.* 2009; Lucherini *et al.* 2009; Sweitzer & Furnas 2016) and suggesting predator behavior to reduce foraging energy expenditure (Foster *et al.* 2013). In their role as mesopredators cats must optimize their use of time not only to encounter prey but also to cope with a sympatric superior predator (Brook, Johnson & Ritchie 2012). The combination of partitioning of habitat, prey size and a 24-hour daily cycle is thought to be a complex mechanism allowing competing felids to coexist in different animal communities (Di Bitetti *et al.* 2010; Foster *et al.* 2013; Silmi, Anggara & Dahlen 2013; Sunarto *et al.* 2015). For example, low overlap in activities has been found between marbled cat (*Pardofelis marmorata*) and clouded leopard (*Neofelis nebulosa*) in Thailand (Lynam *et al.* 2013). Wang & Fisher (2012) also confirmed higher segregation of diel activities of cats with respect to dingoes during wet months. The particularly suppressive effect of an apex carnivore on invasive populations of cats is considered an important conservation issue (Brook, Johnson & Ritchie 2012; Lazenby & Dickman 2013; Doherty, Bengsen & Davis 2015).

We conducted a camera-trap survey on Bohol Island in an attempt to uncover tempo-spatial co-occurrences of terrestrial vertebrate species on regularly used trails and to confirm the presence of cats in the protected primary rainforest (Zone I), a transition zone along the border of the primary rainforest with the agricultural landscape (Zone II), and inside the rural landscape in the proximity of human settlements (Zone III). Our objectives were to: (1) create a general inventory

of camera-trapped taxa; (2) model the species accumulation curve using previous knowledge of the possible number of mammalian, avian and reptile species detectable by camera-traps; and (3) compare the diel activity levels of cats with those of potential prey and competitors.

Materials & Methods

Study site

Our study was conducted under research permit No. 2014-04, issued by DENR, Region VII, Philippines, between July 2nd and December 4th, 2014 in the surroundings of the town of Bilar, Bohol Island, Philippines. The landscape consists of a mixture of distinctive flat rural areas near human settlements, used as rice fields and plantations for various crops, steep karst hills covered by brush and secondary forest, and primary rainforest in protected areas. The town of Bilar lies between two conservation areas, the Rajah Sikatuna Protected Landscape (RSPL) and the Loboc River Watershed Forest Reserve. RSPL is the second largest protected sanctuary on Bohol, covering 11,034 ha of a mostly hilly limestone environment rich in characteristic landforms such as ravines, sinkholes and caves. The altitude in RSPL varies between 300 and 826 m above sea level. The forest canopy is multi-layered, with trees reaching up to 20 m in height. Members of the families Dipterocarpaceae, Moraceae and Melicaceae dominate the canopy. Certain regions of RSPL have been reforested with white teak (*Gmelina arborea*) and Honduras mahogany (*Swietenia macrophylla*) (Barcelona *et al.* 2006). The average annual precipitation reaches 1,600 mm; the rainy season typically lasts from June to December, with an increase in precipitation to 200 mm per month. The driest month is April when approximately 40 mm of rain falls.

Sampling design

We monitored three types of landscape typical of tropical regions and deployed cameras in groups, one camera per location, at eight trapping sites (Fig. 1): Zone I – protected primary rainforest including the Watershed Forest Reserve (site WS), interior of RSPL (site SP) and abandoned farms in the early stages of succession into RSPL (site SF); Zone II – transition zone between the primary rainforest of RSPL and rice fields close to the village of Bulak (site BU), transition zone between RSPL and rice and corn fields close to Logarita Springs (site LS), and transition zone between RSPL and the farms of the village Binantay (site BI); Zone III – mixture of brush and degraded forest and plantations on the edge of the village of Subayon (site SU), and at Bohol Habitat Conservation Center on the edge of the town of Bilar (site HB). Details on camera traps' deployment and duration of sampling are shown in Table 1.

Sampling procedures

We used 41 weatherproof infrared digital camera traps – 29 units of Ltl Acorn 5210MC (Shenzhen Ltl Acorn Electronics Co., Ltd.) and 12 units of SPYPOINT IR7 (SPYPOINT^{MD}, G.G. Telecom). Prior to the study, we tested both types of cameras in a week-long trial which was focused on the difference in detection rates for moving objects. No difference larger than 10 % between numbers of independence events was found. Both types of cameras were also used in every habitat to avoid a bias from site-specific detection rates. Cameras were set up to perform the same delay between recordings – SPYPOINT to take two images with a delay of 10 s between consecutive triggering, and Ltl Acorns to take one picture followed by a 5 s video, with a 5 s delay between triggering. Video sequences served as an additional tool for the identification of species.

We placed all cameras opportunistically on the most frequented trails or their junctions and, according to the expected size of target vertebrates, we fastened cameras with a belt onto the

trunks of trees or bushes nearest to the trail, at a height of up to 0.5 m, with a focal point approximately 2 m from the lens. All cameras were active 24 hours a day; all records in infrared mode were available only in a black-and-white version. No bait was used.

Identification of taxa

Two observers, VB and TJ, independently identified all species visually from images and videos; the results were mutually crosschecked, and disagreeing or unidentifiable records were excluded from the analysis. Based on available databases (Heaney *et al.* 2010; BREO 2015; Lapage 2015), we made a list of terrestrial mammalian and avian ground-dwelling species known or expected to occur on Bohol (Table 2). From reptiles, we included only the largest four-legged taxon, the yellow-headed water monitor (*Varanus cumingi*). Members of the order Chiroptera and the strictly arboreal Philippine colugo (*Cynocephalus volans*) were *a priori* omitted. The conservation status of each species was assessed following IUCN (2015).

Due to the limited nature of the recordings, for the identification process and the calculation of a species accumulation curve all taxa the size of a mouse (*Mus musculus* and also the insectivorous *Crocidura beatus*) were pooled into the group called ‘mice’, and all species of rats (*Rattus* spp. and *Bullimus bagobus*) into the group ‘rats’. In addition, both known species of squirrels (*Exilisciurus concinnus* and *Sundasciurus philippinensis*) were grouped into one taxon: ‘squirrels’. In total, the list consisted of eight taxa of mammals, 13 birds and one reptile. For purposes of *overlap* analyses between cats and their competitors and prey, we pooled both native carnivore species into a group called ‘civets’ and put mice, rats and squirrels into the group ‘rodents’. Ground-dwelling species of birds were the second analyzed group of prey; dogs were accordingly examined as competitors.

Data analysis

Photographs were defined as events (or activity records) when the delay between two consecutive images of an individual exceeded 10 min. The same individual could theoretically trigger more than one camera within 10 min. For each species and Zone, in Table 2 we reported occurrences of species at cameras represented by events (Lazenby & Dickman 2013).

We used a species accumulation curve based on the cumulative number of camera-trapping days, computed in EstimateS Version 9.1.0 (Colwell 2013), to find out if our survey lasted a sufficient number of days to capture the 22 expected terrestrial vertebrate species (including three pooled groups) known from Bohol. We followed Tobler *et al.* (2008) and calculated well-performing estimators of species richness: the non-parametric abundance-based estimator ACE, and the non-parametric incidence-based estimators ICE and Jackknife 1. An abundance-based rarefaction approach with 95% confidence intervals and 1,000 random iterations of sample order was used.

The pair-wise temporal overlap of selected activity patterns was analyzed using the R statistical environment package ‘*overlap*’ (Meredith & Ridout 2014). Following Ridout and Linkie (2009), we applied kernel density estimation on circular data pooled within all study sites. Density of activity (y-axis) uses a von Mises kernel, corresponding to a circular distribution, and is based on recorded time of each event on 24-hour x-axis. The coefficient of overlap (Δ) was calculated with a smoothing parameter of 1.0. We used a smoothed bootstrap of 10,000 resamples to determine standard errors and 95% confidence intervals. We only analyzed combinations of pairs of species, which scored at least 30 events in the activity record (Ridout, pers. comm.) in a given environment. The number of events used for calculation of the activity pattern overlap for each analyzed group of animals and each location is shown in Table 3.

Results

206 *Species inventory*

207 During the whole survey period, lasting 155 days, we accumulated 2,885 trap days and
 208 2,034 events. The combined capture rate across all sites was 73.1 events per 100 trap days. The
 209 list of all 30 animal taxa recorded is shown in Table 2.
 210 The most frequent native species was the barred rail (*Gallirallus torquatus*), captured in 183
 211 independence events. We did not record four expected bird species: *Megapodius cumingii*,
 212 *Coturnix chinensis*, *Turnix sylvaticus* and *Gallinago megala*. On the other hand, we confirmed
 213 the survival of the Philippine warty pig (*Sus philippensis*). Given its size, it was probably a male
 214 individual that was captured, only once, on three images on August 9th (6:35 pm) in a mud
 215 wallow in the interior of RSPL.

216 We found that feral cats most often occurred in the Zone II and III, and were absent
 217 inside the primary forest. A similar trend was found for ground-dwelling birds. Most rats and
 218 other small mammals were recorded in the transition Zone II between the RSPL forest and
 219 agricultural land. Along with feral cats and domestic dogs, we also recorded all three medium-
 220 sized mammals occurring on Bohol – the common palm civet (59 events), Malay civet (16
 221 events) and long-tailed macaque (*Macaca fascicularis*) (7 events). Humans were also captured
 222 but excluded from the analysis.

223 Within all eight sampling sites, we captured 18 of 22 expected target taxa, which
 224 corresponds to a success rate of 81.8% of the species inventory (100% of mammals and reptiles,
 225 69.2% of birds). We used these 18 taxa for calculating the species accumulation curve (Fig. 2).
 226 The mean estimated species richness computed in EstimateS was 19.7 species (ACE = 19.6, ICE
 227 = 19.5 and Jackknife 1 = 20.0). We recorded 15.89 species (72.2% of expected species) in 1,000

trap days. The eight target species of mammals were captured in 1,723 trap days; similarly, nine ground-dwelling birds were recorded within 1,435 trap days.

Temporal overlaps

We recorded cats only in transition Zone II and in the rural landscape close to human settlements (Zone III). Diel activity patterns of cats differed among zones (Fig. 3). Cats showed a decrease in late-afternoon activity near villages, whereas activity in the transition area peaked right before noon. Generally, the activity of cats by daylight was higher in transition zones; in Zone III cats were recorded mainly at night.

The highest overlap in activity patterns between cats and rodents (Table 4) was found in the rural landscape of Zone III, and between cats and ground-dwelling birds in transition Zone II (Fig. 4).

Both categories of potential prey showed shifts in temporal occurrence within sites, based on the presence of cat (Fig. 5). As seen, the peaks of rodent activity decreased in the hours before sunrise and increased after sunset, whereas the activity of ground-dwelling birds peaked about 4 hours sooner at sites where cats were not recorded.

Cats showed the second lowest overlap among all groups with dogs in Zone III (Table 4) where dogs were dominant and active during the day. In Zone II these two animals appeared to peak in their activity at different times: dogs were most active in the morning and late afternoon, whereas cats peaked before noon (Fig. 4).

The overlap between the diel activity patterns of cats and both species of civets is shown in Figure 6. Cats exhibited roughly consistent activity throughout a 24-hour period, with no apparent shift caused by the nocturnal occurrence of sympatric civets.

Discussion

According to our knowledge, to date no study of the behavior and ecology of feral cats has been conducted in the Philippines, nor any camera-trap-based species inventory on Bohol. With the exception of the Philippine pygmy squirrel, *Exilisciurus concinnus*, we were able to capture and identify every non-volant mammalian species recorded as occurring on Bohol larger than a mouse, including an individual of *Sus philippensis*, which is considered to be close to extinction (Oliver 1993), even by local people. Camera traps captured 81.8% of known ground-dwelling mammalian, avian, and reptilian species, similar to the 86 % captured in the Amazon rain forest (Tobler *et al.* 2008) or 89% in the lowland rainforest of Borneo (Bernard *et al.* 2013); both those camera-trapping studies were restricted to mammals. In addition, the initially steep shape of our general species accumulation curve corresponds with studies conducted in tropical ecosystems and confirms the robustness of the approach. Similarly to Rovero *et al.* (2014), we captured the majority of selected species in 1,000 trap days, considered a reliable threshold enabling the detection of rare species (O'Brien 2011).

The absence of cats in the interior of primary rainforest seems not to be driven by distance from the nearest human settlements, given that all three monitored sites were up to approximately 3 km from houses. We suggest that the absence of preferred features and habitats in the rain forest may have resulted in camera traps failing to capture cats. Cats typically use a mixture of vegetation cover at ground level which provides both cover and open space for observing their prey; such habitat may increase hunting success (Doherty, Bengsen & Davis 2015). The habitat heterogeneity hypothesis by Tews *et al.* (2004) predicts that heterogeneous habitats offer a greater diversity and density of potential prey than homogeneous ones, which could be conceivable for cats. Linear features in space (e.g. tree lines, roads and other corridors) are generally considered to maximize cat's detectability (Crooks 2002; Bengsen, Butler &

Masters 2012). We would expect to record cats in primary forest mostly on trails (Trolle & Kéry 2005; Harmsen et al. 2010; Anile et al. 2014) but they could disperse into the undergrowth on paths that are undetected.

The presence of competing, potentially dangerous predators in primary forest is unlikely to explain the absence of cats. Dogs and both species of civets were equally present in all three zones. The common palm civet and Malay civet are omnivorous with a distinctive nocturnal activity pattern (Jennings *et al.* 2009) but they forage in the habitat of cats, and given their size we consider them to be competitors of cats. Nonetheless, cats do not show any temporal avoidance, indicating no interspecies competition, which has evolved during almost a 500-year co-existence (Jubair 1999). For a more comprehensive view of possible niche partitioning, as found for example between felids on Sumatra (Sunarto *et al.* 2015), a camera-trapping study should be conducted on Negros, where the Visayan leopard cat (*Prionailurus bengalensis* ssp. *abori*) occurs as a regional direct competitor (IUCN 2015).

Our results (Tables 2 and 3) show that species richness and availability of both prey categories (rodents and birds) was higher, nearly by orders of magnitude, in both human-altered zones than in primary rainforest. We attribute this to the variety of vertebrate and invertebrate prey, which is more abundant in heterogeneous landscapes. In addition, as suggested by Lozano *et al.* (2003), feral cats use a wide range of habitat components to meet their different activity requirements (e.g. hunting, resting), and this landscape offers a mixture of agricultural features with secondary growth, infrastructure and potential human subsidies (Ferreira et al. 2011). Our data clearly support such a tendency to forage relatively close to human settlements, however we were not able to determine from our records whether a photographed animal was feral or domestic.

Although the diel activity pattern of cats was roughly consistent over 24-h periods (Fig. 6), a detailed analysis of zones revealed a decline in diurnal activity in the vicinity of villages, in contrast to an apparent activity peak before noon in Zone II (Fig. 3). Both Zones II and III offered more-or-less the same number of prey species (Table 2). Compared with Zone II, we hypothesize that more uniform diurnal activity of relatively abundant dogs dissuaded cats from daytime foraging in Zone III (Fig. 4). Also subsidies provided by humans in villages could influence cats to remain inactive in shelters and forage at night. We did not detect any sign of cats being spatially excluded by dogs, but our results support findings that cats optimize their timing of hunting behavior to when dogs are less active, hence avoiding potentially dangerous encounters (Brook, Johnson & Ritchie 2012; Wang & Fisher 2012). Cats were more diurnal in Zone II. This could be explained by the same factor, because the zone-specific relative abundance index of dogs was two times lower than the index of cats than in Zone III (Table 3). So cats could respond both to lower disturbance from dogs and to higher diurnal availability of rodents in Zone II (Fig. 4). Other prey not detected by cameras such as insects or lizards might also be present (Bonnaud *et al.* 2011).

Prey species showed shifts in diel activity patterns between sites where cats were, or were not, present (Fig. 5). When cats were absent, rodents tended to forage visibly by day, while the activity of ground-dwelling birds peaked about 4 hours later. It is difficult to interpret the shift in bird activity; data from sites without cats were considered too scarce to perform a reliable analysis. Rodents shift their activities to become nocturnal if cats are present and more diurnal (Doherty *et al.* 2015). This raised the question of whether almost 500 years of cat presence in the Philippines has driven adaptive mechanisms of prey and competitors to cope with a new predator or not. Our results suggest that this already happened, similarly to the 4000-year history of the

dingo in Australia (Carthey & Banks 2012). Nevertheless, we believe that further research is needed, especially throughout all seasons.

Knowledge of feral cat diet is paradoxically the least researched in tropical habitats with the richest terrestrial biodiversity (Doherty *et al.* 2015; Doherty, Bengsen & Davis 2015). Our findings reveal the first tempo-spatial co-occurrences between feral cats and their potential prey in a typical mixture of Philippine landscapes. We suggest feral cats' temporal avoidance of dogs as the apex predator. We confirm that camera traps are capable of capturing small-bodied fauna, ground-dwelling birds and highly elusive species, such as *Sus philippensis*, as well. Endangered Philippine fauna exposed to invasive species should rapidly become the target of a broad and long-term camera-trapping inventory survey. For an in-depth knowledge of the dietary intake of feral cats in the Philippines, DNA analysis of scat is recommended as a priority for researchers (Nogales *et al.* 2013). In addition, collared and GPS-tracked cats would provide information about habitat use and the size of home ranges. Finally, attention should be paid to the cultural value of cats kept as pets within Philippine society, to inform eradication strategies.

Acknowledgements

We would like to acknowledge the following persons and organizations which made our study possible: Mr. Isabelo R. Montejó, regional director, and Mr. Eusalem S. Quiwag from the Department of Environment and Natural Resources (DENR), Region VII, Philippines; the staff of Rajah Sikatuna PL; Ms. Cristy Burlace and staff from the Habitat Bohol; Dr. Petr Anděl from the Czech University of Life Sciences Prague; Ms. Monika Drimlová for tireless field work; and two anonymous reviewers for comments on this manuscript.

References

Ambal, R.G.R., Duya, M. V, Cruz, M.A., Coroza, O.G., Vergara, S.G., Silva, N. De &

- Molinyawe, N. (2012) Key Biodiversity Areas in the Philippines : Priorities for Conservation. *Journal of Threatened Taxa*, **4**, 2788–2796.
- Anile, S., Ragni, B., Randi, E., Mattucci, F. & Rovero, F. (2014) Wildcat population density on the Etna volcano, Italy: A comparison of density estimation methods. *Journal of Zoology*, **293**, 252–261.
- Barcelona, J.F., Dolotina, N.E., Madroñero, G.S., Granert, W.G. & Sopot, D.D. (2006) The ferns and fern allies of the karst forests of Bohol Island, Philippines. *American Fern Journal*, 1–20.
- Bengsen, A.J., Butler, J.A. & Masters, P. (2012) Applying home-range and landscape-use data to design effective feral-cat control programs. *Wildlife Research*, **39**, 258–265.
- Bernard, H., Ahmad, A.H., Brodie, J., Giordano, A.J., International, L. & York, N. (2013) Camera-Trapping survey of mammals in and around Imbak Canyon Conservation Area in Sabah Malaysian Borneo. *The Raffles Bulletin of Zoology*, **61**, 861–870.
- Di Bitetti, M.S., De Angelo, C.D., Di Blanco, Y.E. & Paviolo, A. (2010) Niche partitioning and species coexistence in a Neotropical felid assemblage. *Acta Oecologica*, **36**, 403–412.
- Bonnaud, E., Medina, F.M., Vidal, E., Nogales, M., Tershy, B., Zavaleta, E., Donlan, C.J., Keitt, B., Le Corre, M. & Horwath, S. V. (2011) The diet of feral cats on islands: A review and a call for more studies. *Biological Invasions*, **13**, 581–603.
- BREO. (2015) Biodiversity Research and Education Outreach—Philippines, [Http://philbreo.lifedesks.org/](http://philbreo.lifedesks.org/)
- Brook, L.A., Johnson, C.N. & Ritchie, E.G. (2012) Effects of predator control on behaviour of an apex predator and indirect consequences for mesopredator suppression. *Journal of*

- 365 *applied ecology*, **49**, 1278–1286.
- 366 Carthey, A.J.R. & Banks, P.B. (2012) When does an alien become a native species? A vulnerable
367 native mammal recognizes and responds to its long-term alien predator. *PLoS One*, **7**,
368 e31804.
- 369 Colwell, R.K. (2013) EstimateS: Statistical estimation of species richness and shared species
370 from samples. Version 9.1.0. User's Guide and application.
- 371 Crooks, K.R. (2002) Relative sensitivities of mammalian carnivores to habitat fragmentation.
372 *Conservation Biology*, **16**, 488–502.
- 373 Doherty, T.S., Bengsen, A.J. & Davis, R.A. (2015) A critical review of habitat use by feral cats
374 and key directions for future research and management. *Wildlife Research*, **41**, 435–446.
- 375 Doherty, T.S., Davis, R.A., Etten, E.J.B., Algar, D., Collier, N., Dickman, C.R., Edwards, G.,
376 Masters, P., Palmer, R. & Robinson, S. (2015) A continental-scale analysis of feral cat diet
377 in Australia. *Journal of Biogeography*, **42**, 964–975.
- 378 Duffy, D.C. & Capece, P. (2012) Biology and Impacts of Pacific Island Invasive Species. 7. The
379 Domestic Cat (*Felis catus*) 1. *Pacific Science*, **66**, 173–212.
- 380 Ferreira, J.P., Leitão, I., Santos-Reis, M. & Revilla, E. (2011) Human-related factors regulate the
381 spatial ecology of domestic cats in sensitive areas for conservation. *PLoS ONE*, **6**, 1–10.
- 382 Foster, V.C., Sarmiento, P., Sollmann, R., Tôrres, N., Jácomo, A.T.A., Negrões, N., Fonseca, C.
383 & Silveira, L. (2013) Jaguar and Puma Activity Patterns and Predator-Prey Interactions in
384 Four Brazilian Biomes. *Biotropica*, **45**, 373–379.
- 385 Harmsen, B.J., Foster, R.J., Silver, S.C., Ostro, L.E.T. & Doncaster, C.P. (2009) Spatial and
386 temporal interactions of sympatric jaguars (*Panthera onca*) and pumas (*Puma concolor*) in a

neotropical forest. *Journal of mammalogy*, **90**, 612–620.

Harmsen, B.J., Foster, R.J., Silver, S., Ostro, L. & Doncaster, C.P. (2010) Differential use of trails by forest mammals and the implications for camera-trap studies: A case study from Belize. *Biotropica*, **42**, 126–133.

Heaney, L.R., Dolar, M.L., Balete, D.S., Esselstyn, J.A., Rickart, E.A. & Sedlock, J.L. (2010) Synopsis of Philippine Mammals, [Http://archive.fieldmuseum.org/philippine_mammals/](http://archive.fieldmuseum.org/philippine_mammals/)

IUCN. (2015) The IUCN Red List of Threatened Species. Version 2015.1., <http://www.iucnredlist.org>

Jennings, A.P., Veron, G., Wilson, D.E. & Mittermeier, R.A. (2009) Family Viverridae (civets, genets and oians). *Handbook of the mammals of the world*, **1**, 174–232.

Jubair, S. (1999) *Bangsamoro, A Nation under Endless Tyranny*. IQ Marin.

Kennedy, R. (2000) *A Guide to the Birds of the Philippines*. Oxford University Press.

Lapage, D. (2015) Avibase – The World Bird Database, [Http://avibase.bsc-eoc.org/](http://avibase.bsc-eoc.org/)

Lazenby, B.T. & Dickman, C.R. (2013) Patterns of Detection and Capture Are Associated with Cohabiting Predators and Prey. *PLoS ONE*, **8**.

Lowe, S., Browne, M., Boudjelas, S. & De Poorter, M. (2000) *100 of the World's Worst Invasive Alien Species: A Selection from the Global Invasive Species Database*. Invasive Species Specialist Group Auckland,, New Zealand.

Lozano, J., Virgós, E., Malo, A.F., Huertas, D.L. & Casanovas, J.G. (2003) Importance of scrub–pastureland mosaics for wild-living cats occurrence in a Mediterranean area: implications for the conservation of the wildcat (*Felis silvestris*). *Biodiversity & Conservation*, **12**, 921–935.

409 Lucherini, M., Reppucci, J.I., Walker, R.S., Villalba, M.L., Wurstten, A., Gallardo, G., Iriarte, A.,
 410 Villalobos, R., Perovic, P., Society, W.C., Paz, L. & Paz, L. (2009) Activity pattern
 411 segregation of carnivores in the high Andes. *Journal of Mammalogy*, **90**, 1404–1409.

412 Lynam, A.J., Jenks, K.E., Tantipisanuh, N., Chutipong, W., Ngoprasert, D., Gale, G.A.,
 413 Steinmetz, R., Sukmasuang, R., Bhumpakphan, N., Grassman, L.I., Cutter, P., Kitamura, S.,
 414 Reed, D.H., Baker, M.C., Mcshea, W., Songsasen, N. & Leimgruber, P. (2013) Terrestrial
 415 activity patterns of wild cats from camera-trapping. *Raffles Bulletin of Zoology*, **61**, 407–
 416 415.

417 MacKinnon, J. & MacKinnon, K. (1980) The behavior of wild spectral tarsiers. *International*
 418 *Journal of Primatology*, **1**, 361–379.

419 Medina, F.M., Bonnaud, E., Vidal, E., Tershy, B.R., Zavaleta, E.S., Josh Donlan, C., Keitt, B.S.,
 420 Corre, M., Horwath, S. V & Nogales, M. (2011) A global review of the impacts of invasive
 421 cats on island endangered vertebrates. *Global Change Biology*, **17**, n/a–n/a.

422 Meredith, M. & Ridout, M.S. (2014) overlap: estimates of coefficients of overlapping for animal
 423 activity patterns. R package version 0.2. 2.

424 Nogales, M., Vidal, E., Medina, F.M., Bonnaud, E., Tershy, B.R. & Campbell, K.J. (2013) Feral
 425 Cats and Biodiversity Conservation: The Urgent Prioritization of Island Management.
 426 *BioScience*, **63**, 804–810.

427 O'Brien, T.G. (2011) Abundance, density and relative abundance: a conceptual framework.
 428 *Camera Traps in Animal Ecology* pp. 71–96. Springer.

429 Oliver, W.L.R. (1993) Pigs, Peccaries, and Hippos Status Survey and Conservation Action Plan.
 430 *IUCN, Gland*, p.219.

- 431 Palomares, F. & Caro, T.M. (1999) Interspecific killing among mammalian carnivores. *The*
432 *American Naturalist*, **153**, 492–508.
- 433 Pearre, S., Maass, R. & Maass, R. (1998) Trends in the prey size-based trophic niches of feral
434 and House Cats *Felis catus* L. *Mammal Review*, **28**, 125–139.
- 435 Ridout, M.S. & Linkie, M. (2009) Estimating overlap of daily activity patterns from camera trap
436 data. *Journal of Agricultural, Biological, and Environmental Statistics*, **14**, 322–337.
- 437 Rovero, F., Martin, E., Rosa, M., Ahumada, J.A. & Spitale, D. (2014) Estimating Species
438 Richness and Modelling Habitat Preferences of Tropical Forest Mammals from Camera
439 Trap Data (ed D Russo). *PLoS ONE*, **9**, e103300.
- 440 Rowcliffe, M.J., Kays, R., Kranstauber, B., Carbone, C. & Jansen, P. a. (2014) Quantifying
441 levels of animal activity using camera-trap data. *Methods in Ecology and Evolution*, n/a–n/a.
- 442 Silmi, M., Anggara, S. & Dahlen, B. (2013) Using leopard cats (*Prionailurus bengalensis*) as
443 biological pest control of rats in a palm oil plantation. *Journal of Indonesia Natural History*,
444 **1**.
- 445 Sunarto, S., Kelly, M.J., Parakkasi, K. & Hutajulu, M.B. (2015) Cat coexistence in central
446 Sumatra: ecological characteristics, spatial and temporal overlap, and implications for
447 management. *Journal of Zoology*, **296**, 104–115.
- 448 Sweitzer, R.A. & Furnas, B.J. (2016) Data from camera surveys identifying co-occurrence and
449 occupancy linkages between fishers (*Pekania pennanti*), rodent prey, mesocarnivores, and
450 larger predators in mixed-conifer forests. *Data in brief*, **6**, 783–792.
- 451 Tambling, C.J., Minnie, L., Meyer, J., Freeman, E.W., Santymire, R.M., Adendorff, J. & Kerley,
452 G.I.H. (2015) Temporal shifts in activity of prey following large predator reintroductions.

Behavioral Ecology and Sociobiology.

Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M.C., Schwager, M. & Jeltsch, F.

(2004) Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *Journal of biogeography*, **31**, 79–92.

Tobler, M.W., Carrillo-Percastegui, S.E., Leite Pitman, R., Mares, R. & Powell, G. (2008) An evaluation of camera traps for inventorying large- and medium-sized terrestrial rainforest mammals. *Animal Conservation*, **11**, 169–178.



Trolle, M. & Kéry, M. (2005) Camera-trap study of ocelot and other secretive mammals in the northern Pantanal. *Mammalia*, **69**, 405–412.

Wang, Y. & Fisher, D.O. (2012) Dingoes affect activity of feral cats, but do not exclude them from the habitat of an endangered macropod. *Wildlife Research*, **39**, 611–620.

Figure 1

Schematized map of the study area in the municipality of Bilar

Figure 1. Schematized map of the study area in the municipality of Bilar, Bohol, Philippines. Circles highlight eight sites surveyed from July to December 2014.



Legend

- + camera traps
- roads
- - - reserve limit
- farmland-plantations
- secondary forest-farms
- primary forest
- farmland-rice paddy
- secondary forest-villages

Figure 2

The species accumulation curves with 95% CIs for species captured in all categories of environment in 2,885 trap days

Figure 2. The species accumulation curves with 95% CIs for species captured in all categories of environment in 2,885 trap days. The dashed-and-dotted line marks the known number of species, while the dotted lines represent species richness estimated by Jackknife 1 in EstimateS (Colwell 2013) .




Figure 3

Overlap between diel activity patterns of cats in transition Zone II and rural Zone III

Figure 3. Overlap between diel activity patterns of cats in transition Zone II (dashed line) and rural Zone III. The number represents the coefficient of overlap (Δ), with standard error in parentheses.




Figure 4

Overlap between the diel activity patterns of cats, rodents, ground-dwelling birds and dogs.

Figure 4. Overlap between the diel activity patterns of cats, rodents, ground-dwelling birds and dogs in transition Zone II (left) and rural Zone III.




Figure 5

Overlap between the diel activity patterns of rodents and ground-dwelling birds at sites with and without the presence of cats

Figure 5. Overlap between the diel activity patterns of rodents (left) and ground-dwelling birds at sites with and without the presence of cats. The numbers represent coefficients of overlap (Δ) with standard errors in parentheses. Arrows indicate shifts in activity levels if cats are present.




Figure 6

Overlap between the diel activity patterns of cats and both species of civets at all sites

Figure 6. Overlap between the diel activity patterns of cats and both species of civets at all sites of their occurrence.




Table 1 (on next page)

Summary of the camera trap deployment

Table 1. Summary of the camera trap deployment in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines.

1 Table 1. Summary of the camera trap deployment in the area of the Rajah Sikatuna Protected
2 Landscape, Bohol, Philippines.

Zone	Site	Date	<i>n</i> camera traps	<i>n</i> trap- days	Range of distances between traps (m)
I	SP	12.7. - 30.7. 2014	12	204	60 – 514
I	SF	30.7. - 14.8. 2014	12	173	38 – 307
I	WS	1.8. - 15.8. 2014	9	125	37 – 265
II	BI	2.8. - 4.12. 2014	10	850	28 – 395
II	LS	12.7. - 4.12. 2014	7	536	48 – 174
II	BU	2.8. - 4.12. 2014	4	383	38 – 44
III	HB	5.7. - 31.7. 2014	10	224	25 – 236
III	SU	2.7. - 31.7. 2014	16	390	23 – 139

3

4

5

Table 2 (on next page)

List of species recorded during a survey in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines.

Table 2. List of species recorded during a survey in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines. Species denoted with ‘*’ are those expected for the species accumulation curve. Values represent number of events of species recorded in each zone.

- 1 Table 2. List of species recorded during a survey in the area of the Rajah Sikatuna Protected
- 2 Landscape, Bohol, Philippines. Species denoted with ‘*’ are those expected for the species
- 3 accumulation curve. Values represent number of events of species recorded in each zone.

Common name	Scientific name	Zone I	Zone II	Zone III	Site
Philippine warty pig*	<i>Sus philippensis</i>	1	0	0	SF
Common palm civet*	<i>Paradoxurus hermaphroditus</i>	10	43	6	Su,Bi,LS,Bu,SP,SF,Ws
Malay civet*	<i>Viverra zangalla</i>	1	8	7	Ha,Bi,LS,SP
Long-tailed macaque*	<i>Macaca fascicularis</i>	3	4	0	LS,Bu,Ws
Philippine tarsier*	<i>Tarsius syrichta</i>	0	1	2	Su,LS
Philippine tree squirrel*	<i>Sundasciurus philippinensis</i>	4	42	1	Su,LS,Bu,SP,SF,Ws
Dog	<i>Canis lupus familiaris</i>	14	39	91	Su,Ha,Bi,LS,Bu,SP
Cat	<i>Felis catus</i>	0	97	83	Su,Ha,Bi,LS
Rat*	<i>Rattus spp.</i>	32	242	47	Su,Ha,Bi,LS,Bu,SP,SF,Ws
Mice*	<i>Mus spp.</i>	5	217	5	Su,Ha,Bi,LS,Bu,SP,SF,Ws
Yellow-headed water monitor*	<i>Varanus cumingi</i>	1	6	5	Su,LS,SF
Hooded pitta*	<i>Pitta sordida</i>	0	0	41	Ha
Red-bellied pitta*	<i>Pitta erythrogaster</i>	0	42	0	Bi,LS
Azure-breasted pitta*	<i>Pitta steerii</i>	1	0	0	Ws
Striated wren-babbler*	<i>Ptilocichla mindanensis</i>	1	9	0	LS,Ws
Red junglefowl*	<i>Gallus gallus</i>	0	5	14	Su,LS,Bu
Barred rail*	<i>Gallirallus torquatus</i>	0	4	179	Su,Ha,Bi,LS,Bu
Slaty-legged crane*	<i>Rallina eurizonoides</i>	0	0	15	Su,Ha
Ruddy-breasted crane*	<i>Zapornia fusca</i>	0	2	1	Su,Bu
Plain bush-hen*	<i>Amaurornis olivacea</i>	0	0	10	Su
Black-faced coucal	<i>Centropus melanops</i>	2	4	2	Su,Bi,LS,Bu,SF,Ws
Philippine coucal	<i>Centropus viridis</i>	0	2	3	Su,Bu
Emerald dove	<i>Chalcophaps indica</i>	7	7	16	Su,Ha,Bi,LS,Bu,SF,Ws
Philippine magpie-robin	<i>Copsychus mindanensis</i>	0	3	1	Ha,Bi
Mindanao bleeding-heart	<i>Gallicolumba crinigera</i>	1	2	0	LS,SP
Hair-crested drongo	<i>Dicrurus hottentottus</i>	1	0	0	SF
Besra	<i>Accipiter virgatus</i>	0	1	1	Su,Bi
Philippine hawk-owl	<i>Ninox philippensis</i>	0	0	1	Ha
Yellow-breasted tailorbird	<i>Orthotomus samarensis</i>	0	1	0	Bi
Domestic chicken	<i>Gallus gallus domesticus</i>	10	254	9	Su,Bi,LS,Bu,SP,SF

Table 3 (on next page)

Number of independent events used for calculation of activity pattern overlap for each analyzed group of animals and each location

Table 3. Number of events used for calculation of activity pattern overlap for each analyzed group of animals and each location in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines. Values in parentheses show the zone-specific relative abundance index (events/total trap days in zone*100). Dashes denote unprocessed entries.

- 1 Table 3. Number of events used for calculation of activity pattern overlap for each analyzed
- 2 group of animals and each location in the area of the Rajah Sikatuna Protected Landscape,
- 3 Bohol, Philippines. Values in parentheses show the zone-specific relative abundance index
- 4 (events/total trap days in zone*100). Dashes denote unprocessed entries.

	Zone I	Zone II	Zone III	All sites	no cats	with cats
cats	0 (0)	67 (4.57)	83 (13.52)	150 (5.81)	-	-
dogs	14 (2.79)	37 (2.53)	90 (14.66)	141 (5.46)	-	-
rodents	41 (8.17)	480 (32.76)	47 (7.65)	568 (22.00)	41 (8.18)	527 (25.35)
ground-dwelling birds	12 (2.39)	242 (16.52)	263 (42.83)	517 (20.03)	12 (2.39)	505 (24.29)
civets	11 (2.19)	16 (1.09)	13 (2.12)	40 (1.39)	-	-

5

Table 4(on next page)

Activity pattern overlaps between cats, their potential prey and competitors in transition Zone II, rural Zone III and among all sites surveyed in the area of Bilar

Table 4. Activity pattern overlaps between cats, their potential prey (rodents and ground-dwelling birds) and competitors (dogs and civets) in transition Zone II, rural Zone III and among all sites surveyed in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines.

- 1 Table 4. Activity pattern overlaps between cats, their potential prey (rodents and ground-
- 2 dwelling birds) and competitors (dogs and civets) in transition Zone II, rural Zone III and among
- 3 all sites surveyed in the area of the Rajah Sikatuna Protected Landscape, Bohol, Philippines.

	site	overlap (Δ)	<i>SE</i>	95% <i>lCI</i>	95% <i>uCI</i>
cats vs. rodents	Zone II	0.48	0.023	0.37	0.58
	Zone III	0.62	0.002	0.52	0.73
cats vs. ground-dwelling birds	Zone II	0.61	0.019	0.50	0.71
	Zone III	0.40	0.041	0.30	0.50
cats vs. dogs	Zone II	0.50	0.052	0.36	0.62
	Zone III	0.45	0.041	0.35	0.56
cats vs. civets	All sites	0.55	0.067	0.45	0.64

4