

On the diversity of the SE Indo-Pacific species of *Terebellides* (Annelida; Trichobranchidae), with the description of a new species (#10561)

1

First submission

Please read the **Important notes** below, and the **Review guidance** on the next page.
When ready [submit online](#). The manuscript starts on page 3.

Important notes

Editor and deadline

Karen Osborn / 20 May 2016

Files

8 Figure file(s)

2 Table file(s)

Please visit the overview page to [download and review](#) the files not included in this review pdf.

Declarations

Describes a new species.



Please in full read before you begin

How to review

When ready [submit your review online](#). The review form is divided into 5 sections. Please consider these when composing your review:


1. BASIC REPORTING

2. EXPERIMENTAL DESIGN

3. VALIDITY OF THE FINDINGS






4. General comments

5. Confidential notes to the editor

 You can also annotate this **pdf** and upload it as part of your review

To finish, enter your editorial recommendation (accept, revise or reject) and submit.





BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standard](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (See [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  Data is robust, statistically sound, & controlled.
-  Conclusion well stated, linked to original research question & limited to supporting results.
-  Speculation is welcome, but should be identified as such.

The above is the editorial criteria summary. To view in full visit <https://peerj.com/about/editorial-criteria/>

On the diversity of the SE Indo-Pacific species of *Terebellides* (Annelida; Trichobranchidae), with the description of a new species

Julio Parapar, Juan Moreira, Daniel Martin

The study of material collected during routine monitoring surveys dealing with oil extraction and aquaculture in waters off Myanmar (North Andaman Sea) and Indonesia (Macasar Strait), respectively, allowed us to analyse the taxonomy and diversity of the polychaete genus *Terebellides* (Annelida). Three species were found, namely *Terebellides* af. **woodlawa**, *Terebellides hutchingsae* **spec. nov.** (a new species fully described and illustrated), and *Terebellides* sp. (likely a new species, but with only one available specimen). The new species is characterised by the combination of some branchial (number, fusion and relative length of lobes and papillation of lamellae), and thoracic (lateral lobes and relative length of notopodia) characters and is compared with all species described or reported in the SW Indo-Pacific area. The taxonomic relevance of the relative length of branchial lobes and different types of ciliature in branchial lamellae for species discrimination in the genus is discussed. A key to all *Terebellides* species described in SE Indo-Pacific waters is presented.

On the diversity of the SE Indo-Pacific species of *Terebellides* (Annelida; Trichobranchidae), with the description of a new species

JULIO PARAPAR¹, JUAN MOREIRA² & DANIEL MARTIN^{3,4}

^{1,4} *Departamento de Biología Animal, Biología Vegetal e Ecología, Universidade da Coruña, 15008 A Coruña, Spain. E-mail: jparapar@udc.es*

² *Departamento de Biología (Zoología), Facultad de Ciencias, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain. E-mail: juan.moreira@uam.es*

³ *Centre d'Estudis Avançats de Blanes, CEAB-CSIC, Blanes, Catalonia, Spain. E-mail: dani@ceab.csic.es*

⁴ *Corresponding author*

Abstract

The study of material collected during routine monitoring surveys dealing with oil extraction and aquaculture in waters off Myanmar (North Andaman Sea) and Indonesia (Macasar Strait), respectively, allowed us to analyse the taxonomy and diversity of the polychaete genus *Terebellides* (Annelida). Three species were found, namely *Terebellides* af. *woodlawa*, *Terebellides hutchingsae* **spec. nov.** (a new species fully described and illustrated), and *Terebellides* sp. (likely a new species, but with only one available specimen). The new species is characterised by the combination of some branchial (number, fusion and relative length of lobes and papillation of lamellae), and thoracic (lateral lobes and relative length of notopodia)

characters and is compared with all species described or reported in the SW Indo-Pacific area. The taxonomic relevance of the relative length of branchial lobes and different types of ciliature in branchial lamellae for species discrimination in the genus is discussed. A key to all *Terebellides* species described in SE Indo-Pacific waters is presented.

Key words

Polychaeta, Myanmar, Indonesia, *Terebellides*, New Species, Branchial morphology, SEM.

Introduction

The genus *Terebellides* is characterised by combination of several characters including the compact appearance of the prostomium, a peristomium forming two lips (upper and lower), a thorax composed by 18 chaetigers, capillary notochaetae, denticulate thoracic neurochaetal hooks and abdominal avicular uncini. Nevertheless, the two most distinctive characters are the single mid-dorsal branchiae composed by 2–5 lamellate lobes, and the geniculate chaetae present in the first 1–2 thoracic neuropodia.

The peculiar shape of the branchiae of the type species (i.e. *T. stroemii* Sars, 1835) led to attribute most subsequent records to this taxon. Therefore, the number of fully described species was relatively low and *T. stroemii* was thought as being cosmopolitan. Prior to the 1980's this species was reported from a wide variety of world areas and depths. In addition to this, the 'Catalogue of World Polychaetes' by Hartman (1959) contributed to this consideration by synonymizing several species with *T. stroemii* (e.g. *T. ypsilon*). However, since Williams (1984), this idea has gradually been changing. Imajima and Williams (1985) and Solís-Weiss *et al.* (1991) further supported to this trend and, thus, a progressively high number of new species have



been ~~(and are being)~~ described (e.g. Hutchings *et al.*, 2015; Parapar & Moreira, 2008; Parapar *et al.*, 2011; 2013; 2016; Schüller & Hutchings, 2010; 2012; 2013). At the same time new characters for the species discrimination have been reported, and those traditionally used (e.g. branchial shape) have increasingly been described in greater detail. As a result, the true diversity of the genus *Terebellides* begins to be revealed.

In the SW Indo-Pacific, ten species of *Terebellides* have been described: four from the Philippine and China Seas (Salazar-Vallejo *et al.*, 2014), namely *T. intoshi* Caullery, 1915, *T. jorgenii* Hutchings, 2007, *T. sieboldi* Kinberg, 1867 and *T. ypsilon* Grube, 1878, and six from the Australian coasts: *T. akares* Hutchings, Nogueira & Carrerette, 2015, *T. jitu* Schüller & Hutchings, 2010, *T. kowinka* Hutchings & Peart, 2000, *T. mundora* Hutchings & Peart, 2000, *T. narribri* Hutchings & Peart, 2000 and *T. woodlawa* Hutchings & Peart, 2000. Additional references to the presence of *T. stroemii* in these waters are found in Caullery (1944), Rullier (1965), Gallardo (1967), Stephenson *et al.* (1970, 1974), Gibbs (1971), Knox & Cameron (1971), Hutchings (1977), Shin (1982), Amoureux (1984), Hutchings & Murray (1984), Hutchings *et al.* (1993) and Tan & Chou (1993). Many reports of *T. stroemii* from Australian and New Zealand waters were summarized by Day & Hutchings (1979), while Hutchings & Peart (2000), by reviewing a high number of references and material of the Australian *Terebellides* (as well as from near the type locality in the SW coast of Norway), described four new species and conclude that *T. stroemii* is not present in southern latitudes. Further papers by Hutchings (2007), Schüller & Hutchings (2010) and Hutchings *et al.* (2015) continued with the reassessment of the diversity of *Terebellides* in Australian-Indonesian coasts.

Our paper addresses the study of the genus in waters off Myanmar and Indonesia, allowing us to describe a new species. We are also reviewing and updating the previous works reporting this

genus in the area, and we present a key to all species recorded in the SE Indo-Pacific. Our study, which is by far not definitive, represents one more contribution for unveiling the hidden diversity of the genus *Terebellides* in world oceans and confirms that the type species is probably absent in the Indo-Pacific area. Furthermore, we provide evidences supporting that the diversity of *Terebellides* is still far to be well known.

Material and Methods

This study is based on 82 specimens of the genus *Terebellides* from 25 samples collected during routine monitoring surveys dealing with oil extraction and aquaculture in waters off Myanmar (North Andaman Sea, 2003) and Indonesia (East of the Borneo Island, North of Macasar Strait, 2004), respectively (Table 1).

The samples were collected by means of a van Veen grab covering about 0.3 m². The grab contents were mixed in a sufficiently large container, and then sieved out on board by pouring the contents through a 1 mm mesh sieve. The retained sediment was then transferred into a plastic bag, fixed with a 10% formaldehyde/seawater solution, stained with “Rose of Bengal” and stored until sorted. An initial sorting was performed under a dissecting stereomicroscope (Zeiss Stemi 2000-C) and the specimens of *Terebellides* were counted and preserved in 70% ethanol.

In Myanmar, a one-liter volume of sediment from one grab was used for physico-chemical analyses (viz. granulometry, organic carbon content). The sediment was taken at each station and transferred into a wide-mouthed double-closing 500 ml polyethylene flasks, which were stored in the dark until transferred to the laboratory. Laser granulometry (% volume) was performed on dry sediment after sifting through a 0.8 mm mesh sieve using a Malvern Mastersizer S laser

granulometer. Sediments were characterized by the percentage of silt and clay (diameter < 63 µm). Estimates of organic carbon have been made according to the European experimental standard NF ISO 14235 (oxidation method, 0.1 % m/m).

Light microscope images were obtained by means of a Olympus SZX12 stereomicroscope equipped with a Olympus C-5050 digital camera. Line drawings were made by means of an Olympus BX40 stereomicroscope equipped with camera lucida. Specimens used for examination with Scanning Electron Microscope (SEM) were prepared by critical point drying, covered with gold and examined and photographed under a JEOL JSM-6400 electron microscope at the Servizos de Apoio á Investigación-SAI (Universidade da Coruña-UDC, Spain).

Most of the obtained material was deposited in the Museo Nacional de Ciencias Naturales (Madrid, Spain; MNCN). Additional paratypes of *T. hutchingsae* **spec. nov.** were deposited in the collections of the Australian Museum (Sydney, Australia; AM) and Göteborgs Naturhistoriska Museum (Göteborg, Sweden; GNM). Type material of *Terebellides gracilis* Malm, 1874 was loaned for study by the Göteborgs Naturhistoriska Museum (Holotype, GNM Polych 641). Type material of *Terebellides sieboldi* Kinberg, 1866 was requested to the Swedish Museum of Natural History for comparison but only one specimen, and badly preserved, could be located (L. Gustavsson, *in litt.*).

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed

through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: 39745D2F-9163-48B2-9FAB-FBF66D3AEFB5. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

Abbreviations used in text and figures: BL—branchial lobes; BT—buccal tentacles; CP = ciliated papillae; CHG = chaetiger with geniculate chaetae; dl—dorsal lobes; gc—geniculate chaeta; go—genital opening; GP = genital papillae; LL—lateral lappets; NACH = number of abdominal chaetigers; npa—nephridial papillae; NRTU = number of rows of frontal rostral teeth in thoracic uncini; PPP = posterior pointed projection; r—rostrum; TC—thoracic chaetiger; TN—thoracic notopodia; tp—terminal projection; TU—thoracic uncini.

Results

Systematics

Family Trichobranchidae Malmgren, 1866

Genus *Terebellides* Sars, 1835, emended by Schüller & Hutchings, 2013

Type species

Terebellides stroemii Sars, 1835, redescribed by Parapar & Hutchings, 2015

Terebellides hutchingsae **spec. nov.**

LSID: 78E96984-41E7-43E6-8E5D-03E9421BE306

(Figs 1–8, Tables 2–3)

139

140 *Material examined*



141 **INDONESIA** (Macasar Strait): **Holotype:** MNCN 16.01/0000 (St. 6). **Paratypes:** MNCN
 142 16.01/0000 (St. 2, 4 specs); MNCN 16.01/0000 (St. 3, 3 specs); MNCN 16.01/0000 (St. 5, 2+1
 143 specs); MNCN 16.01/0000 (St. 5, 1 spec. on SEM stub); MNCN 16.01/0000 (St. 6, 5 specs);
 144 MNCN 16.01/0000 (St. 7, 5 specs); MNCN 16.01/0000 (St. 8, 7 specs); MNCN 16.01/0000 (St.
 145 8, 1 spec. on SEM stub); MNCN 16.01/0000 (St. 15, 2 specs); MNCN 16.01/0000 (St. 16, 6
 146 specs); MNCN 16.01/0000 (St. 23, 1 spec. on SEM stub). **MYANMAR** (North Andaman Sea):
 147 **Paratypes:** MNCN 16.01/0000 (St. E7(2), 1 spec.); MNCN 16.01/0000 (St. E8(3), 1 spec.);
 148 MNCN 16.01/0000 (St. E11B(2), 4 specs); MNCN 16.01/0000 (St. E11B(3), 2 specs); MNCN
 149 16.01/0000 (St. E14(2), 4 specs); MNCN 16.01/0000 (St. E15(2), 10 spec.); MNCN 16.01/0000
 150 (St. E16(1), 2 specs); MNCN 16.01/0000 (St. E16(3), 1 spec.); MNCN 16.01/0000 (St. 17(3), 1
 151 spec.); MNCN 16.01/0000 (St. S2(2), 1 spec.); MNCN 16.01/0000 (St. S3(2), 4 specs); MNCN
 152 16.01/0000 (St. S3(2), 1 spec. on SEM stub); MNCN 16.01/0000 (St. S3(3), 4 specs); MNCN
 153 16.01/0000 (St. S4(2), 2 specs); MNCN 16.01/0000 (St. S4(3), 1 spec.); MNCN 16.01/0000 (St.
 154 WP2(2), 2 specs); MNCN 16.01/0000 (St. WP2(3), 2 specs); MNCN 16.01/0000 (St. WP2(3), 2
 155 specs on SEM stub); MNCN 16.01/0000 (St. WP3(3), 1 spec.).

156

157 *Description* (based on holotype and paratypes)

158 Complete individuals ranging from 9.0 to 14.0 mm in length (14 mm in holotype; Fig. 2A–B)
 159 and 0.7 to 1.5 mm in maximum width at thoracic region (1.3 mm in holotype, excluding
 160 parapodia). Body tapering posteriorly with segments increasingly shorter and crowded towards
 161 pygidium. Prostomium compact; peristomium forming a tentacular membrane with large upper

and lower lips surrounding mouth, sometimes almost devoid of buccal tentacles (Fig. 3A).

Buccal tentacles of two types, short ventral tentacles uniformly cylindrical or slightly expanded at tips, and long dorsal tentacles more expanded at tips (Figs 2B, 4A–B). Lateral lappets on TC1–5 (SGIII–VII), being larger in TC1–3 (Figs 2B, 3A, 4C, 6A). No conspicuous dorsal rounded projection on anterior chaetigers or oval-shaped glandular region in TC3. Both notopodia and notochaetae in TC1 less developed than in following chaetigers (Figs 3A, 4C).

Branchiae arising as single structure from SGII–III, with a single, mid-dorsal, stalk and two pairs of unfused lobes; lower (=ventral) (BL3–4) pair smaller and much shorter than upper (=dorsal) (BL1–2) pair of lobes (Figs 3A–B, 6B–C). Upper and lower lobes with a short terminal pointed projection (although deciduous and sometimes damaged) (Fig. 3C). Dorsal pair of branchial lobes with short anterior projection (fifth lobe; BL5) (Fig. 3D), sometimes hidden behind buccal tentacles (Fig. 2A–B). Loss of any of branchial lobes not observed. One side of branchial lamellae with parallel bent rows of cilia and well-developed ciliated papillae on edge of one side of each branchial lamella (Fig. 3D–F).

Eighteen thoracic chaetigers (SGIII–XX), all with notopodia; neuropodia from SGVIII. Notopodia of TC1 smaller than following ones (Fig. 4C, E); all remaining notopodia similar in size. Thoracic neuropodia as sessile pinnules, from TC6 (SGVIII) to TC18, with uncini in single rows from TC7 (SGIX) throughout. Thoracic notochaetae similar in length, with textured surface (Fig. 4F). Ciliated papilla dorsal to each thoracic notopodia not observed. First thoracic neuropodia (TC6) with 4–7 geniculate acicular chaetae with minute teeth in their upper part forming a *capitium* easily overlooked without SEM (Fig. 6E–F); sharply bend. Subsequent thoracic neuropodia with one row of about 8–10 uncini per torus (Fig. 5A); uncini as shafted denticulate hooks with long, pointed *rostrum* surmounted by 4–5 teeth and an upper crest of

several smaller denticles of different sizes (Fig. 5A–C). One finger-shaped nephridial papilla basal to branchial stem (Fig. 4E); genital openings, dorsal to notopodia in TC4 and TC5 (Figs 4D, 6D).

Twenty seven to 30 abdominal chaetigers (30 in holotype). Abdominal neuropodia as erect pinnules, with about 30 uncini per torus (Fig. 5D). Uncini with 3–4 teeth above main fang (Fig. 5D–E), surmounted by a row of an irregular number of shorter teeth and an upper crest of minute teeth. Pygidium blunt, funnel-like depression. No eggs were observed in body cavity of holotype, but mature females of smaller size were observed (9.0 mm length, 1.0 mm width). Colour in alcohol pale brown.

Type locality

Macasar Strait (Indonesia), muddy bottom with shell fragments at 72 m depth.

Distribution and habitat

Specimens of *T. hutchingsae* **spec. nov.** were found in shallow water bottoms (45.5–51.0 m depth) about 80 Km off the coast of Myanmar (North Andaman Sea) and in slight deeper bottoms (58.0–84.0 m depth) about 16 Km off the mouth of the Mahakam delta in the East coast of the Borneo Island (Indonesia) (North Makassar basin) (Table 1, Fig. 7).

Etymology

The species is named after Dr. Pat Hutchings, for her many contributions to the taxonomy of Terebelliform polychaetes in Australia and SW Pacific waters, and particularly to the genus *Terebellides*, and also for her key role in the study of Australian polychaetes.

208

209 *Remarks*

210 Several species of *Terebellides* were previously described in the Myanmar-Indonesia-
 211 Philippines-North Australia area (Fig 7): *T. intoshi* Caullery, 1915, *T. sieboldi* Kinberg, 1867, *T.*
 212 *ypsilon* Grube, 1878, *T. jorgeni* Hutchings, 2007 and *T. jitu* Schüller & Hutchings, 2010.
 213 *Terebellides intoshi* is characterised by the large size of the notopodia and notochaetae from TC6
 214 onwards (Fig. 8A) and probably by the presence of two chaetigers with geniculate chaetae as
 215 well (see Remarks of *Terebellides* sp.); *T. sieboldi* has geniculate chaetae in TC7 instead of TC6
 216 and *T. ypsilon* is considered undeterminable by Hutchings & Peart (2000) because type material
 217 no longer exists. The two most recently described species, *Terebellides jorgeni* and *T. jitu*, are
 218 the most similar to *T. hutchingsae* **spec. nov.** *Terebellides jorgeni* differs from the new species
 219 in: 1) the presence of glandular and whitish ventral part of anterior segments, SG5 to SG9 (CH3
 220 to CH7) but specially on SG5 to SG7 (absent in *T. hutchingsae* **sp. nov.**), and bearing
 221 pronounced thickening and elevation of dorsal anterior margins forming dorsal crests; 2) genital
 222 pores are present in SG4 and SG5, instead of SG6 and SG7 (TC4 and TC5) as in *T. hutchingsae*
 223 **spec. nov.**; 3) the branchiae are formed by four lobes instead of five. On the other hand, the
 224 overall shape of branchiae is quite similar in both species, being lobes 1–4 unequal sized and
 225 entirely free (not fused), with upper (dorsal) ones larger than lower (ventral) ones, and with
 226 “surface of branchial lamellae weakly papillate” (cfr. p. 78 in Hutchings, 2007); the latter
 227 probably refers to the presence of ciliated papillae, which is a feature difficult to confirm in the
 228 original figures.

Terebellides jitu is also similar to *T. hutchingsae* **spec. nov.** but all branchial lobes are of similar length and fused half of their length instead of the lower ones being much shorter and fused basally as in *T. hutchingsae* **spec. nov.**

Terebellides narribri Hutchings & Peart, 2000 and *T. woolawa* Hutchings & Peart, 2000 were described from the NE Australian coast. Both species share with *T. hutchingsae* **spec. nov.** branchiae with similar shape and composed by five lobes; *Terebellides narribri* differs from the new species by having first thoracic notopodia (TN1) of same size as the following, and TC3 bearing large, white, oval pair of glandular patches. *Terebellides woodlawa* is characterised by the great development of BL5 (see Remarks on *T. af. woodlawa*) and by having anterior thoracic segments with dorsal projections on lateral lappets, which are absent in *T. hutchingsae* **spec. nov.**

The North Atlantic species and type species of the genus *Terebellides*, i. e. *T. stroemii* Sars, 1835, was also widely reported in the area (e. g. Indonesia: Caullery (1944); South Korea: Gallardo (1967); Hong Kong: Shin (1982); Singapore: Tan & Chou (1993); Australian coast: Stephenson *et al.* (1970; 1974), Knox & Cameron (1971), Hutchings (1977), Amoureux (1984), Hutchings & Murray (1984), Hutchings *et al.* (1993); Fig. 7). This species was recently redescribed by Parapar & Hutchings (2015) from Norwegian specimens collected by Michael Sars near the type locality. In the Southern Pacific Ocean, its presence had already been denied by Hutchings & Peart (2000) after examining Norwegian material; indeed, part of this material was already reassigned to other species (see Hutchings & Peart, 2000) while others specimens were not. Among the latter, the material reported by Caullery (1944) and collected during the Siboga expedition might well correspond to more than one species according to the description and illustrations. The shape of the branchiae in specimen from station 271 (fig. 147 in Caullery, 1944; redrawn here in Fig. 8B) and station 311 (fig. 148 in Caullery, 1944; redrawn here in Fig.

8C) sharply differs in BL5 size; the specimen of station 311 is more similar in branchial shape to *T. hutchingsae* **spec. nov.** but differs in the high degree of fusion of dorsal and ventral lobes in Caullery's material (see Fig. 8C). The specimen reported by Gallardo (1967) cannot be properly identified because the description is quite brief (e.g. "The branchia has the typical shape...") and only a lateral view of a thoracic uncinus is illustrated and this is not relevant in species discrimination.

One of the most relevant diagnostic characters of *T. hutchingsae* **spec. nov.** is the presence of ciliated papillae in branchial lamellae. This character was long ignored in *Terebellides* descriptions and was discussed by Parapar *et al.* (2016). In fact, several recently described species from across the world oceans show this feature, namely *T. gracilis* Malm, 1874 *sensu* Parapar *et al.* (2011), off Iceland; *T. jorgeni* Hutchings, 2007, from Indonesia; *T. gracilis* Malm, 1874 *sensu* Parapar *et al.* (2013) and *T. mediterranea* Parapar *et al.*, 2013, from the Adriatic Sea; *T. akares* Hutchings *et al.*, 2015, from the Great Barrier Reef (NE Australia); a new species described by Parapar *et al.* (in press), from the Persian Gulf; and *T. af. woodlawa* Hutchings & Peart, 2000 *sensu* Parapar *et al.* (this work) from South Myanmar. This character is probably much more widespread ~~that was thought previously~~, and shows at least two different morphotypes: 1) low papillae as it was found in *T. gracilis* from Iceland and the Mediterranean, and 2) well developed papillae in the rest of species. The presence of these low ciliated papillae (Parapar *et al.*, 2011; 2013) in Icelandic and Adriatic specimens of *T. gracilis* could not be confirmed yet in the holotype ~~of~~ (see M&M above).

Terebellides af. *woodlawa* Hutchings & Peart, 2000
(Figs 2C–D)

275

276 *Material examined*

277 Two specimens. MNCN 16.01/0000 (St. S4(3), 1 spec.); MNCN 16.01/0000 (St. WP3(3), 1
278 spec.).

279

280 *Distribution and habitat*

281 Both specimens of *T. af. woodlawa* were found in two ~~near~~ shallow water stations (51.0 m depth)
282 about 80 Km off the mouth of the Irawadi river in the coast of Myanmar (North Andaman Sea)
283 (Table 1).

284

285 *Remarks*

286 *Terebellides woolawa* is characterised by the well-developed fifth branchial lobe (BL5) and the
287 presence of dorsal rounded projections on lateral lappets of SG 3–6 (TC1–4). This large species
288 was described from intertidal to shallow water habitats in eastern Australia (Fig. 7) and was
289 found across most of Australian coasts (Hutchings & Peart, 2000). Specimens found in this study
290 are large-sized, and agree fairly well with the original description; in particular, specimen
291 MNCN 16.01/0000 shows the typical shape of the branchiae, which have five lobes, BL1–4 are
292 fused up to half of their length, filamentous tips are short, and BL5 is well developed (Fig.
293 2C–D). Nevertheless, our specimens lack the characteristic dorsal lobes of anterior thoracic
294 lateral lappets: this prevented to fully confirm the identity of our material.

295

296 *Terebellides* sp.

297 (Fig 2E–F, 7, 9)

298

299 *Material examined*

300 One specimen. MNCN 16.01/0000 (St. S4(3), 1 spec.).

301

302 *Distribution and habitat*

303 The specimen was found in shallow water bottom (51.0 m depth) about 16 Km off the coast of
304 Myanmar (North Andaman Sea) (Table 1).

305

306 *Remarks*

307 The specimen differs from *T. hutchingsae* **spec. nov.** and *Terebellides* af. *woolawa* in two
308 features: 1) BL5 is large-sized, about half the length of posterior lobes (BL1–4); and 2) TC5 and
309 TC6 are both provided with acicular geniculate chaetae. Thus, BL5 is longer than in any other
310 described species including *T. woodlawa*; however, this might be due to the preservation state of
311 the specimen, which is slightly deteriorated. Anyway, the combination of the two
312 aforementioned characters may justify the erection of a new species but we prefer to wait for
313 eventual finding of additional specimens to confirm its status.

314 Four species of the genus *Terebellides* were previously described as having geniculate
315 chaetae in two thoracic chaetigers: *T. akares* Hutchings, Nogueira & Carrerette, 2015 (North-
316 East Australia), *T. biaciculata* Hartmann-Schröder, 1992 (French Polynesia), *T. bigeniculatus*
317 Parapar, Moreira & Helgason, 2011 (Iceland) and *T. intoshi* Caullery, 1945 *sensu* Imajima &
318 Williams (1985) (Japan).

319 We follow Parapar *et al.* (2011) in considering that type material of *T. intoshi* from South China
320 Sea (see Figure 8) probably does not have two chaetigers with geniculate chaetae and thus

Japanese material would belong to a different species. Anyway, the latter also differs from *Terebellides* sp. in the branchial shape and the greater development of thoracic notopodia from TC6 (Fig. 8A). In *Terebellides akares*, the branchiae bears a much shorter BL5 and posterior ventral lobes (BL3–4) are completely free from each other; in *Terebellides* sp., these lobes are fused in most of their length (Fig. 2F).

Key of SE Indo-Pacific species of Terebellides

The key here presented has been modified from the previous key of Australian Trichobranchidae (Hutchings & Peart 2000), which was based on a limited number of easy-to-detect characters: 1) number of chaetigers with geniculate chaetae, 2) degree of development of thoracic notopodia, and 3) shape of branchiae, giving special emphasis to the relative size of branchial lobes. *Terebellides ypsilon* Grube, 1878, from the Philippines, was not included because the description is very brief and following Hutchings & Peart (2000), who revised the type material, the taxon should be considered as undeterminable.



1. GC in two TC
 - 2
- GC in one TC
 - 4
2. All TN of similar length
 - 3

342 - TN from TC6 onwards much bigger ~~in size~~ and with more numerous and longer notochaetae
 343 *T. intoshi*
 344 Caullery, 1944
 345 3. TU with GC similar in shape and position *T. akares* Hutchings *et al.*,
 346 2015
 347 - TU with GC different in shape and position *Terebellides*
 348 **sp.**
 349 4. GC in TC7¹ *T. sieboldi* Kinberg,
 350 1867
 351 - GC in TC6
 352 5
 353 5. Branchial lobes 1-4 loosely fused *T. mundora* Hutchings & Peart,
 354 2000
 355 - Branchial lobes 1-4 more or less fused
 356 6
 357 6. Four branchial lobes
 358 7
 359 - Five branchial lobes
 360 8
 361 7. All TN similar in size and well developed *T. kowinka* Hutchings & Peart,
 362 2000
 363 - TN1 and TN2 much smaller than subsequent ones *T. jorgen* Hutchings,
 364 2007

365 8. BL5 about 1/5 length of posterior lobes; thoracic LL without dorsal projections, GC of TC6
 366 sharply bent 9
 367 - BL5 almost 1/2 length of posterior lobes; LL of TC1-4 with dorsal projections, GC of TC6
 368 gently curved *T. woolawa* Hutchings &
 369 Peart, 2000
 370 9. TN1 not reduced; large, white, oval glandular patches in TC3
 371
 372 *T. narribri* Hutchings & Peart,
 373 2000
 374 - TN1 strongly reduced; no glandular patches in TC3
 375 10
 376 10. All branchial lobes of similar length and fused half of their length; BL with transverse ridges
 377 of ciliature *T. jitu* Schüller & Hutchings,
 378 2010
 379 - Ventral (posterior) branchial lobes much shorter than dorsal (anterior) ones and fused basally;
 380 BL with ciliated papillae on border *T. hutchingsae*
 381 **spec. nov.**
 382
 383 ⁽¹⁾ The position of GC in TC7 is very rare in the genus *Terebellides*; this feature is apparently
 384 only shared with *T. pacifica* Kinberg, 1866, a species which has been removed from synonymy
 385 with *T. stroemii* by Garraffoni *et al.* (2005).
 386
 387

Acknowledgements

We are indebted to the staff members of Créocéan, who recovered the material during the processing of benthic macrofaunal samples and explicitly accepted to allow publishing the information contained in this study, and to Andrea Feijoo (UDC) for the initial study of the material. Special thanks are also due to Ada Castro and Cati Sueiro (SAI, UDC) for assisting with the preparation of the specimens and use of the SEM, and to Xela Cunha (Marine Station of A Graña-Ferrol, University of Santiago de Compostela-USC, Spain) for her assistance with the stereomicroscope images. Authors also wish to thank Kennet Lundin (GNM) for sending *T. gracilis* holotype, and Lena Gustavsson (NRM) for her help in trying to locate *T. sieboldi* holotype.

References



- Amoureux, L. (1984) Brief notes on two small collections of Polychaetes from Queensland, with two new records from Australia. *Proceedings of Royal Society of Queensland*, 95, 7–9.
- Bremec, C.S. & Elías, R. (1999) Species of *Terebellides* from South Atlantic Waters off Argentina and Brazil (Polychaeta: Trichobranchidae). *Ophelia*, 5, 177–186.
- Day, J.H. & Hutchings, P.A. (1979) An annotated check-list of Australian and New Zealand Polychaeta, Archiannelida and Myzostomida. *Records of the Australian Museum*, 32(3), 80–161.
- Caullery, M. (1944) Polychètes Sédentaires de l'Expédition du Siboga: Ariciidae, Spionidae, Chaetopteridae, Chlorhaemidae, Opheliidae, Oweniidae, Sabellariidae, Sternaspidae, Amphictenidae, Ampharetidae, Terebellidae. *Siboga Expeditie*, 24(2 bis), 1–204.
- Gallardo, V.A. (1967) Polychaeta from the Bay of Nha Trang, South Viet Nam. *Naga Reports*, 4(3), 35–279.
- Gibbs, P.E. (1971) The polychaete fauna of the Solomon Islands. *Bulletin of the British Museum (Natural History)*, Ser. Zoology, 21, 101–211.

- 414 Garraffoni, A.R.S. & Lana, P.C. (2003) Species of *Terebellides* (Polychaeta, Terebellidae,
415 Trichobranchinae) from the Brazilian coast. *Iheringia, Série Zoológica*, 93, 355–363.
- 416 Hartman, O. (1959) Catalogue of the Polychaetous Annelids of the World. Parts I & II. *Allan*
417 *Hancock Foundation, Occasional Papers*, 23, 1–628.
- 418 Hartman, O. (1974) Polychaetous annelids of the Indian Ocean including an account of species
419 collected by members of the International Indian Ocean Expeditions, 1963–‘64, and a
420 catalogue and bibliography of the species from India. Part II. *Journal of the Marine*
421 *Biological Association of India*, 16 (2), 609–644.
- 422 Hutchings, P.A. (1977) Terebelliform Polychaeta of the families Ampharetidae, Terebellidae and
423 Trichobranchidae from Australia, chiefly from Moreton Bay, Queensland. *Records of the*
424 *Australian Museum*, 31, 1–38.
- 425 Hutchings, P.A. & Murray, A. (1984) Taxonomy of polychaetes from the Hawkesbury River and
426 the southern estuaries of New South Wales, Australia. *Records of the Australian Museum*,
427 36, 1–119.
- 428 Hutchings, P.A., Ward, T.J., Waterhouse, J.H. & Walker, L. (1993) Infauna of marine sediments
429 and seagrass beds of Upper Spencer Gulf near Port Pirie, South Australia. *Transactions of*
430 *the Royal Society of South Australia*, 117, 1–15.
- 431 Hutchings, P. & Peart, R. (2000) A revision of the Australian Trichobranchidae (Polychaeta).
432 *Invertebrate Taxonomy*, 14, 225–272.
- 433 Hutchings, P., Nogueira, J.M.M. & Carrerette, O. (2015) Telothelepodidae, Thelepodidae and
434 Trichobranchidae (Annelida, Terebelliformia) from Lizard Island, Great Barrier Reef,
435 Australia. *Zootaxa*, 4019 (1), 240–274.
- 436 Knox, G. A. & Cameron, D. B. (1971) Port Phillip survey 1057–1963, Victoria, Australia. Part
437 2, No. 4. Polychaeta. *Memoirs of the National Museum, Melbourne*, 32, 21–42.

- 438 Parapar, J. & Hutchings, P. (2015) Redescription of *Terebellides stroemii* (Polychaeta,
439 Trichobranchidae) and designation of a neotype. *Journal of the Marine Biological*
440 *Association of the United Kingdom*, 95, 323–337.
- 441 Parapar, J., Moreira, J. & O'Reilly, M. 2016. A new species of *Terebellides* (Polychaeta:
442 Trichobranchidae) from Scottish waters with an insight into branchial morphology. *Marine*
443 *Biodiversity*, 46 (1), 211–225.
- 444 Parapar, J. & Moreira, J. (2008) Revision of three species of *Terebellides* (Polychaeta:
445 Trichobranchidae) described by C. Hessle in 1917 from the Southern Ocean. *Journal of*
446 *Natural History*, 42 (17–20), 1261–1275.
- 447 Parapar, J., Moreira, J. & Helgason, G.V. (2011) Taxonomy and distribution of *Terebellides*
448 (Polychaeta, Trichobranchidae) in Icelandic waters, with the description of a new species.
449 *Zootaxa*, 2983, 1–20.
- 450 Parapar, J., Mikac, B. & Fiege, D. (2013) Diversity of the genus *Terebellides* (Polychaeta:
451 Trichobranchidae) in the Adriatic Sea with the description of a new species. *Zootaxa*, 3691,
452 333–350.
- 453 Rullier, F. (1965) Contribution à la faune des annélides polychètes de l'Australie. *University of*
454 *Queensland Papers, Department of Zoology*, 2, 163–201.
- 455 Salazar-Vallejo, S.I., Carrera-Parra, L.F., Muir, A.I., De Leon González, J.A., Piotrowski, C. &
456 Sato, M. 2014. Polychaete species (Annelida) described from the Philippine and China Seas.
457 *Zootaxa*, 3842, 1–68.
- 458 Schüller, M. & Hutchings, P.A. (2010) New insights in the taxonomy of Trichobranchidae
459 (Polychaeta) with the description of a new *Terebellides* from Australia. *Zootaxa*, 2395, 1–
460 16.

- 461 Schüller, M. & Hutchings, P.A. (2012) New species of *Terebellides* (Polychaeta:
462 Trichobranchidae) indicate long-distance dispersal between western South Atlantic deep-sea
463 basins. *Zootaxa*, 3254, 1–31.
- 464 Schüller, M. & Hutchings, P.A. (2013) New species of *Terebellides* (Polychaeta:
465 Trichobranchidae) from deep Southern Ocean. *Zootaxa*, 3619, 1–45.
- 466 Shin, K. S. P. (1982) Some polychaetous annelids from Hong Kong waters. In: Morten, B.S.
467 Morten and C.K. Tseng (eds.). *Proceedings of the First International Marine Biological*
468 *Workshop: The Marine Flora and Fauna of Hong Kong and Southern China*, vol. 1, 18
469 April-10 May 1980: 161–172.
- 470 Solís-Weiss, V., Fauchald, K. & Blanckstein, A. (1991) Trichobranchidae (Polychaeta) from
471 shallow warm water areas in the Western Atlantic Ocean. *Proceedings of the Biological*
472 *Society of Washington*, 104 (1), 147–158.
- 473 Stephenson, W., Williams, W.T. & Lance, G.N. (1970) The macrobentos of Moreton Bay.
474 *Ecological Monographs*, 40, 459–494.
- 475 Stephenson, W., Williams, W.T. & Cook, S.D. (1974) The benthic fauna of soft bottoms,
476 Southern Moreton Bay. *Memoirs of the Queensland Museum*, 17 (1), 73–123.
- 477 Tan, L.T. & Chou, L.M. (1993) Checklist of polychaete species from Singapore waters
478 (Annelida). *Raffles Bulletin of Zoology*, 41 (2), 279–295.
- 479 Williams, S.J. (1984) The status of *Terebellides stroemi* (Polychaeta; Trichobranchidae) as a
480 cosmopolitan species, based on a worldwide morphological survey, including description
481 of new species. In: Hutchings, P.A. (Ed.), *Proceedings of the First International*
482 *Polychaete Conference*, Sydney, Australia, 1984. The Linnean Society of New South
483 Wales, Sydney, pp. 118–142.

Table 1(on next page)

Main abiotic characteristics: Silt and Organic Carbon contents


Main abiotic characteristics of the samples where *Terebellides* specimens were collected.

Org. car., organic carbon content (%).

	<i>Station</i>	<i>Date</i>	<i>Longitude N</i>	<i>Latitude E</i>	<i>Depth (m)</i>	$\leq 63 \mu m\%$	<i>Org. car.</i>
Myanmar (N Andaman Sea)	E7	03/12/03	15°07'59.8"	94°46'46.5"	46.0	76.3	1.26
	E8	"	15°06'27.7"	94°46'50.2"	47.0	78.7	1.29
	E11	"	15°07'23.2"	94°46'45.9"	47.0	83.6	1.21
	E14	"	15°07'20.0"	94°46'51.1"	47.4	76.7	1.12
	E15	"	15°07'15.9"	94°46'51.1"	47.0	80.4	1.16
	E16	"	15°07'07.8"	94°46'51.0"	47.6	73.8	1.15
	E17	"	15°07'14.9"	94°45'25.9"	48.0	74.0	1.28
	S2	"	15°02'03.3"	94°45'45.7"	51.0	72.8	0.86
	S3	"	15°02'19.4"	94°46'02.6"	51.0	80.5	0.98
	S4	"	15°03'08.2"	94°46'03.0"	51.0	90.2	0.92
	WP2	"	15°09'06.6"	94°45'26.7"	45.5	86.2	1.19
	WP3	"	15°02'03.0"	94°46'19.2"	51.0	69.6	2.42
	<i>Station</i>	<i>Date</i>	<i>Longitude S</i>	<i>Latitude E</i>	<i>Depth (m)</i>	<i>sediment</i>	
Indonesia (Macasar Strait)	2	23/07/04	0°44'35"	117°45'30"	84	Mud + shell fragments	
	3	"	0°43'20"	117°48'52"	67	Coarse + shell fragments	
	5	26/03/04	0°46'00"	117°47'32"	72	Mud + shell fragments	
	6	"	0°47'02"	117°45'45"	73	"	
	7	25/03/04	0°49'45"	117°44'10"	70	Homogeneous clear mud	
	8	26/03/04	0°48'12"	117°48'15"	78	Mud + shell fragments	
	15	24/03/04	0°55'55"	117°40'52"	58	Clear mud + shell fragments	
	16	"	0°55'41"	117°43'11"	70	"	
	23	29/03/04	0°36'24"	117°35'12"	3	Fine black mud	

Table 2 (on next page)

Comparison of body characters in *Terebellides*

Comparison of several body characters of the species of *Terebellides* described with ciliated papillae in branchial lamellae. 

	Source	GP	LL	Branchiae				CH1	CHG	NRTU	NACH	Distribution
				Lobes	Relative length	CP + presence	PPP					
<i>T. gracilis</i> Malm, 1874	Parapar <i>et al.</i> (2011) ¹	TC4–5	TC1–5 ²	4 ³	Same	Low + both sides	no	Shorter	6	2 ⁴	44	Swedish coast of Skagerrak
<i>T. jorgeni</i> Hutchings, 2007	Hutchings (2007)	TC 2–3 ⁵	SG1–5 and SG7–8	4	Ventral shorter	High ⁶	No ⁷	Shorter ⁸	6	4–6	~48	Bali ⁹
<i>T. akares</i> Hutchings <i>et al.</i> , 2015	Hutchings <i>et al.</i> (2015)	TC4–5	TC1–8	5	Ventral shorter	High + both sides	No ⁷	Shorter	5–6	4 ⁷	17–25	North-east Australia
<i>Terebellides shetlandica</i> Parapar <i>et al.</i> 2016	Parapar <i>et al.</i> (2016)	TC4–5	TC1–6	5	Ventral shorter	High + one side	Short	Shorter	6	4	27–36	Iran
<i>T. hutchingsae</i> spec. nov.	This work	TC4–5	TC1–5	5	Ventral shorter	High + one side	Short	Shorter	6	4	27–30	Indonesia and Myanmar


(1) From a redescription of Malm’s holotype; (2) very low lappets; (3) the fifth lobe is very low and therefore branchiae could be described as with only four lobes; (4) character observed from the study of several specimens under the SEM; it cannot be properly observed in holotype; (5) author mentions SG 3–5; this can be interpreted as NP in SG3 (TC1) and GP in TC2–3; author probably meant TC3–5; (6) author simply mentions “surface of (branchial) lamellae weakly papillate”; (7) not explicitly mentioned in the original description of the species; taken from figures; (8) CH2 also smaller than subsequent ones; (9) this species shows a disjunct distribution, being originally described from Bali but also reported from W and S Africa and Tasman Sea.

Figure 1(on next page)


Line drawings of the species of *Terebellides*

Terebellides hutchingsae **spec. nov.**, Holotype from Indonesia, MNHN 16.01/0000. A, left lateral view; *Terebellides* sp., MNCN 16.01/0000. B, detail of geniculate chaetae in CH5 and CH6 of right side of body.

A




B



2

Stereomicroscope images of *Terebellides* spp.

A-B. *Terebellides hutchingsae* **spec. nov.** Stereomicroscope images. Holotype MNCN 16.01/0000. A, general view; B, anterior end, left lateral view. C-D. *Terebellides* af. *woodlawa* MNCN 16.01/0000. C, anterior end right lateral view; D, detail of branchiae, left lateral view. E-F. *Terebellides* sp. MNCN 16.01/0000. E, anterior end right ventro-lateral view, arrowheads pointing to thoracic chaetigers CH5 and CH6; F, detail of branchiae in ventral view.




3 

SEM micrographs of paratypes of *Terebellides hutchingsae* spec. nov. from Indonesia

Terebellides hutchingsae **spec. nov.**, SEM micrographs of paratypes from Indonesia, MNCN 16.01/0000 and MNCN 16.01/0000. A, anterior end, left lateral view, showing lateral lappets and relative size of branchial lobes; B, detail of branchial lobes showing degree of fusion between dorsal (dl) and ventral (vl) lobes; C, detail of distal end of left dorsal lobe showing ciliary fields and terminal projection (tp); D and E, Detail of ciliated papillae of branchial lobes lamellae.

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*




4

SEM micrographs of paratypes from Indonesia, ~~Cont.~~

Terebellides hutchingsae **spec. nov.**, SEM micrographs of paratypes from Indonesia, MNCN 16.01/0000 and MNCN 16.01/0000. A, several buccal tentacles, one showing ciliated side of distal end (arrowhead); B. detail of ciliated distal end of a buccal tentacle; C, anterior end, left lateral view, showing lateral lappets (ll) from TC1 to TC6 and nephridial papilla (npa) in TC1; D, detail of genital papillae (gp) of TC4 and TC5; ~~D, TC1 and TC2~~; E, TC1 and TC2 thoracic notopodia showing reduction of TC1; F. detail of pilose surface of thoracic notochaetae.

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*




5

SEM micrographs of paratypes from Indonesia, ~~Cont. 2~~



Terebellides hutchingsae **spec. nov.**, SEM micrographs of paratypes from Indonesia, MNCN 16.01/0000 and MNCN 16.01/0000. A, row of uncini in a thoracic uncinigerous torus in lateral view; B, detail of two thoracic uncini in lateral view; C, detail of a thoracic uncinus in frontal view; D, one abdominal neuropodial pinnule; E-F, detail of abdominal uncini in lateral and upper view.

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*



6

SEM micrographs of paratypes from Myanmar

Terebellides hutchingsae **spec. nov.**, SEM micrographs of paratypes from Myanmar, MNCN 16.01/0000 and MNCN 16.01/0000. A, anterior end, left lateral view, showing lateral lappets (ll) from TC1 to TC6 and geniculate chaetae (gc) in TC6; B, lateral view of lower branchial lobes (BL3-4) showing terminal projection (tp); C, ventral view of lower branchial lobes (BL3-4); D, dorsal view of TC4 to TC7 showing genital pores (gp) in TC4-5; E, TC6 geniculate chaetae (gc) in lateral view; F, detail of *capitium* of one geniculate chaeta.

**Note: Auto Gamma Correction was used for the image. This only affects the reviewing manuscript. See original source image if needed for review.*





Figure 7(on next page)

Map of SW Indo-Pacific Ocean

Map of SW Indo-Pacific Ocean showing locations where species of *Terebellides* reported in this paper were found (big circles), along with type localities of other *Terebellides* species previously described (medium circles) or reported (small circles).



Indian Ocean

Figure 8(on next page)

Line drawings of two Terebellides species previously described or reported in SW Pacific Ocean

Two *Terebellides* species previously described or reported in SW Pacific Ocean by Caullery (1915), redrawn from original. *Terebellides intoshi* Caullery, 1915. A. Anterior end in right lateral view showing great size of notopodia and notochaetae of thoracic chaetigers from TC6 onwards; *Terebellides stroemii* Sars, 1835. B. Anterior end in left lateral view of; C. dorsal and ventral view of branchial lobes showing high development of fifth branchial lobe (BL5), small size of branchial ventral lobes (BL3-4) and high degree of fusion of BL1-BL4.

