

Sex differences in pacing during 'Ultraman Hawaii'

Beat Knechtle ^{Corresp., 1, 2}, Pantelis T Nikolaidis ³

¹ Gesundheitszentrum St. Gallen, St. Gallen, Switzerland

² Institute of Primary Care, University of Zurich, Zurich, Switzerland

³ Exercise Physiology Laboratory, Nikaia, Greece

Corresponding Author: Beat Knechtle
Email address: beat.knechtle@hispeed.ch

Background. To date, little is known for pacing in ultra-endurance athletes competing in a non-stop event and in a multi-stage event, and especially, about pacing in a multi-stage event with different disciplines during the stages. Therefore, the aim of the present study was to examine the effect of age, sex and calendaryear on triathlon performance and variation of performance by events (*i.e.* swimming, cycling 1, cycling2 and running) in 'Ultraman Hawaii' held between 1983 and 2015.

Methods. Within each sex, participants were grouped in quartiles (*i.e.* Q1, Q2, Q3 and Q4) with Q1 being the fastest (*i.e.* lowest overall time) and Q4 the slowest (*i.e.* highest overall time). To compare performance among events (*i.e.* swimming, cycling 1, cycling 2 and running), race time in each event was converted in z score and this value was used for further analysis.

Results. A between-within subjects ANOVA showed a large sex×event ($p=0.015$, $\eta^2=0.014$) and a medium performance group×event interaction ($p=0.001$, $\eta^2=0.012$). No main effect of event on performance was observed ($p=0.174$, $\eta^2=0.007$). With regards to the sex×event interaction, three female performance groups (*i.e.* Q2, Q3 and Q4) increased race time from swimming to cycling 1, whereas only one male performance group (Q4) revealed a similar trend. From cycling 1 to cycling 2, the two slower female groups (Q3 and Q4) and the slowest male group (Q4) increased raced time. In women, the fastest group decreased (*i.e.* improved) race time from swimming to cycling 1 and thereafter, maintained performance, whereas in men, the fastest group decreased race time till cycling 2 and increased it in the running.

Conclusion. In summary, women pace differently than men during 'Ultraman Hawaii' where the fastest women decreased performance on day 1 and could then maintain on day 2 and 3, whereas the fastest men worsened performance on day 1 and 2 but improved on day 3.

Sex differences in pacing during ,Ultraman Hawaii'

Short title: Pacing during 'Ultraman Hawaii'

Beat Knechtle ^{1,2}, Pantelis T. Nikolaidis ³

¹ Gesundheitszentrum St. Gallen, St. Gallen, Switzerland

² Institute of Primary Care, University of Zurich, Zurich, Switzerland

³ Exercise Physiology Laboratory, Nikaia, Greece

Corresponding author:

Prof. Dr. med. Beat Knechtle
 Facharzt FMH für Allgemeinmedizin
 Gesundheitszentrum St. Gallen
 Vadianstrasse 26
 9001 St. Gallen
 Switzerland

34 Telefon +41 (0) 71 226 93 00
 35 Telefax +41 (0) 71 226 93 01
 36 E-Mail beat.knechtle@hispeed.ch

37 ABSTRACT

38 **Background.** To date, little is known for pacing in ultra-endurance athletes competing in a non-
 39 stop event and in a multi-stage event, and especially, about pacing in a multi-stage event with
 40 different disciplines during the stages. Therefore, the aim of the present study was to examine the
 41 effect of age, sex and calendar year on triathlon performance and variation of performance by
 42 events (*i.e.* swimming, cycling 1, cycling2 and running) in ‘Ultraman Hawaii’ held between
 43 1983 and 2015.

44 **Methods.** Within each sex, participants were grouped in quartiles (*i.e.* Q1, Q2, Q3 and Q4) with
 45 Q1 being the fastest (*i.e.* lowest overall time) and Q4 the slowest (*i.e.* highest overall time). To
 46 compare performance among events (*i.e.* swimming, cycling 1, cycling 2 and running), race time
 47 in each event was converted in z score and this value was used for further analysis.

48 **Results.** A between-within subjects ANOVA showed a large sex×event ($p=0.015$, $\eta^2=0.014$) and
 49 a medium performance group×event interaction ($p=0.001$, $\eta^2=0.012$). No main effect of event on
 50 performance was observed ($p=0.174$, $\eta^2=0.007$). With regards to the sex×event interaction, three
 51 female performance groups (*i.e.* Q2, Q3 and Q4) increased race time from swimming to cycling
 52 1, whereas only one male performance group (Q4) revealed a similar trend. From cycling 1 to
 53 cycling 2, the two slower female groups (Q3 and Q4) and the slowest male group (Q4) increased
 54 raced time. In women, the fastest group decreased (*i.e.* improved) race time from swimming to
 55 cycling 1 and thereafter, maintained performance, whereas in men, the fastest group decreased
 56 race time till cycling 2 and increased it in the running.

Conclusion. In summary, women pace differently than men during ‘Ultraman Hawaii’ where the fastest women decreased performance on day 1 and could then maintain on day 2 and 3, whereas the fastest men worsened performance on day 1 and 2 but improved on day 3.

Key words: swimming, cycling, running, triathlon, ultra-endurance

73

74

75 **Introduction**

76 ‘Ultraman Hawaii’ is one of the longer distance triathlon lasting three days and consisting of 10
77 km swimming, 165 km cycling (day 1), 261 km cycling (day 2) and 85 km running (day 3).

78 These distances are much longer than the typical Ironman triathlon events (3.8 km swimming,
79 180 km cycling and 42 km running) (Seedhouse et al. 2006). Performance in Ironman triathlon
80 events is related to physiological measures, such as maximal oxygen uptake, anaerobic threshold,
81 economy of movement, hydration and energy homeostasis (Laursen & Rhodes 2001). Another
82 parameter that might play a role on the performance in ‘Ultraman Hawaii’ is pacing. Although
83 pacing has been studied extensively in separate endurance and ultra-endurance events (Knechtle
84 et al. 2015a; Nikolaidis & Knechtle 2016) and in shorter formats of triathlon (Wu et al. 2016;
85 Wu et al. 2015), no information on pacing in a multi-day triathlon such as ‘Ultraman Hawaii’ is
86 available , *i.e.* how the performance in one event (*e.g.* swimming) influences the performance of
87 subsequent events (*e.g.* cycling).

88 Pacing is defined as time per distance, usually expressed in minutes per kilometer or mile
89 (Andrew & Remco 2012). A pacing strategy – or a plan how to distribute an athlete’s potential –
90 is crucial for successful athletic practice (Foster et al. 2005) and has a considerable effect on
91 performance in endurance sports (Abbiss & Laursen 2008; Foster et al. 1993). Abbiss and
92 Laursen (2008) postulated six different pacing strategies such as negative pacing (*i.e.* increase in
93 speed over time), positive pacing (*i.e.* continuous slowing over time), all-out pacing (*i.e.*
94 maximal speed possible), even pacing (*i.e.* same speed over time), parabolic-shaped pacing (*i.e.*

positive and negative pacing in different segments of the race) and variable pacing (*i.e.* pacing with multiple fluctuations). Abbiss and Laursen (2008) stated that athletes in endurance sports often adopt a positive pacing strategy. However, they did not exclude the possibility that an even pacing strategy may be optimal to successfully complete an endurance event. These pacing strategies concern a single race, but their principles can be applied equally in a triathlon where performance might vary from an event to another event.

In a triathlon, the regulation of pacing may be influenced by different intrinsic and extrinsic factors (Wu et al. 2014). Moreover, the role of athletes' physiological state and emotional status has also been recognized (Baron et al. 2011). A complex system has been proposed to be involved in pacing regulation, dealing with elevated temperature, altered oxygen content of air, reduced fuel availability and subjective perception of distance covered (Tucker and Noakes 2009). Most studies investigated pacing strategies for short distances such as sprint distance triathlon (Wu et al. 2016), Olympic distance triathlon (Wu et al. 2015), half-Ironman triathlon (Wu et al. 2015) and Ironman triathlon (Johnson et al. 2016). Only a few studies investigated pacing strategies in very long-distance triathlons (Knechtle et al. 2014). However, no study investigated the pacing strategy in a three-stage ultra-triathlon such as 'Ultraman Hawaii'.

There is evidence that performance in split disciplines has an effect of overall race performance in a triathlon (Le Meur et al. 2011; Vleck et al. 2006; Wu et al. 2016; Wu et al. 2015). In a sprint distance triathlon, faster cycling split times and overall race times were achieved with positive swimming pacing (Wu et al. 2016). For the Olympic distance triathlon, Le Meur *et al.* (Le Meur et al. 2011) investigated the performance during the running split of 107 finishers in an Olympic distance triathlon and stated that elite triathletes should reduce their initial running speed for better performances during competitions. Bernard *et al.* (Bernard et al. 2009) investigated ten elite

triathletes during the cycling split in an Olympic distance triathlon and showed that a progressive decrease in speed combined with an increase in speed variability occurred during the race. A comparison between Olympic distance and Half-Ironman distance triathlon showed pacing strategies during a triathlon race are strongly influenced by both the distance and discipline (Wu et al. 2015). In Ironman triathlon, pacing in downhill segments in the cycling and running split have an influence on overall race performance (Johnson et al. 2016).

There is also evidence that performance decreases during a multi-stage ultra-endurance triathlon (Herbst et al. 2011; Knechtle et al. 2014). In race held over ten times the Ironman distance, performance progressively declined over days (Knechtle et al. 2015a; Knechtle et al. 2014). Additionally, sex has an influence on pacing during triathlon (Le Meur et al. 2009; Vleck et al. 2008). In Olympic distance triathlon, female and male elite triathletes adopted similar positive pacing strategies during swimming and running. Female triathletes were more affected by changes in slope during the cycling and running splits (Le Meur et al. 2009).

To date, little is known for pacing in ultra-endurance athletes competing in a non-stop event (Heidenfelder et al. 2016; Knechtle et al. 2015b) and in a multi-stage event (Herbst et al. 2011). Especially, we have no knowledge about pacing in a multi-stage event with different disciplines during the stages. Such knowledge would be of practical value for both coaches and athletes in order to optimize their pacing strategy according to sex and performance.

Therefore, the aim of the present study was to investigate pacing strategies in ultra-triathletes competing in ‘Ultraman Hawaii’. This race is held since 1983 as a three-stage race with swimming and cycling on day 1, cycling on day 2 and running on day 3 (Meili et al. 2013). We hypothesized that athletes who would start fast on day 1 would slow down more compared to

140 athletes who start slow on day 1. On the contrary, athletes who would start slow in the race
 141 would be able to improve during the race. Furthermore, we hypothesized to find differences in
 142 pacing between women and men.

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

Methods

To test our hypotheses, all women and men who finished ‘Ultraman Hawaii’ between 1983 and 2015 were considered. Sex, age and split times for the three stages were recorded and analyzed. The subjects for this study were obtained from official race website of ‘Ultraman Hawaii’ at <http://ultramanworld.com>. ‘Ultraman Hawaii’ is held as the Ultraman World Championship on the Big Island of Hawaii, USA. The race is limited to 40 athletes on an invitation-only basis and attracts participants from around the world. Racers must have reached their 20th birthday prior to the start of stage one. Each racer must be accompanied by an individual support team of at least two people over the entire course. The first edition of ‘Ultraman Hawaii’ was in 1983. Since then, the race was continuously held until today with exception of 1987 and 1991 where no race was organized. ‘Ultraman Hawaii’ is a three-day, 515 km (320 mile) race. The race is divided into three stages over three days: The first is a 10-km ocean swim from Kailua Bay to Keauhou Bay, followed by a 145-km cross-country bike ride, with vertical climbs that total ~1800 m. Stage two is a 276-km bike ride from Volcanoes National Park to Kohala Village Inn in Hawi, with total vertical climbs of ~1200 m. Stage three is a 84-km double-marathon, which starts at Hawi and finishes on the beach at the Old Kona Airport State Recreation Area. Each stage must be completed within 12 hours or faster. The swim portion of stage one must be completed in 5.5 hours or faster. Participants who do not reach the finish lines within the time limits are disqualified. All procedures used in the study were approved by the Institutional Review Board of Kanton St. Gallen, Switzerland, with a waiver of the requirement for informed consent of the participants given the fact that the study involved the analysis of publicly available data (01/06/2010). Data from 1983 to 2015 were obtained from the

official race website of ‘Ultraman Hawaii’ at <http://ultramanworld.com>. Age, sex, split times for the three stages (i.e. swimming and cycling on day 1, cycling on day 2 and running on day 3) were used for analysis.

Statistical Analyses

All statistical analyses were performed using IBM SPSS v.20.0 (SPSS, Chicago, USA). Descriptive statistics (mean±standard deviations) were calculated for all data. Within each sex, participants were grouped in quartiles (Q1, Q2, Q3 and Q4) with Q1 being the fastest (*i.e.* lowest overall time) and Q4 the slowest (highest overall time). To compare performance among events (*i.e.* swimming, cycling 1, cycling 2 and running), time in each event was converted in z score and this value was used for further analysis. Since race duration and race time variation differed among events, it was hard to compare race times of different events; thus, z score was calculated from the formula score – mean / standard deviation and its use allowed comparing performance in different events. A between-within subjects analysis of variance (ANOVA) examined the effect of sex and overall performance (Q1, Q2, Q3 and Q4), as well as their interaction, on variation among events (*i.e.* swimming, cycling 1, cycling 2 and running). Sex and overall performance were defined as between subjects factors, and performance in events as within subjects factor. Bonferroni post-hoc analysis tested differences among groups. The magnitude of these differences was examined using eta square (η^2), classified as trivial ($\eta^2 < 0.01$), small ($0.01 \leq \eta^2 < 0.06$), medium ($0.06 \leq \eta^2 < 0.14$) and large ($\eta^2 \geq 0.14$). To study differences in performance and age by sex and calendar year from 1983 to 2015, we used a mixed-effects regression model with triathletes as random variable, whereas sex and calendar year were assigned as fixed variables. In addition, we examined the variation of age by calendar year using a mixed-effects regression model with triathletes as random variable, and sex and calendar year

as a fixed variable. We examined interaction effects among these fixed variables. Akaike
information criterion (AIC) was used to select the final model.

Results

Men did not differ from women for age (-0.6 years (-2.1;1.0), mean difference (95% confidence interval), swimming time (-0:04:52 h:min:s (-0:13:32;0:03:46)) and running (-0:15:23 h:min:s (-0:34:32;0:03:46)) (Table 1). Men were faster in the overall race (-1:08:21 h:min:s (-1:51:36;-0:25:07)), cycling 1 (-0:18:06 h:min:s (-0:30:09;-0:06:03)) and cycling 2 (-0:29:59 (-0:45:14;-0:14:43)).

Different trends among events were observed among groups differing for overall performance and between sexes (Figure 1). The between-within subjects ANOVA showed a sex×event ($p=0.015$, $\eta^2=0.014$, small magnitude) and a performance group×event interaction ($p=0.001$, $\eta^2=0.012$, small magnitude). No main effect of event on performance was observed ($p=0.174$, $\eta^2=0.007$, trivial magnitude). With regards to the sex×event interaction, three female performance groups (Q2, Q3 and Q4) increased race time from swimming to cycling 1, whereas only one male performance group (Q4) revealed a similar trend. From cycling 1 to cycling 2, the two slower female groups (Q3 and Q4) and the slowest male group (Q4) increased race time.

In women, the fastest group (Q1) decreased race time from swimming to cycling 1 and thereafter, maintained performance, whereas in men, the fastest group (Q1) decreased race time till cycling 2 and increased it in the running.

With regards to the participation of women and men across calendar years, there was an association between sex and calendar years that was, women and men did not participate similarly: an increased participation of women was observed during the last years (Figure 2).

There was a main effect of calendar year on overall race and all race events' times ($p<0.001$) (Table 2, Figure 3). A sex×calendar year interaction was observed only in cycling 2 ($p=0.013$), in

which the increase in race time was more pronounced in women than in men. Also, there was a main effect of calendar year on age ($p < 0.001$), in which the age of participants increased during the last years (Table 3, Figure 4).

259 Discussion

260 This study investigated pacing trends across stages in ‘Ultraman Hawaii’ held between 1983 and
 261 2015 with the hypothesis that fast starters would slow down (*i.e.* increase race time) compared to
 262 slow starters who would improve (*i.e.* decrease race time). We found, as expected, differences
 263 between performance groups and sexes.

264 The most important finding was that the best women adopted a different pacing strategy during
 265 the three days compared to the best men. In details, the fastest overall women decreased split
 266 times from swimming to cycling 1 (*i.e.* day 1) and maintained performance thereafter (*i.e.* day 2
 267 and day 3), whereas the fastest overall men decreased split times from swimming to cycling 2
 268 (*i.e.* day 1 and day 2) but increased time in running (*i.e.* day 3). Expressed in other words, the best
 269 women improved performance in the beginning of the race (*i.e.* day 1), whereas the best men
 270 improved performance in both day 1 and day 2, but were slower in the end of the race (*i.e.* day
 271 3). Potential explanations for these differences could be sex differences in anthropometry, race
 272 strategy, separate performance in the three split disciplines and the increase in participation in
 273 women in the last years. Therefore, the main sex difference in pacing of the fastest athletes was
 274 highlighted in day 2, when men decreased race time, whereas women maintained a steady
 275 pacing. This pattern in day 2 might explain the sex differences in day 3, when men increased
 276 race time and women maintained steady pacing. These sex differences indicated that
 277 performance in a split might affect following performance.

278 The findings of the present study were in agreement with the existing literature on sex
 279 differences in pacing. For instance, differences in pacing between women and men have been
 280 observed in triathlon (Le Meur et al. 2011; Vleck et al. 2008), but also in other sports disciplines

such as running. Deaner and Lowen (2016) investigated 3,948 performances in 5-km cross country running and showed that women slowed significantly more than men. These authors argued that the sex difference in pacing partly reflected a sex difference in some aspect of decision making, such as over-confidence, risk perception, or willingness to tolerate discomfort. In running, difference in pacing might vary by distance. In marathon running, however, men are more likely to slow down (i.e. increase split times) than women (Deaner et al. 2014). A more recent study, however, found that women, older and faster marathoners are better pacers than men, younger and slower marathoners, respectively (March et al. 2011). These differences in pacing in runners are most likely explained by differences in performance level (Lima-Silva et al. 2010).

A further explanation for the differences in pacing between the best women and men could be the differences in split and overall performances. Considering performance in split and overall race time, men were faster than women in cycling (*i.e.* day 1 and day 2) and overall race time, but not in swimming (*i.e.* day 1) and running (*i.e.* day 3). We found that the fastest men improved performance in cycling 1 and cycling 2, but impaired on day 3 in running. On the contrary, the fastest women improved performance in cycling 1 and maintained it in cycling 2 and running. A potential explanation could be that women saved energy during the cycling for the running split. When men would have completely exhausted got of the bike to run the next day a double marathon, they would not have been able to improve performance. Although women and men suffer an energy deficit during an Ironman triathlon (Kimber et al. 2002), female Ironman triathlons suffer no decrease in body mass (Knechtle et al. 2010b) compared to male Ironman triathletes (Knechtle et al. 2010a). Male Ironman triathletes suffer a decrease in

skeletal muscle mass most probably due to depletion of glycogen stores of the lower limb (Mueller et al. 2013).

Further explanations for the differences in pacing between women and men could be differences in anthropometry, training and previous experience (Knechtle et al. 2010d). Considering predictor variables for Ironman triathlon performance, percent body fat was related to race performance in male athletes and to average weekly training in female athletes. Personal best time in an Ironman triathlon was associated with total race time for both male and female athletes (Knechtle et al. 2010c). The greater height and associated longer limbs of the men might also provide a biomechanical advantage. However, anthropometric characteristics such as body height and limb length were not investigated in Ironman triathletes (Knechtle et al. 2010c, 2010d). Personal best marathon time is a strong predictor for Ironman triathlon race in both women and men (Knechtle et al. 2015c). Also in male marathoners, body fat and running speed during training was a strong predictive variable, not body height or limb length (Barandun et al. 2012). For female marathoners, however, circumference of the calf and running speed during training were the most important predictive variables (Schmid et al. 2012).

Differences in metabolic predominance between men and women might also explain these findings. For example, women tend to use fat metabolism to a higher degree than men (Blaak 2001). Since fat is such a huge energy source, women may be better at preserving energy and maintaining similar pacing with these ultra-races. Men have a greater glycolytic capacity (Lundsgaard & Kiens 2014) which means they have greater anaerobic and high intensity exercise capacities but of course greater metabolite accumulation leading to greater fatigue (Bogdanis 2012).

A further important finding was that the age of the athletes increased across years and their performance impaired. This is in contrast to findings for ‘Ironman Hawaii’ where the age of annual top ten female and male triathletes in the 'Ironman Hawaii' increased between 1983 and 2012 while their performances improved (Gallmann et al. 2014). The most likely explanation for these disparate findings is the fact that in the analysis for ‘Ironman Hawaii’ the annual ten fastest were considered whereas in the present analysis all women and men were included in the analysis. Moreover, a small different sex trend in performance was remarkable during the last years, where race time in cycling 2 increased more in women than in men. The increased overall time during the last years was attributed mostly to increased cycling 2 times in women and to increased cycling 2 and running time in men (Figure 4). To counteract this calendar year trend, this finding implied that coaches should focus on the improvement of cycling 2 in women and on both cycling 2 and running in men.

A limitation of this analysis is the fact that specific characteristics for this race such as alteration in body composition (Baur et al. 2016) and cardiac and pulmonary responses during the race (Seedhouse et al. 2006) were not considered. In addition, since the distance each event is much longer in “Ultraman Hawaii” than other Ironman triathlon events, caution is required in order to generalize the findings of the present study to other triathlon formats. Strength of the present study was that it included all participants to all races of “Ultraman Hawaii”.

Conclusions

In ‘Ultraman Hawaii’, women pace differently than men where the fastest women decreased performance on day 1 and could then maintain on day 2 and 3 whereas the fastest men impaired performance on day 1 and 2 but improved on day 3. For athletes and coaches, women and men should follow sex-tailored training and race tactics in such events.

References

- Abbiss CR, and Laursen PB. 2008. Describing and understanding pacing strategies during athletic competition. *Sports Medicine* 38:239-252.
- Andrew E, and Remco P. 2012. *Pacing in sport and exercise: A psychophysiological perspective*. Hauppauge, NY: Nova Science Publishers.
- Barandun U, Knechtle B, Knechtle P, Klipstein A, Rüst CA, Rosemann T, and Lepers R. 2012. Running speed during training and percent body fat predict race time in recreational male marathoners. *Open Access Journal of Sports Medicine* 3:51-8. doi: 10.2147/OAJSM.S33284
- Baron B, Moullan F, Deruelle F, and Noakes TD. 2011. The role of emotions on pacing strategies and performance in middle and long duration sport events. *British Journal of Sports Medicine* 45:511-517.
- Baur DA, Bach CW, Hyder WJ, and Ormsbee MJ. 2016. Fluid retention, muscle damage, and altered body composition at the Ultraman triathlon. *European Journal of Applied Physiology* 116:447-458. 10.1007/s00421-015-3291-9
- Bernard T, Hausswirth C, Le Meur Y, Bignet F, Dorel S, and Brisswalter J. 2009. Distribution of power output during the cycling stage of a triathlon world cup. *Medicine and Science in Sports and Exercise* 41:1296-1302. 10.1249/MSS.0b013e318195a233
- Blaak E. 2001. Gender differences in fat metabolism. *Current Opinion in Clinical Nutrition and Metabolic Care* 4:499-502.
- Bogdanis GC. 2012. Effects of physical activity and inactivity on muscle fatigue. *Frontiers in Physiology* 3:142. doi: 10.3389/fphys.2012.00142. eCollection 2012.

- Deaner RO, Carter RE, Joyner MJ, and Hunter SK. 2014. Men are more likely than women to slow in the marathon. *Medicine and Science in Sports and Exercise* 47:607-616.
10.1249/MSS.0000000000000432
- Deaner RO, and Lowen A. 2016. Males and Females Pace Differently in High School Cross Country Races. *J Strength Cond Res*. 10.1519/jsc.0000000000001407
- Foster C, Hoyos J, Earnest C, and Lucia A. 2005. Regulation of energy expenditure during prolonged athletic competition. *Medicine and Science in Sports and Exercise* 37:670-675.
10.1249/01.MSS.0000158183.64465.BF
- Foster C, Snyder AC, Thompson NN, Green MA, Foley M, and Schrager M. 1993. Effect of pacing strategy on cycle time trial performance. *Medicine and Science in Sports and Exercise* 25:383-388.
- Gallmann D, Knechtle B, Rüst CA, Rosemann T, and Lepers R. 2014. Elite triathletes in 'Ironman Hawaii' get older but faster. *Age* 36:407-416. 10.1007/s11357-013-9534-y
- Heidenfelder A, Rosemann T, Rüst CA, and Knechtle B. 2016. Pacing strategies of ultracyclists in the "race across America. *Int J Sports Physiol Perform* 11:319-327.
10.1123/ijsp.2015-0051
- Herbst L, Knechtle B, Lopez CL, Andonie JL, Fraire OS, Kohler G, Rüst CA, and Rosemann T. 2011. Pacing strategy and change in body composition during a Deca iron triathlon. *Chinese Journal of Physiology* 54. 10.4077/CJP.2011.AMM115
- Johnson EC, Pryor JL, Casa DJ, Belval LN, Vance JS, DeMartini JK, Maresh CM, and Armstrong LE. 2016. Bike and run pacing on downhill segments predict Ironman triathlon relative success. *Journal of Science and Medicine in Sport* 18:82-87.
10.1016/j.jsams.2013.12.001

- Kimber NE, Ross JJ, Mason SL, and Speedy DB. 2002. Energy balance during an Ironman triathlon in male and female triathletes. *International Journal of Sport Nutrition* 12:47-62.
- Knechtle B, Baumann B, Wirth A, Knechtle P, and Rosemann T. 2010a. Male ironman triathletes lose skeletal muscle mass. *Asia Pacific Journal of Clinical Nutrition* 19:91-97.
- Knechtle B, Bragazzi NL, Rosemann T, and Rüst CA. 2015a. Pacing in a self-paced world record attempt in 24-h road cycling. *SpringerPlus* 4. 10.1186/s40064-015-1445-1
- Knechtle B, Knechtle R, Stiefel M, Zingg MA, Rosemann T, and Rüst CA. 2015c. Variables that influence Ironman triathlon performance - what changed in the last 35 years? *Open Access Journal of Sports Medicine* 6:277-290. doi: 10.2147/OAJSM.S85310. eCollection 2015
- Knechtle B, Rosemann T, Lepers R, and Rüst CA. 2014. A comparison of performance of Deca Iron and Triple Deca Iron ultra-triathletes. *SpringerPlus* 3. 10.1186/2193-1801-3-461
- Knechtle B, Wirth A, Baumann B, Knechtle P, Kohler G, Rosemann T, and Senn O. 2010b. An ironman triathlon does not lead to a change in body mass in female triathletes. *Research in Sports Medicine* 18:115-126. 10.1080/15438621003627059
- Knechtle B, Wirth A, Baumann B, Knechtle P, and Rosemann T. 2010c. Personal best time, percent body fat, and training are differently associated with race time for male and female Ironman triathletes. *Research Quarterly for Exercise and Sport* 81:62-68.
- Knechtle B, Wirth A, Baumann B, Knechtle P, Rosemann T, and Oliver S. 2010d. Differential correlations between anthropometry, training volume, and performance in male and female Ironman triathletes. *Journal of Strength and Conditioning Research* 24:2785-2793. 10.1519/JSC.0b013e3181c643b6

- 440 Knechtle B, Zingg MA, Knechtle P, Rosemann T, and Rüst CA. 2015b. Feet swelling in a
- 441 multistage ultraendurance triathlete: A case study. *International Journal of General*
- 442 *Medicine* 8:325-332. 10.2147/IJGM.S94542
- 443 Laursen PB, and Rhodes EC. 2001. Factors affecting performance in an ultraendurance triathlon.
- 444 *Sports Medicine* 31:195-209.
- 445 Le Meur Y, Bernard T, Dorel S, Abbiss CR, Honnorat G, Brisswalter J, and Hausswirth C. 2011.
- 446 Relationships between triathlon performance and pacing strategy during the run in an
- 447 international competition. *Int J Sports Physiol Perform* 6:183-194.
- 448 Le Meur Y, Hausswirth C, Dorel S, Bignet F, Brisswalter J, and Bernard T. 2009. Influence of
- 449 gender on pacing adopted by elite triathletes during a competition. *European Journal of*
- 450 *Applied Physiology* 106:535-545. 10.1007/s00421-009-1043-4
- 451 Lima-Silva AE, Bertuzzi RCM, Pires FO, Barros RV, Gagliardi JF, Hammond J, Kiss MA, and
- 452 Bishop DJ. 2010. Effect of performance level on pacing strategy during a 10-km running
- 453 race. *European Journal of Applied Physiology* 108:1045-1053. 10.1007/s00421-009-
- 454 1300-6
- 455 Lundsgaard AM, and Kiens B. 2014. Gender differences in skeletal muscle substrate metabolism
- 456 - molecular mechanisms and insulin sensitivity. *Frontiers in Endocrinology (Lausanne)*
- 457 5:195. doi: 10.3389/fendo.2014.00195. eCollection 2014.
- 458 March DS, Vanderburgh PM, Titlebaum PJ, and Hoops ML. 2011. Age, sex, and
- 459 finish time as determinants of pacing in the marathon. *Journal of Strength and*
- 460 *Conditioning Research* 25:386-391. 10.1519/JSC.0b013e3181bffd0f

- Meili D, Knechtle B, Rüst CA, Rosemann T, and Lepers R. 2013. Participation and performance trends in 'Ultraman Hawaii' from 1983 to 2012. *Extreme Physiology and Medicine* 2. 10.1186/2046-7648-2-25
- Mueller SM, Anliker E, Knechtle P, Knechtle B, and Toigo M. 2013. Changes in body composition in triathletes during an Ironman race. *European Journal of Applied Physiology* 113:2343-2352. 10.1007/s00421-013-2670-3
- Nikolaidis PT, and Knechtle B. 2016. Pacing in age group freestyle swimmers - The XV FINA World Masters Championships. *Journal of Sports Sciences* in print. 10.1080/02640414.2016.1213412
- Schmid W, Knechtle B, Knechtle P, Barandun U, Rüst CA, Rosemann T, and Lepers R. 2012. Predictor variables for marathon race time in recreational female runners. *Asian Journal of Sports Medicine* 3:90-98
- Seedhouse ELO, Walsh ML, and Blaber AP. 2006. Heart rate, mean arterial blood pressure, and pulmonary function changes associated with an ultraendurance triathlon. *Wilderness and Environmental Medicine* 17:240-245.
- Vleck VE, Bentley DJ, Millet GP, and Bürgi A. 2008. Pacing during an elite Olympic distance triathlon: Comparison between male and female competitors. *Journal of Science and Medicine in Sport* 11:424-432. 10.1016/j.jsams.2007.01.006
- Vleck VE, Bürgi A, and Bentley DJ. 2006. The consequences of swim, cycle, and run performance on overall result in elite olympic distance triathlon. *International Journal of Sports Medicine* 27:43-48. 10.1055/s-2005-837502
- Wu SS, Peiffer JJ, Brisswalter J, Nosaka K, and Abbiss CR. 2014. Factors influencing pacing in triathlon. *Open Access Journal of Sports Medicine* 5:223-234. 10.2147/oajsm.s44392

Wu SS, Peiffer JJ, Peeling P, Brisswalter J, Lau WY, Nosaka K, and Abbiss CR. 2016. Positive swim pacing improves sprint triathlon performance in trained athletes. *International Journal of Sports Physiology and Performance* 10.1123/ijsp.2015-0580

Wu SSX, Peiffer JJ, Brisswalter J, Nosaka K, Lau WY, and Abbiss CR. 2015. Pacing strategies during the swim, cycle and run disciplines of sprint, Olympic and half-Ironman triathlons. *European Journal of Applied Physiology* 115:1147-1154. 10.1007/s00421-014-3096-2

Table 1 Performance time in participants by event

Total (n=765)	Women (n=120)	Men (n=645)
---------------	---------------	-------------

Age (years)	39.4±8.0	39.9±8.4	39.3±7.9
Overall race (h:min:s)	29:28:50±3:42:48	30:26:25±3:27:52	29:18:06±3:44:00*
Swimming (h:min:s)	3:43:58±0:44:21	3:48:05±0:43:02	3:43:12±0:44:35
Cycling 1 (h:min:s)	6:28:08±1:02:02	6:43:24±1:04:35	6:25:17±1:01:11*
Cycling 2 (h:min:s)	9:53:29±1:18:54	10:18:46±1:22:06	9:48:47±1:17:27†
Running (h:min:s)	9:23:14±1:38:16	9:36:12±1:31:31	9:20:49±1:39:22

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

Table 2 Coefficients (C) and standard errors (SE) from multi-variate regression models for the performance by calendar year in women and men

		C	SE_p	
Swimming	Sex (=female)	9885.21	48918.07	0.840
	Calendar year	121.80	9.26	<0.001
	Interaction sex×calendar year	-5.06	24.42	0.836
Cycling1	Sex (=female)	-58501.28	69593.16	0.401
	Calendar year	148.77	13.18	<0.001
	Interaction sex×calendar year	29.40	34.74	0.398
Cycling2	Sex (=female)	-156860.04	62856.90	0.013
	Calendar year	344.63	11.90	<0.001
	Interaction sex×calendar year	78.40	31.38	0.013
Running	Sex (=female)	118761.88	112339.14	0.291
	Calendar year	233.10	21.27	<0.001
	Interaction sex×calendar year	-59.33	56.08	0.290
Total	Sex (=female)	-86714.23	210194.69	0.680
	Calendar year	848.30	39.80	<0.001
	Interaction sex×calendar year	43.40	104.92	0.679

Table 3 Coefficients (C) and standard errors (SE) from multi-variate regression models for the age of participants by calendar year in women and men

		C	SE	<i>p</i>
Age	Sex (=female)	-195.41	152.82	0.201
	Calendar year	0.29	0.03	<0.001
	Interaction sex×calendar year	0.10	0.08	0.203

552

553

554 **Legends of figures**

555 **Figure 1** Variation of performance in events by overall performance level and sex.

556 **Figure 2** Participation of women and men in the Hawaii Ultraman from 1983 to 2015.

557 Women are depicted in dark and men in bright bars

558 **Figure 3** Variation of female and male participants' performance in the Hawaii Ultraman
559 from 1983 to 2015. Total race time for women (solid line) and men (dashed line)
560 (left), Race time in swimming (S), cycling 1 (C1), cycling 2 (C2) and running (R)
561 for women (middle) and men (right).

562 **Figure 4** Variation of female and male participants' age in the Hawaii Ultraman from 1983
563 to 2015. Women are depicted by solid and men by dashed line.

564

565

566

567

568


569

570


571

572

573 **Figure 1**




575 **Figure 2**



587


588

589 **Figure 3**



590

591 **Figure 4**



592

593