

# Root bacterial endophytes alter plant phenotype, but not physiology

**Jeremiah A Henning** <sup>Corresp., 1</sup>, **David J Weston** <sup>2</sup>, **Dale A Pelletier** <sup>2</sup>, **Collin M Timm** <sup>3</sup>, **Sara S Jawdy** <sup>2</sup>, **Aimée T Classen** <sup>1, 4</sup>

<sup>1</sup> Department of Ecology & Evolutionary Biology, University of Tennessee - Knoxville, Knoxville, Tennessee, United States

<sup>2</sup> Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, United States

<sup>3</sup> Joint Institute for Biological Sciences, University of Tennessee, Oak Ridge, TN, United States

<sup>4</sup> The Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark

Corresponding Author: Jeremiah A Henning

Email address: jhennin2@vols.utk.edu

Plant traits, such as root and leaf area, influence how plants interact with their environment and the diverse, microbiota living within plants can influence plant morphology and physiology. Here, we explored how three bacterial strains isolated from the *Populus* root microbiome, influenced plant phenotype. We chose three bacterial strains that differed in predicted metabolic capabilities, plant hormone production and metabolism, and secondary metabolite synthesis. We inoculated each bacterial strain on a single genotype of *Populus trichocarpa* and measured the response of plant growth related traits (root:shoot, biomass production, root and leaf growth rates) and physiological traits (chlorophyll content, net photosynthesis, net photosynthesis at saturating light -  $A_{sat}$ , and saturating  $CO_2$  -  $A_{max}$ ). Overall, we found that bacterial root endophyte infection increased root growth rate up to 184% and leaf growth rate up to 137% relative to non-inoculated control plants, evidence that plants respond to bacteria by modifying morphology. However, endophyte inoculation had no influence on total plant biomass and photosynthetic traits (net photosynthesis, chlorophyll content). In sum, bacterial inoculation did not significantly increase plant carbon fixation and biomass, but their presence altered where and how carbon was being allocated in the plant host.

Title page

**Root bacterial endophytes alter plant phenotype, but not physiology**

Jeremiah A. Henning<sup>1</sup>, David J. Weston<sup>2</sup>, Dale A. Pelletier<sup>2</sup>, Collin M. Timm<sup>3</sup>, Sara S. Jawdy<sup>2</sup>,  
Aimée T. Classen<sup>1,4</sup>

<sup>1</sup> Ecology & Evolutionary Biology, The University of Tennessee, Knoxville, TN USA

<sup>2</sup> Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN USA

<sup>3</sup> Joint Institute for Biological Sciences, University of Tennessee, Oak Ridge, TN USA

<sup>4</sup> The Natural History Museum of Denmark, The University of Copenhagen København Ø,  
Denmark

Corresponding author:

Jeremiah A. Henning

jhennin2@vols.utk.edu

20 **Abstract (193 words)**

21 Plant traits, such as root and leaf area, influence how plants interact with their environment and  
 22 the diverse, microbiota living within plants can influence plant morphology and physiology.  
 23 Here, we explored how three bacterial strains isolated from the *Populus* root microbiome,  
 24 influenced plant phenotype. We chose three bacterial strains that differed in predicted metabolic  
 25 capabilities, plant hormone production and metabolism, and secondary metabolite synthesis. We  
 26 inoculated each bacterial strain on a single genotype of *Populus trichocarpa* and measured the  
 27 response of plant growth related traits (root:shoot, biomass production, root and leaf growth  
 28 rates) and physiological traits (chlorophyll content, net photosynthesis, net photosynthesis at  
 29 saturating light -  $A_{\text{sat}}$ , and saturating  $\text{CO}_2$  -  $A_{\text{max}}$ ). Overall, we found that bacterial root endophyte  
 30 infection increased root growth rate up to 184% and leaf growth rate up to 137% relative to non-  
 31 inoculated control plants, evidence that plants respond to bacteria by modifying morphology.  
 32 However, endophyte inoculation had no influence on total plant biomass and photosynthetic  
 33 traits (net photosynthesis, chlorophyll content). In sum, bacterial inoculation did not significantly  
 34 increase plant carbon fixation and biomass, but their presence altered where and how carbon was  
 35 being allocated in the plant host.  
 36

# 37 Introduction

38 A recent review exploring microbiome-mediated plant traits found that plant-associated microbes  
 39 can modify fourteen out of thirty commonly measured functional traits (Cornelissen et al., 2003;  
 40 Friesen et al., 2011). For example, inoculation with common root-colonizing bacterial strains  
 41 influenced root and leaf architectural traits, such as specific leaf area and specific root length, as  
 42 well as plant physiological traits such as carbon fixation and chlorophyll content (Harris,  
 43 Pacovsky, & Paul, 1985; Ma et al., 2003; Friesen, 2013). Further, inoculation by different  
 44 members of the plant microbiome may differentially alter plant phenotype (Zamioudis et al.,  
 45 2013; Timm et al., 2016). The presence of unique bacterial strains in legume genotypes  
 46 explained more variation in shoot biomass, root biomass, and plant height than plant genotype  
 47 did (Tan & Tan, 1986). Inoculation of common endophytes can also inhibit primary root  
 48 elongation and promote lateral root formation and root hair production (Zamioudis et al., 2013;  
 49 Weston et al., 2012). Recent breakthroughs in the multitude of the -omics fields have allowed  
 50 for unprecedented mechanistic investigations of microbe-induced changes in host function  
 51 (Verhagen et al., 2004; Walker et al., 2011; Weston et al., 2012; Vandenkoornhuyse et al., 2015;  
 52 Timm et al., 2015; 2016) and have been the subject of multiple recent reviews (Friesen et al.,  
 53 2011; Friesen, 2013; Vandenkoornhuyse et al., 2015; Hacquard & Schadt, 2015; Lebeis, 2015;  
 54 and many others). This work demonstrated that plant growth promoting bacteria elicit numerous  
 55 changes in host gene expression through multiple and simultaneous hormonal and immune  
 56 response pathways (Verhagen et al., 2004; Walker et al., 2011; Weston et al., 2012; Drogue et  
 57 al., 2014; Timm et al., 2016). However, these studies fall short in explaining how changes in  
 58 gene expression influence the overall plant phenotype or plant function. Thus, understanding the  
 59 response of plant traits and overall plant phenotype to microbial strains remains a research gap.

Here, we inoculated three endophytic bacterial strains (*Pseudomonas fluorescens* GM41, *Pseudomonas fluorescens* GM30, and *Burkholderia* sp. BT03), originally isolated from wild *Populus*, on a single genotype of *Populus trichocarpa* and measured plant phenotypic response to bacterial inoculation. We measured a suite of traits commonly measured in the functional trait ecology literature to explore how phenotype is influenced by bacterial strains within the pre-existing functional trait framework. Plant functional trait ecology has largely ignored microbiome contribution to plant phenotype. Bacterial strains belonging to the *Pseudomonas fluorescens* group are common plant growth promoting bacteria that are abundant in the *Populus* microbiome (see Gottel et al., 2011). *Pseudomonas fluorescens* accounted for approximately 34% of the sequences found in the *Populus* endosphere, but only 2-3% of the sequences in the rhizosphere and soil samples originating from the same roots (Gottel et al., 2011). *Pseudomonas* strains can alter plant host function by modifying plant growth (Kloepper et al., 1980; Lugtenberg & Kamilova, 2009; Timm et al., 2015), nutrient allocation (Bisht et al., 2009), hormone signaling (Stearns et al., 2012), up-regulating/down-regulating of gene expression pathways (Timm et al., 2016), and immune function (Verhagen et al., 2004; Weston et al., 2012). Additionally, the *Pseudomonas fluorescens* clade has a large amount of functional diversity (Jun et al., 2016), thus selecting two *Pseudomonas* strains allows us to explore how plant traits and overall phenotype respond to closely related bacterial strain genomes. To contrast with these two strains, we selected a distantly related, but enriched in *Populus* endosphere (Gottel et al., 2011), bacterial strain from the genus *Burkholderia*.

We predicted that aboveground and belowground traits of *Populus trichocarpa* would respond to *Burkholderia* and *Pseudomonas* strains and inoculation of different bacterial strains would result in different plant phenotypes. Further, we predicted that the two *Pseudomonas*

strains would produce a plant phenotype that was more similar to one another than to *Burkholderia* because of phylogenetic relatedness, *i.e.* more shared functionality. To test our predictions, we first conducted a genomic comparison using COG (clusters of orthologous groups) database to predict the functional differences among strains. Next, we inoculated each bacterial strain on *Populus trichocarpa* and measured a suite of physiological and architectural plant traits including the root:shoot, biomass production, root and leaf growth rates, chlorophyll content, net photosynthesis, and net photosynthesis at saturating light -  $A_{\text{sat}}$ , and saturating  $\text{CO}_2$  -  $A_{\text{max}}$ . We chose to measure overall trait response to bacterial endophytes without measuring the pathways involved because we were interested in understanding down-stream consequences of bacterial inoculation on overall plant phenotype.

## Materials and Methods

*Populus trichocarpa* genotype “93-968” (Labbe et al., 2014) was propagated in tissue culture following standard procedures (see Kang et al., 2009). Briefly, *in vitro* cultures were established from actively growing shoot tips collected from greenhouse-grown *Populus* plants. We sterilized shoot tips by soaking fresh cut tips in a 1% Tween 20 solution for 5 min, 70% Ethanol solution for 1 min, a 0.525% sodium hypochlorite solution for 15 min and then rinsed them three times in sterile  $\text{H}_2\text{O}$  for 5 min. Shoot tips were trimmed to 2 cm in length and transferred to a magenta box (Sigma-Aldrich, St. Louis, MO) containing 80 mL of tissue media (1× Murashige & Skoog (MS) basal medium (Murashige & Skoog, 1962) supplemented with MS vitamins (Caisson Labs, North Logan, UT, USA), 0.05% 2-(*N*-morpholino) ethanesulfonic acid (MES hydrate) (Sigma-Aldrich, St. Louis, MO, USA), 3% sucrose, 0.1% PPM™ (plant protective mixture) (Plant Cell Technology, Washington, DC, USA), 0.5% activated charcoal (Sigma-Aldrich, St. Louis, MO,

USA), and 0.15% Gelzan (Plantmedia, bioWORLD, Dublin, OH, USA). Plants were sub-cultured until it was determined, using microscopy and colony formation units with R2A medium, that the plants were axenic.

Plant cultures were rooted in a growth room at 25 °C under a 16 h photoperiod. After root establishment, plants that were similar in size and developmental stage were selected for experimentation. Plants were weighed and scanned to account for initial plant size differences among treatments. To ensure sterility during scanning, plants were placed between two (21.59 x 27.94 cm) sheets of cellulose acetate that were sprayed with 100% ethanol. Scans were performed with a portable scanner (VuPoint Solutions Inc., City of Industry, CA, USA) at 600 × 600 dpi. Scanned images were analyzed in WinRhizo (Regent Instruments, Quebec City, Canada) to determine initial root surface area, root length, stem length, and leaf surface area. After scanning, plants were transferred into experimental microcosms.

## Experimental design

We constructed closed microcosms by interlocking two sterile Magenta boxes (Sigma-Aldrich, St. Louis, MO, USA) with a coupler (Sigma-Aldrich, St. Louis, MO, USA). We added 150 ml calcined clay (Pro's choice Sports Field Products, Chicago, IL, USA) and 70 ml of 1× Hoagland's nutrient solution (Sigma-Aldrich, St. Louis, MO, USA) to each microcosm. We drilled two 7 mm holes on adjacent sides of the upper magenta box and covered the holes with adhesive microfiltration discs (Tissue Quick Plant Laboratories, Hampshire, United Kingdom) to allow air to flow into and out of the microcosms and to prevent outside microbial contamination. Prior to microbial addition, we double sterilized each closed microcosm by autoclaving on a 60 m dry cycle on consecutive days. *Pseudomonas fluorescens* strains (GM30 and GM41) and

*Burkholderia* sp. (BT03), hereafter termed *Pseudomonas* GM30, *Pseudomonas* GM41, and *Burkholderia* BT03 were isolated from *Populus deltoides* endospheres from east Tennessee and western North Carolina, USA (originally described in Brown et al., 2012). For full isolate descriptions, see Brown et al., 2012; Weston et al., 2012; Utturkar et al., 2014; Timm et al., 2015; 2016. We selected these three strains because previous work (*Pseudomonas* GM30 – Weston et al., 2012, Labbe et al., 2014; *Pseudomonas* GM41 – Labbe et al., 2014, Timm et al., 2016; *Burkholderia* Bt03 – Timm et al., 2016) had given us indication that strains were able to influence traits in *Arabidopsis thaliana* (Weston et al., 2012), were able to manipulate plant gene expression and hormonal signaling in *P. deltoides* (Timm et al., 2015; 2016), and were able to influence host interactions with mycorrhizal symbionts (Labbe et al., 2014). Although strains were isolated from *P. deltoides*, strains from *Pseudomonas* and *Burkholderia* readily colonize natural *P. trichocarpa* tissues (Moore et al., 2006; Xin et al., 2009; Knoth et al., 2014; Kahn et al., 2014; Doty et al., 2016). We grew bacterial strains in isolation and at a constant temperature, 25 °C, in 5 ml of R2A medium. After growing overnight they were pelleted and re-suspended in sterile water to an OD600 of 0.01 ( $\sim 1.0 \times 10^7$  cells ml<sup>-1</sup>).

We inoculated each microcosm by adding 10 ml of the bacterial strain ( $10^7$  cells ml<sup>-1</sup>) to the calcined clay substrate and stirring for 30 s to distribute the bacteria. After inoculation, we planted the *Populus* clones within each microcosm. Each *Populus* was grown in an individual microcosm in combination with one of the bacterial strains. Thus, the experiment had four treatment combinations – *Pseudomonas* GM30 inoculation, *Pseudomonas* GM41 inoculation, *Burkholderia* BT03 inoculation, and a bacteria-free control. In total, there were 32 microcosms with four treatments (n = 8). The experiment was divided into three different establishment dates in 2014 (1 March, 3 replicated blocks; 25 March, 2 replicated blocks; and 2 April, 3 replicated



blocks) because microbiome-free plant tissues were difficult to propagate. Plant-bacteria combinations were grown in the microcosms for five weeks with a 16 hr photoperiod, at 21°C and 80% relative humidity.

After 35 days of growth, plants were removed from microcosms, submerged in sterilized deionized H<sub>2</sub>O to remove clay from the root system, weighed, and scanned. Scans were analyzed with WinRhizo to determine final root surface area, total root length, stem length, and leaf surface area. For each plant, the final measurement of root surface area, total root length, stem length, and leaf surface area was subtracted from the initial measurement and divided by the experiment duration to determine tissue growth rates (cm d<sup>-1</sup> or cm<sup>2</sup> d<sup>-1</sup>). Additionally, each plant was dried for 48 hours at 70°C and weighed to measure leaf, shoot (leaf + stem) and root and total dry mass. Specific leaf area and the specific root length of each individual were calculated by dividing leaf area by leaf dry mass or by dividing root length by root dry mass, respectively.

To measure host physiological response to different bacterial strains, leaf gas-exchange was measured and used to estimate leaf photosynthesis on our first replicate block (March 1, n = 3). For each plant, gas exchange of the largest leaf of the plant was measured (Li-Cor model 6400, Li-Cor Biosciences, Lincoln, Nebraska, USA) immediately prior to our experimental harvest. The maximum rate of photosynthesis in saturating light under ambient CO<sub>2</sub> (A<sub>sat</sub>), the maximum rate of photosynthesis in saturating light and saturating CO<sub>2</sub> (A<sub>max</sub>), , and the quantum yield of CO<sub>2</sub> fixation (Φ) were all measured. Finally, average leaf chlorophyll content was measured on three fully opened leaves (Konica Minolta Chlorophyll Meter SPAD-S02, Ramsey, NJ, USA).

## Comparative genomics of microbes

Genomes of *Pseudomonas* GM30 and GM41 and *Burkholderia* BT03 were sequenced at Oak Ridge National Laboratory and genes were identified using Prodigal (Brown et al., 2012, Utturkar et al., 2014) and are available at NCBI (GM41: AKJN000000000.2; GM30: AKJP02000000.2; BT03: NZ\_AKKD000000000.2). Genome annotation, genomes statistics, and annotation comparisons were performed using IMG tools (img.jgi.doe.gov). Genome statistics and COG functional predictions were extracted from Integrated Microbial Genomes (img.jgi.doe.gov) and then they were compared manually for differential inclusion of predicted functions.

# **Bacterial Colonization**

To test for endophytic colonization of *Pseudomonas* GM41, *Pseudomonas* GM30, and *Burkholderia* BT03, we planted cuttings of *P. trichocarpa* into a magenta box using similar methodology and treatments described above (n=3). After 2 weeks of growth, all the plant roots, stems, and 1-2 mature leaves were surface sterilized by dipping them in a ~10% bleach solution, followed by 70% ethanol, and then rinsing in water three times. We recorded wet weight of plant tissues and then separately macerated each plant tissue compartment in a sterile mortar and pestle in 1 ml sterile 1× PBS. We transferred macerated plant tissues to a 24-well plate where we serially diluted each sample by 10% with 1×PBS at 1×, 0.1×, 0.01× of original sample concentration. Each sample was streaked onto R2A media plates and allowed to grow for 48 hours at 20°C. After 48 hours, colony formation was counted. We calculated CFU mg<sup>-1</sup> of plant tissue by multiplying colony number per plate by 10<sup>(dilution factor + 1)</sup> and then dividing that number by the dry tissue mass (mg<sup>1</sup>).

# 198 Statistical analyses

199 We tested all data for normality using the *normalTest* function in the fBasics package (version  
200 3011.87, R metrics core team 2014) for R version 3.0.2 (R development core team, 2013) and  
201 RStudio version 0.98.495 (RStudio, 2013). If data were not normally distributed, we performed  
202 log transformations or square-root transformations to satisfy the normality assumptions of  
203 ANOVA.

204 To explore plant trait response (root dry mass, leaf dry mass, shoot dry mass, total dry  
205 mass, root: shoot, root growth rates, leaf growth rates, change in leaf number, specific root  
206 length, specific leaf area) to bacterial strains, we used linear mixed-effect models using the lme4  
207 package in R (Bates et al., 2014). Bacterial strain was a fixed effect in the model and  
208 experimental block (three establishment dates) was a random factor. For plant dry mass  
209 measures, we incorporated initial measurements of root surface area in the root dry mass model  
210 and initial leaf surface area in the aboveground dry mass model as covariates. To test for  
211 significance of bacterial strain (fixed effects) and covariate (initial growth measure) we  
212 performed a likelihood ratio test to compare models with and without fixed effects and  
213 covariates. If including fixed factors (bacterial strain) was significant an improvement to model  
214 fit ( $p$  value  $< 0.05$  in likelihood ratio test), we calculated least square means and confidence  
215 intervals using the *diffLsmeans* function to calculate differences among strains using the lmerTest  
216 package version 2.0-3 (Kuznetsova, Brockhoff, & Christensen, 2014). We measured host  
217 response to bacterial inoculation by calculating the percent change in trait values ((mean trait  
218 value for *Populus* inoculated with bacterial strain – mean non-inoculated trait value)  $\times 100$ ).

219 To test physiological responses (carboxylase activity,  $A_{\max}$ ,  $A_{\text{sat}}$ ) of plant hosts to  
220 bacterial inoculation, we used one-way analysis of variance (ANOVA) using the *Anova* function

in the CAR package, (version 2.0-22, Fox & Weisberg 2011) because we collected physiology data on only a single sampling date ( $n = 3$ ). All raw data and R code is available in Supplemental Table 2 and 3, respectively.

## Results

### Bacterial strains differ in genomic content

We compared the genomes of *Burkholderia* BT03 and *Pseudomonas* GM30 and GM41 based on predicted enzyme functions using the COG database (Table 1). Overall, our genome comparison demonstrated that the bacterial strains differed in genome size and functional gene content. *Burkholderia* BT03 had a relatively large genome (10.9 Mb) compared to *Pseudomonas* GM30 (6.1 Mb) and *Pseudomonas* GM41 (6.6 Mb) (Table 1). We found all three bacterial strains shared functions that were likely critical for establishment and survival in the plant microbiome including the production of the plant hormone auxin, pili, flagella, chemotaxis, increased signal transduction, and secretion systems. However, we found many functional differences among our strains. The genome of *Burkholderia* encoded multiple pathways predicted to be involved in the metabolism of the plant hormones, salicylate and ethylene (Table 1). Relative to the *Pseudomonas* genomes, the *Burkholderia* genome encoded for numerous secondary metabolite biosynthesis pathways and more carbohydrate and lipid transporters, suggesting increased metabolic capabilities within *Burkholderia* (Table 1).

Even though *Pseudomonas* GM30 and *Pseudomonas* GM41 were classified as the same 16S OTU, their genome size differed as did the predicted functional capabilities of the two strains. The genome of *Pseudomonas* GM41 encoded for phosphorus solubilization and nitrate reduction, which were lacking in the *Pseudomonas* GM30 genome. Additionally, *Pseudomonas*

GM41 contained more secondary metabolite biosynthesis elements compared to *Pseudomonas* GM30. We also found that the genome of *Pseudomonas* GM41 contained more genes coding for carbohydrate metabolism, lipid metabolism, and amino acid transport and metabolism, energy production and conversion, suggesting that *Pseudomonas* GM41 may contain more metabolic breadth than *Pseudomonas* GM30 (Table 1). Taken together, our results demonstrated that these three bacterial strains differ in genome size and their functional gene content.

### **Bacterial colonization of *Populus* root tissue**

All three of the bacterial strains colonized *Populus* hosts. Colony-forming units were enriched in all three bacterial strains relative to the control in the 0.1× and 0.01× dilutions (0.1× dilution  $F = 18.77$ ,  $p < 0.0001$ ; 0.01× dilution  $F = 13.78$ ,  $p < 0.0001$ , Table 2), although CFU number was variable across dilutions, tissue types, and bacterial strain. However, we found no difference in CFUs among non-inoculated control and *Pseudomonas* GM30, GM41, and *Burkholderia* BT03 inoculated host plants at the 1× dilution ( $F = 1.24$ ,  $p = 0.319$  Table 2). Across nearly all tissue types, we found that *Pseudomonas* GM30, *Pseudomonas* GM41, and *Burkholderia* BT03 inoculated plants had 10-10000× more CFUs than did non-inoculated control plants (Table 2). All three bacterial strains colonized leaf and stem tissues, but the highest CFUs across bacterial treatments were consistently observed in roots (Table 2). Inoculated host plants contained 0-28809015 CFU mg<sup>-1</sup> in roots, 0-1166273 CFU mg<sup>-1</sup> in stems, and 0-73537 CFU mg<sup>-1</sup> in leaves compared to 0-400 CFU mg<sup>-1</sup> in root tissues, 0 CFU mg<sup>-1</sup> in stem tissue, 0-1000 CFU mg<sup>-1</sup> in leaf tissue compared to non-inoculated control plants (Table 2).

### **Plant structure is modified by bacterial inoculation**

Overall, we found that plant trait response to bacterial endophytes was strain specific. Specifically, mean root growth rate increased 184% with *Pseudomonas* GM30 colonization ( $t = 3.84, p = 0.001$ ), however root growth rates were unaffected by *Pseudomonas* GM41 ( $t = 1.61, p = 0.12$ ), and *Burkholderia* BT03 ( $t = 1.18, p = 0.25$ ) inoculation (Fig 1, Supplemental Table 1). Similarly, mean leaf growth rate increased 114% and 138% with *Pseudomonas* GM30 ( $t = 2.27, p = 0.03$ ) and *Pseudomonas* GM41 ( $t = 2.86, p = 0.01$ ) inoculation, but leaf growth rate was unaffected by *Burkholderia* inoculation ( $t = 1.02, p = 0.32$ ) (Fig 1, Supplemental Table 1). Inoculation by *Pseudomonas* GM30 increased leaf number by 36% ( $t = 3.34, p = 0.003$ ) but leaf number was unaffected by *Pseudomonas* GM41 ( $t = 0.93, p = 0.36$ ) and *Burkholderia* BT03 ( $t = 1.418, p = 0.17$ ) inoculation (Fig 1). We observed no differences in stem elongation with bacterial inoculation (chisq= 0.06,  $p = 0.97$ , Supplemental Table 1).

Interestingly, we observed no differences in total plant dry mass (chisq = 3.27,  $p = 0.195$ , Fig 2), root dry mass (chisq = 0.00,  $p = 1.00$ , Fig 2), root:shoot ratio (chisq=0.00,  $p = 1.00$ , Supplemental Table 1) or plant height (chisq = 1.99,  $p = 0.158$  Supplemental Table 1) with bacterial inoculation. However, *Pseudomonas* GM30 inoculation increased leaf dry biomass by 86% ( $t = 2.43, p = 0.02$ ) relative to control plants, however leaf biomass was unaffected by *Pseudomonas* GM41 ( $t = 0.97, p = 0.33$ ) and *Burkholderia* BT03 ( $t = 1.70, p = 0.10$ ) (Fig 2, Supplemental Table 1). We observed no differences in specific leaf area with bacterial inoculation (chisq= 2.60,  $p = 0.46$ , Supplemental Table 1). Thus, inoculation of *Pseudomonas* GM30 increased leaf surface area ( $t = 2.27, p = 0.03$ ) and aboveground dry mass ( $t = 2.43, p = 0.02$ ), without changing leaf area:mass ratios. We found no significant differences in root length:dry mass (specific root length, chisq = 1.06,  $p = 0.79$ ) with bacterial inoculation

(Supplemental Table 1). Our results indicate that bacterial strains modify plant resource allocation but not total dry mass production.

## **Plant physiology is not affected by bacterial inoculation**

Bacterial inoculation had no measureable effects on any physiological trait we measured: chlorophyll content (SPAD) ( $\chi^2 = 2.15$ ,  $p = 0.54$ ), quantum yield of photosynthesis ( $\Phi$ ) ( $F = 1.01$ ,  $p = 0.43$ ), net photosynthesis at saturating light conditions ( $A_{\text{sat}}$ ) ( $F = 0.76$ ,  $p = 0.55$ ) or maximum net photosynthesis at saturating light and  $[\text{CO}_2]$  ( $A_{\text{max}}$ ) ( $F = 1.98$ ,  $p = 0.19$ ) (Fig 3). In agreement with the total dry mass data, we did not observe significant changes in the measured photosynthetic parameters. Thus, changes in plant structure were not linked with increased photosynthetic capacity, efficiency, or carbon assimilation rates.

## **Discussion**

The plant root microbiome can have a strong influence on plant production and phenotype (Friesen, 2013; Vandenkoornhuyse, et al., 2015); yet, less is known about how plant trait expression, production, and physiology are influenced by individual endophytic strains. We explored how plant morphological traits, productivity, and cellular physiology in *Populus trichocarpa* responded to inoculation with three bacterial strains, two closely related *Pseudomonas fluorescens* strains (GM30 & GM41) and a more distantly related *Burkholderia* strain (BT03). We selected bacterial strains that were predicted to differ in metabolic capabilities, plant hormone production and metabolism, and secondary metabolite synthesis in an effort to understand how plant phenotype is influenced by inoculation with different strains of common endophytic bacteria (Table 1, Timm et al., 2015; 2016). Our comparative genomic

analysis revealed that while all three strains share many common endophytic functions like plant hormone signal disruption, production of plant hormone auxin, pili, flagella, and chemotaxis, strains potentially differed in their ability to perform these functions. Overall, we found that *Burkholderia* and *Pseudomonas* genomes differed in the carbon substrates they were predicted to degrade, plant hormone production and metabolism, and secondary metabolite synthesis, which led us to predict that plant response to bacterial inoculation would lead to different phenotypes between treatments. All three strains could colonize *Populus* roots, leaves, and stems, however CFU number was highest within root tissues in all three strains (Table 2).

Overall, we found root endophyte inoculation altered plant resource allocation patterns without influencing total plant biomass accumulation (Fig 1). Additionally, we found that plant trait response and overall phenotype differed across bacterial strains in ways that would not have been predicted from our genome analysis. Specifically, *Burkholderia* BT03 was predicted to produce auxin and to metabolize salicylate and ethylene, three plant hormones crucial to plant growth and development (see Yang & Hoffman, 1984; Wasternack & Parthier, 1997; Chen et al., 2009; Dempsey et al., 2011). Additionally, we found the *Burkholderia* genome encoded for multiple transposase elements that degrade poplar-produced aromatics and metabolites (Timm et al., 2015; 2016). Despite the predicted ability of *Burkholderia* to manipulate multiple plant hormonal and signaling pathways, we observed no measurable changes in any traits when *Populus* was inoculated with *Burkholderia* (Figs 1, 2, 3). This was especially surprising since we consistently measured the highest CFU abundance within *Burkholderia* inoculated individuals (Table 2).

In spite of close genetic relatedness and classification under the same 16S OTU profile, our *Pseudomonas* strains differed in key functional capabilities. Specifically, *Pseudomonas*



GM41 encoded for phosphate solubilization and denitrification ability, suggesting these two strains may differentially influence host nutrition, although this remains untested. Our genome analysis revealed that both strains were capable of producing the plant hormone auxin, however another study found that *Pseudomonas* GM41 produced two times more auxin than *Pseudomonas* GM30 (Timm et al., 2015). Auxin synthesis by endophytic bacteria can increase root branching and lateral root formation and decrease overall plant height, leaf number, chlorophyll content and photosynthetic efficiency (Romano, Cooper, & Klee, 1993; Fu & Harberd, 2003; Weston et al., 2012). Thus, we predicted that *Pseudomonas* GM41 would have a strong influence on plant root traits, however we observed no measurable effects of *Pseudomonas* GM41 inoculation on root growth rate or morphology (Fig 1, Supplemental Table 1). Belowground, *Pseudomonas* GM30 inoculation increased root surface area growth rate by 184% (Fig 1) without increasing root biomass (Fig 2), suggesting *Pseudomonas* GM30 inoculation may change root morphology, leading to longer, thinner, highly-branched roots with similar biomass, as we predicted. Similar patterns have been observed when *Pseudomonas* GM30 is inoculated on both *Arabidopsis* (Weston et al., 2012) and *Populus deltoides* (Timm et al., 2015; 2016). Additionally, inoculation of *Pseudomonas* GM30 increased leaf surface area growth rate by 114% (Fig 1), leaf number by 36% (Fig 1), and aboveground biomass by 86% (Fig 2) but did not influence specific leaf area (Supplemental Table 1), whereas closely-related *Pseudomonas* GM41 increased leaf surface area growth rate by 138% (Fig 1) but did not change leaf number (Fig 1) or aboveground biomass (Fig 2). Unlike *Burkholderia*, *Pseudomonas* genomes do not contain the genes to directly metabolize salicylate, however inoculation of *Pseudomonas* GM41 can up-regulate salicylic acid synthesis and degradation in *Populus* (Timm et al., 2016). Taken together, our data suggest that predicting plant phenotypic response to

bacterial inoculation, even in overly simplified systems using fully sequenced bacterial strains, is extremely complex and difficult.

Contrary to our predictions, leaf physiology (Fig 3), plant height (Supplemental Table 1), root:shoot (Supplemental table 1), specific leaf area (Supplemental table 1), specific root length (Supplemental table 1), and total plant dry mass (Fig 2) were not influenced by bacterial inoculation. It is possible that multiple, overlapping plant signaling and gene expression effects induced by bacterial endophyte inoculation may mask a hosts' phenotype response. For example, endophytes simultaneously up- and down-regulate numerous genes and metabolites in plant tissue (see Verhagen et al., 2004; Wang et al., 2005; Walker et al., 2011; Weston et al., 2012; Timm et al., 2016). Thus, counteracting influences among different gene pathways may conceal plant responses to endophyte inoculation when measuring down-stream phenotype and functional traits (Bashan, Holguin, & de-Bashan 2004; Timm et al., 2016). Additionally, host physiological response to endophyte inoculation may vary with bacterial strain (Kandasamy et al., 2009; Weston et al., 2012; Timm et al., 2016), plant host (Smith and Goodman 1999), plant ontogeny (Siddiqui & Shaukat 2003), or plant stress (Dimkpa, Weinand, & Asch 2009; Yang, Kloepper, & Ryu 2009; Lau & Lennon 2012). For example, Root colonization by *Pseudomonas* can reduce chlorophyll content and net photosynthesis ( $A_{sat}$ ) in a variety of plant hosts (Zou et al., 2005; Weston et al., 2012). However *Pseudomonas* colonization can also increase photosynthetic activity and chlorophyll content (Kandasamy et al., 2009, Timm et al., 2016). Thus, biotic and abiotic contexts may drive the phenotypic response of hosts to endophyte inoculation, however this idea requires further testing.

Our study focused on the response of plant functional traits to monoculture associations of common endosphere bacteria, however future studies should focus on exploring how plant

phenotype responds to diverse microbiome communities. With a few well-known exceptions (Tan & Tan 1986; Harris, Pacovsky, & Paul 1985; Ma et al., 2003, Lau & Lennon 2011; 2012), bacterial community composition in roots has been ignored in studies exploring what drives natural variation in plant traits (Friesen et al., 2011; Friesen 2013; Timm et al., 2016). We propose a multifaceted approach to investigate linkages among the plant microbiome and natural plant trait variation. First, incorporation of microbiome composition into studies that currently investigate host identity/genotype and environmental parameters may be important for finding patterns in natural trait variation – especially when conducted across a variety of environmental gradients. Second, once correlations between microbiomes and plant traits are observed in the field, detailed work constructing communities in the lab and greenhouse would enable a mechanistic understanding of what is underlying the observed patterns. These studies could be especially fruitful when conducted across natural biotic and abiotic environmental gradients in the laboratory, greenhouse, and field settings (Classen et al., 2015).

## Conclusions

Our study demonstrates that bacteria living in plant roots can influence plant morphological traits. Increasingly, ecologists are using plant functional traits to explore how changing environments might alter plant function (Wright et al., 2004; Reich 2014). Plant traits, such as specific leaf area and specific root length, are often significantly correlated with important plant functions such as carbon fixation and nutrient uptake (Diaz & Cabido 2001). Researchers are using correlations between plant traits and function to extrapolate how plants and ecosystems will respond to global changes (Reich et al., 1999; Wright et al., 2004; Reich 2014). While interactions between plant genotype and environment undoubtedly influence plant phenotypic

plasticity (Bradshaw 1965; Schlichting 1986; Sultan 2000; Des Marais, Hernandez, & Juenger 2013), phenotype is also heavily influenced by biotic factors, like the microbiome bacterial endophytes (Lau & Lennon 2011; 2012; Wagner et al., 2014; Hacquard & Schadt 2015). Given that plant-microbial studies, including ours, have observed strong linkages between microbiome and plant phenotype (reviewed in Friesen et al., 2011; Friesen 2013) interactions among global change drivers, plant genotypes, and plant microbiomes, should be considered in trait-based approaches to ecological questions (Classen et al., 2015).

# **Acknowledgements**

We thank Lee Gunter, Tse-Yuan Lu, Kelsey Carter, Jesse Labbe, and W. Nathan Cude for their assistance in data collection and plant propagation. The Classen Ecosystem Ecology group and three reviewers provided valuable input on earlier reviews of this manuscript.

# References

- Bashan Y, Holguin G, de-Bashan LE. 2004. *Azospirillum*-plant relationships: physiological, molecular, agricultural, and environmental advances (1997-2003). *Canadian Journal of Microbiology*. 50:521-577.
- Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*. 67:1-48.
- Bisht R, Chaturvedi S, Srivastava R, Sharma A, Johri B. 2009. Effect of arbuscular mycorrhizal fungi, *Pseudomonas fluorescens* and *Rhizobium leguminosarum* on the growth and nutrient status of *Dalbergia sissoo* Roxb. *Tropical Ecology*. 50:231.
- Bradshaw AD. 1965. Evolutionary significance of phenotypic plasticity in plants. *Advances in Genetics*. 13:115-155.
- Brown SD, Utturkar SM, Klingeman DM, Johnson CM, Martin SL, Land ML, Lu TYS, Schadt CW, Doktycz MJ, Pelletier DA. 2012. Twenty-one genome sequences from *Pseudomonas* species and 19 genome sequences from diverse bacteria isolated from the rhizosphere and endosphere of *Populus deltoides*. *Journal of Bacteriology*. 194:5991-5993.
- Chen Z, Zheng Z, Huang J, Lai Z, Fan B. 2009. Biosynthesis of salicylic acid in plants. *Plant Signaling & Behavior*. 4:493-496.
- Classen AT, Sundqvist M, Henning JA, Newman GS, Moore JAM, Cregger M, Moorhead LC, Patterson CM. 2015. Direct and indirect effects of climate change on soil microbial and soil microbial-plant interactions: What lies ahead? *Ecosphere*. 6(8):130. doi: 10.1890/ES15-00217.1.
- Cornelissen JHC, Lavorel S, Garnier E, Diaz S, Buchmann N, Gurvich DE, Reich PB, ter Steege H, Morgan HD, van der Heijden MGA, Pausas JG, Poorter H. 2003. A handbook of protocols

441 for standardised and easy measurement of plant functional traits worldwide. *Australian*  
 442 *Journal of Botany*. 51:335-380.

443 Dempsey DMA, Vlot AC, Wildermuth MC, and Klessig DF. 2011. Salicylic Acid Biosynthesis  
 444 and Metabolism. *The Arabidopsis Book*. 9: e0156 doi: 10.1199/tab.0156.

445 Des Marais DL, Hernandez KM, Juenger TE. 2013. Genotype-by-environment interaction and  
 446 plasticity: exploring genomic responses of plants to the abiotic environment. *Annual Review*  
 447 *of Ecology, Evolution, and Systematics*. 44:5-29.

448 Díaz S, Cabido M. 2001. Vive la différence: plant functional diversity matters to ecosystem  
 449 processes. *Trends in Ecology & Evolution*. 16:646-655.

450 Dimkpa C, Weinand T, Asch F. 2009. Plant–rhizobacteria interactions alleviate abiotic stress  
 451 conditions. *Plant, cell & Environment*. 32:1682-1694.

452 Doty SL, Sher AW, Fleck ND, Khorasani M, Bumgarner RE, Khan Z, Ko AWK, Kim S-H,  
 453 DeLuca TH. 2016. Variable Nitrogen Fixation in Wild *Populus*. *PLoS ONE*.  
 454 doi:11:e0155979.

455 Drogue B, Sanguin H, Chamam A, Mozar M, Llauro C, Panaud O, Prigent-Combaret C, Picault  
 456 N, Wisniewski-Dyé F. 2014. Plant root transcriptome profiling reveals a strain-dependent  
 457 response during *Azospirillum*-rice cooperation. *Frontiers in Plant Science*. 5:607  
 458 doi:10.3389/fpls.2014.00607.

459 Fox J, Weisberg S. 2011. *An {R} Companion to Applied Regression*, Second Edition. Thousand  
 460 Oaks CA: Sage.

461 Friesen ML. 2013. *Microbially Mediated plant functional traits molecular microbial ecology of*  
 462 *the rhizosphere*. John Wiley and Sons, Inc., pp 87-102.

463 Friesen ML, Porter SS, Stark SC, von Wettberg EJ, Sachs JL, Martinez-Romero E. 2011.  
464 Microbially mediated plant functional traits. *Annual Review of Ecology, Evolution, and*  
465 *Systematics*. 42:23-46.

466 Fu X, Harberd NP. 2003. Auxin promotes Arabidopsis root growth by modulating gibberellin  
467 response. *Nature*. 421:740-743.

468 Gottel NR, Castro HF, Kerley M, Yang Z, Pelletier DA, Podar M, Karpinets T, Uberbacher E,  
469 Tuskan GA, Vilgalys R, Doktycz MJ, Schadt CW. 2011. Distinct microbial communities  
470 within the endosphere and rhizosphere of *Populus deltoides* roots across contrasting soil  
471 types. *Applied and Environmental Microbiology*. 77:5934-5944.

472 Hacquard S, Schadt CW. 2015. Towards a holistic understanding of the beneficial interactions  
473 across the *Populus* microbiome. *New Phytologist*. 205:1424-1430.

474 Harris D, Pacovsky RS, Paul EA. 1985. Carbon economy of soybean-Rhizobium-Glomus  
475 associations. *New Phytologist*. 101:427.

476 Jun SR, Wassenaar TM, Nookaew I, Hauser L, Wanchai V, Land M, Timm CM, Lu TYS, Schadt  
477 CW, Doktycz MJ, Pelletier DA, Ussery DW. 2016. Diversity of *Pseudomonas* Genomes,  
478 Including *Populus*-Associated Isolates, as Revealed by Comparative Genome Analysis.  
479 *Applied and Environmental Microbiology*. 82:375-383.

480 Kang B, Osburn L, Kopsell D, Tuskan G, Cheng ZM. 2009. Micropropagation of *Populus*  
481 *trichocarpa* ‘Nisqually-1’: the genotype deriving the *Populus* reference genome. *Plant Cell,*  
482 *Tissue and Organ Culture*. 99:251-257.

483 Khan Z, Roman D, Kintz T, delas Alas M, Yap R, Doty S. 2014. Degradation, phytoprotection  
484 and phytoremediation of phenanthrene by endophyte *Pseudomonas putida*, PD1.  
485 *Environmental science & technology*. 48:12221-12228.

486 Kloepper JW, Leong J, Teintze M, Schroth MN. 1980 Enhanced plant growth by siderophores  
487 produced by plant growth-promoting rhizobacteria. *Nature*. 286:885-886.

488 Knoth JL, Kim S-H, Ettl GJ, Doty SL. 2014. Biological nitrogen fixation and biomass  
489 accumulation within poplar clones as a result of inoculations with diazotrophic endophyte  
490 consortia. *New Phytologist*. 201:599-609.

491 Kandasamy S, Loganathan K, Muthuraj R, Duraisamy S, Seetharaman S, Thiruvengadam R,  
492 Ponnusamy B, Ramasamy S. 2009. Understanding the molecular basis of plant growth  
493 promotional effect of *Pseudomonas fluorescens* on rice through protein profiling. *Proteome*  
494 *Science* 7:1-8.

495 Kuznetsova A, Brockhoff P, Christensen R. 2014. *LmerTest: tests for random and fixed effects*  
496 *for linear mixed effect models*. R package, version 2.0-3.

497 Labbé JL, Weston DJ, Dunkirk N, Pelletier DA, Tuskan GA. 2014. Newly identified helper  
498 bacteria stimulate ectomycorrhizal formation in *Populus*. *Frontiers of Plant Science*. 5:579  
499 doi: 10.3389/fpls.2014.00579.

500 Lau JA, Lennon JT. 2011. Evolutionary ecology of plant-microbe interactions: soil microbial  
501 structure alters selection on plant traits. *New Phytologist*. 192:215-224.

502 Lau JA, Lennon JT. 2012. Rapid responses of soil microorganisms improve plant fitness in novel  
503 environments. *Proceedings of the National Academy of Sciences*. 109:14058-62.

504 Lebeis SL. 2015. Greater than the sum of their parts: characterizing plant microbiomes at the  
505 community-level. *Current Opinion in Plant Biology*. 24:82-86.

506 Lugtenberg B, Kamilova F. 2009. Plant-Growth-Promoting Rhizobacteria. *Annual Review of*  
507 *Microbiology*. 63:541-556.



508 Ma W, Sebestianova S, Sebestian J, Burd G, Guinel F, Glick B. 2003. Prevalence of 1-  
509 aminocyclopropane-1-carboxylate deaminase in *Rhizobium* spp. *Antonie van Leeuwenhoek*.  
510 83:285-291.

511 Moore FP, Barac T, Borremans B, Oeyen L, Vangronsveld J, van der Lelie D, Campbell CD,  
512 Moore ERB. 2006. Endophytic bacterial diversity in poplar trees growing on a BTEX-  
513 contaminated site: The characterisation of isolates with potential to enhance  
514 phytoremediation. *Systematic and Applied Microbiology*. 29:539-556.

515 Murashige T, Skoog F. 1962. A revised medium for rapid growth and bio assays with tobacco  
516 tissue cultures. *Physiologia Plantarum*. 15:473-497.

517 Reich PB. 2014. The world-wide ‘fast–slow’ plant economics spectrum: a traits manifesto.  
518 *Journal of Ecology*. 102:275-301.

519 Reich PB, Ellsworth DS, Walters MB, Vose JM, Gresham C, Volin JC, Bowman WD. 1999.  
520 Generality of leaf trait relationships: a test across six biomes. *Ecology*. 80:1955-1969.

521 Romano CP, Cooper ML, Klee HJ. 1993. Uncoupling auxin and ethylene effects in transgenic  
522 tobacco and arabidopsis plants. *Plant Cell*. 5:181-189.

523 Schlichting CD. 1986. The evolution of phenotypic plasticity in plants. *Annual Review of*  
524 *Ecology, Evolution, and Systematics*. 17:667-693.

525 Siddiqui IA, Shaukat SS. 2003. Plant species, host age and host genotype effects on *Meloidogyne*  
526 *incognita* biocontrol by *Pseudomonas fluorescens* strain CHA0 and its genetically-modified  
527 derivatives. *Journal of Phytopathology*. 151:231-238.

528 Smith KP, Goodman RM. 1999. Host variation for interactions with beneficial plant-associated  
529 microbes. *Annual Review of Phytopathology*. 37:473-491.

530 Stearns JC, Woody OZ, McConkey BJ, Glick BR. 2012. Effects of bacterial ACC deaminase on  
531 *Brassica napus* gene expression. *Molecular Plant Microbe Interactions*. 25:668-676 doi:  
532 10.1094/MPMI-08-11-0213.

533 Sultan SE. 2000. Phenotypic plasticity for plant development, function and life history. *Trends in*  
534 *Plant Science*. 5:537-542.

535 Tan G, Tan W. 1986. Interaction between alfalfa cultivars and *Rhizobium* strains for nitrogen  
536 fixation. *Theoretical and Applied Genetics*. 71:724-729.

537 Timm CM, Campbell AG, Utturkar, SM, Jun SR, Paraless RE, Tan W, Robeson MS, Lu TYS,  
538 Jawdy S, Brown SD, Ussery DW, Schadt CW, Tuskan GA, Doktycz MJ, Weston DJ, Pelletier  
539 DA. 2015. Metabolic functions of *Pseudomonas fluorescens* strains from *Populus deltoides*  
540 depend on rhizosphere or endosphere isolation compartment. *Frontiers in Microbiology*. 6  
541 doi: [10.3389/fmicb.2015.01118](https://doi.org/10.3389/fmicb.2015.01118).

542 Timm CM, Pelletier DA, Jawdy S, Gunter LE, Henning JA, Engle N, Aufrecht J, Gee E,  
543 Nookaew I, Yang Z, Lu TYS, Tschaplinski TJ, Doktycz MJ, Tuskan GA, Weston DJ. 2016.  
544 Two poplar-associated bacterial isolates induce additive favorable responses in a constructed  
545 plant-microbiome system. *Frontiers in Plant Science*. 7 doi: [10.3389/fpls.2016.00497](https://doi.org/10.3389/fpls.2016.00497).

546 Utturkar SM, Klingeman DM, Land ML, Schadt CW, Doktycz MJ, Pelletier DA, Brown SD.  
547 2014. Evaluation and validation of de novo and hybrid assembly techniques to derive high-  
548 quality genome sequences. *Bioinformatics*. 30:2709-2716.

549 Vandenkoornhuyse P, Quaiser A, Duhamel M, Le Van A, Dufresne A. 2015. The importance of  
550 the microbiome of the plant holobiont. *New Phytologist*. 206:1196-1206.

Verhagen BWM, Glazebrook J, Zhu T, Chang HS, van Loon LC, Pieterse CMJ. 2004. The transcriptome of rhizobacteria-induced systemic resistance in *Arabidopsis*. *Molecular Plant Microbe Interactions*. 17:895-908.

Wagner MR, Lundberg DS, Coleman-Derr D, Tringe SG, Dangl JL, Mitchell-Olds T. 2014. Natural soil microbes alter flowering phenology and the intensity of selection on flowering time in a wild *Arabidopsis* relative. *Ecology Letters*. 17:717-726.

Walker V, Bertrand C, Bellvert F, Moënne-Loccoz Y, Bally R, Comte G. 2011. Host plant secondary metabolite profiling shows a complex, strain-dependent response of maize to plant growth-promoting rhizobacteria of the genus *Azospirillum*. *New Phytologist*. 189:494-506.

Wang Y, Ohara Y, Nakayashiki H, Tosa Y, Mayama S. 2005. Microarray analysis of the gene expression profile induced by the endophytic plant growth-promoting rhizobacteria, *Pseudomonas fluorescens* FPT9601-T5 in *Arabidopsis*. *Molecular Plant Microbe Interactions*. 18:385-396.

Wasternack C, Parthier B. 1997. Jasmonate-signalled plant gene expression. *Trends in Plant Science*. 2: 302-307.

Weston DJ, Pelletier DA, Morrell-Falvey JL, Tschaplinski TJ, Jawdy SS, Lu TYS, Allen SM, Melton SJ, Martin MZ, Schadt CW, Karve AA, Chen JG, Yang X, Doktycz MJ, Tuskan GA. 2012. *Pseudomonas fluorescens* induces strain-dependent and strain-independent host plant responses in defense networks, primary metabolism, photosynthesis, and fitness. *Molecular Plant Microbe Interactions*. 25:765-778.

Wright IJ, Reich, PB, Westoby M, Ackerly DD, Baruch Z. 2004. The worldwide leaf economics spectrum. *Nature*. 428:821-827.

573 Xin G, Zhang G, Kang JW, Staley JT, Doty SL. 2009. A diazotrophic, indole-3-acetic acid-  
574 producing endophyte from wild cottonwood. *Biology and Fertility of Soils*. 45:669-674.

575 Yang J, Kloepper JW, Ryu C-M. 2009. Rhizosphere bacteria help plants tolerate abiotic stress.  
576 *Trends in Plant Science*. 14:1-4.

577 Yang SF, Hoffman NE. 1984. Ethylene biosynthesis and its regulation in higher plants. *Annual*  
578 *Review of Plant Physiology and Plant Molecular Biology*. 35:155-189.

579 Zamioudis C, Mastranesti P, Dhonukshe P, Blilou I, Pieterse CMJ. 2013. Unraveling Root  
580 Developmental Programs Initiated by Beneficial *Pseudomonas* spp. Bacteria. *Plant*  
581 *Physiology*. 162:304-318.

582 Zou J, Rodriguez-Zas S, Aldea M, L, M, Zhu J, Gonzalez DO, Vodkin LO, DeLucia E, Clough  
583 SJ. 2005. Expression Profiling Soybean Response to *Pseudomonas syringae* Reveals New  
584 Defense-Related Genes and Rapid HR-Specific Downregulation of Photosynthesis. *Molecular*  
585 *Plant Microbe Interactions*. 18:1161-1168.

# **Table 1**(on next page)

Predicted plant-interaction pathways in bacterial strains *Burkholderia* sp. BT03, *Pseudomonas fluorescens* GM30, and *Pseudomonas fluorescens* GM4.

Predicted plant-interaction pathways in bacterial strains *Burkholderia* sp. BT03, *Pseudomonas fluorescens* GM30, and *Pseudomonas fluorescens* GM41. Genome size, relevant pathways, and COG category statistics were identified using IMG tools. Where applicable, gene loci indicating predicted functions in genomes (individual genes or pathways) were included. NA = not applicable.

1

	<i>Burkholderia</i> BT03	<i>Pseudomonas</i> GM30	<i>Pseudomonas</i> GM41
Genome size (Mb)	10.9	6.1	6.6
ACC deaminase	PMI06_0002752	PMI25_02765	PMI27_01478
salicylate metabolism	PMI06_001931	NA	PMI27_05197
auxin biosynthesis	PMI06_005275	PMI25_03791	PMI27_00952
pili, fimbriae	PMI06_00372-3373	PMI25_00378-0372	NA
flagella	PMI06_009483-9498	PMI25_03624 -3649	PMI27_02843-2866
chemotaxis	PMI06_009463-9475	PMI25_05665-5658	PMI27_05395-5382
type 2 secretion system	PMI06_001352-1341	PMI25_00837-00844	NA
type 3 secretion system	PMI06_000607-0617	NA	NA
type 4 secretion system	PMI06_009642-9622	NA	NA
type 6 secretion system	PMI06_001813-1833	PMI25_012011220	PMI27_02378-2397
carbohydrate metabolism (# of genes)	582	222	291
secondary metabolite metabolism (# of genes)	337	113	148

2

## Table 2 (on next page)

Colony forming units found in leaf, root, and stem tissue of *Populus trichocarpa* genotypes inoculated with *Pseudomonas* GM30, *Pseudomonas* GM41, or *Burkholderia* BT03 across three dilution factors.

Colony forming units found in leaf, root, and stem tissue of *Populus trichocarpa* genotypes inoculated with *Pseudomonas* GM30, *Pseudomonas* GM41, or *Burkholderia* BT03 across three different dilution factors: 1×, 0.1×, 0.01× concentrations of the original sample.

*Pseudomonas* GM41 and *Burkholderia* BT03 data were first published in Timm et al. (2016).

1

Treatment	Tissue	Dilution	mean CFU	St dev		Sum Sq	Df	F	<i>p</i>
Control	leaf	1.0E+01	1080.5	1871.5	Bact.	1.3E+12	3	1.24	0.319
GM30	leaf	1.0E+01	19574.7	30672.7	Tissue	1.4E+12	2	1.92	0.1699
GM41	leaf	1.0E+01	1141.3	1809.3	B × T	1.6E+12	6	0.74	0.6264
BT03	leaf	1.0E+01	41175.9	45063.1	Resid.	8.1E+12	24		
Control	root	1.0E+01	110.2	131.5					
GM30	root	1.0E+01	170447.1	212977.7					
GM41	root	1.0E+01	2438.9	1563.8					
BT03	root	1.0E+01	309628.0	106958.6					
Control	stem	1.0E+01	0.0	0.0					
GM30	stem	1.0E+01	1166273.0	1872593.0					
GM41	stem	1.0E+01	1510.2	2135.8					
BT03	stem	1.0E+01	654513.2	688365.7					
Control	leaf	1.0E-01	1044.4	1809.0	Bact.	1.2E+13	3	18.77	<b>&gt;0.001</b>
GM30	leaf	1.0E-01	16643.8	28827.9	Tissue	3.8E+12	2	9.21	<b>0.001</b>
GM41	leaf	1.0E-01	566.2	980.7	B × T	1.1E+13	6	8.91	<b>&gt;0.001</b>
BT03	leaf	1.0E-01	60745.9	54910.0	Resid.	5.0E+12	24		
Control	root	1.0E-01	402.7	377.6					
GM30	root	1.0E-01	120591.5	111174.4					
GM41	root	1.0E-01	2851.9	3319.7					
BT03	root	1.0E-01	3096279.7	1069585.6					
Control	stem	1.0E-01	0.0	0.0					
GM30	stem	1.0E-01	289189.7	330089.7					
GM41	stem	1.0E-01	0.0	0.0					



BT03	stem	1.0E-01	904314.7	1099508.6					
Control	leaf	1.0E-02	0.0	0.0	Bact.	6.9E+14	3	13.78	>0.001
GM30	leaf	1.0E-02	0.0	0.0	Tissue	4.0E+14	2	11.79	>0.001
GM41	leaf	1.0E-02	0.0	0.0	B × T	1.2E+15	6	11.47	>0.001
BT03	leaf	1.0E-02	73537.1	80004.4	Resid.	4.1E+14	24		
Control	root	1.0E-02	0.0	0.0					
GM30	root	1.0E-02	368195.0	510398.0					
GM41	root	1.0E-02	20595.2	35671.9					
BT03	root	1.0E-02	28809015.5	14126689.6					
Control	stem	1.0E-02	0.0	0.0					
GM30	stem	1.0E-02	227127.9	252544.5					
GM41	stem	1.0E-02	0.0	0.0					
BT03	stem	1.0E-02	1805855.2	1567125.7					


2

3

# Figure 1

Structural traits of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) or were inoculated with *Pseudomonas* GM30, *Pseudomonas* GM4, or *Burkholderia* BT03.


Figure 1: Structural traits of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) (n = 8), were inoculated with *Pseudomonas* GM30 (n = 7), *Pseudomonas* GM41 (n = 8), or *Burkholderia* BT03 (n = 7). a) Change in leaf number from the first to last day of the experiment. Negative values indicate that leaves senesced during the experiment. b) Leaf surface area growth rates ( $\text{cm}^2 \text{d}^{-1}$ ), c) Stem growth rate ( $\text{cm}^1 \text{d}^{-1}$ ). d) Root surface area growth rates ( $\text{cm}^2 \text{d}^{-1}$ ). Letters represent significant differences of post-hoc least squares means among bacterial treatments. Boxplots display median, first and third quartiles, and vertical lines represent 1.5×inner quartile range of our dataset. The dots represent raw data values.



# Figure 2

Biomass allocation of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) or were inoculated with *Pseudomonas* GM30, *Pseudomonas* GM41, or *Burkholderia* BT03.

Figure 2: Biomass allocation of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) (n = 8) , were inoculated with *Pseudomonas* GM30 (n = 7), *Pseudomonas* GM41 ( n = 8), or *Burkholderia* BT03 (n = 7). a) Total dry mass (mg). b) Leaf biomass (mg), c) Root biomass (mg). Letters represent significant differences of post-hoc least squares means among bacterial treatments. Boxplots display median, first and third quartiles, and vertical lines represent 1.5×inner quartile range of our dataset. The dots represent raw data values.



# Figure 3

Physiology traits of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) or were inoculated with *Pseudomonas* GM30, *Pseudomonas* GM41, or *Burkholderia* BT03.

Figure 3: Physiology traits of *Populus trichocarpa* that were not inoculated with bacteria (no microbe control) (n = 8), were inoculated with *Pseudomonas* GM30 (n = 7), *Pseudomonas* GM41 (n = 8), or *Burkholderia* BT03 (n = 7). a) Plant chlorophyll content (SPAD), b)  $\Phi\text{CO}_2$  (expressed as the slope of carboxylase activity across different light levels), and c) carboxylase activity under maximum light level and  $\text{CO}_2$  concentration ( $A_{\text{max}}$ ). Letters represent significant differences of post-hoc least squares means among bacterial treatments. Boxplots display median, first and third quartiles, and vertical lines represent 1.5×inner quartile range of our dataset. The dots represent raw data values.

