

Pests, diseases and crop protection practices in the smallholder sweetpotato production system of the highlands of Papua New Guinea

Geoff M Gurr^{Corresp., 1,2,3}, Jian Liu^{1,4}, Anne C Johnson³, Deane N Woruba⁵, Gunnar Kirchhof⁶, Ryosuke Fujinuma⁶, William Sirabis⁷, Yapo Jeffery⁷, Ramakrishna Akkinapally⁸

¹ State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, Fujian Agriculture & Forestry University, Fuzhou, Fujian, China

² Institute of Applied Ecology, Fujian Agriculture & Forestry University, Fuzhou, Fujian, China

³ Graham Centre for Agricultural Innovation, Charles Sturt University, Orange, New South Wales, Australia

⁴ Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou, Fujian, China

⁵ Elizabeth Macarthur Agricultural Institute, NSW Department of Primary Industries, Menangle, New South Wales, Australia

⁶ School of Agriculture and Food Sciences, The University of Queensland, St Lucia, Queensland, Australia

⁷ Highlands Regional Centre, National Agricultural Research Institute, Aiyura, Eastern Highlands Province, Papua New Guinea

⁸ National Agricultural Research Institute, Lae, Morobe Province, Papua New Guinea

Corresponding Author: Geoff M Gurr

Email address: ggurr@csu.edu.au

Sweetpotato (*Ipomea batatans*) is a food crop of global significance. The storage roots and foliage of crop are attacked by a wide range of pests and diseases. Whilst these are generally well controlled in developed countries using approaches such as clean planting material and monitoring with pheromone traps to guide insecticide use, research into methods suitable for developing countries has lagged. In Papua New Guinea (PNG), sweetpotato is grown extensively as a subsistence crop and commercial production as a cash crop is developing. We report results from a survey of 33 smallholder producers located in the Highlands of PNG where the crop is of particular importance. Surveys of interviewees' crops showed high levels of pest and disease impact to foliage, stems and storage roots, especially in crops that were several years old. Weevils (Curculionidae) were reportedly the most damaging pests and scab (caused by the fungus *Elisnoe batatus*) the most damaging disease. Most producers reported root damage from the former and foliar damage from the latter but the general level of knowledge of pest and disease types was low. Despite the apparency of pest and disease signs and symptoms and recognition of their importance by farmers, a large majority of producers reported practiced no active pest or disease management. This was despite low numbers of farmers reporting use of traditional cultural practices including phytosanitary measures and insecticidal plants that had the scope for far wider use. Only one respondent reported use of insecticide though pesticides were available in nearby cities. This low level of pest and disease management in most cases, likely due to paucity in biological and technical knowledge among growers,

hampers efforts to establish food security and constrains the development of sweetpotato as a cash crop.

Pests, diseases and crop protection practices in the smallholder sweetpotato production system of the highlands of Papua New Guinea

Geoff M Gurr^{1, 2, 3 *}, Jian Liu^{1, 2}, Anne C Johnson³, Deane N Woruba⁴, Gunnar Kirchhof⁵,
Ryosuke Fujinuma⁵, William Sirabis⁶, Yapo Jeffery⁷, Ramakrishna Akkinapally⁷

¹State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, Fujian
Agriculture and Forestry University, Fuzhou 350002, China; email: ggurr@csu.edu.au

²Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou 350002,
China

³Graham Centre for Agricultural Innovation (Charles Sturt University & NSW Department of
Primary Industries), PO Box 883, Orange, NSW 2800. Australia. ggurr@csu.edu.au

⁴NSW Department of Primary Industries, Elizabeth Macarthur Agricultural Institute,
Woodbridge Road, Menangle, NSW 2568, Australia. nabre3@gmail.com

⁵School of Agriculture and Food Sciences, The University of Queensland, St Lucia Queensland
4072, Australia. g.kirchhof1@uq.edu.au; R.fujinuma@uq.edu.au

⁶National Agricultural Research Institute, Highlands Regional Centre, Aiyura, P.O. Box 210,

21 Ukarumpa 444, Eastern Highlands Province, Papua New Guinea. william.sirabis@nari.org.pg;
 22 yapo.jeffery@nari.org.pg.

23 ⁷ National Agricultural Research Institute, P O Box 4415, Lae 411, Morobe Province, Papua
 24 New Guinea. a.ramakrishna@nari.org.pg

25 * Corresponding author

27 Abstract

28 Sweetpotato (*Ipomea batatas*) is a food crop of global significance. The storage roots and foliage
 29 of the crop are attacked by a wide range of pests and diseases. Whilst these are generally well
 30 controlled in developed countries using approaches such as clean planting material and
 31 monitoring with pheromone traps to guide insecticide use, research into methods suitable for
 32 developing countries has lagged. In Papua New Guinea (PNG), sweetpotato is grown extensively
 33 as a subsistence crop and commercial production as a cash crop is developing. We report results
 34 from a survey of 33 smallholder producers located in multiple sites in the Highlands of PNG
 35 where the crop is of particular importance. Surveys of interviewees' crops showed high levels of
 36 pest and disease impact to foliage, stems and storage roots, especially in areas where many
 37 successive crops had been grown. Weevils (Curculionidae) were reportedly the most damaging
 38 pests and were present in crops. Symptoms of scab (caused by the fungus *Elisnoe batatus*) were
 39 the most common foliar symptoms and this was the disease of most concern to farmers. Most
 40 producers reported root damage from the former and foliar damage from the latter but the
 41 general level of knowledge of pest and disease types was low. Despite the apparency of pest and
 42 disease signs and symptoms and recognition of their importance by farmers, a large majority of
 43 producers reported practicing no active pest or disease management. This was despite the
 44 practice among some farmers of traditional cultural methods including phytosanitary measures
 45 and insecticidal plants that had the scope for wider use. Only one respondent reported use of
 46 insecticide though pesticides were available in nearby towns. This low level of pest and disease
 47 management, likely due to paucity in biological and technical knowledge among growers,
 48 hampers efforts to establish food security and constrains the development of sweetpotato as a
 49 cash crop.

50

51

52 **Introduction**

53 Among globally important food crops, sweetpotato (*Ipomoea batatas*) ranks number seven (Clark
54 et al. 2013) but has been the subject of far less research than other staples such as potato (*Solanum*
55 *tuberosum*) and wheat (*Triticum aestivum*) (Clark et al. 2013). This reflects the fact sweetpotato is
56 a relatively minor crop in most developed countries in contrast to its widespread production in
57 many tropical and sub-tropical, developing regions such as Africa, southern Asia and the Pacific
58 where it is important for local consumption in subsistence communities (Woolfe 1992, Bourke
59 2009, Loebenstein and Thottapilly 2009, Zhang et al. 2009). In these areas, sweetpotato is critical
60 for food security as it is often a major source of calories as well as vitamins such as carotenoids
61 which are vital in preventing malnutrition in children (Lebot 2010, Woolfe 1992, Kismul, Van den
62 Broeck, and Lunde 2014).

63 The storage roots of sweetpotato have high sugar and water content making them highly
64 susceptible to biotic threats, especially during storage and if roots have been damaged by
65 harvesting or pest attack (Woolfe 1992). In developed country production systems, losses are
66 prevented by the availability of infrastructure such as cool storage facilities and rapid
67 transportation systems. In subsistence production systems, however, post-harvest losses are
68 avoided only by progressive harvest on-demand for immediate use (Okonya et al. 2014); with the
69 general lack of infrastructure otherwise leading to high levels of damage (Johnson and Gurr 2016).
70 This slows the development of commercial production and the livelihood benefits that value chains
71 and processing potentially offer to impoverished rural communities.

Sweetpotato is attacked by around 300 species of arthropods (Talekar 1991) that can cause severe to complete crop loss, as well as at least 30 diseases (Clark et al. 2013). Johnson and Gurr (2016) provide a recent, comprehensive review of those most common in smallholder production. The fact that sweetpotato is vegetatively propagated, either by storage root fragments (slips) or by stem cuttings means that there is high scope for transfer of pest and pathogen inoculum from old to new crops. For example, eggs and larvae of the sweetpotato weevil *Cylas formicarius* (Fabricius), an especially important pest, can be found in these propagules (Hartemink et al. 2000). Still more difficult for subsistence farmers to manage is the fact that plant pathogen inoculum, especially of viruses, is readily multiplied and distributed in slips and cuttings (Clark et al. 2012). Pests and diseases of sweetpotato are generally well controlled in developed countries by the use of pathogen-tested (clean) planting material, pheromone trapping and pesticides (Clark et al. 2013, Jansson and Raman 1991). In developing countries, however, these technologies are less available, particularly in outlying areas, and often unaffordable, making subsistence growers more reliant on traditional practices. These cultural practices include ‘slash and burn’ production in which crops are established on newly-cleared land. However, population growth and associated land shortage makes it increasingly difficult to continue these cultural practices resulting in more intense production with shorter fallow periods (Bourke 2001). A further factor that exacerbates the potential impact of pests and diseases in developing countries is that sweetpotato is often grown in a small production unit (garden) as a series of consecutive crops for multiple years rather than as an annual crop rotated among multiple fields as in developed countries. This increases the time period over which pest densities and pathogen inoculum and infection levels can reach damaging levels, potentially compounded by depletion of nutrients from the soil resulting from repeated

harvest of storage roots (Bailey 2009, Hughes et al. 2009, Kirchhof, Taraken, Ramakrishna, et al. 2009).

Overall, sweetpotato production in developing countries is critical for food security but threatened - in a general sense - by pests and diseases; and effective management is difficult because well-studied technologies that are used in developed countries are not appropriate. Further, traditional practices that have allowed production for many generations are becoming less viable because of land shortages whilst research on management approaches that can be implemented has lagged because these regions are often lack funding and capacity for agricultural research. To address this situation, the aim of this study was to capture data that would identify the major biotic threats to sweetpotato production as a guide to future investment of research funding. The geographical focus of the study was the Highlands of PNG where this crop is the main food staple and where there are currently efforts to establish sweetpotato as a commercial cash crop. Whilst agronomic and soil management issues in this region have been the subject of some earlier research (Kirchhof, Taraken, Ratsch, et al. 2009, Wegener, Kirchhof, and Wilson 2009), no information has been available on pests, diseases and their management. A group of the authors visited 33 farmers spanning the major sweetpotato growing areas of the Highlands, conducting an extended interview with each and collecting data from their crops. Retail outlets in the two major towns of the region were also visited to determine the availability of pesticides.

Methods

Sweetpotato farmer surveys were conducted in the Highlands region of PNG in 2014 covering the same sites used in a 2005 survey of farming systems and soil management (Kirchhof,

116 Taraken, Ratsch, et al. 2009, Wegener, Kirchhof, and Wilson 2009). The survey covered the five
 117 population centres of Asaro and Lufa in the Eastern Highlands Province, Gumni and Sinasina in
 118 the Simbu Province, and Mount Hagen in the Western Highlands Province. The Highlands
 119 region experiences sporadic outbreaks of inter-tribal conflicts and armed violence is common.
 120 Significant areas of potentially productive land sited between population centres is either
 121 uncultivated or is being overgrown with revegetation because it is considered too dangerous for
 122 people to regularly cultivate. Reflecting these hazards, local officials, village extension workers
 123 and police were used to facilitate an initial visit to population centres for the purposes of this
 124 study. Armed police accompanied the research team for one centre. At each centre, a preliminary
 125 meeting was held with the community in which authors able to speak the local dialect explained
 126 the nature of the survey and sought their participation. Thereafter, six to seven farmers from each
 127 village were surveyed, a total of 33. Conditions did not permit detailed assessments and
 128 replicated destructive sampling for each site so the survey consisted of a rapid rural appraisal
 129 (RRA) (Kirchhof, Taraken, Ratsch, et al. 2009). Responses of interviewees were recorded on a
 130 standardised form in English. Interviewees were then asked to take the research team (4-5
 131 persons depending on date) to a representative ‘new garden’ in which few successive
 132 sweetpotato crops had been grown and a representative ‘old garden’ in which many successive
 133 sweetpotato crops had been grown and that was planned to be placed into fallow or planted to a
 134 non-sweetpotato crop in the near future. Gardens of both categories were made available on most
 135 sites. A total of 27 varieties were reported from these gardens with I Don’t Care (7), Wagi Besta
 136 (4), Susan’s Black Eye (2) and Carrot Kaukau (2) being the only varieties present in more than
 137 one garden. All gardens were well established and producing storage roots at the time of
 138 inspection. Yield data for the sites and the region in general are not available because

sweetpotato is grown as a subsistence crop that is harvested in a progressive manner. Gardens varied in size from approximately 50 m² to 200 m². This small size allowed the whole garden to be visually assessed for presence/absence of foliar symptoms. Permission was sought to harvest two, randomly selected sweetpotato plants from each garden. This was granted in a majority of cases (more readily for old than newly-planted gardens). The base of the stems was split to assess incidence of weevil larvae and their feeding tunnels and all of the storage roots beneath sampled plants were inspected for the presence of holes smaller than 3 mm in diameter and holes with greater diameter. Chi square analyses using the Quantpsy tool (Preacher 2001) were used to compare old and new gardens, and compare the distribution of farmer responses within garden ages.

Concurrent with the farmer survey, the senior author visited all rural supply retailers in the major townships in the region, Goroka and Mount Hagen, to determine the availability of insecticide and fungicide products.

Results

The 33 farmers made available for inspection a total of 27 newly planted gardens and 28 old gardens.

Crop Inspections

The incidence of crops that were free of foliar symptoms was significantly ($P < 0.05$) lower for old than new gardens (Figure 1). Deformities of the young leaves symptomatic of scab disease, caused by the fungus *Elisnoe batatus* Viégas & Jenkins, were the most common symptoms in old and new gardens. This was distinct from more general stunting of leaf size and discolouration (including mosaic) characteristic of viral diseases which was observed as frequently as scab

symptoms in the old gardens. Viral symptoms were significantly ($P < 0.05$) less frequently observed among new than old gardens (Figure 1). Splitting stem bases of sweetpotato vines was possible for only some gardens because growers tended to be concerned about destructive inspection of even a single plant because of the small size of the gardens but growers more inclined to approve this in old gardens (Figure 2). Weevil larvae were detected in six of the 14 old gardens but only one of the 10 new gardens, however the small sample size meant that this difference was not significantly different ($P > 0.05$) (Figure 2). For gardens of both ages, crops in which holes were consistently absent from all storage roots sampled from both randomly selected plants were in the minority (Figure 3). The storage roots in most of the old gardens had small (< 3 mm diameter) holes typical of sweetpotato weevil *C. formicarius*. Larger (> 3 mm diameter) holes that may have been caused by the gregariously-feeding West Indian sweetpotato weevil *Euscepes postfasciatus* (Fairmaire) as well as other pests such as molluscs and rats was less common than smaller holes for gardens of both ages. For neither category of hole did the incidence differ significantly between old and new gardens.

Farmer responses

New gardens reportedly had an average of 2.9 successive plantings (including the current crop) with an average fallow period between crops of 11.40 months compared with 25.8 successive plantings for old gardens with just 2.45 months between crops. Prior to the establishment of these gardens, the new ones had an average of 7.56 years of fallow with responses as high as “more than 50 years”, whilst the old gardens were in fallow for 7.39 years with responses extending to “too long ago to remember”. Farmers’ expectation of storage root yield were most commonly high for new gardens and low for old gardens with differences between garden ages very highly significant ($P < 0.001$) (Figure 4).

Very low number of farmers reported that their crops tended not to be attacked by pests and diseases (Figure 5). Damage from these biotic factors was very much the norm. Chi square analysis comparing the null hypothesis of uniform pest attack across all plant parts with the farmers' reports of which plant parts were attacked showed significant ($p < 0.05$) differences for new gardens such that storage roots (the harvestable portion) were most attacked and roots least attacked (Figure 5 a). The same trend across plant parts was apparent among old gardens but the distribution of pest attack did not differ significantly from the null hypothesis. For diseases, stems and leaves were reportedly most commonly attacked and roots least attacked, a trend that was consistent across both garden ages and significantly different from the null hypothesis ($p < 0.05$) within each age (Figure 5 b). Caterpillars were considered a particular problem at the 5-6 month stage and gall mites and scab at harvest time.

Sweetpotato weevil (species unspecified) was ranked by the farmers as the crop protection issue of greatest concern and for which they most wanted a solution. Chi square analysis comparing the null hypothesis of all pest types reported with equal frequency with the farmers' reports showed very highly significant ($p < 0.001$) differences within new and old gardens (Figure 6). This applied to the extent that weevils ranked more highly than all other biotic threat responses combined. Gall mite was the second highest ranked pest priority for gardens of both ages whilst grasshoppers and crickets were also specific concerns. Scab was the highest-ranked sweetpotato disease problem, again in gardens of both ages. 'Nematode', 'tuber rot', 'rust' and other, unknown diseases were also mentioned as biotic issues of concern. When asked to specify the times of year pest were most problematic the responses were varied. For sweetpotato weevil, attack was reported by farmers at widely varying times of the year and plant growth period but

was mostly associated with the dry season. For crickets, planting and wet seasons were periods of reported risk. Gall mites and scab were of greatest concern at harvest time.

Despite all farmers noticing pests and diseases (Figure 5) and considering pest damage, particularly by weevils, as a concern (Figure 6) very few reported taking action to prevent or control pest attack. The great majority of farmers reported taking no action to manage pests (Figure 7 a). Chi square analysis comparing the null hypothesis of all pest management approaches (including no control) being reported with equal frequency with the farmers' reports showed very highly significant ($p < 0.001$) differences for new and old gardens (Figure 7a). No more than four farmers each used the soil management approaches of mounding-up over storage roots or breaking up mounds to expose roots to heat; biological control with ants or chickens, mulching with plant materials such as 'fish-kill' (*Tephrosia* spp.) or other insecticidal plants. One farmer mentioned use of insecticide, Karate® (lambda-cyhalothrin) in his new garden. Only one grower reported the use of a combination of methods, soil management with rogueing (removal of infested stems), for pest management.

An equivalent lack of intervention was evident for disease management (Figure 7 b). Chi square analysis comparing the null hypothesis of all disease management approaches (including no control) being reported with equal frequency with the farmers' reports showed very highly significant ($p < 0.001$) differences for new and old gardens (Figure 7b). One grower reported the use of 'clean planting material' but this was sourced from their own gardens rather than from a pathogen-tested planting material scheme. In a separate question specifically about use of planting material that was 'certified or disease tested', all farmers reported no such use. One grower each reported rogueing (removal of symptomatic stems), fallowing and use of an unspecified resistant variety.

229 *Survey of pesticide availability*

230 A survey of the seven rural supply shops in the two major townships of Mount Hagen and
 231 Goroka found that a small range of pesticides was available (Table 1). Of the eight insecticides
 232 available, only lambda-cyhalothrin was sold in most shops. Chlorothalonil was the only
 233 fungicide available in the two cities but on sale in most of the shops. Retailers reported these
 234 were usually purchased for use on cash crops such as Irish potato (*Solanum tuberosum*), allowing
 235 the cost of the input to be recouped, and rarely for use in sweetpotato since this was principally
 236 for consumption by the extended family. In some stores, the pesticides were repackaged into
 237 smaller, unlabelled packs for sale at low prices. More generally, labelling practices were not
 238 stringent, with packs of one chlorothalonil product carrying the contradictory wording
 239 ‘protective fungicide’ and ‘group Y herbicide’ (Figure 8).

240

241 **Discussion**

242 Developing country pest and disease issues tend to receive less attention than those in developed
 243 countries and this is compounded in regions where studies are made more difficult because of
 244 instability and violence. Thus, though agricultural research in PNG has been the subject of
 245 significant effort in recent years, there is a relative dearth of information to inform priorities and
 246 investment. The present study of smallholder sweetpotato growers in the region of PNG, where
 247 this crop is the main staple, provides strong evidence that pests and diseases are having a large
 248 impact on production and that current management efforts are inadequate.

249 Among the biotic threats that farmers reported to be of high concern, weevils were paramount.

250 This was evident also in the assessment of damage to storage roots and inspections of stems in

which weevils were frequently present. Internationally, the sweetpotato weevil is consistently ranked as the most problematic pest in sweetpotato production (Ebregt et al. 2004, Fielding and van Crowder 1995, Nsibande and McGeoch 1999, Okonya et al. 2014, Parr, Ntonifor, and Jackai 2014, Placide et al. 2015) though the damage can be confused with that from other pests such as millipedes (Diplopoda) (Ebregt et al. 2004). *Euscepes postfasciatus* is present in PNG (Hughes 2013) and this causes some forms of damage similar to that of the sweetpotato weevil (*C. formicarius*). Though the adults of these two weevils are dissimilar in appearance, the immatures look very similar. No farmers mentioned either species specifically so the relative importance of these two species as pests remains to be determined. Certainly, both are potentially serious pests. Weevil attack was reported by farmers at widely varying times of the year but was mostly associated with the dry season, reflecting the fact that storage roots are more exposed to attack if soil cracks as a result of dry conditions (Lutulele, 2001; Parr *et al.*, 2014a) and this suggests that impact could be more severe under climate change conditions (Okonya and Kroschel, 2013).

Native to the Indian subcontinent and eastwards to Malaysia, *C. formicarius* is a serious pest in the south west Pacific, the southern USA, Caribbean and South America (Chalfant et al. 1990, Sherman and Tamashiro 1954, Waterhouse and Norris 1987). Austin, Jansson, and Wolfe (1991) and Horton and Ewell (1991) considered this pest of great importance in causing pre-harvest damage. *Euscepes postfasciatus* originated from the Caribbean and is now a pest in the Pacific region and South America (Katsuki et al. 2012, Raman and Alleyne 1991, Sherman and Tamashiro 1954). An important mode of dispersal for both species is as immatures within storage roots or stem cuttings (Hartemink et al. 2000, Ray, Mishra, and Mishra 1983). Larvae of both weevil species feed on the storage root or within stems causing tunnelling packed with frass. Adult *E. postfasciatus* tend to feed on storage roots gregariously, causing relatively few

large holes. In contrast *C. formicarius* adults tend to feed individually causing smaller wounds (Sherman and Tamashiro 1954). Accordingly, our classification of observed holes on storage roots into <3mm and >3mm diameter provides an approximate indication that *C. formicarius* may be the dominant weevil species. Clearly storage root holes could also be caused by other pests, such as molluscs and rats, especially in the case of larger holes, so these results are tentative. Studies based on rearing-out adults from infested storage roots or identifying immatures (potentially aided by the development of molecular diagnostic tools) are necessary in order to discriminate the incidence and impact of these two weevil species and plan appropriate research and management priorities and such studies are currently underway.

Gall mite, *Eriophyes gastrotrichus* Nalepa (Acari: Eriophyidae), causes erinose, a foliar disease characterised by blister-like galls on the stems of sweetpotato plants in the Philippines, and PNG where it is has previously been reported to be a problem of increasing concern in the Highlands (Ames et al. 1996, Hughes et al. 2009). This pest was the second most highly-rated concern among growers. Since it infests the foliage, it is readily spread by stem cuttings which are commonly used in the region. The use of slips or, more especially, pathogen-tested planting material would allow crops to be established in a ‘clean’ state and allow production for some time before field infection occurs. The Australian Centre for International Agricultural Research has invested in establishing a pathogen-tested planting material program in the region. Whilst the principal focus of this is control of viruses (see below) it would also benefit crop protection more widely including for gall mite. In the present study, however, none of the farmers reported prior use of planting material that was pathogen tested, certified or disease tested. Some reported use of ‘clean planting material’ but this was sourced from their own or nearby gardens and illustrates that they were aware of this infection pathway and the need to manage carryover of inoculum.

Symptoms of scab, caused by *Elsinoe batatas* Viégas & Jenkins, was the most commonly observed form of foliar symptoms in both old and new gardens and was also the disease considered of highest priority by farmers. Though the symptoms of this disease are characteristic and unlikely to be confused with those of other diseases, pathogen isolation in future studies is required to confirm identity. Throughout tropical regions, scab is considered the most serious fungal disease of sweetpotato (Clark et al. 2013, Coleman et al. 2009). Though the storage roots can be infected this tends to cause little impact; though foliar damage can be so severe that photosynthetic area is reduced leading to storage root yield reductions as high as 34% (Coleman et al. 2009). Pathogen inoculum survives on crop residues and can be transmitted readily by stem cuttings so is chiefly a problem when sweetpotato is grown continuously (Clark et al. 2013, Coleman et al. 2009). It is noteworthy, then, that its incidence was high even in the new crop gardens and this reflects the fact that no farmers had accessed pathogen-tested clean planting material.

Viruses are widely considered to be of great economic importance in sweetpotato production (Clark et al. 2012, Gibson and Kreuze 2014). A survey of scientists from less developed countries rated viruses as the top priority (Fuglie 2007). Notably, however, no farmers in the present study mentioned viruses though a large proportion of old gardens showed foliar symptoms consistent with viral infection. As noted above, pathogen identification is required in future work to confirm the precise cause of these symptoms. The apparent lack of concern amongst growers about viral diseases likely reflects the fact that symptoms of viral infection can be subtle and develop over a prolonged period with little or no direct symptoms on the storage roots other than yield decline which is likely to be attributed to pests because of their greater apparency. Related to this, the concept of a plant pathogenic virus, that has no signs, is relatively

unfamiliar to many farmers so it not being mentioned is likely to reflect this fact. The availability of molecular detection methods has led to rapid advances in sweetpotato virus knowledge and at least 30 viruses of sweetpotato are known (Clark et al. 2012), some with multiple strains (Dolores, Yebron, and Laurena 2012). Yields of virus-infected sweetpotato plants are often severely affected, reduced by as much as 80-90% (Carey et al. 1999, Clark et al. 2012, Davis and Ruabete 2010). Though insects such as aphids such as *Aphis gossypii* and whiteflies including *Bemisia tabaci* can transmit viruses (Clark et al. 2012, Byamukama et al. 2004), propagation material is the chief means of viral spread (Gibson et al. 1997, Moyer and Larsen 1991, Mbanzibwa et al. 2014). Foliar symptoms of virus infection include leaf distortion, strapping and crinkling, mosaics, vein clearing, brown blotches and general stunting and chlorosis (Mbanzibwa et al. 2014). These symotoms were significantly more frequently seen in old rather than new gardens reflecting the time available for plant-to-plant transmission and build up of infection levels.

These differences in pest and disease apparency between old and new gardens underscore the importance of political action to establish peaceful rural communities in order to allow potentially productive farmlands to be used. Prior to the establishment of these gardens, the new ones had 7.56 years of fallow whilst the old gardens were in fallow for 7.39 year, less than half as long as the 16.8 (SE=2.4) year reported for a 2005 survey of the same sites (Kirchhof, Taraken, Ramakrishna, et al. 2009). This shortening of fallows reflects land shortages resulting from rapidly increasing human population densities (Bourke, 2001) and is likely to allow pest and disease pressure to increase because fallowing has been demonstrated to increase yields via benefits to crop nutrition (Hartemink 2003, Hartemink et al. 2000). Accordingly, if farming communities in the Highlands of New Guinea felt sufficiently safe to extend their cropping

activities back into areas that has fallen out of production because of fear of inter-tribal violence, this would alleviate both biotic and abiotic (nutritional) stress on crops.

A striking finding about pest and disease management practices among the surveyed growers is the very large majority who reported not practicing any active management. This is despite the existence of a potentially large number of methods that could be employed in this setting. Small numbers of farmers reported using insecticidal plants, basic phytosanitation methods and simple forms of biological control using ants or livestock. The makum system is a traditional PNG practice for production of taro on mounds in which the ant, *Pheidole megacephala* (Fabricius), has nested, and has been adapted for use in sweetpotato production (Sar et al. 2009). Ants are also employed in a system in Cuba involving green tree ants being transported into sweetpotato fields from banana plantations within their rolled banana leaf nests (Lagnaoui et al. 2000). Ants can provide sweetpotato weevil control in a more cost effective than insecticides (Chalfant et al. 1990), so merits more attention as a method that could be readily adopted in smallholder systems. It is not possible to determine from the present study why such low rates of pest and disease management were apparent in the present study but the most likely explanation – based on general interactions with the farming communities – is lack of knowledge. In particular, though farmers recognised a range of pest and disease types, their knowledge of lifecycles and essential concepts such as microscopic disease causing agents was rudimentary. Further, though expectations of storage tuber yield from old gardens was lower than from new gardens, there was a tendency to associate this with nutrient depletion. Associated with this, the adoption of strategies to manage nutrition, such as not burning crop residues (Bailey 2009), could exacerbate carryover of pests and pathogen inocula.

A survey of sweetpotato growers in Tanzania found that although farmers could identify diseased plants they could not distinguish the different types of disease (Adam, Sindi, and Badstue 2015). Though those African farmers had a very limited knowledge of pests and pathogens, they took at active precautions to manage them (Adam, Sindi, and Badstue 2015, Nsibande and McGeoch 1999). For example, they identified plants that looked healthy and free of pests for use in planting material (Adam, Sindi, and Badstue 2015). This was not widely reported as a pest or disease management practice in the present survey though farmers are likely to select relatively healthy cuttings on the basis of these being likely to root readily and grow vigorously. The closer a village was to a main town or main road with passing traffic the more likely the farmers in the Tanzanian study were to be able to identify diseases that affect sweetpotato (Adam, Sindi, and Badstue 2015). Sites with easier access also tended to facilitate the use of higher quality planting material. In the present study, all sites were accessible by roads (Kirchhof, Taraken, Ratsch, et al. 2009) is it is likely that levels of knowledge and active pest and disease management are still lower in the more remote areas of the PNG Highlands. Farmer to farmer interactions are an important source of information sharing on pest management (Adam, Sindi, and Badstue 2015, Pouratashi and Iravani 2012) but this communication channel is impeded in PNG by tribal conflict and this underscores the importance of extension efforts and initiatives such as the development of a pathogen-tested planting material scheme. Among the challenges for such a scheme is that many dozens of sweetpotato varieties are grown in the Highlands of PNG so the scheme would need to ‘clean-up’ and make available a wide range of cultivars to meet farmers’ needs.

Conclusion

Like many developing countries, PNG is experiencing rapid population growth and government policies are seeking to establish greater food security and livelihood development, the latter by developing cash crops and value adding to agricultural commodities by processing and marketing. Sweetpotato potentially can contribute strongly to both these objectives because it is widely grown and culturally integral to traditional diets, yet strongly impacted by pests and diseases that are not well managed. The recent IPES-Food (2016) ‘Uniformity to Diversity’ Report highlighted the multiple negative outcomes from intensive agriculture in developed countries. These include loss of biodiversity and reliance on non-renewable and environmentally hazardous inputs including pesticides. Accordingly, the development trajectory of countries such as PNG need to be cognizant of the negative aspects of simply following practices already established in developed nation agricultural systems. For example, making pesticides more readily available and promoting their use are not logical from the sustainability perspective and would also complicate the common practice of feeding sweetpotato foliage to pigs. Production needs to be increased to meet human needs but achieving this by becoming reliant on non-renewable inputs and eroding the natural resource base of agriculture will lead to unsustainability (Godfray 2011). As an alternative, ecological intensification (in which ecosystem services such as biological pest control and nutrient cycling are key) offers viable benefits (Bommarco, Kleijn, and Potts 2013). If wider use of pesticides is to be avoided, the need for alternative approaches is clear but traditional practices of ancient agricultural systems, such as ants and livestock for biological control, and insecticidal plants, can underpin this if their efficacy and utility are better understood and appropriate extension efforts are made. Parallel with such technological efforts, however, advances are necessary in the political and policy arena to make rural communities safer and more sustainable. Recent human population growth and inter-tribal conflict over ever-

more-scarce land has resulted in more intensive cropping in areas close to villages exacerbating pest and disease build-up.

Acknowledgement

Mr. Kai Lali (NARI) drove and provided field support. Sargent Simon Wakala (PNG Royal Constabulary) assisted with security.

References

- Adam, Rahma Isaack, Kirimi Sindi, and Lone Badstue. 2015. "Farmers' knowledge, perceptions and management of diseases affecting sweet potatoes in the Lake Victoria Zone region, Tanzania." *Crop Protection* 72 (0):97-107. doi: 10.1016/j.cropro.2015.02.010.
- Ames, T, NEJM Smit, Ann R. Braun, JN O Sullivan, and Skoglund L.G. 1996. *Sweetpotato: Major pests, diseases, and nutritional disorders*. Peru: International Potato Center.
- Austin, DF, RK Jansson, and GW Wolfe. 1991. "Convolvulaceae and *Cylas*: a proposed hypothesis on the origins of this plant/insect relationship." *Tropical Agriculture* 68 (2):162-170.
- Bailey, J. 2009. "An evaluation of nutritional constraints on sweetpotato production in Papua New Guinea highlands using the diagnosis and recommendation inegration System (DRIS)." In *Soil fertility in sweetpotato based cropping systems in the highlands of Papua New Guinea*, edited by G. Kirchhof, 7-11. Canberra, ACT: Australian Centre for International Agricultural Research.
- Bommarco, Riccardo, David Kleijn, and Simon G Potts. 2013. "Ecological intensification: harnessing ecosystem services for food security." *Trends in Ecology & Evolution* 28 (4):230-238. doi: 10.1016/j.tree.2012.10.012.
- Bourke, R. M. 2009. "Sweetpotato in Oceania." In *The Sweetpotato*, edited by G. Loebenstein and G. Thottapilly, 498-502. The Netherlands: Springer.
- Bourke, R. Michael. 2001. "Intensification of Agricultural Systems in Papua New Guinea." *Asia Pacific Viewpoint* 42 (2-3):219-235. doi: 10.1111/1467-8373.00146.

- Byamukama, E., R. W. Gibson, V. Aritua, and E. Adipala. 2004. "Within-crop spread of sweet potato virus disease and the population dynamics of its whitefly and aphid vectors." *Crop Protection* 23 (2):109-116. doi: 10.1016/j.cropro.2003.07.003.
- Carey, EE, RW Gibson, S Fuentes, M Machmud, ROM Mwanga, G Turyamureeba, L Zhang, D Ma, F Abo El-Abbas, and R El-Bedewy. 1999. The causes and control of virus diseases of sweetpotato in developing countries: Is sweetpotato virus disease the main problem? In *Impact on changing world. 1997-98 Program Report*. Lima, Peru: International Potato Centre.
- Chalfant, R B, R K Jansson, D R Seal, and J M Schalk. 1990. "Ecology and management of sweet potato insects." *Annual Review of Entomology* 35 (1):157-180. doi: doi:10.1146/annurev.en.35.010190.001105.
- Clark, C. A., DM Ferrin, TP Smith, and GJ Holmes, eds. 2013. *Compendium of sweetpotato diseases, pests and disorders*. Second ed. Minnesota, USA: APS Press.
- Clark, Christopher A., Jeffrey A. Davis, Jorge A. Abad, Wilmer J. Cuellar, Segundo Fuentes, Jan F. Kreuze, Richard William Gibson, Settumba B. Mukasa, Arthur K. Tugume, Fred Donati Tairo, and Jari P. T. Valkonen. 2012. "Sweetpotato Viruses: 15 Years of Progress on Understanding and Managing Complex Diseases." *Plant Disease* 96 (2):168-185. doi: 10.1094/pdis-07-11-0550.
- Coleman, E., M. Hughes, G. Jackson, B. Komolong, and E. Guaf. 2009. "Genetics and disease as factors in the yield decline of sweetpotato in the Papua New Guinea highlands." In *Soil fertility in sweetpotato based cropping systems in the highlands of Papua New Guinea*, edited by G. Kirchhof, 33-42. Canberra, ACT: Australian Centre for International Agricultural Research.
- Davis, R. I., and T. K. Ruabete. 2010. "Records of plant pathogenic viruses and virus-like agents from 22 Pacific island countries and territories: a review and an update." *Australasian Plant Pathology* 39 (3):265-291. doi: 10.1071/AP10047.
- Dolores, L. M., M. G. N. Yebron, Jr., and A. C. Laurena. 2012. "Molecular and biological characterization of selected Sweet potato feathery mottle virus (SPFMV) strains in the Philippines." *Philippine Journal of Crop Science* 37 (2):29-37.
- Ebregt, E., P. C. Struik, P. E. Abidin, and B. Odongo. 2004. "Farmers' information on sweet potato production and millipede infestation in north-eastern Uganda. I. Associations between spatial and temporal crop diversity and the level of pest infestation." *NJAS - Wageningen Journal of Life Sciences* 52 (1):47-68. doi: 10.1016/S1573-5214(04)80029-0.
- Fielding, W., and L. van Crowder. 1995. "Sweet potato weevils in Jamaica: Acceptable pests?" *Journal of Sustainable Agriculture* 5 (4):105-117.
- Fuglie, KO. 2007. "Priorities for sweetpotato in developing countries: results of a survey." *HortScience* 42 (5):1200-1206.
- Gibson, R. W., and J. F. Kreuze. 2014. "Degeneration in sweetpotato due to viruses, virus-cleaned planting material and reversion: a review." *Plant Pathology* 64 (1):1-15. doi: 10.1111/ppa.12273.

- 474 Gibson, R. W., R. O. M. Mwanga, S. Kasule, I. Mpembe, and E. E. Carey. 1997. "Apparent absence of
475 viruses in most symptomless field-grown sweet potato in Uganda." *Annals of Applied Biology* 130
476 (3):481-490. doi: 10.1111/j.1744-7348.1997.tb07676.x.
- 477 Godfray, H Charles J. 2011. "Ecology, food and biodiversity." *Science* 333 (6047):1231-1232. doi:
478 10.1126/science.1211815.
- 479 Hartemink, AE. 2003. "Sweet potato yields and nutrient dynamics after short term fallows in the humid
480 lowlands of Papua New Guinea." *NJAS - Wageningen Journal of Life Sciences* 50:297-319.
- 481 Hartemink, Alfred E., S. Poloma, M. Maino, K. S. Powell, J. Egenae, and J. N. O'Sullivan. 2000. "Yield
482 decline of sweet potato in the humid lowlands of Papua New Guinea." *Agriculture, Ecosystems &
483 Environment* 79 (2-3):259-269. doi: 10.1016/S0167-8809(00)00139-0.
- 484 Horton, D.E., and P.T Ewell. 1991. "Sweet potato pest management: A social science perspective." In
485 *Sweet potato pest management: A global perspective*, edited by R K Jansson and Kandukuri V.
486 Raman, 407-428. San Francisco: Westview Press.
- 487 Hughes, M. 2013. Identifying appropriate strategies for reducing virus and weevil losses in sweetpotato
488 production systems in Papua New Guinea and Australia. In *Final report HORT/2011/053*. Canberra,
489 ACT: Australian Centre of International Agricultural Research.
- 490 Hughes, M., E. Coleman, Issac T. Taraken, and P. Igua. 2009. "Sweet potato agronomy in Papua New
491 Guinea." In *Soil fertility in sweetpotato based cropping systems in the highlands of Papua New
492 Guinea*, edited by G. Kirchhof, 7-11. Canberra: Australian Centre for International Agricultural
493 Research.
- 494 IPES-Food. 2016. From uniformity to diversity: a paradigm shift from industrial agriculture to diversified
495 agroecological systems. International Panel of Experts on Sustainable Food systems. [www.ipes-](http://www.ipes-food.org)
496 [food.org](http://www.ipes-food.org).
- 497 Jansson, R K, and Kandukuri V. Raman, eds. 1991. *Sweet potato pest management: A Global perspective*.
498 San Francisco: Westview Press
- 499 Johnson, A. C., and G. M. Gurr. 2016. "Invertebrate pests and diseases of sweetpotato (*Ipomoea*
500 *batatas*): a review and identification of research priorities for smallholder production." *Annals of*
501 *Applied Biology* 168 (3):291-320. doi: 10.1111/aab.12265.
- 502 Katsuki, Masako, Yusuke Omae, Kensuke Okada, Toru Kamura, Takashi Matsuyama, Dai Haraguchi,
503 Tsuguo Kohama, and Takahisa Miyatake. 2012. "Ultraviolet light-emitting diode (UV LED) trap the
504 West Indian sweet potato weevil, *Euscepes postfasciatus* (Coleoptera: Curculionidae)." *Applied*
505 *Entomology and Zoology* 47 (3):285-290. doi: 10.1007/s13355-012-0113-y.
- 506 Kirchhof, G., Issac T. Taraken, A. Ramakrishna, Rainer Ratsch, and P. Igua. 2009. "Biophysical constraints
507 of sweetpotato-based cropping systems in the Papua New Guinea highlands." In *Soil fertility in*
508 *sweetpotato based cropping systems in the highlands of Papua New Guinea*, edited by G. Kirchhof,
509 95-109. Canberra, ACT: Australian Centre for International Agricultural Research.

- Kirchhof, G., Issac T. Taraken, Rainer Ratsch, D. Kapal, and P. Igua. 2009. "Survey methodology to assess socioeconomic and biophysical constraints - lessons learnt in the highlands of Papua New Guinea." In *Soil fertility in sweetpotato based cropping systems in the highlands of Papua New Guinea*, edited by G. Kirchhof, 70-78. Canberra, ACT: Australian Centre for International Agricultural Research.
- Kismul, H., J. Van den Broeck, and T. M. Lunde. 2014. "Diet and kwashiorkor: a prospective study from rural DR Congo." *PeerJ* 2:e350. doi: 10.7717/peerj.350.
- Lagnaoui, A, F Cisneros, J Alcazar, and F Morales. 2000. *A sustainable pest management strategy for sweetpotato weevil in Cuba: A success story*: Food Fertilizer Technology Center.
- Lebot, V. . 2010. "Sweetpotato." In *Root and tuber crops*, edited by J. E. Bradshaw, 97-125. New York, USA: Springer Science & Business Media.
- Loebenstein, G., and G. Thottapilly, eds. 2009. *The Sweetpotato*. Netherlands: Springer.
- Mbanzibwa, D. R., A. K. Tugume, E. Chiunga, D. Mark, and F. D. Tairo. 2014. "Small RNA deep sequencing-based detection and further evidence of DNA viruses infecting sweetpotato plants in Tanzania." *Annals of Applied Biology* 165 (3):329-339. doi: 10.1111/aab.12136.
- Moyer, J.W., and R.C Larsen. 1991. "Management of insect vectors of viruses infecting sweet potato." In *Sweet potato pest management: A Global perspective*, edited by R K Jansson and Kandukuri V. Raman, 341-358. San Francisco: Westview Press
- Nsibande, M. L., and M. A. McGeoch. 1999. "Sweet potato, *Ipomoea batatas* (L), cropping practices and perceived production constraints in Swaziland: implications for pest management." *International Journal of Pest Management* 45 (1):29-33. doi: 10.1080/096708799228012.
- Okonya, JS, R. Mwanga, K Syndikus, and J Kroschel. 2014. "Insect pests of sweetpotato in Uganda: farmers' perceptions of their importance and control practices." *SpringerPlus* 3 (303):1-10. doi: 10.1186/2193-1801-3-303.
- Parr, MC, NN Ntonifor, and LE Jackai. 2014. "Effect of planting dates on the population dynamics of *Cylas puncticollis* and sweet potato storage roots damage in South Western Cameroon." *Journal of Biology, Agriculture and Healthcare* 4 (18):41-48.
- Placide, R, H. Shimelis, M Laing, and D Gahakwa. 2015. "Farmers' perceptions, production and productivity constraints, preferences and breeding priorities of sweetpotato in Rwanda." *HortScience* 50 (1):36-43.
- Pouratashi, Mahtab, and Hooshang Iravani. 2012. "Farmers' knowledge of integrated pest management and learning style preferences: Implications for information delivery." *International Journal of Pest Management* 58 (4):347-353. doi: 10.1080/09670874.2012.724468.
- Preacher, K. 2001. "Calculation for the chi-square test: An interactive calculation tool for chi-square tests of goodness of fit and independence. [Computer software]." <http://quantpsy.org>.

- Raman, Kandukuri V., and E. H. Alleyne. 1991. "Biology and management of the West Indian sweet potato weevil, *Euscepes postfasciatus*." In *Sweet potato pest management: A Global perspective*, edited by R K Jansson and Kandukuri V. Raman, 263-281. San Francisco: Westview Press.
- Ray, P. K., S. Mishra, and S. S. Mishra. 1983. "Sweet-potato productivity as affected by recurrent use of vines as planting-material." *Scientia Horticulturae* 20 (4):319-322. doi: 10.1016/0304-4238(83)90145-0.
- Sar, Sim A, Christine King, Elske van de Fliert, Redley Opasa, Michael Atoai, Ana Appa, and Triya Papaya. 2009. "Engaging stakeholders through participatory research: Farmer innovations in the use of predatory ants for pest management in Papua New Guinea." *Innovation Asia Pacific Symposium*, Nepal, 4-7 May.
- Sherman, Martin, and Minoru Tamashiro. 1954. The sweetpotato weevils in Hawaii: Their biology and control. In *Technical Bulletin No. 23*, edited by University of Hawaii Hawaii Agricultural Experimental Station. Hawaii: University of Hawaii.
- Talekar, N.S. 1991. "Integrated Control of *Cylas formicarius*." In *Sweet potato pest management: A Global perspective*, edited by R K Jansson and Kandukuri V. Raman, 139-156. San Francisco: Westview Press.
- Waterhouse, D. F., and K. R. Norris. 1987. *Biological control, Pacific prospects*. Edited by Research Australian Centre for International Agricultural. Melbourne: Melbourne : Inkata Press.
- Wegener, M., G. Kirchhof, and T. Wilson. 2009. "An analysis of village garden management in the Papua New Guinea highlands." In *Soil fertility in sweetpotato based cropping systems in the highlands of Papua New Guinea*, edited by G. Kirchhof, 79-87. Canberra, ACT: Australian Centre for International Agricultural Research.
- Woolfe, Jennifer A. 1992. *Sweet potato: An untapped food resource*. New York: Cambridge University Press.
- Zhang, L., Q. Wang, Q. Liu, and Q. Wang. 2009. "Sweetpotato in China." In *The Sweetpotato*, edited by Gad Loebenstein and George Thottappilly, 325-358. Netherlands: Springer

Table 1: Insecticide and fungicide availability in retail outlets in the Papua New Guinea Highlands region townships of Goroka and Mount Hagen. (The anonymity of the retail suppliers is protected by de-identification and the use of lettering)

City	Retail Supplier	Product name and active constituent	Type
Goroka	A	Karate® 25g/L; lambda-cyhalothrin	Insecticide
		Eko® 720 g/L; chlorothalonil	Fungicide
		Barrek® 500g/L; chlorothalonil	Fungicide
	B	Lambda® C2.5EC; lambda-cyhalothrin	Insecticide
		Malathion®; malathion	Insecticide
		Eko® 720 g/L; chlorothalonil	Fungicide
	C	Permethrin® 250 EC; permethrin	Insecticide
		Lambda® C2.5EC; lambda-cyhalothrin	Insecticide
		Bifenthrin® 10%; bifenthrin	Insecticide
		Eko® 720 g/L; chlorothalonil	Fungicide
	D	Confidor®; imidacloprid	Insecticide
Mount Hagen	E	Permethrin®; permethrin	Insecticide
		Carbofuran®; carbofuran	Insecticide
		Acephate® 75%wv; acephate	Insecticide
		Bifenthrin®; bifenthrin	Insecticide
		Chlorpyrifos® 480EC; chlorpyrifos	Insecticide
		Barrek® 500g/L; chlorothalonil	Fungicide
	F	Eko® 720 g/L; chlorothalonil	Fungicide




Figure 1. Incidence of foliar symptoms (viral infection and scab infection and symptom-free) among sweetpotato crops (n=25 new and 20 old). (Symptoms were non-mutually, exclusive; some crops had symptoms of more than one type. (Chi-square tests compared old and new gardens: viral, $X^2 = 4.543$, $df=1$, $p=0.033$; scab, $X^2=1.635$, $df=1$, $p=0.202$; no symptoms, $X^2=4.021$, $df=1$, $p=0.044$).





Figure 2. Incidence of weevils in the base of the stems among sweetpotato crops (n=10 new and 14 old). (Chi-square test compared old and new gardens $X^2=3.048$, $df=1$, $p=0.081$)

596



597

598

599

600

601

602

603

604

Figure 3. Incidence of pest damage holes in sweetpotato storage roots (n=10 new and 14 old). (One crop had holes of both sizes.) (Chi-square tests compared old and new gardens: <3mm: $X^2= 1.143$, df=1, p=0.285; >3mm: $X^2= 2.194$, df=1, p=0.138)





Figure 4. Yield expectation of farmers for new and old gardens. (Chi-square test compared distribution of responses between garden ages: $X^2=24.316$, $df=3$, $p<0.001$)

a



b




Figure 5. Farmers' responses on whether and where they observe damage by pests (a) and diseases (b). (Means are the number of farmers mentioning a given concern and are non-mutually exclusive, some

farmers mentioning one, and some multiple plant portions.) (Chi-square tests compared plant portions within each garden age: PESTS, new gardens, $X^2=7.849$, $df=3$, $p=0.049$; old gardens, $X^2=4.524$, $df=3$, $p=0.201$. DISEASES, old gardens: $X^2=8.544$, $df=3$, $p=0.036$; old gardens, $X^2=9.9444$, $df=3$, $p=0.0190$)






Figure 6. Plant protection issues cited in the top three concerns by farmers' for pest and disease problems. (Means are the number of farmers mentioning a given concern and are non-mutually exclusive, some farmers mentioning one, and some up to three issues. (Chi-square test compared pest types within each garden age: new gardens, priority is used by times been listed without giving any points. Weevil: $X^2=16.448$, $df=4$, $p=0.002$; old gardens, $X^2=23.836$, $df=4$, $p<0.001$)

637 **a**



638

639 **B**




640

641

642 **Figure 7.** Reported actions taken to control pests (a) and diseases (b) on sweetpotato crops. (Chi-square
643 test compared management approaches within each garden age: PESTS, new gardens, $X^2=41.989$, $df=6$,

644 $p < 0.001$; old gardens, $X^2 = 52.738$, $df = 6$, $p < 0.001$. DISEASES, new gardens, $X^2 = 38.338$, $df = 4$, $p < 0.001$;
 645 old gardens, $X^2 = 52.277$, $df = 4$, $p < 0.001$)
 646

647
648
649
650



651
652
653
654
655
656

Figure 8: Example of pesticide labelling anomaly. Photograph from pesticide label on product for sale in Goroka.